



ANNALES

DE

L'INSTITUT FOURIER

Markus SZYMIK

The rational stable homology of mapping class groups of universal nil-manifolds

Article à paraître, mis en ligne le 5 février 2019, 21 p.



© Association des Annales de l'institut Fourier, 2019,
Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

THE RATIONAL STABLE HOMOLOGY OF MAPPING CLASS GROUPS OF UNIVERSAL NIL-MANIFOLDS

by Markus SZYMIK

ABSTRACT. — We compute the rational stable homology of the automorphism groups of free nilpotent groups. These groups interpolate between the general linear groups over the ring of integers and the automorphism groups of free groups, and we employ functor homology to reduce to the abelian case. As an application, we also compute the rational stable homology of the outer automorphism groups and of the mapping class groups of the associated aspherical nil-manifolds in the TOP, PL, and DIFF categories.

RÉSUMÉ. — Nous calculons l'homologie rationnelle stable des groupes d'automorphismes de groupes nilpotents libres. Ces groupes s'intercalent entre les groupes généraux linéaires sur l'anneau des entiers et les groupes d'automorphismes de groupes libres, et nous employons l'homologie de foncteurs pour nous réduire au cas abélien. A titre d'application, nous calculons également l'homologie rationnelle stable des groupes d'automorphismes extérieurs et des groupes modulaires des variétés asphériques associées dans les catégories TOP, PL, et DIFF.

1. Introduction

For any integer $r \geq 1$, let F_r denote the free group on r generators. The automorphism groups $\text{Aut}(F_r)$ of these groups have been the subject of plenteous research. Nielsen began his investigations in that direction about a century ago, and more recent work has led to substantial progress in the computation of their homology. This reckoning applies in particular after stabilization: We can extend an automorphism of F_r to F_{r+1} by sending the new generator to itself, and in the (co)limit $r \rightarrow \infty$, we get the stable automorphism group $\text{Aut}(F_\infty)$. Note that this notation does *not* refer to an automorphism group of a group F_∞ . The homology of the group $\text{Aut}(F_\infty)$ is

Keywords: stable homology, automorphism groups, nilpotent groups, functor categories, Hochschild homology, stable K-theory, spectral sequences.

2010 Mathematics Subject Classification: 20J05, 20E36, 19M05, 18A25, 18G40.

the colimit of the homologies of the groups $\text{Aut}(F_r)$. This stable homology has been computed by Galatius [14]. In particular, it is rationally trivial, as had been conjectured by Hatcher and Vogtmann [18].

The descending sequence of normal subgroups of the free group F_r that is given by $\Gamma_1(F_r) = F_r$ and $\Gamma_{c+1}(F_r) = [F_r, \Gamma_c(F_r)]$ is the lower central series. The quotient groups $N_c^r = F_r/\Gamma_{c+1}(F_r)$ are the free nilpotent groups of class $c \geq 1$.

For instance, the first case gives the free abelian groups $N_1^r \cong \mathbb{Z}^r$. Their automorphism groups are the general linear groups $\text{GL}_r(\mathbb{Z})$ over the ring \mathbb{Z} of integers, and their homology is related to the algebraic K-theory of the ring \mathbb{Z} . In contrast to the case of the groups $\text{Aut}(F_r)$, we only know very few of their stable homology groups integrally. This changes when we are willing to work rationally. In that case, a complete computation of the stable homology has been achieved by Borel [6] using analytic methods. There is an isomorphism

$$(1.1) \quad H^*(\text{GL}_\infty(\mathbb{Z}); \mathbb{Q}) \cong \Lambda_{\mathbb{Q}}(r_5, r_9, r_{13}, \dots)$$

between the rational stable cohomology ring and a rational exterior algebra on generators in degrees $5, 9, 13, \dots, 4n + 1, \dots$. The methods that we will use here are entirely algebraic and topological. They only provide new information relative to the homology of the general linear groups, to which we have nothing to add. We could, in principle, also attempt to use them to obtain information on torsion in the relative homology. At present, such computations are out of reach, however.

In this writing, we address the general (and therefore non-abelian) free nilpotent groups N_c^r of class $c \geq 2$ and rank $r \geq 2$. Among these is the famous Heisenberg group of unit upper-triangular $(3, 3)$ -matrices with integral entries as the smallest example N_2^2 . Its classifying space is the 3-dimensional Heisenberg manifold X_2^2 . In general, the groups N_c^r arise as fundamental groups of certain aspherical nil-manifolds X_c^r that we can present as iterated torus bundles over tori.

We are interested in the symmetries of these groups and manifolds, beginning with the automorphism groups $\text{Aut}(N_c^r)$ of the groups N_c^r , and their homology. Again, we can stabilize by passing from r to $r + 1$, extending an automorphism by sending the new generator to itself. In the (co)limit $r \rightarrow \infty$, we get the stable automorphism group $\text{Aut}(N_c^\infty)$. Again, this notation is *not* used to refer to the automorphism group of a group N_c^∞ . The homology of the group $\text{Aut}(N_c^\infty)$ is the colimit of the homologies of the groups $\text{Aut}(N_c^r)$.

The homology of a group with constant coefficients is arguably the most fundamental case to consider. However, when we are dealing with families of groups, it turns out that it is often helpful to set up the computations to function in the much broader context of *polynomial* (or *finite degree*) coefficients. See [4], [9], and [33], for instance, and the proofs of Theorem 1.1 and Theorem 1.3 below. The more comprehensive setting often allows to feed inductions, and whenever possible, it is, therefore, appropriate to give the results in that generality.

THEOREM 1.1. — *Let V be a polynomial functor from the category \mathbf{ab} of finitely generated free abelian groups to the category of rational vector spaces. For every integer $c \geq 1$ the canonical homomorphism*

$$H_*(\text{Aut}(\mathbb{N}_c^\infty); V_\infty) \longrightarrow H_*(\text{GL}_\infty(\mathbb{Z}); V_\infty)$$

is an isomorphism.

By 2-out-of-3, it follows that for every integer $c \geq 2$ the canonical homomorphism

$$H_*(\text{Aut}(\mathbb{N}_c^\infty); V_\infty) \longrightarrow H_*(\text{Aut}(\mathbb{N}_{c-1}^\infty); V_\infty)$$

is an isomorphism.

In the case of the constant functor V with $V(\mathbb{Z}^r) = \mathbb{Q}$, we see that the natural homomorphism $\text{Aut}(\mathbb{N}_c^\infty) \rightarrow \text{GL}_\infty(\mathbb{Z})$ is a rational homology equivalence for every integer $c \geq 1$:

COROLLARY 1.2. — *For all integers $c \geq 2$ the canonical homomorphisms*

$$\cdots \longrightarrow \text{Aut}(\mathbb{N}_c^r) \longrightarrow \text{Aut}(\mathbb{N}_{c-1}^r) \longrightarrow \cdots \longrightarrow \text{GL}_r(\mathbb{Z})$$

induce isomorphisms in rational stable homology.

As a geometric application, we consider the automorphism groups (in various geometric categories) of the aspherical nil-manifolds X_c^r with fundamental groups isomorphic to \mathbb{N}_c^r .

The group $\pi_0 G(X_c^r)$ of components of the topological monoid $G(X_c^r)$ of homotopy self-equivalences of the classifying space X_c^r of the group \mathbb{N}_c^r is isomorphic to the group $\text{Out}(\mathbb{N}_c^r)$ of outer automorphism of the group \mathbb{N}_c^r . The following result is Theorem 6.1 in the main text.

THEOREM 1.3. — *For every class $c \geq 1$ the canonical projection induces an isomorphism*

$$H_d(\text{Aut}(\mathbb{N}_c^r); V(\mathbb{Z}^r)) \longrightarrow H_d(\text{Out}(\mathbb{N}_c^r); V(\mathbb{Z}^r))$$

in the stable range of $\text{Aut}(\mathbb{N}_c^r)$ for all finite degree functors V on \mathbf{ab} with values in rational vector spaces.

In particular, for all integers $c \geq 1$ the canonical homomorphisms

$$\mathrm{Aut}(\mathbb{N}_c^r) \rightarrow \mathrm{Out}(\mathbb{N}_c^r)$$

of groups induce isomorphisms in rational stable homology.

The following result is Theorem 7.1 in the main text. Starting from the isomorphisms $\mathrm{Out}(\mathbb{N}_c^r) \cong \pi_0 \mathbb{G}(X_c^r)$ of groups, it allows us to extend the preceding results from the homotopy category to the mapping class groups (groups of components of the automorphism groups) in the TOP, PL, and DIFF categories.

THEOREM 1.4. — *For all integers $c \geq 1$, if the nil-manifold X_c^r is of dimension at least 5, the canonical homomorphisms*

$$\pi_0 \mathrm{DIFF}(X_c^r) \longrightarrow \pi_0 \mathrm{PL}(X_c^r) \longrightarrow \pi_0 \mathrm{TOP}(X_c^r) \longrightarrow \pi_0 \mathbb{G}(X_c^r)$$

of discrete groups induce isomorphisms in rational homology.

Taken together, Theorem 1.1, Theorem 1.3, and Theorem 1.4 compute the rational stable homology of the mapping class groups of the universal nil-manifolds X_c^r , justifying the title.

Stable homology computations such as those in the present paper produce most impact in the presence of homological stability: This guarantees that the stable homology determines infinitely many unstable values. Luckily, such results are already in place for the groups of interest here: Rational homological stability for the general linear groups over the ring \mathbb{Z} of integers is due to Borel [6]. With integral coefficients, the result is due to Charney [8], Maazen (unpublished), van der Kallen [22], and Suslin [32]. Homological stability results for the automorphism groups of free groups are more recent than those for the general linear groups, see [16] and [17]. They have been revisited by Bestvina [3] lately. For the automorphism groups of free nilpotent groups of an arbitrary class c , the author has proven homological stability in [33]. In none of these cases is the exact stable range known, and we will not reproduce the estimates from the references. Our goal is the computation of the stable homology, and the mere existence of bounds for homological stability is enough to reach it.

An outline of this paper is as follows. Section 2 contains the necessary background from homological algebra in functor categories and provides for a basic rational vanishing result, Proposition 2.5. Section 3 is about stable K-theory and recalls Scorichenko's theorem. In Section 4 we review the free nilpotent groups and their automorphisms. Section 5 provides a proof of Theorem 1.1. Sections 6 and 7 contain the applications to the outer

automorphism groups and the TOP, PL, and DIFF mapping class groups of the associated aspherical nil-manifolds, respectively.

2. Functor homology

In this section, we briefly review some homological algebra in functor (and bifunctor) categories to the extent that we will need in the following sections. We also prove Proposition 2.5, a rational vanishing result that enters fundamentally into the proof of our main result.

2.1. Mac Lane homology

If \mathbf{C} is a small (or essentially small) category, then we write $\mathcal{F}(\mathbf{C})$ for the category of functors $\mathbf{C} \rightarrow \mathbf{Ab}$ from the category \mathbf{C} to the category \mathbf{Ab} of all abelian groups. This is an abelian category with enough projective objects to do homological algebra in. Note that we are looking at *all* functors, even if the category \mathbf{C} is such that it would make sense for us to consider only additive functors $\mathbf{C} \rightarrow \mathbf{Ab}$. For instance, we will be particularly interested in the case when $\mathbf{C} = \mathbf{ab}$ is the essentially small category \mathbf{ab} of finitely generated free abelian groups. Then we have the inclusion functor $I: \mathbf{ab} \rightarrow \mathbf{Ab}$ at our disposal.

A pair of functors $E: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Ab}$ and $F: \mathbf{C} \rightarrow \mathbf{Ab}$ has an external tensor product, defined by

$$(2.1) \quad (E \boxtimes F)(X, Y) = E(X) \otimes F(Y),$$

which gives a bifunctor $E \boxtimes F: \mathbf{C}^{\text{op}} \times \mathbf{C} \rightarrow \mathbf{Ab}$. Of course, not every bifunctor has this form. But every bifunctor $\mathbf{C}^{\text{op}} \times \mathbf{C} \rightarrow \mathbf{Ab}$ has a coend, and in the case of $E \boxtimes F$ this is the internal tensor product $E \otimes F$, or sometimes $E \otimes_{\mathbf{C}} F$ for emphasis. This is just an abelian group. The derived functors will be denoted by $\text{Tor}_*^{\mathcal{F}(\mathbf{C})}$ or just $\text{Tor}_*^{\mathbf{C}}$.

The following definition is due to Jibladze and Pirashvili, compare [21, Def. 1.2].

DEFINITION 2.1. — *The Mac Lane homology of the ring \mathbb{Z} of integers with coefficients in a functor $F \in \mathcal{F}(\mathbf{ab})$ is defined as*

$$\text{HML}_*(\mathbb{Z}; F) = \text{Tor}_*^{\mathcal{F}(\mathbf{ab})}(\mathbb{I}^\vee, F),$$

where $\mathbb{I}^\vee: \mathbf{ab}^{\text{op}} \rightarrow \mathbf{Ab}$ is the dual of the inclusion I , so that $\mathbb{I}^\vee(X) = \text{Hom}(X, \mathbb{Z})$.

Remark 2.2. — A functor $F \in \mathcal{F}(\mathbf{ab})$ is additive if and only if it is given (up to isomorphism) by tensoring with an abelian group [21, p. 254]. In that case, the Mac Lane homology is the Hochschild homology of the \mathbb{Q} -construction $\mathbb{Q}(\mathbb{Z})$ with coefficients in that abelian group, thought of as a (\mathbb{Z}, \mathbb{Z}) -bimodule, see [21, 2.1 and 6.2]. Also, in that case, the Mac Lane homology agrees with the topological Hochschild homology (THH) of the Eilenberg–Mac Lane spectrum $\mathbb{H}\mathbb{Z}$ with coefficients in the associated bimodule spectrum, see [30, Thm. 3.2] and [10, Thm. 2.10].

If \mathbb{Z} also denotes the constant functor $\mathbf{C}^{\text{op}} \rightarrow \mathbf{Ab}$ with value the group \mathbb{Z} of integers, and $F: \mathbf{C} \rightarrow \mathbf{Ab}$ is any functor, the homology of the category \mathbf{C} with coefficients in the functor F is defined as

$$(2.2) \quad H_*(\mathbf{C}; F) = \text{Tor}_*^{\mathcal{F}(\mathbf{C})}(\mathbb{Z}, F).$$

Both the Mac Lane homology as well as the homology of categories are special cases of the Hochschild (bifunctor) homology, as we shall now see.

2.2. Hochschild (bifunctor) homology

Let $D: \mathbf{C}^{\text{op}} \times \mathbf{C} \rightarrow \mathbf{Ab}$ be a bifunctor. For instance, we might have $D = E \boxtimes F$ as in (2.1). Or we might have $D = J$, where $J(X, Y) = \mathbb{Z}\mathbf{C}(X, Y)$ is the free abelian group on the set $\mathbf{C}(X, Y)$ of morphisms $X \rightarrow Y$ in \mathbf{C} . The bifunctor J is useful even when \mathbf{C} is already enriched in abelian groups: The Hochschild homology of the category \mathbf{C} with coefficients in the bifunctor D is defined as

$$\text{HH}_*(\mathbf{C}; D) = \text{Tor}_*^{\mathcal{F}(\mathbf{C}^{\text{op}} \times \mathbf{C})}(J, D).$$

See Loday [26, App. C.10] and Franjou–Pirashvili [13, 2.5]. They use the notation H for this, but we shall reserve it for the homology of categories with coefficients in a plain functor (rather than a bifunctor) as in (2.2). The latter is a special case:

PROPOSITION 2.3. — *For all functors $F \in \mathcal{F}(\mathbf{C})$ there is an isomorphism*

$$(2.3) \quad H_*(\mathbf{C}; F) \cong \text{HH}_*(\mathbf{C}; F \circ \text{pr}_2),$$

where $\text{pr}_2: \mathbf{C}^{\text{op}} \times \mathbf{C} \rightarrow \mathbf{C}$ is the projection.

Proof. — Note that we can write the composition of functors as $F \circ \text{pr}_2 = \mathbb{Z} \boxtimes F$, so that both sides of (2.3) are isomorphic to $\text{Tor}_*^{\mathcal{F}(\mathbf{C})}(\mathbb{Z}, F)$ by [13, Prop. 2.10 on p. 115] which says $\text{Tor}_*^{\mathcal{F}(\mathbf{C})}(E, F) \cong \text{HH}_*(\mathbf{C}; E \boxtimes F)$ in our notation, when E takes values in free abelian groups. \square

PROPOSITION 2.4. — *For all functors $F \in \mathcal{F}(\mathbf{ab})$ there is an isomorphism*

$$\mathrm{HML}_*(\mathbb{Z}; F) \cong \mathrm{HH}_*(\mathbf{ab}; \mathrm{I}^\vee \boxtimes F),$$

where $\mathrm{I}^\vee \boxtimes F$ is the bifunctor with $(\mathrm{I}^\vee \boxtimes F)(X, Y) \cong \mathrm{Hom}(X, FY)$.

Proof. — Both sides are isomorphic to $\mathrm{Tor}_*^{\mathcal{F}(\mathbf{ab})}(\mathrm{I}^\vee, F)$: the left hand side by Definition 2.1, and the right hand side by [13, Prop. 2.10 on p. 115] again. Alternatively, see [26, 13.1.7, 13.2.17]. \square

2.3. A rational vanishing result

The following rational vanishing result will be the main computational input in our proof of Theorem 1.1. It refers to the Lie functors $\mathrm{Lie}^b: \mathbf{ab} \rightarrow \mathbf{Ab}$, that is the degree b homogeneous part of the free Lie algebra functor. We will abbreviate

$$\mathrm{Lie}^{[a,c]} = \bigoplus_{b=a}^c \mathrm{Lie}^b,$$

so that $\mathrm{Lie}^{[1,c]}(\mathbb{Z}^r)$ is nothing but the free nilpotent Lie algebra of class c on r generators.

PROPOSITION 2.5. — *For all $c \geq 2$ and $q \geq 1$, the Hochschild homology groups*

$$\mathrm{HH}_*(\mathbf{ab}; (\mathrm{I}^\vee \boxtimes \mathrm{Lie}^{[2,c]})^{\otimes q} \otimes V)$$

vanish whenever V is a functor on \mathbf{ab} with values in rational vector spaces.

Proof. — We will assume that the functor V is constant, so that we are dealing with the groups $\mathrm{HH}_*(\mathbf{ab}; (\mathrm{I}^\vee \boxtimes \mathrm{Lie}^{[2,c]})^{\otimes q})$ rationally. The same argument gives the result in the general case, inserting $\otimes V$ in suitable places.

We can check the first case $q = 1$ directly, because

$$\mathrm{HH}_*(\mathbf{ab}; \mathrm{I}^\vee \boxtimes \mathrm{Lie}^{[2,c]}) \cong \mathrm{HML}_*(\mathbb{Z}; \mathrm{Lie}^{[2,c]})$$

is isomorphic to the Mac Lane homology of the ring \mathbb{Z} of integers with coefficients in the functor $\mathrm{Lie}^{[2,c]}$ by Proposition 2.4. The Lie functors Lie^b are rationally retracts of the tensor powers \bigotimes^b : There are morphisms

$$\mathrm{Lie}^b \longrightarrow \bigotimes^b \longrightarrow \mathrm{Lie}^b$$

such that their composition is multiplication with b [31, Prop. 3.3]. Since the functor \otimes^b has vanishing Mac Lane homology by Pirashvili's Lemma (compare [12, Lem. 2.2]), so have the functors Lie^b , and therefore also their direct sum

$$\text{Lie}^{[2,c]} = \bigoplus_{b=2}^c \text{Lie}^b.$$

This proves the claim in the case $q = 1$.

To finish the proof, we reduce the case $q \geq 2$ to the previous one: There is an obvious isomorphism

$$(\mathbb{I}^\vee \boxtimes \text{Lie}^{[2,c]})^{\otimes q} \cong (\mathbb{I}^\vee)^{\otimes q} \boxtimes (\text{Lie}^{[2,c]})^{\otimes q},$$

and by another invocation of [13, Prop. 2.10 on p. 115] we have

$$\text{HH}_*(\mathbf{ab}; (\mathbb{I}^\vee)^{\otimes q} \boxtimes (\text{Lie}^{[2,c]})^{\otimes q}) \cong \text{Tor}_*^{\mathcal{F}(\mathbf{ab})}((\mathbb{I}^\vee)^{\otimes q}, (\text{Lie}^{[2,c]})^{\otimes q}).$$

The right hand side can be computed, using the Künneth theorem for Tor , from

$$\text{Tor}_*^{\mathcal{F}(\mathbf{ab})}(\mathbb{I}^\vee, \text{Lie}^{[2,c]})^{\otimes q},$$

and it therefore suffices to show that this is zero. But, working backwards, we already know that this is isomorphic to

$$\text{HH}_*(\mathbf{ab}; \mathbb{I}^\vee \boxtimes \text{Lie}^{[2,c]})^{\otimes q} \cong \text{HML}_*(\mathbb{Z}; \text{Lie}^{[2,c]})^{\otimes q},$$

and this vanishes as explained for $q = 1$. □

Remark 2.6. — In the case $c = 2$ and $q = 1$ of the preceding result, the Mac Lane cohomology of the ring of integers with coefficients in the functor $\text{Lie}^2 = \Lambda^2$ has been computed integrally by Franjou and Pirashvili [12, Cor. 2.3]. It is all 2-torsion.

3. Stable K-theory

There is a simple trick (due to Waldhausen, compare [35, Sec. 6]) that helps us compute the stable homology

$$(3.1) \quad \text{H}_*(\text{GL}_\infty(\mathbb{Z}); D_\infty) = \text{colim}_r \text{H}_*(\text{GL}_r(\mathbb{Z}); D(\mathbb{Z}^r, \mathbb{Z}^r))$$

of the general linear groups with coefficients in a module D_∞ that comes from a bifunctor $D: \mathbf{ab} \times \mathbf{ab}^{\text{op}} \rightarrow \mathbf{Ab}$: We can use the Serre spectral sequence for the fibration

$$X \longrightarrow \text{BGL}_\infty(\mathbb{Z}) \longrightarrow \text{BGL}_\infty(\mathbb{Z})^+,$$

where the space X is the homotopy fiber of the indicated Quillen plus construction. The groups on the E^2 page take the form $H_*(\mathrm{BGL}_\infty(\mathbb{Z})^+; H_*(X; D_\infty))$, where the coefficients $H_*(X; D_\infty)$ are the homology of the fiber X with coefficients in the $\pi_1(X)$ -module $D_\infty = \mathrm{colim}_r D(\mathbb{Z}^r, \mathbb{Z}^r)$. The fundamental group $\pi_1(X)$ acts on D_∞ via the homomorphism

$$\pi_1(X) \rightarrow \pi_1(\mathrm{BGL}_\infty(\mathbb{Z})) \cong \mathrm{GL}_\infty(\mathbb{Z}).$$

Remark 3.1. — The action of the fundamental group of $\mathrm{BGL}_\infty(\mathbb{Z})^+$ on these coefficients $H_*(X; D_\infty)$ is now trivial, see [23, Thm. 3.1] and [24, proof of Thm. 2.16]. We will not need this fact, since we will show that in our situation the coefficients itself will be trivial.

Since the Quillen plus construction is an acyclic map, it induces homology isomorphisms with respect to all coefficients, twisted and untwisted. Therefore, we can remove the plus to get a spectral sequence

$$(3.2) \quad E_{s,t}^2 \implies H_{s+t}(\mathrm{GL}_\infty(\mathbb{Z}); D_\infty),$$

with

$$(3.3) \quad E_{s,t}^2 = H_s(\mathrm{GL}_\infty(\mathbb{Z}); K_t^{\mathrm{st}}(\mathbb{Z}; D))$$

where, by definition,

$$(3.4) \quad K_*^{\mathrm{st}}(\mathbb{Z}; D) = H_*(X; D_\infty)$$

is the *stable K-theory* of the ring of integers with coefficients in the bifunctor D .

Remark 3.2. — Bökstedt pointed out that this spectral sequence often degenerates. See also [5, proof of Thm. 2]. Again, this will be obvious in our situation, because the entire E^2 page will be trivial.

The definition (3.4) of stable K-theory is topological in nature. We will need the following result that gives an algebraic description of it.

THEOREM 3.3 (Scorichenko). — *For any ring R , let \mathbf{mod}_R be the category of finitely generated projective left R -modules, and let D be a bifunctor on it. If D has finite degree with respect to each of the variables, then there is an isomorphism between Waldhausen’s stable K-theory and the Hochschild homology of the category \mathbf{mod}_R :*

$$K_*^{\mathrm{st}}(R; D) \cong \mathrm{HH}_*(\mathbf{mod}_R; D).$$

See the exposition [13] by Franjou and Pirashvili. The cited result is stated as Theorem 1.1 there, and the proof given is complete for rings R with the property that submodules of finitely generated projective left R -modules are still finitely generated and projective. This property is satisfied for $R = \mathbb{Z}$, when $\mathbf{mod}_{\mathbb{Z}} = \mathbf{ab}$, which is the only case that we will be using here. See also [9, 5.2] for an enlightening discussion, and [11, App.].

4. Free nilpotent groups and their automorphisms

In this section, we present some basic results about the free nilpotent groups and their automorphisms. Most of this must be well-known, and we can refer to the exposition in [33], for instance. We will, however, prove the two Propositions 4.1 and 4.3 which will later be used.

Let G be a (discrete) group. For integers $n \geq 1$, the subgroups $\Gamma_n(G)$ are defined inductively by $\Gamma_1(G) = G$ and $\Gamma_{n+1}(G) = [G, \Gamma_n(G)]$. We also set $\Gamma_\infty(G)$ to be the intersection of all the $\Gamma_n(G)$. This gives a series

$$G = \Gamma_1(G) \geq \Gamma_2(G) \geq \cdots \geq \Gamma_\infty(G)$$

of normal subgroups, the *descending/lower central series* of G . The associated graded group is abelian, and the commutator bracket induces the structure of a graded Lie algebra $\mathrm{Lie}(G)$ on it.

Let F_r denote a free group on a set of r generators. In this case, an old theorem of Magnus says that the subgroup $\Gamma_\infty(F_r)$ is trivial, so that the free groups F_r are *residually nilpotent*. By Witt's theorem, the canonical homomorphism from a free Lie algebra on a set of r generators to the associated graded Lie algebra of F_r is an isomorphism. In particular

$$(4.1) \quad \Gamma_n(F_r)/\Gamma_{n+1}(F_r) \cong \mathrm{Lie}^n(\mathbb{Z}^r)$$

is a free abelian group, the degree n homogeneous part of the free Lie algebra on a set of r generators. For instance, we have $\mathrm{Lie}^1 = \mathrm{Id}$ and $\mathrm{Lie}^2 = \Lambda^2$.

4.1. Free nilpotent groups

The universal examples of nilpotent groups of class $c \geq 1$ are the quotients $N_c^r = F_r/\Gamma_{c+1}(F_r)$. As the two extreme cases, we obtain $N_1^r \cong \mathbb{Z}^r$ and $N_\infty^r \cong F_r$. It follows from (4.1) that there are extensions

$$(4.2) \quad 0 \longrightarrow \mathrm{Lie}^c(\mathbb{Z}^r) \longrightarrow N_c^r \longrightarrow N_{c-1}^r \longrightarrow 1$$

of groups.

4.2. Rational homology

We will later need the following structural result on the rational homology of the groups N_c^r .

PROPOSITION 4.1. — *For every class $c \geq 1$ and every degree d , there is a polynomial functor P , depending on c and d , and of degree at most cd , from **ab** to rational vector spaces, such that $H_d(N_c^r; \mathbb{Q}) \cong P(\mathbb{Z}^r)$ for all r .*

Proof. — Since we are only interested in the rational homology of nilpotent groups, we may just as well consider the rationalizations/Malcev completions $N_c^r \otimes \mathbb{Q}$, see [25, Sec. 4]. We have natural isomorphisms

$$H_d(N_c^r; \mathbb{Q}) \cong H_d(N_c^r \otimes \mathbb{Q}; \mathbb{Q}).$$

Then, the category of uniquely divisible nilpotent groups is equivalent to the category of nilpotent Lie algebras over \mathbb{Q} , and we get a natural isomorphism

$$H_d(N_c^r; \mathbb{Q}) \cong H_d(\text{Lie}(N_c^r \otimes \mathbb{Q}); \mathbb{Q}),$$

as demonstrated by Nomizu [28, Thm. 1] and, more generally and explicitly, by Pickel [29, Thm. 10]. As a consequence of Witt’s theorem, the Lie algebra $\text{Lie}(N_c^r \otimes \mathbb{Q})$ of the group of interest is nothing else than the free nilpotent Lie algebra $\text{Lie}^{[1,c]}(\mathbb{Q}^r)$ on r generators, and this is clearly functorial in \mathbb{Q}^r (and in particular in \mathbb{Z}^r) of degree c . The d -th homology of a Lie algebra \mathfrak{g} is the homology of the Chevalley–Eilenberg resolution

$$\cdots \longleftarrow \Lambda^d \mathfrak{g} \longleftarrow \cdots .$$

As a subquotient of the functor $\Lambda^d \text{Lie}(N_c^r \otimes \mathbb{Q})$, which is of degree cd , the homology is then also functorial of degree at most cd . \square

4.3. Automorphisms

More than in the free nilpotent groups N_c^r themselves, we are interested in the groups $\text{Aut}(N_c^r)$ of their automorphisms. In the case $c = 1$, these are the general linear groups $\text{GL}_r(\mathbb{Z})$, and in the limiting case $c = \infty$, these are the automorphism groups $\text{Aut}(F_r)$ of free groups.

PROPOSITION 4.2. — *There are extensions*

$$(4.3) \quad 0 \longrightarrow \text{Hom}(\mathbb{Z}^r, \text{Lie}^c(\mathbb{Z}^r)) \longrightarrow \text{Aut}(N_c^r) \longrightarrow \text{Aut}(N_{c-1}^r) \longrightarrow 1$$

of groups.

See again [33, Prop. 3.1], for instance. Note that any element of the quotient group $\text{Aut}(N_{c-1}^r)$ acts on the kernel group via restriction along the canonical projection $\text{Aut}(N_{c-1}^r) \rightarrow \text{Aut}(N_1^r) = \text{GL}_r(\mathbb{Z})$. For that reason, it is sometimes more efficient to study the extensions

$$(4.4) \quad 1 \longrightarrow \text{IA}_c^r \longrightarrow \text{Aut}(N_c^r) \longrightarrow \text{GL}_r(\mathbb{Z}) \longrightarrow 1$$

of groups. The kernels IA_c^r are known to be nilpotent of class $c - 1$. See for instance Andreadakis [1, Cor. 1.3.]. By what has already been said, it is clear that they differ by free abelian groups in the sense that there are extensions

$$(4.5) \quad 0 \longrightarrow \text{Hom}(\mathbb{Z}^r, \text{Lie}^c(\mathbb{Z}^r)) \longrightarrow \text{IA}_c^r \longrightarrow \text{IA}_{c-1}^r \longrightarrow 1.$$

4.4. On the Andreadakis–Johnson module

We can now put this together to get the following result that will later help us to estimate homology.

PROPOSITION 4.3. — *The $\text{GL}_r(\mathbb{Z})$ -module $\text{H}_q(\text{IA}_c^r) \otimes \mathbb{Q}$ is isomorphic to a direct summand of the $\text{GL}_r(\mathbb{Z})$ -module $\Lambda^q(\text{Hom}(\mathbb{Z}^r, \text{Lie}^{[2,c]}(\mathbb{Z}^r)) \otimes \mathbb{Q}$.*

Proof. — Our starting point are the Lyndon–Hochschild–Serre spectral sequences associated with the extensions (4.5). We have

$$\text{E}_{s,t}^2 \implies \text{H}_{s+t}(\text{IA}_c^r)$$

with

$$(4.6) \quad \text{E}_{s,t}^2 = \text{H}_s(\text{IA}_{c-1}^r; \text{H}_t \text{Hom}(\mathbb{Z}^r, \text{Lie}^c(\mathbb{Z}^r))).$$

Since we are working rationally, and GL_r is reductive, every short exact sequence splits. It therefore suffices to show that the filtration quotients $\text{E}_{s,q-s}^\infty$ of the homology $\text{H}_q(\text{IA}_c^r)$ form a direct summand of $\Lambda^q(\text{Hom}(\mathbb{Z}^r, \text{Lie}^{[2,c]}(\mathbb{Z}^r)))$. In turn, this will follow once we know the same for the terms $\text{E}_{s,q-s}^2$ on the E^2 page, because the groups $\text{E}_{s,q-s}^\infty$ are subquotients (and rationally direct summands) thereof.

As for the groups (4.6) on the E^2 page, it is evident by construction that the actions of the kernels IA_{c-1}^r of the canonical projections $\text{Aut}(N_c^r) \rightarrow \text{GL}_r(\mathbb{Z})$ on the coefficient groups $\text{Hom}(\mathbb{Z}^r, \text{Lie}^c(\mathbb{Z}^r))$ are trivial. Therefore, we can write the E^2 term as a graded tensor product

$$\text{E}_{*,*}^2 \cong \text{H}_*(\text{IA}_{c-1}^r) \otimes \Lambda^* \text{Hom}(\mathbb{Z}^r, \text{Lie}^c(\mathbb{Z}^r)),$$

using that the homology of the free abelian groups is given by the exterior powers. We see that $H_q(\mathbf{IA}_c^r)$ is a direct summand of the degree q part of

$$H_*(\mathbf{IA}_{c-1}^r) \otimes \Lambda^* \text{Hom}(\mathbb{Z}^r, \text{Lie}^c(\mathbb{Z}^r)).$$

By induction on c , this is isomorphic to a direct summand of the degree q part of

$$\begin{aligned} \Lambda^* \text{Hom}(\mathbb{Z}^r, \text{Lie}^2(\mathbb{Z}^r)) \otimes \cdots \otimes \Lambda^* \text{Hom}(\mathbb{Z}^r, \text{Lie}^c(\mathbb{Z}^r)) \\ \cong \Lambda^* \text{Hom}(\mathbb{Z}^r, \text{Lie}^2(\mathbb{Z}^r) \oplus \cdots \oplus \text{Lie}^c(\mathbb{Z}^r)), \end{aligned}$$

which is $\Lambda^q \text{Hom}(\mathbb{Z}^r, \text{Lie}^{[2,c]}(\mathbb{Z}^r))$, as claimed. \square

Tensoring with another rational representation $V(\mathbb{Z}^r)$ that originates in a functor V from the category of finitely generated free abelian groups to the category of rational vector spaces, we obtain:

COROLLARY 4.4. — *For every functor V from the category **ab** of the finitely generated free abelian groups to the category of rational vector spaces, the $\text{GL}_r(\mathbb{Z})$ -module $H_q(\mathbf{IA}_c^r) \otimes V(\mathbb{Z}^r)$ is a direct summand (up to isomorphism) of the $\text{GL}_r(\mathbb{Z})$ -module $\Lambda^q(\text{Hom}(\mathbb{Z}^r, \text{Lie}^{[2,c]}(\mathbb{Z}^r)) \otimes V(\mathbb{Z}^r))$.*

5. A proof of Theorem 1.1

We now have enough ingredients to prove Theorem 1.1.

Proof. — The homomorphism

$$(5.1) \quad H_*(\text{Aut}(\mathbb{N}_c^\infty); V_\infty) \longrightarrow H_*(\text{GL}_\infty(\mathbb{Z}); V_\infty)$$

in question is the edge homomorphism in the Lyndon–Hochschild–Serre spectral sequence

$$E_{p,q}^2 \implies H_{p+q}(\text{Aut}(\mathbb{N}_c^\infty); V_\infty)$$

for the colimit of the extensions (4.4), with E^2 page

$$(5.2) \quad E_{p,q}^2 = H_p(\text{GL}_\infty(\mathbb{Z}); H_q(\mathbf{IA}_c^\infty; V_\infty)).$$

Since \mathbf{IA}_c^∞ maps trivially to the general linear groups, we can pull the coefficients out:

$$E_{p,q}^2 \cong H_p(\text{GL}_\infty(\mathbb{Z}); H_q(\mathbf{IA}_c^\infty) \otimes V_\infty).$$

In fact, the 0-line is just the homology of the stable general linear group with coefficients in V_∞ . We will show that the other groups (i.e. the lines with $q \geq 1$) on the E^2 page vanish rationally:

$$(5.3) \quad H_*(\text{GL}_\infty(\mathbb{Z}); H_q(\mathbf{IA}_c^\infty) \otimes V_\infty) = 0$$

for all $c \geq 2$ and $q \geq 1$. This will imply the result.

We will prove (5.3) using some very crude estimates. (Here these always originate from spectral sequences.) The first one is Corollary 4.4, which implies that it suffices to be shown that

$$(5.4) \quad H_*(GL_\infty(\mathbb{Z}); \Lambda^q(I^\vee \boxtimes \text{Lie}^{[2,c]})_\infty \otimes V_\infty) = 0$$

for all $c \geq 2$ and $q \geq 1$. As explained in Section 3, such twisted homology can be computed from the stable K-theory $K_*^{\text{st}}(\mathbb{Z}; \Lambda^q(I^\vee \boxtimes \text{Lie}^{[2,c]}) \otimes V)$ using the spectral sequence (3.2) and (3.3) for the bifunctor $D = \Lambda^q(I^\vee \boxtimes \text{Lie}^{[2,c]}) \otimes V$:

$$E_{s,t}^2 \implies H_{s+t}(GL_\infty(\mathbb{Z}); \Lambda^q(I^\vee \boxtimes \text{Lie}^{[2,c]})_\infty \otimes V_\infty)$$

with

$$E_{s,t}^2 = H_s(GL_\infty(\mathbb{Z}); K_t^{\text{st}}(\mathbb{Z}; \Lambda^q(I^\vee \boxtimes \text{Lie}^{[2,c]}) \otimes V)).$$

The bifunctor $\Lambda^q(I^\vee \boxtimes \text{Lie}^{[2,c]}) \otimes V$ at hand is of finite degree in each variable, so that we can use Scorichenko's Theorem 3.3 to the effect that stable K-theory is bifunctor homology,

$$K_*^{\text{st}}(\mathbb{Z}; \Lambda^q(I^\vee \boxtimes \text{Lie}^{[2,c]}) \otimes V) \cong \text{HH}_*(\mathbf{ab}; \Lambda^q(I^\vee \boxtimes \text{Lie}^{[2,c]}) \otimes V).$$

Thus, it suffices to see that the right hand side is zero rationally. Since the functor Λ^q is a retract of \otimes^q , this follows from Proposition 2.5. \square

Remark 5.1. — By Betley's work [4, Thm.4.2] on the homology of the general linear groups with twisted coefficients, the right hand side of (5.1) (and therefore also the left hand side) vanishes if the functor V is reduced, that is if we have $V(0) = 0$.

6. Outer automorphism groups

The outer automorphism group $\text{Out}(G)$ of a group G is the quotient in an extension

$$1 \longrightarrow \text{Inn}(G) \longrightarrow \text{Aut}(G) \longrightarrow \text{Out}(G) \longrightarrow 1,$$

with inner automorphism group $\text{Inn}(G) \cong G/Z(G)$ and the projection gives a factorization $\text{Aut}(G) \rightarrow \text{Out}(G) \rightarrow \text{Aut}(H_*G)$ of the canonical homomorphism to the automorphism group of the homology/abelianization: Inner automorphisms act trivially on homology.

In the case when $G = N_c^r$ is a free nilpotent group, the center is the kernel of the canonical projection $N_c^r \rightarrow N_{c-1}^r$, so that we have

$$Z(N_c^r) \cong \text{Lie}^c(\mathbb{Z}^r)$$

and

$$\text{Inn}(\mathbb{N}_c^r) \cong \mathbb{N}_{c-1}^r.$$

Consequently, we get an extension

$$(6.1) \quad 1 \longrightarrow \mathbb{N}_{c-1}^r \longrightarrow \text{Aut}(\mathbb{N}_c^r) \longrightarrow \text{Out}(\mathbb{N}_c^r) \longrightarrow 1.$$

In parallel with (4.4), we define subgroups $\text{IO}_c^r \leq \text{Out}(\mathbb{N}_c^r)$ as the kernels in the extensions

$$(6.2) \quad 1 \longrightarrow \text{IO}_c^r \longrightarrow \text{Out}(\mathbb{N}_c^r) \longrightarrow \text{GL}_r(\mathbb{Z}) \longrightarrow 1.$$

This gives extensions

$$(6.3) \quad 1 \longrightarrow \mathbb{N}_{c-1}^r \longrightarrow \text{IA}_c^r \longrightarrow \text{IO}_c^r \longrightarrow 1$$

of nilpotent groups of class $c - 1$.

This finishes our description of the outer automorphism groups $\text{Out}(\mathbb{N}_c^r)$ up to extensions. We can proceed to compute their rational homology.

Let V be a polynomial functor (of finite degree) from the category **ab** of finitely generated free abelian groups to the category of rational vector spaces. This defines twisted coefficients for both of the groups $\text{Aut}(\mathbb{N}_c^\infty)$ and $\text{Out}(\mathbb{N}_c^\infty)$ via their homomorphisms to the general linear groups.

THEOREM 6.1. — *For every class $c \geq 1$ the canonical projection induces an isomorphism*

$$(6.4) \quad \text{H}_d(\text{Aut}(\mathbb{N}_c^r); V(\mathbb{Z}^r)) \longrightarrow \text{H}_d(\text{Out}(\mathbb{N}_c^r); V(\mathbb{Z}^r))$$

*in the stable range of $\text{Aut}(\mathbb{N}_c^r)$ for all finite degree functors V on **ab** with values in rational vector spaces.*

Remark 6.2. — The existence of a stable range for the homology of the automorphism groups (with polynomial coefficients) has been established in [33, Thm. 4.5], even integrally. The proof of Theorem 6.1 given below uses that stability result. It says that there exists an integer $R = R(c, d, V)$ such that the homology $\text{H}_d(\text{Aut}(\mathbb{N}_c^r); V(\mathbb{Z}^r))$ is independent (in a precise sense) of r as soon as $r \geq R$. At the time of writing, the best possible value is not known.

Remark 6.3. — The theorem implies the existence of a stable range for the homology of the *outer* automorphism groups as well, at least rationally. We note that, in particular, the rational homology vanishes in the stable range if the functor V is reduced, by Theorem 1.1 and Remark 5.1 following it.

Proof of Theorem 6.1. — We prove this by induction on the homological degree d .

We start the induction in degree $d = 0$. This case concerns the co-invariants of the actions on $V(\mathbb{Z}^r)$. Since the groups $\text{Aut}(\mathbb{N}_c^r)$ surject onto the groups $\text{Out}(\mathbb{N}_c^r)$, the co-invariants are the same.

Let us now assume that the degree d is positive, and that the result has already been proven for all smaller degrees than that. We consider the Lyndon–Hochschild–Serre spectral sequence

$$E_{s,t}^2(r) = H_s(\text{Out}(\mathbb{N}_c^r); H_t(\mathbb{N}_{c-1}^r; V(\mathbb{Z}^r))) \implies H_{s+t}(\text{Aut}(\mathbb{N}_c^r); V(\mathbb{Z}^r))$$

for the extension (6.1) in the stable range, that is for r so large that the homology groups under consideration represent the stable values. (There are only finitely many of them relevant at each given time.) We would like to show that the rows with $t \neq 0$ vanish, so that the edge homomorphism (which is the homomorphism (6.4) in question) is an isomorphism.

Since the action on the coefficients $V(\mathbb{Z}^r)$ factors through the quotient $\text{Out}(\mathbb{N}_c^r)$, the coefficients for the homology $H_t(\mathbb{N}_{c-1}^r; V(\mathbb{Z}^r))$ are actually untwisted.

$$H_t(\mathbb{N}_{c-1}^r; V(\mathbb{Z}^r)) \cong H_t(\mathbb{N}_{c-1}^r) \otimes V(\mathbb{Z}^r)$$

From Proposition 4.1 we deduce that this is a polynomial functor (of finite degree). And if $t \neq 0$, then this functor is reduced: Since the group \mathbb{N}_{c-1}^0 is trivial, so is the homology $H_t(\mathbb{N}_{c-1}^0)$ for $t \neq 0$. It follows by induction (and Remark 6.3) that the groups $E_{s,t}^2(r)$ vanish for $t \neq 0$, and we are left with an isomorphism

$$H_d(\text{Aut}(\mathbb{N}_c^r); V(\mathbb{Z}^r)) \cong E_{d,0}^2(r) = H_d(\text{Out}(\mathbb{N}_c^r); V(\mathbb{Z}^r)),$$

as desired. □

Remark 6.4. — The basic set-up for this proof is the same as in [33, Sec. 1]. There it was used to prove homological stability, whereas here it is used to compute stable homology.

Combining the specializations of both Theorems 1.1 and 6.1 to the case of constant coefficients we get:

COROLLARY 6.5. — *The canonical projection induces an isomorphism*

$$H_*(\text{Out}(\mathbb{N}_c^\infty); \mathbb{Q}) \longrightarrow H_*(\text{GL}_\infty(\mathbb{Z}); \mathbb{Q})$$

in rational homology for all classes $c \geq 1$.

7. Mapping class groups

The free nilpotent groups N_c^r of class c on r generators arise as fundamental groups of aspherical nil-manifolds: Upon passage to classifying spaces, the extensions (4.2) gives rise to iterated torus bundles over tori. These manifolds will be denoted by X_c^r here. We can now spell out some implications of our preceding results for the mapping class groups of the X_c^r for the various categories of automorphisms. Note that the dimension $\dim(X_c^r)$ of the manifolds X_c^r increases with r , so that for “large” enough r , which is the situation that we are interested in here, surgery and concordance/pseudo-isotopy theory apply and will compute the mapping class groups in the TOP, PL, and DIFF categories. For basic information about the automorphism groups of manifolds in these categories, we refer to the survey articles by Burghelea [7], Hatcher [15], Balcerak–Hajduk [2], and Weiss–Williams [37].

We begin with the homotopy category. For any discrete group G , the group $\text{Out}(G)$ is isomorphic to the group of components of the topological monoid $G(X)$ of homotopy self-equivalences of the classifying space X of G . For the groups $G = N_c^r$ with classifying spaces X_c^r , we therefore have an isomorphism

$$(7.1) \quad \pi_0 G(X_c^r) \cong \text{Out}(N_c^r)$$

of groups, and Corollary 6.5 directly gives the stable rational homology of the symmetries of the manifolds X_c^r in the homotopy category.

We now explain how to transfer this to the TOP, PL, and DIFF categories:

THEOREM 7.1. — *For all integers $c \geq 1$, if the nil-manifold X_c^r is of dimension at least 5, the canonical homomorphisms*

$$\pi_0 \text{DIFF}(X_c^r) \longrightarrow \pi_0 \text{PL}(X_c^r) \longrightarrow \pi_0 \text{TOP}(X_c^r) \longrightarrow \pi_0 G(X_c^r)$$

of discrete groups induce isomorphisms in rational homology of discrete groups.

Remark 7.2. — The first case $c = 1$ of this result is already known as a consequence of the (independent) work of Hsiang–Sharpe [19] and Hatcher [15]. In order to generalize their arguments, we need to know the vanishing of surgery obstructions for manifolds with free nilpotent fundamental groups. Since the latter are poly- \mathbb{Z} groups by (4.1), we can refer to Wall [36, 15B]. Waldhausen [34, Thms. 17.5, 19.4] has shown that the Whitehead spaces vanish for a class of groups that contains all poly- \mathbb{Z} groups.

Proof of Theorem 7.1. — We start with the homeomorphisms. It is clear that the canonical homomorphism $\pi_0 \text{TOP}(X_c^r) \rightarrow \pi_0 \text{G}(X_c^r)$ is surjective. For $\dim(X_c^r) \geq 5$ this follows from surgery theory, see for instance [36, Thm. 15B.1]. In our situation, it can also be seen explicitly, and for all X_c^r , because the projection $\text{Aut}(N_c^r) \rightarrow \text{Out}(N_c^r)$ is evidently surjective, and the group $\text{Aut}(N_c^r)$ tautologically acts on N_c^r , on the completion $N_c^r \otimes \mathbb{R}$ of $N_c^r \otimes \mathbb{Q}$, and therefore also on the quotient $X_c^r \cong N_c^r \otimes \mathbb{R}/N_c^r$. This action is algebraic, hence smooth.

We need to estimate the kernel of the homomorphism from $\pi_0 \text{TOP}(X_c^r)$ to $\pi_0 \text{G}(X_c^r)$, which is isomorphic to the fundamental group $\pi_1(\text{G}(X_c^r)/\text{TOP}(X_c^r))$. It follows from surgery theory (see again [36, Sec. 15B, 17A] and the remark preceding this proof) that $\text{G}(X_c^r)/\widetilde{\text{TOP}}(X_c^r)$ is contractible, where the notation $\widetilde{\text{TOP}}$ as usually refers to the block homeomorphism group. We thus have an isomorphism

$$\pi_1(\text{G}(X_c^r)/\text{TOP}(X_c^r)) \cong \pi_1(\widetilde{\text{TOP}}(X_c^r)/\text{TOP}(X_c^r)).$$

The right hand side can be determined from concordance/pseudo-isotopy theory as follows: Work of Hatcher and Igusa ([15, Sec. 2, 3] and [20, Sec. 8]) shows that this group is dominated by π_0 of the concordance space of X_c^r . Therefore, the only possible contribution to π_0 is a quotient of the 2-torsion group $\mathbb{F}_2[N_c^r]/\mathbb{F}_2$. We see that this vanishes rationally (in fact, away from 2), so that

$$\pi_0 \text{TOP}(X_c^r) \longrightarrow \pi_0 \text{G}(X_c^r)$$

is a rational homology isomorphism if $\dim(X_c^r) \geq 5$. This settles the TOP case.

Since the group $\pi_1(\widetilde{\text{CAT}}(X_c^r)/\text{CAT}(X_c^r))$ is independent of the category CAT of symmetries considered [15], we see that it vanishes rationally for the remaining two cases $\text{CAT} = \text{PL}$ and $\text{CAT} = \text{DIFF}$ as well. For this reason, the kernels of the homomorphisms

$$(7.2) \quad \pi_0 \text{DIFF}(X_c^r) \longrightarrow \pi_0 \text{PL}(X_c^r) \longrightarrow \pi_0 \text{TOP}(X_c^r)$$

between the mapping class groups in the various categories of structures are the same as the kernels between the corresponding blocked groups $\pi_0 \widetilde{\text{CAT}}(X_c^r)$, and it remains to be checked that these kernels are rationally trivial.

From surgery theory, see the references for [36, Sec. 17A], there exists an equivalence between the quotient space $\widetilde{\text{TOP}}(X_c^r)/\widetilde{\text{PL}}(X_c^r)$, which measures the difference between the two block automorphism groups, and the function space of maps $X_c^r \rightarrow \text{TOP}/\text{PL}$. The work of Kirby–Siebenmann (see [27, Rem. 2.25] for instance) shows that the target space TOP/PL is

an Eilenberg–Mac Lane space of type $(\mathbb{Z}/2, 3)$, which is rationally trivial. Therefore, the function space is also rationally trivial.

We similarly have an equivalence between $\widetilde{\text{PL}}(X_c^r)/\widetilde{\text{DIFF}}(X_c^r)$ and the function space of maps $X_c^r \rightarrow \text{PL}/\text{O}$. By work of Cerf and Kervaire–Milnor, the target PL/O also has finite homotopy groups, essentially the groups of exotic spheres, see [27, Rem. 4.21]. We can therefore argue as in the paragraph before. \square

Acknowledgements. I thank A. Djament for his constructive comments on early drafts of this text. His and a referee’s suggestions led to substantial improvements. I also thank B. I. Dundas, W. G. Dwyer, T. Pirashvili, A. Putman, C. Vespa, and N. Wahl. This research has been supported by the Danish National Research Foundation through the Centre for Symmetry and Deformation (DNRF92), and parts of this paper were written while I was visiting the Hausdorff Research Institute for Mathematics, Bonn.

BIBLIOGRAPHY

- [1] S. ANDREADAKIS, “On the automorphisms of free groups and free nilpotent groups”, *Proc. Lond. Math. Soc.* **15** (1965), p. 239-268.
- [2] W. BALCERAK & B. HAJDUK, “Homotopy type of automorphism groups of manifolds”, *Colloq. Math.* **45** (1981), no. 1, p. 1-33.
- [3] M. BESTVINA, “Homological stability of $\text{Aut}(F_n)$ revisited”, in *Hyperbolic geometry and geometric group theory*, Advanced Studies in Pure Mathematics, vol. 73, Mathematical Society of Japan, 2017, p. 1-11.
- [4] S. BETLEY, “Homology of $\text{GL}(R)$ with coefficients in a functor of finite degree”, *J. Algebra* **150** (1992), no. 1, p. 73-86.
- [5] S. BETLEY & T. PIRASHVILI, “Stable K-theory as a derived functor”, *J. Pure Appl. Algebra* **96** (1994), no. 3, p. 245-258.
- [6] A. BOREL, “Stable real cohomology of arithmetic groups”, *Ann. Sci. Éc. Norm. Supér.* **7** (1974), p. 235-272.
- [7] D. BURGHELEA, “Automorphisms of manifolds”, in *Algebraic and geometric topology (Stanford, 1976), Part 1*, Proceedings of Symposia in Pure Mathematics, vol. 32, American Mathematical Society, 1978, p. 347-371.
- [8] R. M. CHARNEY, “Homology stability for GL_n of a Dedekind domain”, *Invent. Math.* **56** (1980), no. 1, p. 1-17.
- [9] A. DJAMENT, “Sur l’homologie des groupes unitaires à coefficients polynomiaux”, *J. K-Theory* **10** (2012), no. 1, p. 87-139.
- [10] Z. FIEDOROWICZ, T. PIRASHVILI, R. SCHWÄNZL, R. VOGT & F. WALDHAUSEN, “Mac Lane homology and topological Hochschild homology”, *Math. Ann.* **303** (1995), no. 1, p. 149-164.
- [11] V. FRANJOU, E. M. FRIEDLANDER, A. SCORICHENKO & A. A. SUSLIN, “General linear and functor cohomology over finite fields”, *Ann. Math.* **150** (1999), no. 2, p. 663-728.

- [12] V. FRANJOU & T. PIRASHVILI, “On the Mac Lane cohomology for the ring of integers”, *Topology* **37** (1998), no. 1, p. 109-114.
- [13] ———, “Stable K -theory is bifunctor homology (after A. Scorichenko)”, in *Rational representations, the Steenrod algebra and functor homology*, Panoramas et Synthèses, vol. 16, Société Mathématique de France, 2003, p. 107-126.
- [14] S. GALATIUS, “Stable homology of automorphism groups of free groups”, *Ann. Math.* **173** (2011), no. 2, p. 705-768.
- [15] A. E. HATCHER, “Concordance spaces, higher simple-homotopy theory, and applications”, in *Algebraic and geometric topology (Stanford, 1976), Part 1*, Proceedings of Symposia in Pure Mathematics, American Mathematical Society, 1978, p. 3-21.
- [16] ———, “Homological stability for automorphism groups of free groups”, *Comment. Math. Helv.* **70** (1995), no. 1, p. 39-62.
- [17] A. E. HATCHER & K. VOGTMANN, “Cerf theory for graphs”, *J. Lond. Math. Soc.* **58** (1998), no. 3, p. 633-655.
- [18] ———, “Rational homology of $\text{Aut}(F_n)$ ”, *Math. Res. Lett.* **5** (1998), no. 6, p. 759-780.
- [19] W.-C. HSIANG & R. W. SHARPE, “Parametrized surgery and isotopy”, *Pac. J. Math.* **67** (1976), no. 2, p. 401-459.
- [20] K. IGUSA, “What happens to Hatcher and Wagoner’s formulas for $\pi_0 C(M)$ when the first Postnikov invariant of M is nontrivial?”, in *Algebraic K-theory, number theory, geometry and analysis (Bielefeld, 1982)*, Lecture Notes in Mathematics, vol. 1046, Springer, 1984, p. 104-172.
- [21] M. JIBLADZE & T. PIRASHVILI, “Cohomology of algebraic theories”, *J. Algebra* **137** (1991), no. 2, p. 253-296.
- [22] W. VAN DER KALLEN, “Homology stability for linear groups”, *Invent. Math.* **60** (1980), no. 3, p. 269-295.
- [23] C. KASSEL, “La K -théorie stable”, *Bull. Soc. Math. Fr.* **110** (1982), no. 4, p. 381-416.
- [24] ———, “Calcul algébrique de l’homologie de certains groupes de matrices”, *J. Algebra* **80** (1983), no. 1, p. 235-260.
- [25] M. LAZARD, “Sur les groupes nilpotents et les anneaux de Lie”, *Ann. Sci. Éc. Norm. Supér.* **71** (1954), p. 101-190.
- [26] J.-L. LODAY, *Cyclic homology*, second ed., Grundlehren der Mathematischen Wissenschaften, vol. 301, Springer, 1998, xx+513 pages.
- [27] I. MADSEN & R. J. MILGRAM, *The classifying spaces for surgery and cobordism of manifolds*, Annals of Mathematics Studies, vol. 92, Princeton University Press; University of Tokyo Press, 1979, xii+279 pages.
- [28] K. NOMIZU, “On the cohomology of compact homogeneous spaces of nilpotent Lie groups”, *Ann. Math.* **59** (1954), p. 531-538.
- [29] P. F. PICKEL, “Rational cohomology of nilpotent groups and Lie algebras”, *Commun. Algebra* **6** (1978), no. 4, p. 409-419.
- [30] T. PIRASHVILI & F. WALDHAUSEN, “Mac Lane homology and topological Hochschild homology”, *J. Pure Appl. Algebra* **82** (1992), no. 1, p. 81-98.
- [31] J. W. SCHLESINGER, “The semi-simplicial free Lie ring”, *Trans. Am. Math. Soc.* **122** (1966), p. 436-442.
- [32] A. A. SUSLIN, “Stability in algebraic K -theory”, in *Algebraic K-theory, Part I (Oberwolfach, 1980)*, Lecture Notes in Mathematics, vol. 966, Springer, 1982, p. 304-333.
- [33] M. SZYMIK, “Twisted homological stability for extensions and automorphism groups of free nilpotent groups”, *J. K-Theory* **14** (2014), no. 1, p. 185-201.
- [34] F. WALDHAUSEN, “Algebraic K -theory of generalized free products. III, IV”, *Ann. Math.* **108** (1978), no. 2, p. 205-256.

- [35] ———, “Algebraic K-theory of topological spaces. I”, in *Algebraic and geometric topology (Stanford, 1976), Part 1*, Proceedings of Symposia in Pure Mathematics, American Mathematical Society, 1978, p. 35-60.
- [36] C. T. C. WALL, *Surgery on compact manifolds*, vol. 1, Academic Press Inc., 1970, London Mathematical Society Monographs, x+280 pages.
- [37] M. WEISS & B. WILLIAMS, “Automorphisms of manifolds”, in *Surveys on surgery theory, Vol. 2*, Annals of Mathematics Studies, vol. 149, Princeton University Press, 2001, p. 165-220.

Manuscrit reçu le 19 juillet 2017,
révisé le 6 avril 2018,
accepté le 26 avril 2018.

Markus SZYMIK
Department of Mathematical Sciences
NTNU Norwegian University of Science and
Technology
7491 Trondheim (Norway)
markus.szymik@ntnu.no