



ANNALES

DE

L'INSTITUT FOURIER

John Alexander CRUZ MORALES, Anton MELLIT,
Nicolas PERRIN & Maxim SMIRNOV

On quantum cohomology of Grassmannians of isotropic lines, unfoldings of A_n -singularities, and Lefschetz exceptional collections

Article à paraître, mis en ligne le 18 février 2019, 37 p.



© Association des Annales de l'institut Fourier, 2019,
Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

ON QUANTUM COHOMOLOGY OF GRASSMANNIANS OF ISOTROPIC LINES, UNFOLDINGS OF A_n -SINGULARITIES, AND LEFSCHETZ EXCEPTIONAL COLLECTIONS

by John Alexander CRUZ MORALES, Anton MELLIT,
Nicolas PERRIN & Maxim SMIRNOV

With an appendix by Alexander KUZNETSOV (*)

Dedicated to Yuri Ivanovich Manin, with admiration and gratitude

ABSTRACT. — The subject of this paper is the big quantum cohomology rings of symplectic isotropic Grassmannians $IG(2, 2n)$. We show that these rings are regular. In particular, by “generic smoothness”, we obtain a conceptual proof of generic semisimplicity of the big quantum cohomology for $IG(2, 2n)$. Further, by a general result of Hertling, the regularity of these rings implies that they have a description in terms of isolated hypersurface singularities, which we show in this case to be of type A_{n-1} . By the homological mirror symmetry conjecture, these results suggest the existence of a very special full exceptional collection in the derived category of coherent sheaves on $IG(2, 2n)$. Such a collection is constructed in the appendix by Alexander Kuznetsov.

RÉSUMÉ. — Dans cet article, nous nous intéressons au gros anneau de cohomologie quantique de $IG(2, 2n)$, la grassmannienne symplectique des droites isotropes. Nous montrons que cet anneau est régulier et en déduisons par « lissité générique » une preuve conceptuelle de la semi-simplicité générique du gros anneau de cohomologie quantique de $IG(2, 2n)$. Par ailleurs, par un résultat général de Hertling, cette régularité donne une description de cet anneau en termes de singularités isolées d’hypersurfaces et nous montrons que les singularités qui apparaissent sont de type A_{n-1} . La conjecture de symétrie miroir homologique prédit l’existence de suites exceptionnelles très spéciales dans la catégorie dérivée des faisceaux cohérents de $IG(2, 2n)$. L’existence de telles collections est démontrée en appendice par Alexander Kuznetsov.

Keywords: semisimplicity of quantum cohomology, unfoldings of singularities, Lefschetz exceptional collections.

2010 *Mathematics Subject Classification:* 14N35, 53D45.

(*) A.K. was partially supported by the Russian Academic Excellence Project “5-100”, and by the Program of the Presidium of the Russian Academy of Sciences No. 01 “Fundamental Mathematics and its Applications” under grant PRAS-18-01.

1. Introduction

1.1. Dubrovin's conjecture

Quantum cohomology rings of smooth projective varieties have been an object of intensive study ever since they were introduced at the beginning of 1990s. A particular question that attracted a lot of attention is a conjecture formulated by Boris Dubrovin in [7]. This conjecture provides a beautiful relation between two a priori seemingly unrelated objects — the quantum cohomology ring of a variety X and its bounded derived category of coherent sheaves $D^b(X)$. Dubrovin's conjecture has several parts, and its first part claims that the generic semisimplicity of the quantum cohomology of X is equivalent to the existence of a full exceptional collection in $D^b(X)$. Though there are no general approaches to this conjecture, it has been tested in many examples and the original formulation of Dubrovin has been made more precise (see [2, 7, 8, 11, 16] and references therein).

1.2. Quantum cohomology: big vs. small

The quantum cohomology ring of X is a (formal) deformation family of rings provided by the genus zero Gromov–Witten theory, that specializes to the ordinary cohomology ring $H^*(X, \mathbb{Q})$, if one sets all deformation parameters to zero. There exist two such deformation families that one often meets in the literature: the small quantum cohomology (involves only 3-point GW invariants) and the big quantum cohomology (involves GW invariants with arbitrary many insertions). The former family is a specialization of the latter (i.e. some of the deformation parameters are set to zero). It is the big quantum cohomology ring that Dubrovin used in [7] to formulate the conjecture.

In this paper we are concerned with generic semisimplicity of quantum cohomology. Therefore, we only consider the above deformation families in the neighbourhood of the generic point of the small quantum cohomology. In this way the small quantum cohomology (denote it $\mathrm{QH}(X)$) becomes a commutative, associative, finite dimensional algebra over some field K , and the big quantum cohomology (denote it $\mathrm{BQH}(X)$) is a formal deformation family of $\mathrm{QH}(X)$. This will be made more precise in Section 2.

In Dubrovin's conjecture we are interested in the semisimplicity of the generic member of the deformation family $\mathrm{BQH}(X)$. So it could happen that the special fiber $\mathrm{QH}(X)$ is not semisimple, whereas the generic fiber

of the full family $\text{BQH}(X)$ is. This is exactly what happens in the case of symplectic isotropic Grassmannians $\text{IG}(2, 2n)$. Indeed, it was shown in [4, 5] that the small quantum cohomology of $\text{IG}(2, 2n)$ is not semisimple. Further, in [12, 26] jointly with Sergey Galkin, we have proved that the big quantum cohomology of $\text{IG}(2, 2n)$ is generically semisimple. Since, according to [20, 28], the derived category of coherent sheaves on $\text{IG}(2, 2n)$ has a full exceptional collection, the first part of Dubrovin's conjecture holds.

1.3. Generic semisimplicity via generic smoothness

Our first result is the following (see Theorem 6.4).

THEOREM A. — *The ring $\text{BQH}(\text{IG}(2, 2n))$ is regular.*

As an easy consequence of the above theorem and generic smoothness (see Corollary 6.5) we recover generic semisimplicity of the big quantum cohomology of $\text{IG}(2, 2n)$ proved in [12, 26]. Note that in [12] the authors only considered the case of $\text{IG}(2, 6)$ and the proof was computer-assisted. The proof in [26] works for all $\text{IG}(2, 2n)$ but needs many lengthy computations. Our proof is more conceptual and clarifies the situation. This is the content of Sections 2–6.

Our approach via generic smoothness gives a new perspective towards a proof of generic semisimplicity of quantum cohomology for more general Grassmannians $\text{IG}(m, 2n)$ or even rational homogeneous spaces G/P . We plan to address this in a future work.

1.4. F -manifolds with smooth spectral cover

In [16, Problem 2.8] the authors formulated the following question: *characterise those varieties X for which the quantum cohomology has smooth spectral cover.* In our terminology this question translates into the problem of characterising those varieties X for which the big quantum cohomology $\text{BQH}(X)$ is a regular ring. According to the above theorem, isotropic Grassmannians $\text{IG}(2, 2n)$ provide non-trivial examples. In fact, it is natural to expect that this property holds for any rational homogeneous space G/P .

The regularity of $\text{BQH}(X)$ implies, by a beautiful result of C. Hertling, that the F -manifold defined by the quantum cohomology of X has a description in terms of unfoldings of isolated hypersurface singularities. In our situation we prove the following (see Section 7.5).

THEOREM B. — *Assume that the genus zero Gromov–Witten potential of $\mathrm{IG}(2, 2n)$ has non-trivial convergence radius. Then the F -manifold of $\mathrm{BQH}(\mathrm{IG}(2, 2n))$ decomposes into the product of the unfolding of an A_{n-1} -singularity and $(2n - 1)(n - 1)$ copies of the unfolding of an A_1 -singularity.*

1.5. Derived category of coherent sheaves

The above theorem and mirror symmetry suggest that $\mathrm{IG}(2, 2n)$ should have a Landau–Ginzburg model with one degenerate critical point of type A_{n-1} and $(2n - 1)(n - 1)$ non-degenerate critical points. Therefore, the Fukaya–Seidel category of this LG model should have a semiorthogonal decomposition

$$\langle \mathcal{C}_{n-1}, E_1, \dots, E_{(2n-1)(n-1)} \rangle,$$

where E_i are exceptional objects given by the non-degenerate critical points and \mathcal{C}_{n-1} is the Fukaya–Seidel category of an A_{n-1} -singularity. By homological mirror symmetry conjecture, the bounded derived category of coherent sheaves $D^b(\mathrm{IG}(2, 2n))$ should be equivalent to the Fukaya–Seidel category of the LG model. Hence, the derived category $D^b(\mathrm{IG}(2, 2n))$ should also have a decomposition of this form. A more detailed account is contained in Section 8.

The above discussion is highly conjectural. Nonetheless, the aforementioned semiorthogonal decomposition of $D^b(\mathrm{IG}(2, 2n))$ is constructed directly in the Appendix by Alexander Kuznetsov.

THEOREM C (Kuznetsov). — *There exists a semiorthogonal decomposition*

$$D^b(\mathrm{IG}(2, 2n)) = \langle \mathcal{A}_{n-1}, E_1, \dots, E_{(2n-1)(n-1)} \rangle,$$

where the E_i are some exceptional objects and the subcategory \mathcal{A}_{n-1} is equivalent to the bounded derived category of representations of the quiver of type A_{n-1} .

This theorem confirms the conjectural picture described above, as the bounded derived category of representations of the quiver of type A_{n-1} is equivalent to the Fukaya–Seidel category of an A_{n-1} -singularity by [29]. Note that the result in the appendix is stronger. Namely, the objects $E_1, \dots, E_{(2n-1)(n-1)}$ form a rectangular Lefschetz exceptional collection.

One can view these results as an enhanced version of Dubrovin’s conjecture, i.e. a prediction of a more subtle relation between quantum cohomology rings and derived categories of coherent sheaves.

Acknowledgements. We are indebted to Yuri Ivanovich Manin for his support over the years and for drawing our attention to results of Claus Hertling indispensable for this paper. Further we are very grateful to Alexander Kuznetsov for providing the appendix, and numerous remarks on the main body of the paper. Special thanks go to Mohammed Abouzaid, Sheel Ganatra, and Ailsa Keating for the helpful email correspondence about Fukaya–Seidel categories. Last but not least we would like to thank our friends and colleagues Tarig Abdelgadir, Erik Carlsson, Roman Fedorov, Sergey Galkin, Christian Lehn, Sina Türelı and Runako Williams for valuable discussions and comments.

We would like to thank institutions that supported us at various stages of this project. Namely, we are very grateful to the International Centre for Theoretical Physics (ICTP) in Trieste, the Institute for Algebraic Geometry and the Riemann Center for Geometry and Physics at the Leibniz Universität Hannover, and the Max Planck Institute for Mathematics (MPIM) in Bonn. The third author was supported by a public grant as part of the Investissement d’avenir project, reference ANR-11-LABX-0056-LMH, LabEx LMH.

2. Conventions and notation for quantum cohomology

Here we briefly recall the definition of the quantum cohomology ring for a smooth projective variety X . To simplify the exposition and avoid introducing unnecessary notation we impose from the beginning the following conditions on X : it is a Fano variety of Picard rank 1 and $H^{\text{odd}}(X, \mathbb{Q}) = 0$. For a thorough introduction we refer to [24].

2.1. Definition

Let us fix a graded basis $\Delta_0, \dots, \Delta_s$ in $H^*(X, \mathbb{Q})$ and dual linear coordinates t_0, \dots, t_s . It is customary to choose $\Delta_0 = 1$. For cohomology classes we use the Chow grading, i.e. we divide the topological degree by two. Further, for variables t_i we set $\deg(t_i) = 1 - \deg(\Delta_i)$.

Let R be the ring of formal power series $\mathbb{Q}[[q]]$, k its field of fractions, and K an algebraic closure of k . We set $\deg(q) = \text{index}(X)$, which is the largest integer n such that $-K_X = nH$ for some ample divisor H on X , where K_X is the canonical class of X .

The genus zero Gromov–Witten potential of X is an element $\Phi \in R[[t_0, \dots, t_s]]$ defined by the formula

$$(2.1) \quad \Phi = \sum_{(i_0, \dots, i_s)} \langle \Delta_0^{\otimes i_0}, \dots, \Delta_s^{\otimes i_s} \rangle \frac{t_0^{i_0} \dots t_s^{i_s}}{i_0! \dots i_s!},$$

where

$$\langle \Delta_0^{\otimes i_0}, \dots, \Delta_s^{\otimes i_s} \rangle = \sum_{d=0}^{\infty} \langle \Delta_0^{\otimes i_0}, \dots, \Delta_s^{\otimes i_s} \rangle_d q^d,$$

and $\langle \Delta_0^{\otimes i_0}, \dots, \Delta_s^{\otimes i_s} \rangle_d$ are rational numbers called Gromov–Witten invariants of X of degree d . With respect to the grading defined above Φ is homogeneous of degree $3 - \dim X$.

Using (2.1) one defines the *big quantum cohomology ring* of X . Namely, let us endow the $K[[t_0, \dots, t_s]]$ -module

$$\text{BQH}(X) = H^*(X, \mathbb{Q}) \otimes_{\mathbb{Q}} K[[t_0, \dots, t_s]]$$

with a ring structure by setting

$$(2.2) \quad \Delta_a \star \Delta_b = \sum_c \frac{\partial^3 \Phi}{\partial t_a \partial t_b \partial t_c} \Delta^c,$$

on the basis elements and extending to the whole $\text{BQH}(X)$ by $K[[t_0, \dots, t_s]]$ -linearity. Here $\Delta^0, \dots, \Delta^s$ is the basis dual to $\Delta_0, \dots, \Delta_s$ with respect to the Poincaré pairing. It is well known that (2.2) makes $\text{BQH}(X)$ into a commutative, associative, graded $K[[t_0, \dots, t_s]]$ -algebra with the identity element Δ_0 .

The algebra $\text{BQH}(X)$ is called the *big quantum cohomology algebra* of X to distinguish it from a simpler object called the *small quantum cohomology algebra* which is the quotient of $\text{BQH}(X)$ with respect to the ideal (t_0, \dots, t_s) . We will denote the latter $\text{QH}(X)$ and use \star_0 instead of \star for the product in this algebra. It is a finite dimensional K -algebra. Equivalently one can say that

$$\text{QH}(X) = H^*(X, \mathbb{Q}) \otimes_{\mathbb{Q}} K$$

as a vector space, and the K -algebra structure is defined by putting

$$\Delta_a \star_0 \Delta_b = \sum_c \langle \Delta_a, \Delta_b, \Delta_c \rangle \Delta^c.$$

Remark 2.1. — We are using a somewhat non-standard notation $\text{BQH}(X)$ for the big quantum cohomology and $\text{QH}(X)$ for the small quantum cohomology to stress the difference between the two. Note that this notation is different from the one used in [12] and is closer to the notation of [26].

Remark 2.2. — The above definitions look slightly different from the ones given in [24]. The differences are of two types. The first one is that $\text{QH}(X)$ and $\text{BQH}(X)$ are in fact defined already over the ring R and not only over K . We pass to K from the beginning, since in this paper we are only interested in generic semisimplicity of quantum cohomology. The second difference is that in some papers on quantum cohomology one unifies the coordinate q with the coordinate t_i which is dual to $H^2(X, \mathbb{Q})$, but the resulting structures are equivalent.

2.2. Deformation picture

The small quantum cohomology, if considered over the ring R (cf. Remark 2.2), is a deformation of the ordinary cohomology algebra, i.e. if we put $q = 0$, then the quantum product becomes the ordinary cup-product. Similarly, the big quantum cohomology is an even bigger deformation family of algebras. Since we work not over R but over K , we lose the point of classical limit but still retain the fact that $\text{BQH}(X)$ is a deformation family of algebras with the special fiber being $\text{QH}(X)$.

In this paper we view $\text{Spec}(\text{BQH}(X))$ as a deformation family of zero-dimensional schemes over $\text{Spec}(K[[t_0, \dots, t_s]])$. In the base of the deformation we consider the following two points: the origin (the closed point given by the maximal ideal (t_0, \dots, t_s)) and the generic point η . The fiber of this family over the origin is the spectrum of the small quantum cohomology $\text{Spec}(\text{QH}(X))$. The fiber over the generic point will be denoted by $\text{Spec}(\text{BQH}(X)_\eta)$. It is convenient to summarize this setup in the diagram

$$(2.3) \quad \begin{array}{ccccc} \text{Spec}(\text{QH}(X)) & \longrightarrow & \text{Spec}(\text{BQH}(X)) & \longleftarrow & \text{Spec}(\text{BQH}(X)_\eta) \\ \downarrow & & \downarrow \pi & & \downarrow \pi_\eta \\ \text{Spec}(K) & \longrightarrow & \text{Spec}(K[[t_0, \dots, t_s]]) & \longleftarrow & \eta \end{array}$$

where both squares are Cartesian.

By construction $\text{BQH}(X)$ is a free module of finite rank over $K[[t_0, \dots, t_s]]$. Therefore, it is a noetherian semilocal K -algebra which is flat and finite over $K[[t_0, \dots, t_s]]$. Note that neither $K[[t_0, \dots, t_s]]$ nor $\text{BQH}(X)$ are finitely generated over the ground field K . Therefore, some extra care is required in the standard commutative algebra (or algebraic geometry) constructions. For example, the notion of smoothness is one of such concepts.

2.3. Semisimplicity

Let A be a finite dimensional algebra over a field F of characteristic zero. It is called *semisimple* if it is a product of fields. Equivalently, the algebra A is semisimple if the scheme $\text{Spec}(A)$ is reduced. Another equivalent condition is to require the morphism $\text{Spec}(A) \rightarrow \text{Spec}(F)$ to be smooth.

DEFINITION 2.3. — We say that $\text{BQH}(X)$ is generically semisimple if $\text{BQH}(X)_\eta$ is a semisimple algebra.

3. Geometry of $\text{IG}(2, 2n)$

Let V be a complex vector space endowed with a symplectic form ω . In this case the dimension of V has to be even and we denote it by $2n$. For any $1 \leq m \leq n$ there exists an algebraic variety $\text{IG}(m, V)$ that parametrizes m -dimensional isotropic subspaces of V . For $m = 2$, and this is the case we are considering in this paper, it has the following explicit description. Consider the ordinary Grassmannian $\text{G}(2, V)$ with its Plücker embedding into $\mathbb{P}(\Lambda^2 V)$. The symplectic form ω defines a hyperplane $H_\omega = \mathbb{P}(\text{Ker } \omega) \subset \mathbb{P}(\Lambda^2 V)$ and the intersection of $\text{G}(2, V)$ with H_ω is $\text{IG}(2, V)$. Thus, we have inclusions

$$(3.1) \quad \text{IG}(2, V) \subset \text{G}(2, V) \subset \mathbb{P}(\Lambda^2 V).$$

As for different symplectic forms on V we obtain isomorphic isotropic Grassmannians, it is unambiguous to write $\text{IG}(2, 2n)$. We will use this notation starting from Paragraph 3.2.

3.1. Two sets of cohomology classes

As for ordinary Grassmannians, one considers the short exact sequence of vector bundles on $X = \text{IG}(2, V)$

$$(3.2) \quad 0 \rightarrow \mathcal{U} \rightarrow \mathcal{V} \rightarrow \mathcal{V}/\mathcal{U} \rightarrow 0,$$

where \mathcal{V} is the trivial vector bundle with fiber V , \mathcal{U} is the subbundle of isotropic subspaces, and \mathcal{V}/\mathcal{U} is the quotient bundle. Usually one refers to \mathcal{U} and \mathcal{V}/\mathcal{U} as *tautological subbundle* and *tautological quotient bundle* respectively.

One also defines a vector bundle \mathcal{U}^\perp as the kernel of the composition $\mathcal{V} \xrightarrow{\omega} \mathcal{V}^* \rightarrow \mathcal{U}^*$, where the first morphism is the isomorphism induced by

the symplectic form, and the second one is the dual of the natural inclusion $\mathcal{U} \rightarrow \mathcal{V}$. From the definition of \mathcal{U}^\perp we immediately obtain an isomorphism

$$(3.3) \quad \mathcal{V}/\mathcal{U}^\perp \simeq \mathcal{U}^*.$$

Further, we have inclusions of vector bundles $\mathcal{U} \subset \mathcal{U}^\perp \subset \mathcal{V}$, and can also consider the short exact sequence

$$(3.4) \quad 0 \rightarrow \mathcal{U}^\perp/\mathcal{U} \rightarrow \mathcal{V}/\mathcal{U} \rightarrow \mathcal{V}/\mathcal{U}^\perp \rightarrow 0.$$

By taking Chern classes of vector bundles in (3.2) and (3.4) we obtain two sets of cohomology classes which generate the cohomology ring:

(a) Chern classes of \mathcal{V}/\mathcal{U} . The vector bundle \mathcal{V}/\mathcal{U} is of rank $2n - 2$ and we denote its Chern classes by $\sigma_k = c_k(\mathcal{V}/\mathcal{U})$. Cycles representing these cohomology classes can be explicitly described as follows. Let

$$(3.5) \quad Z(E_{2n-k-1}) = \{V_2 \in X \mid \dim(V_2 \cap E_{2n-k-1}) \geq 1\}.$$

Then $\sigma_k = [Z(E_{2n-k-1})] \in H^{2k}(X, \mathbb{Z})$. In the above, E_{2n-k-1} is a subspace of dimension $2n - k - 1$ such that the rank of $\omega|_{E_{2n-k-1}}$ is minimal. Note that $\sigma_0 = 1$ is the fundamental class of X . These cohomology classes are usually called *special Schubert classes*.

(b) Chern classes of \mathcal{U} and $\mathcal{U}^\perp/\mathcal{U}$. The vector bundle \mathcal{U} is of rank 2, so it only has two non-vanishing Chern classes $a_i = c_i(\mathcal{U})$ for $i = 1, 2$. The vector bundle $\mathcal{U}^\perp/\mathcal{U}$ is self-dual of rank $2n - 4$, therefore it has only $n - 2$ non-vanishing Chern classes $b_i = c_{2i}(\mathcal{U}^\perp/\mathcal{U})$ for $i \in [1, n - 2]$.

3.2. Cohomology ring of $\text{IG}(2, 2n)$

The cohomology ring of $X = \text{IG}(2, 2n)$ can be described in terms of generators and relations. We will give two presentations using the two sets of special cohomology classes defined above.

PROPOSITION 3.1 ([3, Theorem 1.2]). — *The cohomology ring $H^*(X, \mathbb{Q})$ is isomorphic to the quotient of the ring $\mathbb{Q}[\sigma_1, \dots, \sigma_{2n-2}]$ by the ideal generated by the elements*

$$(3.6) \quad \det(\sigma_{1+j-i})_{1 \leq i, j \leq r}, \quad \text{with } r \in [3, 2n - 2]$$

and the two elements

$$(3.7) \quad \sigma_{n-1}^2 + 2 \sum_{i=1}^{n-1} (-1)^i \sigma_{n-1+i} \sigma_{n-1-i} \quad \text{and} \quad \sigma_n^2 + 2 \sum_{i=1}^{n-2} (-1)^i \sigma_{n+i} \sigma_{n-i}.$$

The dimension of $H^*(X, \mathbb{Q})$ is equal to $2^2 \binom{n}{2} = 2n(n - 1)$.

PROPOSITION 3.2. — *The cohomology ring $H^*(X, \mathbb{Q})$ is isomorphic to the quotient of the ring $\mathbb{Q}[a_1, a_2, b_1, \dots, b_{n-2}]$ by the ideal generated by*

$$(3.8) \quad (1 + (2a_2 - a_1^2)x^2 + a_2^2x^4)(1 + b_1x^2 + \dots + b_{n-2}x^{2n-4}) = 1$$

The last equality is viewed as an equality of polynomials in the variable x and gives a concise way to write a system of equations in the variables a_i, b_i .

Proof. — This result is well known to specialists but we include a short proof for the convenience of the reader.

Let us start by checking that (3.8) holds in the cohomology ring. Define $P(x) = 1 + a_1x + a_2x^2$ and $Q(x) = 1 + b_1x^2 + \dots + b_{n-2}x^{2n-4}$ and rewrite (3.8) as $P(x)P(-x)Q(x) = 1$. We interpret the polynomial $P(x)$ as the total Chern class of \mathcal{U} and $Q(x)$ as the total Chern class of $\mathcal{U}^\perp/\mathcal{U}$. Now by using basic properties of Chern classes, short exact sequences (3.2), (3.4), and the isomorphism (3.3), it is easy to see that the above relation does hold.

The above discussion shows that we have a natural homomorphism of \mathbb{Q} -algebras

$$(3.9) \quad \psi: \mathbb{Q}[a_1, a_2, b_1, \dots, b_{n-2}]/(P(x)P(-x)Q(x) - 1) \rightarrow H^*(X, \mathbb{Q})$$

sending a_i 's to a_i 's and b_i 's to b_i 's. To prove the proposition it is enough to establish two facts: i) σ_i 's can be expressed in terms of a_i 's and b_i 's, ii) the dimensions of both algebras in (3.9) are equal. To prove the first fact one can use again simple properties of Chern classes and the exact sequence (3.4). For the second fact, we first need to show that $\mathbb{Q}[a_1, a_2, b_1, \dots, b_{n-2}]/(P(x)P(-x)Q(x) - 1)$ is finite dimensional. This can be done similarly to the finite-dimensionality part of the proof of [3, Theorem 1.2]. Then, we need to compute the dimension of this algebra. For this we can proceed similarly to the proof of [3, Lemma 1.1], which is based on [30], and get the desired dimension $2n(n-1)$. \square

4. Small quantum cohomology of $\text{IG}(2, 2n)$

4.1. Two presentations

As described in Section 3.2, the ordinary cohomology ring of the Grassmannian $\text{IG}(2, 2n)$ is generated by the special Schubert classes $\sigma_1, \dots, \sigma_{2n-2}$ with relations (3.6) and (3.7). To pass to the small quantum cohomology, informally speaking, we need to adjoin a new variable q to the σ_i 's and

adjust relations (3.6)–(3.7) in such a way that they remain homogeneous and specialize to the original ones when setting $q = 0$. The degree of the variable q in the case of $\text{IG}(2, 2n)$ is equal to $2n - 1$ (cf. Section 2.1). Thus, the only relation that needs to be modified is the second equation in (3.7). Moreover, up to a constant factor, this modification is unique for degree reasons. The complete answer for arbitrary Grassmannians $\text{IG}(m, 2n)$ was given in [3, Theorem 1.5] which we reproduce here in the special case of $m = 2$.

THEOREM 4.1 ([3, Theorem 1.5]). — *The small quantum cohomology ring $\text{QH}(X)$ is isomorphic to the quotient of the ring $K[\sigma_1, \dots, \sigma_{2n-2}]$ by the ideal generated by the elements*

$$\det(\sigma_{1+j-i})_{1 \leq i, j \leq r}, \quad \text{with } r \in [3, 2n - 2]$$

and the two elements

$$\sigma_{n-1}^2 + 2 \sum_{i=1}^{n-1} (-1)^i \sigma_{n-1+i} \sigma_{n-1-i}$$

and

$$\sigma_n^2 + 2 \sum_{i=1}^{n-2} (-1)^i \sigma_{n+i} \sigma_{n-i} + (-1)^{n+1} q \sigma_1.$$

Combining the above theorem with Proposition 3.2 we arrive at the following statement.

COROLLARY 4.2. — *The small quantum cohomology ring $\text{QH}(X)$ is isomorphic to the quotient of the ring $K[a_1, a_2, b_1, \dots, b_{n-2}]$ by the ideal generated by*

$$(4.1) \quad (1 + (2a_2 - a_1^2)x^2 + a_2^2x^4)(1 + b_1x^2 + \dots + b_{n-2}x^{2n-4}) = 1 - qa_1x^{2n}.$$

Proof. — We need to find the quantum deformation of (3.8). Since the index of X is $2n - 1$, there are no quantum corrections except possibly in degrees $2n - 1$ and $2n$. Furthermore, as (3.8) has no terms of degree $2n - 1$, we only need to check the deformation in degree $2n$. Explicitly we need to check the relation $a_2^2 b_{n-2} = -qa_1$. Note that since $a_2^2 b_{n-2}$ has degree $2n$ and since $a_2^2 b_{n-2}$ vanishes in $H^*(X, \mathbb{Q})$ (see Proposition 3.2), we have $a_2^2 b_{n-2} = \lambda a_1 q$ with $\lambda = -\langle a_2, a_2 b_{n-2}, \ell \rangle_1$ and ℓ the class of a line in X .

By the definition of a_i 's we have $a_2 = c_2(\mathcal{U})$, and so a_2 is a Schubert class (in fact, we have $a_2 = \sigma_{1,1}$). Further, using basic properties of Chern

classes, and Formulas (3.2)–(3.4) one obtains the equality of cohomology classes

$$a_2 b_{n-2} = \sigma_{2n-2}.$$

Therefore, $a_2 b_{n-2}$ is also a Schubert class. As ℓ is yet again a Schubert class, all insertions in the GW invariant $\langle a_2, a_2 b_{n-2}, \ell \rangle_1$ are Schubert classes in X . One can compute such an invariant either by using methods of Section 5, or the quantum-to-classical principle for degree 1 invariants from [5, Section 4], and obtains $\langle a_2, a_2 b_{n-2}, \ell \rangle_1 = 1$. \square

4.2. Structure of $\mathrm{QH}(X)$

In this paragraph we decompose the K -algebra $\mathrm{QH}(X)$ as a direct product, or, equivalently, we decompose the K -scheme $\mathrm{Spec}(\mathrm{QH}(X))$ into connected components.

PROPOSITION 4.3. — *The scheme $\mathrm{Spec}(\mathrm{QH}(X))$ is the disjoint union of $(2n-1)(n-1)$ reduced points $\mathrm{Spec}(K)$ and one fat point $\mathrm{Spec}(K[\varepsilon]/(\varepsilon^{n-1}))$.*

Proof. — By the presentation of $\mathrm{QH}(X)$ described in Corollary 4.2, the K -scheme $\mathrm{Spec}(\mathrm{QH}(X))$ is given as a closed subscheme of the affine space $\mathbb{A}^n = \mathrm{Spec}(K[a_1, a_2, b_1, \dots, b_{n-2}])$ defined by the equations

$$\begin{aligned}
 & 2a_2 - a_1^2 + b_1 = 0 \\
 & a_2^2 + b_1(2a_2 - a_1^2) + b_2 = 0 \\
 & b_1 a_2^2 + b_2(2a_2 - a_1^2) + b_3 = 0 \\
 & \dots \\
 (4.2) \quad & b_{i-1} a_2^2 + b_i(2a_2 - a_1^2) + b_{i+1} = 0 \\
 & \dots \\
 & b_{n-4} a_2^2 + b_{n-3}(2a_2 - a_1^2) + b_{n-2} = 0 \\
 & b_{n-3} a_2^2 + b_{n-2}(2a_2 - a_1^2) = 0 \\
 & b_{n-2} a_2^2 + q a_1 = 0.
 \end{aligned}$$

It is clear that the origin of \mathbb{A}^n is a solution of this system. Moreover, this solution corresponds to a fat point of $\mathrm{Spec}(\mathrm{QH}(X))$. Indeed, it is easy to see that the Zariski tangent space of (4.2) at the origin is one-dimensional. Thus, the origin is a fat point of $\mathrm{Spec}(\mathrm{QH}(X))$. Let A be the corresponding factor of $\mathrm{QH}(X)$. Thus, we have the direct product decomposition

$$(4.3) \quad \mathrm{QH}(X) = A \times B,$$

where B corresponds to components of $\text{Spec}(\text{QH}(X))$ supported outside the origin.

Setting $a_1 = 0$ in (4.2) we obtain $b_i = (-1)^i(i+1)a_2^i$. Therefore, eliminating the b_i 's, we get an isomorphism of K -algebras

$$\text{QH}(X)/(a_1) \simeq A/(a_1) \simeq K[a_2]/(a_2^{n-1}).$$

In particular, we have a surjective homomorphism of K -algebras

$$(4.4) \quad A \rightarrow K[a_2]/(a_2^{n-1}).$$

Thus, the dimension of A is at least $n-1$. In fact, below we will see that $\dim_K A = n-1$.

Now let us examine the structure of B , i.e. we need to study solutions of (4.2) different from the origin. It is convenient to rewrite (4.1) as

$$(z^4 + (2a_2 - a_1^2)z^2 + a_2^2)(z^{2n-4} + b_1z^{2n-6} + \cdots + b_{n-2}) = z^{2n} - a_1,$$

where we set $q = 1$ for convenience. Making the substitution $a_1 = z_1 + z_2$, $a_2 = z_1z_2$, and putting $Q(z) = z^{2n-4} + b_1z^{2n-6} + \cdots + b_{n-2}$ we arrive at

$$(4.5) \quad (z^2 - z_1^2)(z^2 - z_2^2)Q(z) = z^{2n} - (z_1 + z_2).$$

In geometric terms this manipulation corresponds to pulling back our system (4.2) with respect to the morphism

$$\text{Spec}(K[z_1, z_2, b_1, \dots, b_{n-2}]) \rightarrow \text{Spec}(K[a_1, a_2, b_1, \dots, b_{n-2}])$$

defined by

$$a_1 \mapsto z_1 + z_2, \quad a_2 \mapsto z_1z_2, \quad b_i \mapsto b_i.$$

It is a double cover unramified outside of the locus $z_1 = z_2$.

Let us count solutions of (4.5) for which $z_1 \neq z_2$ and both of them are non-zero. This reduces to counting pairs z_1, z_2 satisfying

$$(4.6) \quad \begin{aligned} z_1^{2n} &= z_1 + z_2 \\ z_2^{2n} &= z_1 + z_2. \end{aligned}$$

Eliminating z_2 using the first equation we obtain that z_1 must be a solution of

$$(z_1^{2n} - z_1)^{2n} = z_1^{2n}.$$

Now it is straightforward to count that there are $(2n-2)(2n-1)$ distinct pairs (z_1, z_2) , with $z_1 \neq z_2 \neq 0$, satisfying the system (4.6).

In terms of the original system (4.2) the above computation means that there are at least $(n-1)(2n-1)$ distinct solutions outside of the origin. Note that we have divided the number of solutions by 2, since the initial

count was on the double cover. In other words the dimension of B is at least $(n-1)(2n-1)$.

Up to now we have shown that

$$\dim_K(A) \geq n-1 \quad \text{and} \quad \dim_K(B) \geq (n-1)(2n-1).$$

Since $\dim_K(\mathrm{QH}(X)) = 2n(n-1)$, this implies that $\dim_K(A) = n-1$ and $\dim_K(B) = (n-1)(2n-1)$. Hence, (4.4) is an isomorphism. \square

Remark 4.4. — The above proposition implies that $\mathrm{QH}(\mathrm{IG}(2, 2n))$ is not semisimple, as was already mentioned in the introduction. Moreover, we have an explicit description of the non-semisimple factor as $K[\varepsilon]/(\varepsilon^{n-1})$. Note that the latter is the Milnor algebra of the A_{n-1} -singularity.

5. Four-point Gromov–Witten invariants

In Section 6 we will study the deformation of $\mathrm{QH}(\mathrm{IG}(2, 2n))$ in the big quantum cohomology $\mathrm{BQH}(\mathrm{IG}(2, 2n))$ in the direction of the special Schubert class σ_2 . In particular, we will need the values of Gromov–Witten invariants

$$\langle \mathrm{pt}, \sigma_2, \sigma_i, \sigma_j \rangle_1,$$

which we compute in this section. Note that for dimension reasons these invariants vanish unless $i+j = 2n-2$.

Morally a GW invariant $\langle \Delta_1, \dots, \Delta_n \rangle_d$ counts the number of rational curves of degree d on X incident to general representatives of the cohomology classes Δ_i . Even though for a general variety X this enumerative meaning fails, in our case $X = \mathrm{IG}(2, 2n)$ we have the following fact.

LEMMA 5.1. — *The Gromov–Witten invariant $\langle \mathrm{pt}, \sigma_2, \sigma_i, \sigma_j \rangle_1$ is the number of lines meeting general representatives of the cohomology classes pt , σ_2 , σ_i and σ_j . Furthermore, given any open dense subset of the set of lines, the above lines can be chosen in this open subset.*

Proof. — It follows from Kleiman–Bertini’s Theorem [18] (see also [10, Lemma 14]). \square

5.1. Lines on $\mathrm{IG}(2, 2n)$

We recall the description of the Hilbert scheme of lines on $\mathrm{IG}(2, 2n)$ from [22, Theorem 4.3] and [31, Proposition 3]. First start with the Hilbert

scheme of lines on the ordinary Grassmannian $G(2, 2n)$. The Hilbert scheme of lines on a variety X and the universal line are denoted $F_1(X)$ and $Z_1(X)$ respectively.

A line on $G(2, 2n)$ is given by a pair (W_1, W_3) of nested subspaces of \mathbb{C}^{2n} of dimension 1 and 3 respectively. Geometrically the corresponding line is given by

$$(5.1) \quad \ell(W_1, W_3) = \{V_2 \in G(2, 2n) \mid W_1 \subset V_2 \subset W_3\}.$$

Extending the above description to families one can show that there exist isomorphisms

$$\begin{aligned} F_1(G(2, 2n)) &\simeq \text{Fl}(1, 3; 2n) \\ Z_1(G(2, 2n)) &\simeq \text{Fl}(1, 2, 3; 2n). \end{aligned}$$

The natural projections from $Z_1(G(2, 2n)) \subset G(2, 2n) \times F_1(G(2, 2n))$ to $G(2, 2n)$ and $F_1(G(2, 2n))$ are given by forgetting a part of the flag

$$\begin{array}{ccc} \text{Fl}(1, 2, 3; 2n) & \xrightarrow{f} & G(2, 2n) \\ \downarrow p & & \\ \text{Fl}(1, 3; 2n) & & \end{array}$$

Now if the line $\ell(W_1, W_3)$ is contained in $\text{IG}(2, 2n)$, we have that any two-dimensional subspace V_2 with $W_1 \subset V_2 \subset W_3$ is isotropic. This is easily seen to be equivalent to the fact that $W_1 \subset W_3^\perp$ or equivalently $W_1 \subset \ker(\omega|_{W_3})$ (note that in [31, Proposition 3], this last condition is missing). In particular we have

$$F_1(\text{IG}(2, 2n)) = \{\ell(W_1, W_3) \in F_1(G(2, 2n)) \mid \omega(W_1, W_3) = 0\}.$$

Schematically, this can be described as the zero locus of $\omega : \mathcal{W}_1 \otimes \mathcal{W}_3 \rightarrow \mathcal{O}_{F_1(G(2, 2n))}$ in $F_1(G(2, 2n))$, where \mathcal{W}_1 and \mathcal{W}_3 are the tautological subbundles. In particular a general element $\ell(W_1, W_3) \in F_1(\text{IG}(2, 2n))$ is determined by W_3 since $W_1 = \ker(\omega|_{W_3})$. We make this more precise. Let $G(3, 2n)$ be the Grassmannian of 3-dimensional vector subspaces in \mathbb{C}^{2n} and $\text{IG}(3, 2n)$ the closed subset representing subspaces isotropic for the form ω .

PROPOSITION 5.2. — *The Hilbert scheme $F_1(\text{IG}(2, 2n))$ is the blow-up of $\text{IG}(3, 2n)$ in $G(3, 2n)$.*

Proof. — Since $F_1(\text{IG}(2, 2n))$ is a subscheme of $F_1(G(2, 2n)) = \text{Fl}(1, 3; 2n)$, we have a morphism $\pi : F_1(\text{IG}(2, 2n)) \rightarrow G(3, 2n)$ defined by $(W_1, W_3) \mapsto W_3$ by forgetting W_1 . We prove that this is the desired

blow-up. Denote by \mathcal{W}_1 and \mathcal{W}_3 the tautological rank 1 and 3 bundles on $\mathrm{Fl}(1, 3; 2n)$ and $\mathrm{G}(3, 2n)$. Recall that $\mathrm{IG}(3, 2n)$ is defined by the vanishing of $\omega : \Lambda^2 \mathcal{W}_3 \rightarrow \mathcal{O}_{\mathrm{G}(3, 2n)}$. Pulling back via π and using the fact that $\mathcal{W}_1 \subset \ker(\omega|_{\mathcal{W}_3})$, we get that $\pi^* \omega$ factors as follows:

$$\begin{array}{ccc} \Lambda^2 \mathcal{W}_3 & \longrightarrow & \Lambda^2(\mathcal{W}_3/\mathcal{W}_1) \\ & \searrow \pi^* \omega & \downarrow s \\ & & \mathcal{O}_{F_1(\mathrm{IG}(2, 2n))}. \end{array}$$

Since the top map is a morphism of vector bundles and never vanishes, the vanishing of $\pi^* \omega$ is equivalent to the vanishing of s . This just means that the transform of the ideal defining $\mathrm{IG}(2, 2n)$ in $\mathrm{G}(2, 2n)$ to $F_1(\mathrm{IG}(2, 2n))$ is $\Lambda^2(\mathcal{W}_3/\mathcal{W}_1)$ and therefore invertible. By the universal property of blow-up, we deduce that π factors through the blow-up $Bl_{\mathrm{IG}(3, 2n)}(\mathrm{G}(3, 2n))$ of $\mathrm{IG}(3, 2n)$ in $\mathrm{G}(3, 2n)$:

$$\begin{array}{ccc} F_1(\mathrm{IG}(2, 2n)) & \xrightarrow{f} & Bl_{\mathrm{IG}(3, 2n)}(\mathrm{G}(3, 2n)) \\ & \searrow \pi & \downarrow p \\ & & \mathrm{G}(3, 2n). \end{array}$$

Since $\mathrm{IG}(3, 2n)$ and $\mathrm{G}(3, 2n)$ are smooth, we only need to check that f is bijective since Zariski's Main Theorem will imply that f is an isomorphism. Note that all the diagram is Sp_{2n} -equivariant and is bijective on the complement of $\mathrm{IG}(3, 2n)$: the subspace W_1 is determined by $W_1 = \ker(\omega|_{W_3})$. Now the fiber $\pi^{-1}(W_3)$ for $W_3 \in \mathrm{IG}(3, 2n)$ is isomorphic to $\mathbb{P}(W_3) \simeq \mathbb{P}^2$ and is homogeneous under the stabiliser of W_3 in Sp_{2n} . Since the fiber of p over W_3 is also isomorphic to \mathbb{P}^2 (the codimension of $\mathrm{IG}(3, 2n)$ in $\mathrm{G}(3, 2n)$ is 3), the map f must be bijective on the fiber. The result follows. \square

The group of symplectic automorphisms Sp_{2n} acts on $\mathrm{IG}(2, 2n)$ and $F_1(\mathrm{IG}(2, 2n))$. In terms of (5.1) there are two types of lines on $\mathrm{IG}(2, 2n)$ corresponding to the two Sp_{2n} -orbits on $F_1(\mathrm{IG}(2, 2n))$:

- (i) If W_3 is not isotropic, then W_1 is the kernel of the symplectic form ω on W_3 . Therefore, W_1 is completely determined by W_3 . A line of this type will be denoted by $\ell(W_3)$. These lines form the open orbit of the Sp_{2n} -action.
- (ii) If W_3 is isotropic, then W_1 is any one-dimensional subspace of W_3 . These lines form the closed orbit of the Sp_{2n} -action.

Note that in terms of the blow-up description the open orbit is $\pi^{-1}(\mathbb{G}(3, 2n) \setminus \text{IG}(3, 2n))$ and the closed orbit is the exceptional divisor of the blow-up which is isomorphic to the isotropic flag variety $\text{IFl}(1, 3; 2n)$.

5.2. Computation of the invariant

Let X be $\text{IG}(2, 2n)$. Denote by Y the Hilbert scheme of lines on X described above, and by \mathring{Y} the open orbit of the Sp_{2n} -action. In order to compute $\langle \text{pt}, \sigma_2, \sigma_i, \sigma_j \rangle_1$, according to Lemma 5.1, it is enough to consider lines in the open orbit \mathring{Y} .

For a closed subvariety $Z \subset X$ we define

$$\mathring{Y}(Z) = \{\ell \in \mathring{Y} \mid \ell \cap Z \neq \emptyset\},$$

which is a closed subvariety of \mathring{Y} .

LEMMA 5.3.

(1) *There exist identifications*

$$\begin{aligned} \mathring{Y}(\{E_2\}) &\stackrel{\text{def}}{=} \{\ell(W_3) \in \mathring{Y} \mid E_2 \subset W_3\} \simeq \mathbb{P}(\mathbb{C}^{2n}/E_2) \setminus \mathbb{P}(E_2^\perp/E_2) \\ \mathring{Y}(Z(E_{2n-k-1})) &= \{\ell(W_3) \in \mathring{Y} \mid \dim(W_3 \cap E_{2n-k-1}) \geq 1\}, \end{aligned}$$

where $\{E_2\}$ is the one-point subvariety of X corresponding to an isotropic subspace $E_2 \subset \mathbb{C}^{2n}$, and $Z(E_{2n-k-1})$ was defined in (3.5).

(2) *If E_2 and E_{2n-k-1} are in general position, the intersection $\mathring{Y}(\{E_2\}) \cap \mathring{Y}(Z(E_{2n-k-1}))$ is isomorphic to*

$$\mathbb{P}((E_2 + E_{2n-k-1})/E_2) \setminus \mathbb{P}(((E_2 + E_{2n-k-1}) \cap E_2^\perp)/E_2),$$

which we view as a subvariety of $\mathbb{P}(\mathbb{C}^{2n}/E_2) \setminus \mathbb{P}(E_2^\perp/E_2)$.

Proof.

(1) The first isomorphism just maps W_3 to W_3/E_2 and uses the fact that W_3 is non-isotropic iff W_3/E_2 is not contained in E_2^\perp/E_2 .

The second equality works as follows. Consider a point $V_2 \in (\ell(W_3) \cap Z(E_{2n-k-1})) \subset X$. Then, by definition of $\ell(W_3)$ and $Z(E_{2n-k-1})$, we have $V_2 \subset W_3$ and $\dim(V_2 \cap E_{2n-k-1}) \geq 1$. Thus, we get $\dim(W_3 \cap E_{2n-k-1}) \geq 1$. Conversely, for W_3 satisfying $\dim(W_3 \cap E_{2n-k-1}) \geq 1$, one can find a 2-dimensional isotropic subspace $V_2 \subset W_3$ meeting E_{2n-k-1} non-trivially.

- (2) First, assume that $k = 0$. Then, we have $E_2 + E_{2n-1} = \mathbb{C}^{2n}$, as E_2 and E_{2n-1} are assumed to be in general position. Further, since we clearly have $\dot{Y}(Z(E_{2n-1})) = \dot{Y}$, the claim follows.

Now assume that $k \geq 1$. Since E_2 and E_{2n-k-1} are in general position, we have $E_2 \cap E_{2n-k-1} = 0$. If $\ell(W_3)$ is a point of $\dot{Y}(\{E_2\}) \cap \dot{Y}(Z(E_{2n-k-1}))$, then we have the inclusion $W_3 \subset E_2 + E_{2n-k-1}$. Therefore, we have $W_3/E_2 \subset (E_2 + E_{2n-k-1})/E_2$, which proves the claim. \square

COROLLARY 5.4. — $\langle \text{pt}, \sigma_2, \sigma_i, \sigma_j \rangle_1 = \delta_{i+j, 2n-2}$.

Proof. — As was mentioned at the beginning of this section, the invariant vanishes unless $i + j = 2n - 2$. So we assume $i + j = 2n - 2$. In that case, $\langle \text{pt}, \sigma_2, \sigma_i, \sigma_j \rangle_1$ is the number of lines meeting $\{E_2\}$, $Z(E_{2n-3})$, $Z(E_{2n-i-1})$ and $Z(E_{2n-j-1})$, where the subspaces $E_k \subset \mathbb{C}^{2n}$ are in general position with $\omega|_{E_k}$ of minimal rank.

The locus of such lines is the intersection in $\mathbb{P}(\mathbb{C}^{2n}/E_2) \setminus \mathbb{P}(E_2^\perp/E_2)$ of three linear spaces

$$\mathbb{P}((E_2 + E_{2n-3})/E_2), \quad \mathbb{P}((E_2 + E_{2n-i-1})/E_2) \quad \text{and} \quad \mathbb{P}((E_2 + E_{2n-j-1})/E_2).$$

Since these linear spaces are in general position, and of respective codimensions 1, $i - 1$ and $j - 1$ that add up to the dimension of $\mathbb{P}(\mathbb{C}^{2n}/E_2)$, they meet in exactly one point. \square

6. Big quantum cohomology of $\text{IG}(2, 2n)$

In this section we show the regularity of $\text{BQH}(\text{IG}(2, 2n))$ and deduce its generic semisimplicity, which was proved previously in [12, 26].

6.1. Deformation

As before we let $X = \text{IG}(2, 2n)$ and consider the deformation $\text{BQH}_\tau(X)$ of the small quantum cohomology $\text{QH}(X)$ inside the big quantum cohomology $\text{BQH}(X)$ in the direction of the Schubert class $\tau = \sigma_2$. Thus, the ring $\text{BQH}_\tau(X)$ is the quotient of $\text{BQH}(X)$ with respect to the ideal generated by those elements of $(H^*(X, \mathbb{Q}))^*$ which vanish on σ_2 .

Explicitly $\text{BQH}_\tau(X)$ can be described in the following way. For any cohomology classes $a, b \in H^*(X)$ the product is of the form

$$a \star_\tau b = a \star_0 b + t \sum_{\sigma} \left(\sum_{d \geq 1} \langle a, b, \sigma, \sigma_2 \rangle_{dq^d} \right) \sigma^\vee + O(t^2),$$

where $a \star_0 b$ is the small quantum product, σ runs over the basis of the cohomology consisting of Schubert classes, σ^\vee is the dual basis, and t is the deformation parameter of degree -1 .

Remark 6.1. — The variable t can only appear together with a positive power of q so that $q^d t$ can only occur in elements of degree at least $\deg(q) + \deg(t) = 2n - 2$.

Let us have a closer look at the products of the form $\sigma_i \star_\tau \sigma_j$. According to the dimension axiom for GW invariants $\langle \sigma_i, \sigma_j, \sigma, \sigma_2 \rangle_d$ vanishes unless

$$i + j + \deg(\sigma) + 2 = (2n - 1)d + 2(2n - 2).$$

Therefore, applying Corollary 5.4, we have

$$(6.1) \quad \sigma_i \star_\tau \sigma_j = \sigma_i \star_0 \sigma_j + \delta_{i+j, 2n-2} q t + O(t^2),$$

for $i + j \leq 2n - 2$.

6.2. Presentation for $\text{BQH}_\tau(X)$

Consider the following elements in $\text{BQH}_\tau(X)$

$$\begin{aligned} \Delta_r &= \det(\sigma_{1+j-i})_{1 \leq i, j \leq r} \quad \text{for } r \in [3, 2n - 2] \\ \Sigma_{2n-2} &= \sigma_{n-1} \star_\tau \sigma_{n-1} + 2 \sum_{i=1}^{n-1} (-1)^i \sigma_{n-1+i} \star_\tau \sigma_{n-1-i} \\ \Sigma_{2n} &= \sigma_n \star_\tau \sigma_n + 2 \sum_{i=1}^{n-2} (-1)^i \sigma_{n+i} \star_\tau \sigma_{n-i} + (-1)^{n+1} q \sigma_1, \end{aligned}$$

where all products are taken in $\text{BQH}_\tau(X)$. Note that these are “the same” elements as the relations in the presentation of $\text{QH}(X)$ in Theorem 4.1. The lower indices indicate the degree of the respective element in $\text{BQH}_\tau(X)$.

LEMMA 6.2. — We have $\Delta_r = O(t^2)$ for all $r \in [3, 2n - 2]$, $\Sigma_{2n-2} = (-1)^n q t + O(t^2)$ and $\Sigma_{2n} = O(t)$.

Proof.

- (1) Here we prove the statement about Δ_r 's. From Theorem 4.1 and Remark 6.1 we know that $\Delta_r = O(t^2)$ for $r \in [3, 2n - 3]$. Thus, we only need to consider Δ_{2n-2} . Inductively developing the determinant with respect to the first column we have

$$\begin{aligned} \Delta_{2n-2} &= \sum_{s=1}^{2n-2} (-1)^{s-1} \sigma_s \star_\tau \Delta_{2n-2-s} \\ &= O(t^2) - \sigma_{2n-4} \star_\tau \Delta_2 + \sigma_{2n-3} \star_\tau \Delta_1 - \sigma_{2n-2}, \end{aligned}$$

where we used the fact that $\Delta_r = O(t^2)$ for $r \in [3, 2n - 3]$. Thus, we need to prove that

$$(6.2) \quad \sigma_{2n-4} \star_\tau \sigma_2 - \sigma_{2n-4} \star_\tau \sigma_1 \star_\tau \sigma_1 + \sigma_{2n-3} \star_\tau \sigma_1 - \sigma_{2n-2}$$

is of the form $O(t^2)$. We will see this by reducing everything to products of special Schubert classes. First, let us look at the term $\sigma_{2n-4} \star_\tau \sigma_1 \star_\tau \sigma_1$. Since $\sigma_{2n-4} \star_0 \sigma_1 = \sigma_{2n-3} + \sigma'_{2n-3}$, where σ'_{2n-3} is a Schubert class of degree $2n - 3$ different from σ_{2n-3} (see [5]), and for degree reasons, we have

$$(6.3) \quad (\sigma_{2n-4} \star_\tau \sigma_1) \star_\tau \sigma_1 = (\sigma_{2n-3} + \sigma'_{2n-3}) \star_\tau \sigma_1.$$

By (6.1) we have $\sigma_{2n-3} \star_\tau \sigma_1 = \sigma_{2n-3} \star_0 \sigma_1 + qt + O(t^2)$. Thus, we only need to take care of the term $\sigma'_{2n-3} \star_\tau \sigma_1$. For degree reasons we have

$$\sigma'_{2n-3} \star_\tau \sigma_1 = \sigma'_{2n-3} \star_0 \sigma_1 + \langle \sigma'_{2n-3}, \sigma_1, \text{pt}, \sigma_2 \rangle_1 qt + O(t^2).$$

Applying the divisor axiom for Gromov–Witten invariants we see that the 4-point invariant $\langle \sigma'_{2n-3}, \sigma_1, \text{pt}, \sigma_2 \rangle_1$ is equal to the 3-point invariant $\langle \sigma'_{2n-3}, \text{pt}, \sigma_2 \rangle_1$. The latter can be computed using results in [5] and we get $\langle \sigma'_{2n-3}, \text{pt}, \sigma_2 \rangle_1 = 1$. Therefore, we obtain

$$(6.4) \quad \sigma'_{2n-3} \star_\tau \sigma_1 = \sigma'_{2n-3} \star_0 \sigma_1 + qt + O(t^2).$$

Plugging (6.3)–(6.4) into (6.2), using (6.1) and the fact that Δ_{2n-2} vanishes modulo t , we obtain the required statement.

- (2) To prove the statement about Σ_{2n-2} we just apply (6.1) to each summand of Σ_{2n-2} , and use the fact that Σ_{2n-2} vanishes modulo t .
- (3) The statement about Σ_{2n} is evident, as it just claims that there might be a non-trivial deformation along t . \square

By a standard argument (see e.g. [10, Proposition 10]) we obtain the following.

COROLLARY 6.3. — *The ring $\text{BQH}_\tau(X)$ is the quotient of $(K[[t]])[\sigma_1, \dots, \sigma_{2n-2}]$ modulo relations of the form*

$$(6.5) \quad \Delta_r + O(t^2) \quad \text{for all } r \in [3, 2n - 2]$$

$$(6.6) \quad \Sigma_{2n-2} + (-1)^{n+1} qt + O(t^2)$$

$$(6.7) \quad \Sigma_{2n} + O(t).$$

6.3. Structure of $\text{BQH}_\tau(X)$

Recall from Section 4.2 that for $\text{QH}(X)$ we have the direct product decomposition

$$\text{QH}(X) = A \times B = A \times \prod_{i \in I} B_i$$

where $A = K[\varepsilon]/(\varepsilon^{n-1})$ and each B_i is the ground field K . By Hensel's lemma this decomposition lifts to $\text{BQH}_\tau(X)$ and we have

$$(6.8) \quad \text{BQH}_\tau(X) = \mathcal{A} \times \prod_{i \in I} \mathcal{B}_i$$

where \mathcal{A} and \mathcal{B}_i are algebras over $K[[t]]$.

By construction the $K[[t]]$ -algebra $\text{BQH}_\tau(X)$ is a free module over $K[[t]]$ of finite rank equal to $\dim H^*(X)$. Therefore, \mathcal{A} and \mathcal{B}_i 's are also free $K[[t]]$ -modules of finite rank. Reducing modulo t one sees that the rank of \mathcal{A} is $n - 1$ and the ranks of \mathcal{B}_i 's are all equal to 1. Similarly we obtain that the natural algebra homomorphism $K[[t]] \rightarrow \mathcal{B}_i$ is an isomorphism of K -algebras.

THEOREM 6.4. — *The ring $\text{BQH}(X)$ is regular.*

Proof. — Let us first prove that $\text{BQH}_\tau(X)$ is a regular ring, which amounts to showing that \mathcal{A} is regular. Consider the presentation of $\text{BQH}_\tau(X)$ given in Corollary 6.3. Taking linear terms one computes the Zariski tangent space of \mathcal{A} at the maximal ideal (it corresponds to the origin in coordinates $\sigma_1, \dots, \sigma_{2n-2}, t$). We have: (6.5) contribute σ_r for $r \in [3, 2n - 2]$, the relation (6.6) contributes $2\sigma_{2n-2} + qt$, and the relation (6.7) contributes σ_1 (note that the deformation along t has not changed the linear part of this relation). Therefore, the dimension of the tangent space is 1. Since the Krull dimension of \mathcal{A} is also equal to 1, we obtain regularity.

The regularity of $\text{BQH}(X)$ follows from an identical argument. The only difference is the number of the deformation parameters. Note that no additional GW invariants are necessary. \square

COROLLARY 6.5. — *$\text{BQH}(X)$ is generically semisimple.*

Proof. — This is a simple corollary of the above Theorem 6.4. Let us give a short proof for completeness. The coordinate ring of the generic fiber

$$\text{BQH}(X)_\eta = \text{BQH}(X) \otimes_{K[[t_0, \dots, t_s]]} K((t_0, \dots, t_s))$$

is a localization of $\text{BQH}(X)$. Therefore, it is also a regular ring, since $\text{BQH}(X)$ was regular. This implies that $\text{BQH}(X)_\eta$ is a product of finite field extensions of $K((t_0, \dots, t_s))$ which was our definition of semisimplicity. \square

7. Relation to the unfolding of A_{n-1} -singularity

In this section we establish a relation between the quantum cohomology of $\text{IG}(2, 2n)$ and the unfolding of an A_{n-1} -singularity. The comparison takes place at the level of F -manifolds. The concept of an F -manifold was introduced in [15] as a weakening of that of a Frobenius manifold. Every Frobenius manifold gives rise to an F -manifold after forgetting a part of the structure. Both F -manifolds and Frobenius manifolds arise naturally in quantum cohomology and singularity theory. We begin by briefly recalling relevant concepts and refer to [17] for details.

7.1. F -manifolds: generalities

Let M be a complex manifold. Endow \mathcal{T}_M (the holomorphic tangent sheaf) with an \mathcal{O}_M -bilinear, commutative, associative, and unital product $\circ: \mathcal{T}_M \times \mathcal{T}_M \rightarrow \mathcal{T}_M$, and denote e the global unit vector field. The triple (M, \circ, e) is called an F -manifold, if the identity

$$(7.1) \quad \text{Lie}_{X \circ Y}(\circ) = X \circ \text{Lie}_Y(\circ) + Y \circ \text{Lie}_X(\circ)$$

is satisfied for any local vector fields X and Y .

Given two F -manifolds (M_1, \circ_1, e_1) and (M_2, \circ_2, e_2) one can endow the product $M_1 \times M_2$ with a natural structure of an F -manifold (see [17, Proposition 2.10]). The resulting F -manifold is denoted $(M_1 \times M_2, \circ_1 \boxplus \circ_2, e_1 \boxplus e_2)$.

If (M, \circ, e) is an F -manifold and $P \in M$ a point, then by (M, P, \circ, e) we will denote the germ of (M, \circ, e) at P . A germ (M, P, \circ, e) is called *irreducible* if it cannot be written as a product of germs of F -manifolds.

Let (M, P, \circ, e) be a germ of an F -manifold. The tangent space $T_P M$ is a finite dimensional commutative \mathbb{C} -algebra, and as such it uniquely decomposes into the product of local algebras $T_P M = B_1 \times \dots \times B_r$. Moreover, this decomposition extends to an analytic neighbourhood of P , i.e. we have the decomposition

$$(M, P, \circ, e) = \prod_{i=1}^r (M_i, P_i, \circ_i, e_i)$$

of germs of irreducible F -manifolds (see [17, Theorem 2.11]).

7.2. F -manifold of $\text{BQH}(X)$

In Section 2 we have defined big quantum cohomology ring $\text{BQH}(X)$ as an algebra over $K[[t_0, \dots, t_s]]$, where K is the algebraic closure of $\mathbb{Q}((q))$. Another way of packaging the ingredients appearing in $\text{BQH}(X)$ is in the language of formal Frobenius manifold and formal F -manifolds (see [24]).

For reasons that will become clear later, the setting of formal F -manifolds is not suitable for our needs. Therefore, we will describe how to pass to the analytic setting by making some convergence assumption. As was already mentioned in Remark 2.2, $\text{BQH}(X)$ can be defined as an algebra over $R[[t_0, \dots, t_s]]$, where $R = \mathbb{Q}[[q]]$. Moreover, since X is a Fano variety, one can further replace R with $\mathbb{Q}[[q]]$. Further, we can specialize to $q = 1$ and replace \mathbb{Q} with \mathbb{C} . This way we get $\text{BQH}(X)$ defined over $\mathbb{C}[[t_0, \dots, t_s]]$ which we will denote $\text{BQH}(X)_{\mathbb{C}}$.

Another way to formulate the above procedure is to say that for a Fano variety X , if we put $q = 1$, the Gromov–Witten Potential (2.1) gives an element of the power series ring $\mathbb{C}[[t_0, \dots, t_n]]$. Let us denote this power series by $\Phi_{q=1}$.

A priori it is not clear that $\Phi_{q=1}$ has a non-trivial domain of convergence. This is one of the standard expectations in Gromov–Witten theory, but by no means an established result. Therefore, we will make the following assumption.

CONVERGENCE ASSUMPTION. — *The power series $\Phi_{q=1}$ converges in some open neighbourhood M of the origin.*

Repeating the definition of quantum multiplication we now obtain an analytic F -manifold (even Frobenius) structure on M . Namely, basis elements Δ_i appearing in (2.2) form a basis of sections of the holomorphic tangent sheaf of M . Thus, Formula (2.2) endows \mathcal{T}_M with a multiplication, and one can further check that (7.1) is satisfied.

7.3. F -manifolds for isolated hypersurfaces singularities

An *isolated hypersurface singularity* is a holomorphic function germ $f: (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}, 0)$ with an isolated singularity at the origin, i.e. the origin is an isolated critical point of f . One defines the *Milnor algebra* of f as $\mathcal{O}_{\mathbb{C}^m, 0} / (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_m})$, which is the algebra of functions on the critical locus of f .

An *unfolding* of an isolated hypersurface singularity $f: (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}, 0)$ is a holomorphic function germ $F: (\mathbb{C}^m, 0) \times (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$ satisfying $F|_{\mathbb{C}^m \times \{0\}} = f$. The germ $(\mathbb{C}^n, 0)$ is called the *base* of the unfolding and is often denoted $(M, 0)$. The *relative critical locus* $C \subset (\mathbb{C}^m, 0) \times (M, 0)$ is the closed subspace defined by the ideal $J_F = (\frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_m})$. Note that C is not necessarily reduced. This setup can be conveniently described by the diagram

$$(7.2) \quad \begin{array}{ccc} C & \xrightarrow{i} & (\mathbb{C}^m \times M, 0) \xrightarrow{F} (\mathbb{C}, 0) \\ & \searrow p & \downarrow \pi \\ & & (M, 0) \end{array}$$

Define a morphism of sheaves

$$(7.3) \quad \mathcal{T}_{(M,0)} \rightarrow p_* \mathcal{O}_C$$

by sending a local vector field X to $\tilde{X}(F)|_C$, where \tilde{X} is a lift of X to $(\mathbb{C}^m \times M, 0)$. One says that unfolding F is *semiuniversal* if (7.3) is an isomorphism (cf. [17, Theorem 5.1]). In this case one can transport the product structure from $p_* \mathcal{O}_C$ to \mathcal{T}_M . Moreover, this multiplication endows $(M, 0)$ with an F -manifold structure (cf. [17, Theorem 5.3]).

7.3.1. Example: A_n -singularity

We illustrate the above construction in the simplest case of $f = x^{n+1}$, which is also the relevant case for our considerations. In this case a semiuniversal unfolding is given by

$$F = x^{n+1} + a_{n-1}x^{n-1} + \dots + a_1x + a_0,$$

where a_0, \dots, a_{n-1} are coordinates on the base of the unfolding $M = \mathbb{C}^n$. The fact that f is a function of one variable is a special feature of this example. Among isolated hypersurface singularities it is the only one that can be realised in this way.

7.4. Spectral cover of an F -manifold

Let (M, \circ, e) be an F -manifold. The *spectral cover* of (M, \circ, e) is defined as the relative analytic spectrum $\text{Specan}(\mathcal{T}_M, \circ)$. By construction there exists a canonical morphism $\text{Specan}(\mathcal{T}_M, \circ) \rightarrow M$. In general $\text{Specan}(\mathcal{T}_M, \circ)$ is only an analytic space, i.e. it could be singular and/or

non-reduced. The canonical morphism $\text{Specan}(\mathcal{T}_M, \circ) \rightarrow M$ is finite, and in the reduced case can be seen as a ramified covering of M .

In the quantum cohomology setting the spectral cover is given by the spectrum of the big quantum cohomology ring. In the singularity theory setting the spectral cover is nothing else but the relative critical locus C that appears in (7.2).

An F -manifold (M, \circ, e) is called generically semisimple if there is an open subset $U \subset M$, so that for every point in $P \in U$ the algebra structure on $T_P M$ is semisimple.

One can nicely characterise generic semisimplicity of an F -manifold in terms of its spectral cover. Namely, we have the following

THEOREM 7.1 ([17, Theorem 3.2]). — *An F -manifold (M, \circ, e) is generically semisimple if and only if its spectral cover is reduced.*

Note that, the “if” part of the statement is an immediate consequence of the generic smoothness in characteristic zero.

The next theorem can be viewed as a refinement of the above one and gives a beautiful characterisation of (germs of) F -manifolds with smooth spectral covers.

THEOREM 7.2 ([17, Theorem 5.6]). — *Any germ of an irreducible generically semisimple F -manifold with smooth spectral cover is isomorphic to the base space of a semiuniversal unfolding of an isolated hypersurface singularity.*

7.5. F -manifold of $\text{BQH}(\text{IG}(2, 2n))$

Let (M, O, \circ, e) be the germ at the origin of the F -manifold for the big quantum cohomology of $\text{IG}(2, 2n)$ as defined in Section 7.2. The decomposition of $\text{QH}(\text{IG}(2, 2n))$ described in Section 4.2 gives rise to the decomposition of the algebra $(T_O M, \circ_O) = \text{QH}(\text{IG}(2, 2n))_{\mathbb{C}}$ into the product of local algebras. Thus, according to Section 7.1, we obtain the product decomposition

$$(M, O, \circ, e) = (M_0, P_0, \circ_0, e_0) \times \prod_{i=1}^{(2n-1)(n-1)} (M_i, P_i, \circ_i, e_i),$$

where (M_i, P_i, \circ_i, e_i) are one-dimensional F -manifolds and (M_0, P_0, \circ_0, e_0) is an irreducible $(n - 1)$ -dimensional F -manifold.

Computing the rank of the Jacobian matrix, similar to the proof of Theorem 6.4, implies that the spectral cover of (M, O, \circ, e) and, hence, of

(M_i, P_i, \circ_i, e_i) is smooth. Thus, by Theorem 7.2 each germ (M_i, P_i, \circ_i, e_i) is isomorphic to the base space of a semiuniversal unfolding of an isolated hypersurface singularity. Moreover, we can identify the type of this singularity.

THEOREM 7.3. — *If the convergence assumption of Section 7.2 is satisfied for $\text{IG}(2, 2n)$, then the following statements hold.*

- (1) *The F -manifold (M_0, P_0, \circ_0, e_0) is isomorphic to the base space of a semiuniversal unfolding of an isolated hypersurface singularity of type A_{n-1} .*
- (2) *For $i \geq 1$ the F -manifolds (M_i, P_i, \circ_i, e_i) are isomorphic to the base space of a semiuniversal unfolding of an isolated hypersurface singularity of type A_1 .*

Proof. — We only prove the first assertion. The second one is proved in a similar way.

The statement follows from the following observation. Proposition 4.3 implies that the tangent space to $\text{Spec } T_{P_0} M_0$ at its unique point is of dimension one. Now, assuming that (M_0, P_0, \circ_0, e_0) is given by an unfolding of f , we get an identification between the Milnor algebra of f and $T_{P_0} M_0$. In singularity theory the dimension of the tangent space to the spectrum of the Milnor algebra is known as *corank of f at the critical point*. Now, the statement follows from the fact that the only singularity of corank 1 and Milnor number $n - 1$ is the A_{n-1} -singularity (see [1, Section 11.1]). \square

Remark 7.4. — Note that the convergence assumption is classically expected (for example, see [6] and references therein). Further, it may be possible that Theorem 7.2 holds in the formal setting [14].

8. LG model: expectations

In this paragraph we will see how our previous results fit into the general picture of homological mirror symmetry for Fano varieties. Note that we are not going to prove a mirror symmetry type statement, but rather speculate about properties of a Landau–Ginzburg mirror for $\text{IG}(2, 2n)$.

8.1. Mirror symmetry for Fano varieties

Let X be a smooth projective Fano variety and (Y, w) a Landau–Ginzburg model, i.e. Y is a smooth variety and w a regular function on it. To be able

to formulate mirror-symmetry-type statements we also endow X and Y with symplectic forms compatible with respective complex structures but our notation will not reflect the symplectic forms explicitly. For simplicity we assume that w has only isolated, but possibly degenerate, critical points.

To X one attaches two triangulated categories — the category of D-branes of type A and the category of D-branes of type B. The category of D-branes of type A is the derived Fukaya category $D\text{Fuk}(X)$, which encodes the symplectic geometry of X and is independent of the complex structure. The category of D-branes of type B is $D^b\text{Coh}(X)$, the bounded derived category of coherent sheaves on X , which encodes the complex geometry of X and is independent of the symplectic structure.

Further one defines categories of D-branes of type A and B for the pair (Y, w) . The category of D-branes of type A is the derived Fukaya–Seidel category $D\text{FS}(Y, w)$ and the category of D-branes of type B is the triangulated category of matrix factorisations $\text{MF}(Y, w)$. As before the category of D-branes of type A is independent of the complex structure and the category of D-branes of type B of the symplectic one.

One says that X and (Y, w) are mirror to each other if there exist triangulated equivalences of categories

$$(8.1) \quad D\text{Fuk}(X) \simeq \text{MF}(Y, w)$$

$$(8.2) \quad D^b\text{Coh}(X) \simeq D\text{FS}(Y, w).$$

As it is not essential for our purposes, we refrain from recalling precise definitions of the above categories and instead refer to [25] and references therein.

8.2. Quantum cohomology and mirror symmetry

Up to now we viewed the quantum cohomology of X as a purely algebro-geometric object. Historically, Gromov–Witten invariants and quantum cohomology were first defined in symplectic geometry [27], and in mirror symmetry they appear on the symplectic side. Namely, conjecturally there exists an isomorphism of algebras

$$\text{QH}(X) \rightarrow HH^*(D\text{Fuk}(X)),$$

known as closed-open map (e.g. see [13]), from the small quantum cohomology to the Hochschild cohomology of $D\text{Fuk}(X)$.

Let us assume that X and (Y, w) are mirror to each other. Then (8.1) gives rise to the isomorphism

$$HH^*(D \text{Fuk}(X)) \simeq HH^*(\text{MF}(Y, w)).$$

According to [23, Theorem 3.1], the Hochschild cohomology $HH^*(\text{MF}(Y, w))$ is isomorphic to the Milnor algebra of (Y, w) . Therefore, reduced points of $\text{Spec } \text{QH}(X)$ correspond to non-degenerate critical points of w , whereas fat points correspond to degenerate ones.

8.3. LG model for $\text{IG}(2, 2n)$

Let us assume that $\text{IG}(2, 2n)$ has a mirror LG model (Y, w) with only isolated critical points. Then the above discussion combined with Proposition 4.3 imply that the critical locus of w has only one degenerate critical point. Moreover, Theorem 7.3 predicts that the degenerate critical point is of type A_{n-1} .

The above should imply (see [25, 29]) that $D \text{FS}(Y, w)$ has a semiorthogonal decomposition of the form

$$(8.3) \quad D \text{FS}(Y, w) = \langle \mathcal{C}_{n-1}, E_1, \dots, E_{(2n-1)(n-1)} \rangle$$

where E_i are exceptional objects given by vanishing thimbles attached to non-degenerate critical points and \mathcal{C}_{n-1} is the subcategory generated by vanishing thimbles attached to the degenerate one.

Further, it should also follow that subcategory \mathcal{C}_{n-1} is the Fukaya–Seidel category of an isolated hypersurface singularity of type A_{n-1} . The latter was studied by P. Seidel in [29] and shown to be equivalent to the bounded derived category of representations of the quiver of type A_{n-1} . Note that \mathcal{C}_{n-1} also has a full exceptional collection.

The above discussion combined with equivalence (8.2) suggest that $D^b \text{Coh}(\text{IG}(2, 2n))$ should have a semiorthogonal decomposition (in fact, a full exceptional collection)

$$D^b \text{Coh}(\text{IG}(2, 2n)) = \langle \mathcal{A}_{n-1}, E_1, \dots, E_{(2n-1)(n-1)} \rangle,$$

where subcategory \mathcal{A}_{n-1} is equivalent to the bounded derived category of representations of the quiver of type A_{n-1} . Such a decomposition is constructed explicitly in Theorem 9.6 of the Appendix by Alexander Kuznetsov.

9. Appendix (by Alexander Kuznetsov)

9.1. Lefschetz exceptional collections

Recall the notion of a Lefschetz semiorthogonal decomposition ([19, Definition 4.1] and [21, Definition 3.1]). In this paper we will need its special case, a Lefschetz exceptional collection ([20, Definition 2.1]). We will remind this notion for the readers convenience.

DEFINITION 9.1. — A Lefschetz exceptional collection in $D^b(X)$ with respect to a line bundle $\mathcal{O}_X(1)$ is an exceptional collection in $D^b(X)$ which has a block structure

$$\left(\underbrace{(E_1, E_2, \dots, E_{\lambda_0})}_{\text{block 1}}, \underbrace{(E_1(1), E_2(1), \dots, E_{\lambda_1}(1))}_{\text{block 2}}, \dots, \underbrace{(E_1(m-1), E_2(m-1), \dots, E_{\lambda_{m-1}}(m-1))}_{\text{block } m} \right)$$

where $\lambda = (\lambda_0 \geq \lambda_1 \geq \dots \geq \lambda_{m-1} \geq 0)$ is a non-increasing sequence of positive integers (the support partition of the Lefschetz collection). A Lefschetz exceptional collection is full if it generates the derived category $D^b(X)$. It is called rectangular if $\lambda_0 = \lambda_1 = \dots = \lambda_{m-1}$ (i.e., if the Young diagram of its support partition is a rectangle).

Clearly, to specify a Lefschetz exceptional collection one needs to give its first block and its support partition. We will denote by (E_\bullet, λ) the corresponding Lefschetz exceptional collection. Let us list several interesting examples.

Example 9.2. — Let $X = G(2, m)$. Set $k = \lfloor m/2 \rfloor$, $E_i = S^{i-1}\mathcal{U}^*$ for $1 \leq i \leq k$, and

$$\lambda = \begin{cases} (k^{2k+1}), & \text{if } m = 2k + 1 \text{ is odd,} \\ (k^k, (k-1)^k), & \text{if } m = 2k \text{ is even} \end{cases}$$

(here the exponents stand for the multiplicities of the entries, thus the first line denotes the partition with $2k + 1$ parts equal to k , and the second line means the partition with k parts equal k and k parts equal $k - 1$).

Example 9.3. — Let $X = IG(2, 2k)$. Set $E_i = S^{i-1}\mathcal{U}^*$ for $1 \leq i \leq k$, and $\lambda = (k^{k-1}, (k-1)^k)$ with the same convention about the exponents as in the previous example.

The collections listed in these examples are full [20, Theorems 4.1, 5.1]. With the conventions we took they can be rewritten as

$$(9.1) \quad D^b(G(2, 2k+1)) = \langle S^{\bullet-1}\mathcal{U}^*, (k^{2k+1}) \rangle,$$

$$(9.2) \quad D^b(G(2, 2k)) = \langle S^{\bullet-1}\mathcal{U}^*, (k^k, (k-1)^k) \rangle,$$

$$(9.3) \quad D^b(\mathrm{IG}(2, 2k)) = \langle S^{\bullet-1}\mathcal{U}^*, (k^{k-1}, (k-1)^k) \rangle.$$

The first of these collections is rectangular, while the last two are not.

It is known that $G(k, n)$ has a rectangular Lefschetz exceptional collection of length n if and only if k and n are coprime [9]. Lefschetz exceptional collections on general isotropic Grassmannians are not yet known, it is a good question to construct such a collection.

DEFINITION 9.4. — *Given a Lefschetz exceptional collection (E_\bullet, λ) with the length of λ equal to m , the subcollection*

$$(E_\bullet, (\lambda_{m-1})^m) = (E_1, \dots, E_{\lambda_{m-1}}, E_1(1), \dots, E_{\lambda_{m-1}}(1), \dots, \\ E_1(m-1), \dots, E_{\lambda_{m-1}}(m-1))$$

is called the rectangular part of (E_\bullet, λ) . The subcategory orthogonal to the rectangular part of a Lefschetz exceptional collection is called the residual subcategory. It is zero if and only if the original collection is full and rectangular.

The main result of this section is the following two theorems describing the residual subcategories of (9.2) and (9.3). Note that the residual subcategory of (9.1) is zero.

THEOREM 9.5. — *The residual subcategory of (9.2) is generated by k completely orthogonal exceptional objects. In particular, it is equivalent to the derived category of the union of k disjoint points.*

THEOREM 9.6. — *The residual subcategory of (9.3) is equivalent to the derived category of representations of A_{k-1} quiver.*

We prove Theorem 9.5 in Section 9.2, and Theorem 9.6 in Section 9.3 below.

9.2. Usual Grassmannian

Denote the block of the rectangular part of the collection (9.2) by \mathcal{A} , so that

$$\mathcal{A} = \langle \mathcal{O}, \mathcal{U}^*, \dots, S^{k-2}\mathcal{U}^* \rangle.$$

The Lefschetz decomposition of $G(2, 2k)$ then can be rewritten as

$$(9.4) \quad D^b(G(2, 2k)) = \langle \mathcal{A}, S^{k-1}\mathcal{U}^*, \mathcal{A}(1), S^{k-1}\mathcal{U}^*(1), \dots, \\ \mathcal{A}(k-1), S^{k-1}\mathcal{U}^*(k-1), \mathcal{A}(k), \mathcal{A}(k+1), \dots, \mathcal{A}(2k-1) \rangle.$$

We denote the corresponding residual subcategory of $G(2, 2k)$ by \mathcal{R}_k , so that

$$\mathcal{R}_k = \langle \mathcal{A}, \mathcal{A}(1), \dots, \mathcal{A}(2k-1) \rangle^\perp.$$

Using mutations of exceptional collections it is easy to deduce from (9.4) that

$$(9.5) \quad \mathcal{R}_k = \langle \mathbb{L}_{\mathcal{A}}(S^{k-1}\mathcal{U}^*), \mathbb{L}_{\langle \mathcal{A}, \mathcal{A}(1) \rangle}(S^{k-1}\mathcal{U}^*(1)), \\ \dots \mathbb{L}_{\langle \mathcal{A}, \mathcal{A}(1), \dots, \mathcal{A}(k-1) \rangle}(S^{k-1}\mathcal{U}^*(k-1)) \rangle,$$

where $\mathbb{L}_{\mathcal{A}}$ is the left mutation through \mathcal{A} , $\mathbb{L}_{\langle \mathcal{A}, \mathcal{A}(1) \rangle}$ is the left mutation through $\langle \mathcal{A}, \mathcal{A}(1) \rangle$, and so on. So, it is enough to show that these objects (they are automatically exceptional) are completely orthogonal.

For this we will use an alternative description of the objects, obtained by using the following long exact sequence (constructed in [20, Proof of Lemma 4.3]):

$$(9.6) \quad 0 \rightarrow S^{k-1}\mathcal{U}^*(-k) \rightarrow V \otimes S^{k-2}\mathcal{U}^*(1-k) \rightarrow \dots \\ \rightarrow \Lambda^{k-2}V \otimes \mathcal{U}^*(-2) \rightarrow \Lambda^{k-1}V \otimes \mathcal{O}_X(-1) \rightarrow \Lambda^{k-1}V^* \otimes \mathcal{O}_X \\ \rightarrow \Lambda^{k-2}V^* \otimes \mathcal{U}^* \rightarrow \dots \rightarrow V^* \otimes S^{k-2}\mathcal{U}^* \rightarrow S^{k-1}\mathcal{U}^* \rightarrow 0.$$

The complex formed by the last k terms of (9.6) is the Koszul complex, and its cohomology is isomorphic to $\Lambda^{k-1}\mathcal{U}^\perp$. The complex formed by the first k terms of (9.6) is dual to the above complex twisted by $\mathcal{O}_X(-1)$ taking into account isomorphisms $S^i\mathcal{U} \cong S^i\mathcal{U}^*(-i)$ (obtained by taking the symmetric powers of the isomorphism $\mathcal{U} \cong \mathcal{U}^*(-1)$). Consequently, the cohomology of the complex formed by the first k terms of (9.6) is isomorphic to $\Lambda^{k-1}(V/\mathcal{U}) \otimes \mathcal{O}_X(-1)$. So, the canonical isomorphism

$$\Lambda^{k-1}\mathcal{U}^\perp \cong \Lambda^{k-1}(V/\mathcal{U}) \otimes \mathcal{O}_X(-1)$$

allows to glue the two halves of (9.6) into a single long exact sequence.

We denote by F_i the cohomology of the subcomplex of (9.6) consisting of its first i terms. In other words, F_i are defined by the following exact sequences

$$(9.7) \quad 0 \rightarrow S^{k-1}\mathcal{U}^*(-k) \rightarrow V \otimes S^{k-2}\mathcal{U}^*(1-k) \rightarrow \dots \\ \rightarrow \Lambda^{i-1}V \otimes S^{k-i}\mathcal{U}^*(i-k-1) \rightarrow F_i \rightarrow 0.$$

LEMMA 9.7. — For every $1 \leq i \leq k$ there is an isomorphism

$$\mathbb{L}_{\langle \mathcal{A}, \dots, \mathcal{A}(k-i) \rangle}(S^{k-1}\mathcal{U}^*(k-i)) \cong F_i(k-i)[2k-i-1].$$

Proof. — Since (9.6) is exact, we have yet another exact sequence for F_i :

$$(9.8) \quad 0 \rightarrow F_i \rightarrow \Lambda^i V \otimes S^{k-i-1}\mathcal{U}^*(i-k) \rightarrow \dots \rightarrow \Lambda^{k-1} V \otimes \mathcal{O}_X(-1) \rightarrow \\ \rightarrow \Lambda^{k-1} V^* \otimes \mathcal{O}_X \rightarrow \Lambda^{k-2} V^* \otimes \mathcal{U}^* \rightarrow \dots \rightarrow V^* \otimes S^{k-2}\mathcal{U}^* \rightarrow S^{k-1}\mathcal{U}^* \rightarrow 0.$$

Twisting it by $\mathcal{O}_X(k-i)$ we note that all its terms (except for the leftmost $F_i(k-i)$ and the rightmost $S^{k-1}\mathcal{U}^*(k-i)$) lie in the subcategory $\langle \mathcal{A}, \dots, \mathcal{A}(k-i-1), \mathcal{A}(k-i) \rangle$. In other words, the cone of the morphism

$$S^{k-1}\mathcal{U}^*(k-i) \rightarrow F_i(k-i)[2k-i-1]$$

represented by the extension class of this exact sequence, is contained in the subcategory $\langle \mathcal{A}, \dots, \mathcal{A}(k-i-1), \mathcal{A}(k-i) \rangle$. So, to prove the claim of the lemma, it is enough to show that $F_i(k-i) \in \langle \mathcal{A}, \dots, \mathcal{A}(k-i) \rangle^\perp$. For this we use the twist by $\mathcal{O}_X(k-i)$ of (9.7). The terms of this twist clearly belong to the first half $\langle \mathcal{A}, S^{k-1}\mathcal{U}^*, \dots, \mathcal{A}(k-1), S^{k-1}\mathcal{U}^*(k-1) \rangle$ twisted by $\mathcal{O}_X(-k)$, hence to the orthogonal of the second half of (9.4) twisted by $\mathcal{O}_X(-k)$, i.e., to $\langle \mathcal{A}, \mathcal{A}(1), \dots, \mathcal{A}(k-1) \rangle^\perp$. This gives the required embedding. \square

Combining this lemma with (9.5) we deduce an exceptional collection

$$(9.9) \quad \mathcal{R}_k = \langle F_k, F_{k-1}(1), \dots, F_1(k-1) \rangle.$$

To prove the theorem it remains to show it is completely orthogonal.

LEMMA 9.8. — We have $\text{Ext}^\bullet(F_i(k-i), F_j(k-j)) = 0$ for all $1 \leq j < i \leq k$.

Proof. — To show this we use for $F_i(k-i)$ resolution (9.8) twisted by $\mathcal{O}_X(k-i)$, and for $F_j(k-j)$ resolution (9.7) twisted by $\mathcal{O}_X(k-j)$. The terms of the first are contained in the subcategory

$$\langle \mathcal{A}, \dots, \mathcal{A}(k-i-1), \mathcal{A}(k-i), S^{k-1}\mathcal{U}^*(k-i) \rangle,$$

and the terms of the second are contained in the subcategory

$$\langle S^{k-1}\mathcal{U}^*(-j), \mathcal{A}(1-j), \dots, \mathcal{A}(-1) \rangle.$$

All the components of these subcategories are clearly semiorthogonal. This follows immediately from the twist of (9.4) by $\mathcal{O}_X(-j)$, since $-j < k-i \leq k-1-j$ by the assumption on i and j taken in the lemma. \square

This lemma together with the exceptional collection (9.9) proves Theorem 9.5.

9.3. Isotropic Grassmannian

Denote the block of the rectangular part of the collection (9.3) by $\bar{\mathcal{A}}$, so that

$$\bar{\mathcal{A}} = \langle \mathcal{O}, \mathcal{U}^*, \dots, S^{k-2}\mathcal{U}^* \rangle.$$

The Lefschetz decomposition of $\mathrm{IG}(2, 2k)$ then can be rewritten as

$$(9.10) \quad D^b(\mathrm{IG}(2, 2k)) = \langle \bar{\mathcal{A}}, S^{k-1}\mathcal{U}^*, \bar{\mathcal{A}}(1), S^{k-1}\mathcal{U}^*(1), \dots, \bar{\mathcal{A}}(k-2), S^{k-1}\mathcal{U}^*(k-2), \bar{\mathcal{A}}(k-1), \bar{\mathcal{A}}(k), \dots, \bar{\mathcal{A}}(2k-2) \rangle.$$

We denote the corresponding residual subcategory of $\mathrm{IG}(2, 2k)$ by $\bar{\mathcal{R}}_k$, so that

$$\bar{\mathcal{R}}_k = \langle \bar{\mathcal{A}}, \bar{\mathcal{A}}(1), \dots, \bar{\mathcal{A}}(2k-2) \rangle^\perp.$$

Using mutations of exceptional collections it is easy to deduce from (9.10) that

$$(9.11) \quad \bar{\mathcal{R}}_k = \langle \mathbb{L}_{\bar{\mathcal{A}}}(S^{k-1}\mathcal{U}^*), \mathbb{L}_{\langle \bar{\mathcal{A}}, \bar{\mathcal{A}}(1) \rangle}(S^{k-1}\mathcal{U}^*(1)), \dots, \mathbb{L}_{\langle \bar{\mathcal{A}}, \bar{\mathcal{A}}(1), \dots, \bar{\mathcal{A}}(k-2) \rangle}(S^{k-1}\mathcal{U}^*(k-2)) \rangle.$$

The first step of the proof in this case is to show that these objects are given by (the restrictions to $\mathrm{IG}(2, 2k)$) of the same sheaves F_i as before.

LEMMA 9.9. — *For every $2 \leq i \leq k$ there is an isomorphism*

$$\mathbb{L}_{\langle \bar{\mathcal{A}}, \dots, \bar{\mathcal{A}}(k-i) \rangle}(S^{k-1}\mathcal{U}^*(k-i)) \cong F_i(k-i)[2k-i-1].$$

Proof. — The same proof as in Lemma 9.7 applies. We restrict exact sequences (9.7) and (9.8) to $X = \mathrm{IG}(2, 2k)$. Twisting the restriction of (9.8) by $\mathcal{O}_X(k-i)$ we note that all its terms (except for the leftmost $F_i(k-i)$ and the rightmost $S^{k-1}\mathcal{U}^*(k-i)$) lie in the subcategory $\langle \bar{\mathcal{A}}, \dots, \bar{\mathcal{A}}(k-i-1), \bar{\mathcal{A}}(k-i) \rangle$. In other words, the cone of the morphism

$$S^{k-1}\mathcal{U}^*(k-i) \rightarrow F_i(k-i)[2k-i-1]$$

represented by the extension class of this exact sequence, is contained in the subcategory $\langle \bar{\mathcal{A}}, \dots, \bar{\mathcal{A}}(k-i-1), \bar{\mathcal{A}}(k-i) \rangle$. So, to prove the claim of the lemma, it is enough to show that $F_i(k-i) \in \langle \bar{\mathcal{A}}, \dots, \bar{\mathcal{A}}(k-i) \rangle^\perp$. For this we use the twist by $\mathcal{O}_X(k-i)$ of the restriction of (9.7). The terms of this twist belong to the first part $\langle \bar{\mathcal{A}}, S^{k-1}\mathcal{U}^*, \dots, \bar{\mathcal{A}}(k-2), S^{k-1}\mathcal{U}^*(k-2), \bar{\mathcal{A}}(k-1) \rangle$ of (9.10) twisted by $\mathcal{O}_X(-k)$, hence to the orthogonal of the second part of (9.10) twisted by $\mathcal{O}_X(-k)$, i.e., to $\langle \bar{\mathcal{A}}, \bar{\mathcal{A}}(1), \dots, \bar{\mathcal{A}}(k-2) \rangle^\perp$. This gives the required embedding. \square

Combining this lemma with (9.11) we deduce an exceptional collection

$$(9.12) \quad \bar{\mathcal{R}}_k = \langle F_k, F_{k-1}(1), \dots, F_2(k-2) \rangle.$$

Remark 9.10. — Note that the object $F_1(k-1) = S^{k-1}\mathcal{U}^*(-1)$ also belongs to $\bar{\mathcal{R}}_k$. This follows easily by mutating the first component $\bar{\mathcal{A}}$ of (9.10) to the right (it gets twisted by the anticanonical class $\mathcal{O}_X(2k-1)$), and then twisting the resulting decomposition by $\mathcal{O}_X(-1)$.

To prove the theorem it remains to compute Ext-spaces between its objects.

LEMMA 9.11. — *Assume $1 \leq j < i \leq k$. We have*

$$\mathrm{Ext}^\bullet(F_i(k-i), F_j(k-j)) = \begin{cases} \mathbb{C}, & \text{if } i = j + 1, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. — To show this we use for $F_i(k-i)$ (the restriction of) resolution (9.8) twisted by $\mathcal{O}_X(k-i)$, and for $F_j(k-j)$ (the restriction of) resolution (9.7) twisted by $\mathcal{O}_X(k-j)$. The terms of the first are contained in the subcategory

$$\langle \bar{\mathcal{A}}, \dots, \bar{\mathcal{A}}(k-i-1), \bar{\mathcal{A}}(k-i), S^{k-1}\mathcal{U}^*(k-i) \rangle,$$

and the terms of the second are contained in the subcategory

$$\langle S^{k-1}\mathcal{U}^*(-j), \bar{\mathcal{A}}(1-j), \dots, \bar{\mathcal{A}}(-1) \rangle.$$

In case of $i \geq j + 2$ all the components of these subcategories are clearly semiorthogonal. This follows immediately from the twist of (9.10) by $\mathcal{O}_X(-j)$, since $-j < k-i \leq k-2-j$ by the assumption on i and j . If, however, $i = j + 1$ we do not have a semiorthogonality between $S^{k-1}\mathcal{U}^*(k-i)$ and $S^{k-1}\mathcal{U}^*(-j)$. On a contrary, a direct Borel–Bott–Weil calculation shows that

$$\begin{aligned} \mathrm{Ext}^\bullet(S^{k-1}\mathcal{U}^*(k-i), S^{k-1}\mathcal{U}^*(-j)) \\ &\cong \mathrm{Ext}^\bullet(S^{k-1}\mathcal{U}^*, S^{k-1}\mathcal{U}^*(i-j-k)) \\ &\cong \mathrm{Ext}^\bullet(S^{k-1}\mathcal{U}^*, S^{k-1}\mathcal{U}^*(1-k)) \cong \mathbb{C}[3-2k]. \end{aligned}$$

Finally, using exact sequences (9.7) and (9.8) we deduce that

$$\begin{aligned} \mathrm{Ext}^\bullet(F_i(k-i), F_j(k-j)) \\ &\cong \mathrm{Ext}^\bullet(S^{k-1}\mathcal{U}^*(k-i)[i+1-2k], S^{k-1}\mathcal{U}^*(-j)[j-1]) \\ &\cong \mathrm{Ext}^\bullet(S^{k-1}\mathcal{U}^*(k-i), S^{k-1}\mathcal{U}^*(-j))[2k-3] \cong \mathbb{C}. \quad \square \end{aligned}$$

Remark 9.12. — A chain of morphisms

$$F_k \rightarrow F_{k-1}(1) \rightarrow \cdots \rightarrow F_2(k-2) \rightarrow F_1(k-1)$$

computed in Lemma 9.11 can be described as follows. Consider the bicomplex of [20, Proposition 5.3]. Replacing m by k and W by V (according to the difference in the notation), and identifying V with V^* via the symplectic form, we note that the columns of the obtained bicomplex coincide with the sequences (9.7) twisted by $\mathcal{O}(2k-i)$. Since the totalization of the bicomplex is exact, we conclude that the cohomology sheaves of the vertical differential of the bicomplex form an exact sequence

$$0 \rightarrow F_k(k) \rightarrow F_{k-1}(k+1) \rightarrow \cdots \rightarrow F_2(2k-2) \rightarrow F_1(2k-1) \rightarrow 0,$$

from which the required chain can be obtained by a twist.

Lemma 9.11 together with the exceptional collection (9.9) proves Theorem 9.6. Indeed, we can define the functor $D^b(A_{k-1}) \rightarrow \bar{\mathcal{R}}_k$ by the rule

$$S_i \mapsto F_{k+1-i}(i-1)[1-i], \quad 1 \leq i \leq k-1$$

where S_i stand for the simple representations of the quiver. Since

$$\mathrm{Ext}^\bullet(S_i, S_j) = \begin{cases} \mathbb{C}[-1], & \text{if } i = j - 1, \\ 0, & \text{otherwise,} \end{cases}$$

it follows that the functor is fully faithful, and by (9.12) it is essentially surjective.

Remark 9.13. — Under this equivalence the extra object $F_1(k-1) \in \bar{\mathcal{R}}_k$ corresponds (up to a shift) to the projective module of the first vertex of the quiver.

BIBLIOGRAPHY

- [1] V. I. ARNOLD, S. M. GUSEIN-ZADE & A. N. VARCHENKO, *Singularities of differentiable maps. Volume 1. Classification of critical points, caustics and wave fronts*, Modern Birkhäuser Classics, Birkhäuser, 2012, translated from the Russian by Ian Porteous based on a previous translation by Mark Reynolds, reprint of the 1985 edition, xii+382 pages.
- [2] A. BAYER, “Semisimple quantum cohomology and blowups”, *Int. Math. Res. Not.* (2004), no. 40, p. 2069-2083.
- [3] A. S. BUCH, A. KRESCH & H. TAMVAKIS, “Quantum Pieri rules for isotropic Grassmannians”, *Invent. Math.* **178** (2009), no. 2, p. 345-405.
- [4] P.-E. CHAPUT, L. MANIVEL & N. PERRIN, “Quantum cohomology of minuscule homogeneous spaces III. Semi-simplicity and consequences”, *Can. J. Math.* **62** (2010), no. 6, p. 1246-1263.
- [5] P.-E. CHAPUT & N. PERRIN, “On the quantum cohomology of adjoint varieties”, *Proc. Lond. Math. Soc.* **103** (2011), no. 2, p. 294-330.

- [6] T. COATES & H. IRITANI, “On the convergence of Gromov-Witten potentials and Givental’s formula”, *Mich. Math. J.* **64** (2015), no. 3, p. 587-631.
- [7] B. DUBROVIN, “Geometry and analytic theory of Frobenius manifolds”, *Doc. Math.* (1998), p. 315-326, Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998).
- [8] ———, “Quantum cohomology and isomonodromic deformation”, Lecture at “Recent Progress in the Theory of Painlevé Equations: Algebraic, asymptotic and topological aspects”, Strasbourg., 2013.
- [9] A. V. FONARĚV, “Minimal Lefschetz decompositions of the derived categories for Grassmannians”, *Izv. Math.* **77** (2013), no. 5, p. 203-224.
- [10] W. FULTON & R. PANDHARIPANDE, “Notes on stable maps and quantum cohomology”, in *Algebraic geometry—Santa Cruz 1995*, Proceedings of Symposia in Pure Mathematics, vol. 62, American Mathematical Society, 1997, p. 45-96.
- [11] S. GALKIN, V. GOLYSHEV & H. IRITANI, “Gamma classes and quantum cohomology of Fano manifolds: gamma conjectures”, *Duke Math. J.* **165** (2016), no. 11, p. 2005-2077.
- [12] S. GALKIN, A. MELLIT & M. SMIRNOV, “Dubrovin’s conjecture for $IG(2, 6)$ ”, *Int. Math. Res. Not.* (2015), no. 18, p. 8847-8859.
- [13] S. GANATRA, “Automatically generating Fukaya categories and computing quantum cohomology”, <https://arxiv.org/abs/1605.07702>, 2016.
- [14] C. HERTLING, Email communication dated 04.04.2016.
- [15] C. HERTLING & YU. I. MANIN, “Weak Frobenius manifolds”, *Int. Math. Res. Not.* (1999), no. 6, p. 277-286.
- [16] C. HERTLING, YU. I. MANIN & C. TELEMAN, “An update on semisimple quantum cohomology and F -manifolds”, *Tr. Mat. Inst. Steklova* **264** (2009), p. 69-76.
- [17] C. HERTLING, *Frobenius manifolds and moduli spaces for singularities*, Cambridge Tracts in Mathematics, vol. 151, Cambridge University Press, 2002, x+270 pages.
- [18] S. L. KLEIMAN, “The transversality of a general translate”, *Compos. Math.* **28** (1974), p. 287-297.
- [19] A. KUZNETSOV, “Homological projective duality”, *Publ. Math., Inst. Hautes Étud. Sci.* (2007), no. 105, p. 157-220.
- [20] ———, “Exceptional collections for Grassmannians of isotropic lines”, *Proc. Lond. Math. Soc.* **97** (2008), no. 1, p. 155-182.
- [21] ———, “Semiorthogonal decompositions in algebraic geometry”, in *Proceedings of the International Congress of Mathematicians, Vol. II (Seoul, 2014)*, KM Kyung Moon Sa, 2014, p. 635-660.
- [22] J. M. LANDSBERG & L. MANIVEL, “On the projective geometry of rational homogeneous varieties”, *Comment. Math. Helv.* **78** (2003), no. 1, p. 65-100.
- [23] K. H. LIN & D. POMERLEANO, “Global matrix factorizations”, *Math. Res. Lett.* **20** (2013), no. 1, p. 91-106.
- [24] YU. I. MANIN, *Frobenius manifolds, quantum cohomology, and moduli spaces*, Colloquium Publications, vol. 47, American Mathematical Society, 1999, xiv+303 pages.
- [25] D. ORLOV, “Landau-Ginzburg models, D-branes and mirror symmetry”, *Mat. Contemp.* **41** (2012), p. 75-112.
- [26] N. PERRIN, “Semisimple quantum cohomology of some Fano varieties”, <https://arxiv.org/abs/1405.5914v1>, 2014.
- [27] Y. RUAN & G. TIAN, “A mathematical theory of quantum cohomology”, *J. Differ. Geom.* **42** (1995), no. 2, p. 259-367.
- [28] A. SAMOKHIN, “Some remarks on the derived categories of coherent sheaves on homogeneous spaces”, *J. Lond. Math. Soc.* **76** (2007), no. 1, p. 122-134.

- [29] P. SEIDEL, “More about vanishing cycles and mutation”, in *Symplectic geometry and mirror symmetry (Seoul, 2000)*, World Scientific, 2001, p. 429-465.
- [30] R. P. STANLEY, “Hilbert functions of graded algebras”, *Adv. Math.* **28** (1978), no. 1, p. 57-83.
- [31] E. STRICKLAND, “Lines in G/P ”, *Math. Z.* **242** (2002), no. 2, p. 227-240.

Manuscrit reçu le 23 juin 2017,
 accepté le 14 septembre 2017.

John Alexander CRUZ MORALES
 Universidad Nacional de Colombia
 Departamento de Matemáticas
 Carrera 45 No. 26–85 Edificio Uriel Gutiérrez
 Bogotá D.C. (Colombia)
jacruzmo@unal.edu.co

Anton MELLIT
 Faculty of Mathematics
 University of Vienna
 Oskar-Morgenstern-Platz 1
 1090 Vienna (Austria)
anton.mellit@univie.ac.at

Nicolas PERRIN
 Laboratoire de Mathématiques de Versailles,
 UVSQ, CNRS, Université Paris–Saclay
 78035 Versailles (France)
nicolas.perrin@uvsq.fr

Maxim SMIRNOV
 Universität Augsburg, Institut für Mathematik
 Universitätsstr. 14
 86159 Augsburg (Germany)
maxim.smirnov@math.uni-augsburg.de

Alexander KUZNETSOV
 Algebraic Geometry Section
 Steklov Mathematical Institute
 of Russian Academy of Sciences
 8 Gubkin str.
 Moscow 119991 (Russia)
and
 Interdisciplinary Scientific Center J.-V. Poncelet
 (CNRS UMI 2615)
 Moscow (Russia)
and
 Laboratory of Algebraic Geometry
 National Research University Higher School of
 Economics
 Moscow (Russia)
akuznet@mi.ras.ru