



ANNALES DE L'INSTITUT FOURIER

François MAUCOURANT

**Unique ergodicity of asynchronous rotations, and
application**

Article à paraître, mis en ligne le 17 juin 2025, 27 p.

Article mis à disposition par son auteur selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE



<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les *Annales de l'Institut Fourier* sont membres du
Centre Mersenne pour l'édition scientifique ouverte

www.centre-mersenne.org

e-ISSN : 1777-5310

UNIQUE ERGODICITY OF ASYNCHRONOUS ROTATIONS, AND APPLICATION

by François MAUCOURANT

ABSTRACT. — The main result of this paper is an analogue for a continuous family of tori of Kronecker–Weyl’s unique ergodicity of irrational rotations. We show that the notion corresponding in this setup to irrationality, namely asynchronicity, is satisfied in some homogeneous dynamical systems. This is used to prove the ergodicity of naturals lifts of invariant measures.

RÉSUMÉ. — Nous étudions sur une famille continue de tores les rotations dites asynchrones, analogues aux rotations irrationnelles sur les tores classiques. Le résultat principal est l’unique ergodicité de ces rotations sur un monoïde adapté. Nous prouvons que la condition d’asynchronicité est vérifiée dans une famille d’exemples issue de la dynamique homogène, ce qui nous permet de déduire l’ergodicité de relevés de certaines transformations dans des fibrés en tores.

1. Introduction

1.1. Motivations

The original motivation of this study is the inquiry of ergodic properties of torus extension of homogeneous dynamical systems. Such dynamics have drawn some attention recently — see for example [7, 10, 13] for unipotent actions, and [2, 12] for diagonal actions.

As an informal example, consider a diagonal element $a \in \mathrm{SL}(d, \mathbb{R}) = G_0$ ($d \geq 2$) with positive diagonal entries, acting by left multiplication on the homogeneous space $\mathrm{SL}(d, \mathbb{R})/\mathrm{SL}(d, \mathbb{Z}) = G_0/\Gamma_0$, and let μ be an a -invariant and ergodic probability measure on G_0/Γ_0 . The main interesting cases for our purpose occur when the measure μ is not algebraic (i.e. not a H -invariant probability measure on a closed homogeneous submanifold

$Hx\Gamma_0$, where $H \subset G_0$ is a closed subgroup containing a). This dynamical system is a factor of the action of (a, α) , where $\alpha \in \mathbb{R}^d$ is arbitrary, by left multiplication on $(\mathrm{SL}(d, \mathbb{R}) \ltimes \mathbb{R}^d)/(\mathrm{SL}(d, \mathbb{Z}) \ltimes \mathbb{Z}^d) = G/\Gamma$. This latter space is a torus bundle above G_0/Γ_0 . Amongst the possible (a, α) -invariant measures that projects onto μ , there is a particular one, denoted by λ , which decomposes into the Haar measure of tori on each fiber. It is natural to ask about its ergodicity with respect to (a, α) .

Following the classical Hopf argument (see [5]), one is naturally led to inquire about ergodic properties of the strong stable foliation of (a, α) . It turns out that this foliation contains the orbits of another action, namely the multiplication by (e, β) on G/Γ , where $\beta \in \mathbb{R}^d$ is an eigenvector for a associated to an eigenvalue < 1 (Lemma 6.1). This action is a unipotent action, but since it is “vertical” (in the sense trivial in the factor G_0/Γ_0), Ratner’s theory yield in this case no more information than Kronecker–Weyl’s uniform distribution on the torus.

To visualize this action of (e, β) on each fiber, one may think of it as a rotation by a fixed vector β on a varying torus depending on the base-point. Here, we will prefer to think of it as the rotation by a varying vector $f_\beta(x)$ depending on the base-point x , on a fixed torus \mathbb{T}^d . One may hope in this situation that $f_\beta(x) \in \mathbb{T}^d$ is irrational for almost every x . It turns out that under appropriate assumptions, the rotations defined by f_β satisfy a stronger property, namely *asynchronicity*.

As we will see shortly, such rotations on torus bundle above a measured space like $(G_0/\Gamma_0, \mu)$ enjoy strong ergodic properties, enabling us in this setting to prove a unique ergodicity result. In some sense, this can be considered as a weak analogue of Furstenberg’s unique ergodicity of horocyclic flow.

Finally, we will return to the question of the ergodicity of λ with respect to (a, α) , and related mixing properties.

These kind of fiber-wise system were also investigated independently by Damien Thomine [14], using another point of view.

1.2. Asynchronous rotations

Let $(I, \mathcal{B}(I), \mathcal{L})$ be a standard probability space without atoms, and let $\mathbb{T}^d = (\mathbb{R}/\mathbb{Z})^d$ be a torus, for some integer $d \geq 1$. We think of a measurable map $f : I \rightarrow \mathbb{T}^d$ as the data, for each $x \in I$, of a rotation adding the angle $f(x)$ in a torus indexed by x . We will call such a map an *angle map*, and

two angle maps will be identified if they coincide \mathcal{L} -almost everywhere. Despite what the above motivational example might suggest, in this abstract setting, the case $d = 1$ of rotations on a family of circles above a probability space is already interesting, and contains most of the difficulties.

The set of angle maps $\{f : I \rightarrow \mathbb{T}^d\}$, is naturally an abelian group under pointwise addition of functions. We denote by $((\mathbb{T}^d)^I, +)$ this group, by a slight abuse of notation. We would like to study the translation by f in $(\mathbb{T}^d)^I$, but as it lacks a nice topology, we consider a compactification of the group $((\mathbb{T}^d)^I, +)$, which will be a monoid, as follows.

Let $\text{Prob}_{\mathcal{L}}$ be the space of probability measures on $I \times \mathbb{T}^d$ which project to \mathcal{L} on the first factor. To an angle map $f \in (\mathbb{T}^d)^I$, we can associate the probability measure \mathcal{D}_f on $I \times \mathbb{T}^d$, supported by its graph, which is the pushforward of \mathcal{L} by the map $x \in I \mapsto (x, f(x)) \in I \times \mathbb{T}^d$. This defines an embedding of $(\mathbb{T}^d)^I$ into $\text{Prob}_{\mathcal{L}}$. It is not hard to see that the group law $+$ on $(\mathbb{T}^d)^I$ correspond to a fiber-wise convolution product $*$ on $\text{Prob}_{\mathcal{L}}$, which turns $(\text{Prob}_{\mathcal{L}}, *)$ into an abelian monoid, with neutral \mathcal{D}_0 , where $0 : I \rightarrow \mathbb{T}^d$ is the zero map.

The space $\text{Prob}_{\mathcal{L}}$ is equipped naturally with a weak- $*$ topology, for which it is a compact metric space. A tricky fact is that the convolution product $(\mu, \nu) \mapsto \mu * \nu$ is *not* continuous of the two variables, but is of each variable separately. A more detailed description of these objects, and explanations of the implied claims, are given in Section 2.

There is a particular element in $\text{Prob}_{\mathcal{L}}$, the measure $\lambda = \mathcal{L} \otimes \text{Haar}_{\mathbb{T}^d}$. It satisfies the relation: $\forall \mu \in \text{Prob}_{\mathcal{L}}, \mu * \lambda = \lambda$.

We are interested in studying the dynamics of convolution $\mathcal{D}_f *$ on the monoid $\text{Prob}_{\mathcal{L}}$. Unsurprisingly, we now need a kind of irrationality condition.

DEFINITION. — *The angle map $f : I \rightarrow \mathbb{T}^d$ is said to be asynchronous if the image measure $f_*\mathcal{L}$ gives zero mass to any translate of any proper closed subgroup of \mathbb{T}^d .*

See Lemma 3.1 for equivalent definitions of asynchronicity. Intuitively, for $d = 1$, this means that one looks at an action by rotation on a family of circles indexed by $x \in I$, by angles $f(x)$, which are different from one another if picked randomly following the probability \mathcal{L} . For $d \geq 1$, it means that for almost every couple (x, y) , $f(x)$ and $f(y)$ do not belong to the same coset modulo any closed, strict subgroup of \mathbb{T}^d .

THEOREM 1.1. — *The following are equivalent.*

- (1) *The angle map $f : I \rightarrow \mathbb{T}^d$ is asynchronous.*

(2) The closure $\overline{\{\mathcal{D}_{nf}\}_{n \in \mathbb{Z}}}$ contains λ .

(3) The convolution action of \mathcal{D}_f on $\text{Prob}_{\mathcal{L}}$ is uniquely ergodic.

If these are true, the only invariant probability measure on $\text{Prob}_{\mathcal{L}}$ under $\mu \mapsto \mathcal{D}_f * \mu$ is the Dirac measure δ_λ .

The fact that the invariant measure is a Dirac measure implies (see Proposition 3.4) that there exists a subset of the integers $E \subset \mathbb{Z}$, of natural density 1, such that for any $\mu \in \text{Prob}_{\mathcal{L}}$,

$$\lim_{n \rightarrow \pm\infty, n \in E} \mathcal{D}_{nf} * \mu = \lambda.$$

The question whether λ is an attracting point of the dynamic, that is if

$$(1.1) \quad \lim_{n \rightarrow \pm\infty} \mathcal{D}_{nf} * \mu = \lambda,$$

or if this fails along some subsequence of zero density, is more delicate, and its answer depends on f . Note (1.1) is in fact equivalent to

$$\lim_{n \rightarrow \pm\infty} \mathcal{D}_{nf} = \lambda,$$

see Section 3.2.

If $I = [0, 1]$ equipped with Lebesgue measure, $d = 1$, and f is a C^2 map with non-vanishing derivative, then λ is an attracting point (Proposition 3.2), and there is no exceptional subsequence. The C^2 regularity condition is not optimal, as Thomine obtained similar results for C^1 maps [14].

However, for an angle map f which is only measurable, the convolution action of \mathcal{D}_f might behave more like an intermittent map with the neutral fixed point λ . An example of this phenomenon is the following. Again, let $I = [0, 1]$ endowed with the Lebesgue measure \mathcal{L} . Let ν be the (probability) Hausdorff measure of dimension $\log 2 / \log 3$ on the usual Cantor set C , viewed as a subset of \mathbb{T}^1 by identifying 0 and 1. Define $f : [0, 1] \rightarrow \mathbb{T}^1$, by

$$f(x) = \inf\{t \geq 0 : \nu(0, t) \geq x\} \bmod 1.$$

Then $f_*\mathcal{L} = \nu$, and $f(I) \subset C$. Alternatively, f can be defined as the reciprocal, outside of dyadic rationals, of the usual devil's staircase, modulo 1. Since ν does not have any atom, f is an asynchronous angle map. We claim that the sequence $(\mathcal{D}_{3^k f})_{k \geq 1}$ does not intersect a fixed neighborhood of λ . Indeed, since C is invariant by the multiplication $\times 3$ on the circle, the graph of $3^k f$ is contained in $I \times C$, so the measure $\mathcal{D}_{3^k f}$ is supported on $I \times C$, a proper compact subset of $I \times \mathbb{T}^d$. This forbids $\mathcal{D}_{3^k f}$ to be close to λ . Still, by Theorem 1.1, subsequences like $(\mathcal{D}_{3^k f})_{k \geq 0}$ are scarce, as the points $(\mathcal{D}_{nf})_{n \in \mathbb{Z}}$ tend to λ for a subset of \mathbb{Z} of density one.

1.3. Main example, and a related ergodicity result

As hinted in the motivational paragraph, asynchronous rotations occur naturally in the context of homogeneous dynamics on torus bundle.

More precisely, let \mathbb{G}_0 be a connected, semisimple algebraic linear group defined over \mathbb{Q} , $G_0 = \mathbb{G}_0(\mathbb{R})$ the group of its \mathbb{R} -points and $\Gamma_0 = \mathbb{G}_0(\mathbb{Z})$ its integer points. By the Borel–Harish-Chandra Theorem, Γ_0 is a lattice in G_0 . We will consider invariant measures on G_0/Γ_0 under some elements $a \in G_0$, under the following assumptions.

DEFINITION. — *An element $a \in G_0 - \{e\}$ is said triangularizable with positive eigenvalues if for every finite dimensional representation α of \mathbb{G}_0 defined over \mathbb{Q} , $\alpha(a)$ has only real, positive eigenvalues.*

It is the case, for example, when G_0 is the real split form of \mathbb{G}_0 , meaning the real rank equals the complex rank, and if a is the exponential of a non-zero element of a Cartan subalgebra. It also happens when a is unipotent, but this case is less interesting for our purpose, since by Ratner’s Theory (see e.g. [9, 11]), a -invariant ergodic measures (or their extensions to semidirect product as considered below) are classified and well understood in this case. Existence of such an element a implicitly rules out the case where G_0 is the real compact form of \mathbb{G}_0 .

DEFINITION. — *Let μ be a probability measure on G_0/Γ_0 , invariant and ergodic under the action of a . Such a measure is said to be non-concentrated if for every $H \subset G_0$ closed algebraic, strict subgroup containing a , and every $x \in G_0/\Gamma_0$ such that $Hx\Gamma_0$ is closed, then $\mu(Hx\Gamma_0) < 1$.*

We now consider a fiber bundle over the probability space $(G_0/\Gamma_0, \mu)$, whose fibers are tori.

Let $\rho : \mathbb{G}_0 \rightarrow \mathrm{GL}(V)$ a representation defined over \mathbb{Q} on a finite-dimensional space $V = \mathbb{R}^d$ endowed with the \mathbb{Z} -structure \mathbb{Z}^d . We will always assume that $d \geq 2$, and that ρ is irreducible over \mathbb{Q} . The semidirect product $G = G_0 \times_{\rho} V$ is endowed with the group law

$$\forall (g, v) \in G, \forall (h, w) \in G, (g, v)(h, w) = (gh, v + \rho(g)w).$$

Up to replacing Γ_0 with a subgroup of finite index in a way such that $\rho(\Gamma) \subset \mathrm{GL}(d, \mathbb{Z})$, the set $\Gamma = \Gamma_0 \times_{\rho} \mathbb{Z}^d$ is a subgroup of G , and the map

$$\pi : G/\Gamma \longrightarrow G_0/\Gamma_0,$$

is a torus bundle. Indeed, the lattice of $\{e\} \times \mathbb{R}^d$ (for the action of multiplication on the left) stabilizing a point $(x, v)\Gamma$ is precisely $\{e\} \times \rho(x)\mathbb{Z}^d$,

thus the fiber of π over $x\Gamma_0$ is the torus $\mathbb{R}^d/\rho(x)\mathbb{Z}^d$. It will be convenient to have measurable coordinates where this fiber bundle is a direct product.

Let $I \subset G_0$ be a measurable fundamental domain for the action of Γ_0 , and put \mathcal{L} the restriction to I of the Γ_0 -invariant lift of μ . As previously, we denote by \mathbb{T}^d the d -dimensional torus $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$. The map

$$\begin{aligned} I \times \mathbb{T}^d &\longrightarrow G/\Gamma, \\ (x, \bar{v})_{I \times \mathbb{T}^d} &\longmapsto (x, \rho(x)v)\Gamma, \end{aligned}$$

is a measurable bijection, that we will use as an identification between $I \times \mathbb{T}^d$ and G/Γ , the subscript $I \times \mathbb{T}^d$ indicating the coordinates we are using. Likewise, we will identify G_0/Γ_0 with I and μ with \mathcal{L} .

For $\beta \in V$, the action of (e, β) by multiplication on G/Γ on the left, can be read in the $I \times \mathbb{T}^d$ coordinates as the map

$$(x, \bar{v})_{I \times \mathbb{T}^d} \longmapsto (x, \bar{v} + \rho(x)^{-1}\beta)_{I \times \mathbb{T}^d},$$

i.e. it is a rotation by an angle map $f_\beta : I \rightarrow \mathbb{T}^d$, with

$$f_\beta(x) = \rho(x)^{-1}\beta \bmod \mathbb{Z}^d.$$

We prove:

THEOREM 1.2. — *Assume that $a \in G_0$ is triangularizable with positive eigenvalues, that μ is an a -invariant, non-concentrated, ergodic probability on G_0/Γ_0 , that ρ is irreducible over \mathbb{Q} , and $\dim(V) > 1$. Assume also that $\beta \in V - \{0\}$ is an eigenvector for $\rho(a)$. Then the angle map*

$$\begin{aligned} f_\beta : I &\longrightarrow \mathbb{T}^d \\ x &\longmapsto \rho(x)^{-1}\beta \bmod \mathbb{Z}^d, \end{aligned}$$

is asynchronous.

Using the identification of G/Γ with $I \times \mathbb{T}^d$, we still denote by λ the measure on G/Γ such that $\pi_*\lambda = \mu$, whose disintegration along each fiber of μ are the Haar measures on each tori.

Direct application of Theorem 1.1 gives:

COROLLARY 1.3. — *We assume the same hypotheses as Theorem 1.2. Let Prob_μ be the set of probabilities on G/Γ projecting onto μ . The action on Prob_μ induced by the left multiplication by (e, β) on G/Γ is uniquely ergodic, with invariant measure δ_λ .*

Now choose any $\alpha \in V$. The action of left multiplication by (a, α) on G/Γ admits the action of a on G_0/Γ_0 as a factor. A natural question is if the measure λ , which is invariant, is ergodic with respect to this action.

If β is any eigenvector of $\rho(a)$, multiplication by (e, β) is in some sense moving in some part of the stable, unstable or neutral direction (depending on the eigenvalue) of the action of (a, α) . Theorem 1.2 and Hopf's argument allows us to prove the following ergodicity result:

THEOREM 1.4. — *Assume that $a \in G_0$ is triangularizable with positive eigenvalues, that μ is an a -invariant, non-concentrated, ergodic probability on G_0/Γ_0 , and that ρ is irreducible over \mathbb{Q} , of dimension > 1 . Choose $\alpha \in V$, then the action by left multiplication by (a, α) on G/Γ is ergodic with respect to the invariant measure λ .*

If we assume moreover that $\rho(a)$ is not unipotent, then the action of (a, α) on $(G/\Gamma, \lambda)$ is weakly mixing if and only if the action of a on $(G_0/\Gamma_0, \mu)$ is weakly mixing, and the same property holds for strong mixing.

1.4. Plan of the paper

The first part of the paper deals with abstract ergodic theory. In Section 2, we collect some facts about the topology of $\text{Prob}_{\mathcal{L}}$.

In Section 3, after dealing with the toy example where f is a C^2 map, we prove Theorem 1.1, mainly using Fourier series.

In the second part of the paper, we turn to a setting in homogeneous dynamics; the reader new to this might benefit from the survey [9].

In Section 4, we prove that in the algebraic setting, the smallest algebraic subgroup of G_0 containing the elements $\gamma \in \Gamma_0$ induced by Poincaré recurrence of the a -action, is G_0 itself. This result (Theorem 4.1) is the main ingredient of the proof of Theorem 1.2.

In Section 5, we prove the asynchronicity of the rotation obtained by the construction in homogeneous dynamics (Theorem 1.2).

In Section 6, we prove the ergodicity of the extension of the action of a on G_0/Γ_0 (Theorem 1.4), and related mixing properties. This mainly relies on the following easy observation: if an angle map $f : I \rightarrow \mathbb{T}^d$ is asynchronous, then for almost every x , the action of translation by $f(x)$ on the torus \mathbb{T}^d is (uniquely) ergodic.

1.5. Acknowledgements

I would like to thank to Jean-Pierre Conze, Serge Cantat, Sébastien Gouëzel, Barbara Schapira, Damien Thomine and the referee for their feedback and comments on the subject.

2. The space $\text{Prob}_{\mathcal{L}}$

We recall briefly the setting. Let $(I, \mathcal{B}(I), \mathcal{L})$ be a standard probability space without atoms, \mathbb{T}^d be the d -dimensional torus. The group $((\mathbb{T}^d)^I, +)$ is the set of equivalence classes of measurable maps $f : I \rightarrow \mathbb{T}^d$ where two maps are identified if they agree on a set of full \mathcal{L} -measure, endowed with the usual addition. Such a (class of) map is called an *angle map*. The set $\text{Prob}_{\mathcal{L}}$ is the set of probability measures on $I \times \mathbb{T}^d$ which projects to \mathcal{L} on I . Our goal here is first to define a topology on $\text{Prob}_{\mathcal{L}}$, and then a convolution that extends the group law of addition of angle maps.

2.1. Topology of $\text{Prob}_{\mathcal{L}}$

We recall that $(I, \mathcal{B}(I))$ be a standard measurable space means that I can be endowed with a complete, separable distance d_I , such that $\mathcal{B}(I)$ is the σ -algebra of its Borel sets. The facts about standard probability spaces we will use are summarized in [1, Chapter 1.1]. Choosing such a distance on I defines a topology on the space of probability measures on $I \times \mathbb{T}^d$, and hence on $\text{Prob}_{\mathcal{L}}$.

Although the weak- $*$ topology of the space of measures on $I \times \mathbb{T}^d$ depends in a strong way on the choice of topology on I , it turns out that:

LEMMA 2.1. — *The topology induced on $\text{Prob}_{\mathcal{L}}$ does not depend on the choice of topology on I .*

Proof. — Let I_1, I_2 two complete, separable metric space endowed with probabilities \mathcal{L}_i , with a map $\varphi : I_1 \rightarrow I_2$ an isomorphism such that $\varphi_*\mathcal{L}_1 = \mathcal{L}_2$. The topologies induced on measures on $I_i \times \mathbb{T}^d$ are generated by the open sets:

$$\mathcal{U}^i(F, \delta, \mu) = \left\{ \nu : \left| \int_{I_i \times \mathbb{T}^d} F d\mu - \int_{I_i \times \mathbb{T}^d} F d\nu \right| < \delta \right\},$$

where $F : I_i \times \mathbb{T}^d \rightarrow \mathbb{C}$ is continuous with compact support for the relevant topology on $I_i \times \mathbb{T}^d$.

Denote by $\tilde{\varphi}$ the map $I_1 \times \mathbb{T}^d \rightarrow I_2 \times \mathbb{T}^d$, $\tilde{\varphi}(x, y) = (\varphi(x), y)$. To show that $\tilde{\varphi}_* : \text{Prob}_{\mathcal{L}_1} \rightarrow \text{Prob}_{\mathcal{L}_2}$ is a homeomorphism, it is sufficient by symmetry to show its continuity.

We fix a neighborhood $\mathcal{U}^2(F, \epsilon, \tilde{\varphi}_*\mu)$, and wish to show that its preimage contains some neighborhood of the initial point $\mathcal{U}^1(G, \delta, \tilde{\varphi}_*\mu)$, for some G, δ .

The map φ from I_1 to I_2 is measurable. By Lusin's Theorem, for every $\delta > 0$, there is a compact set $J \subset I_1$, such that $\mathcal{L}_1(J) > 1 - \delta$, on which φ is continuous. By Tietze–Urysohn's Theorem, there exists a continuous function $G_\delta : I_1 \times \mathbb{T}^d \rightarrow \mathbb{C}$ which extends the continuous map $F \circ \tilde{\varphi} : J \times \mathbb{T}^d \rightarrow \mathbb{C}$. Moreover, since F is bounded, G_δ can be chosen such that $\|G_\delta\|_\infty = \|F\|_\infty$. If $\nu \in \text{Prob}_{\mathcal{L}_1}$,

$$\left| \int_{I_2 \times \mathbb{T}^d} F d\tilde{\varphi}_* \mu - \int_{I_2 \times \mathbb{T}^d} F d\tilde{\varphi}_* \nu \right| \leq 2\delta \|F\|_\infty + \left| \int_{J \times \mathbb{T}^d} G_\delta d\mu - \int_{J \times \mathbb{T}^d} G_\delta d\nu \right|,$$

because $\mu((I_2 - \varphi J) \times \mathbb{T}^d) = \mathcal{L}_2(I_2 - \varphi J) = \mathcal{L}_1(I_1 - J) < \delta$ and the same holds for ν . If we choose $\delta > 0$ such that $2\delta \|F\|_\infty < \epsilon/2$, we have:

$$\left| \int_{I_2 \times \mathbb{T}^d} F d\tilde{\varphi}_* \mu - \int_{I_2 \times \mathbb{T}^d} F d\tilde{\varphi}_* \nu \right| \leq \epsilon/2 + \left| \int_{I_1 \times \mathbb{T}^d} G_\delta d\mu - \int_{I_1 \times \mathbb{T}^d} G_\delta d\nu \right|,$$

and therefore, provided that $\delta < \epsilon/2$,

$$\text{Prob}_{\mathcal{L}_1} \cap \mathcal{U}^1(G_\delta, \delta, \mu) \subset \tilde{\varphi}_*^{-1}(\mathcal{U}^2(F, \epsilon, \tilde{\varphi}_* \mu) \cap \text{Prob}_{\mathcal{L}_2}).$$

As any neighborhood of $\varphi_* \mu$ contains finite intersections of sets of the form $\mathcal{U}^2(F, \epsilon, \varphi_* \mu)$, this implies that $\tilde{\varphi}_* : \text{Prob}_{\mathcal{L}_1} \rightarrow \text{Prob}_{\mathcal{L}_2}$ is continuous, as required. \square

A corollary of this discussion is that we can assume for example that $I = [0, 1]$ and \mathcal{L} is the Lebesgue measure on this interval, endowed with its usual topology. Since in this case, the set of probability measures on $I \times \mathbb{T}^d$ is a compact, separable metric space, it follows that $\text{Prob}_{\mathcal{L}}$ is also compact, separable and metric.

2.2. Graphs, Disintegration and convolutions

For an angle map $g : I \rightarrow \mathbb{T}^d$, we define the *graph measure* \mathcal{D}_g , of g as the direct image of \mathcal{L} by the map $x \in I \mapsto (x, g(x)) \in I \times \mathbb{T}^d$. Two angle maps $I \rightarrow \mathbb{T}^d$ define the same graph measure if and only if they are equal (\mathcal{L} -almost everywhere).

Let \mathcal{G} be the set of graph measures, this is a subset of $\text{Prob}_{\mathcal{L}}$.

Any $\mu \in \text{Prob}_{\mathcal{L}}$ can be disintegrated as a family of measures $(\mu^x)_{x \in I}$, such that for any continuous test-function with compact support $F : I \times \mathbb{T}^d \rightarrow \mathbb{C}$,

$$\mu(F) = \int_I \left(\int_{\mathbb{T}^d} F(x, y) d\mu^x(y) \right) d\mathcal{L}(x).$$

Moreover, the map $x \mapsto \mu^x$ is measurable, and uniquely defined modulo zero sets. See e.g. [8, Theorem 5.8].

For μ_1, μ_2 two measures in $\text{Prob}_{\mathcal{L}}$, we define the fiberwise convolution product of μ_1, μ_2 by

$$\mu_1 * \mu_2(F) = \int_I \left(\int_{(\mathbb{T}^d)^2} F(x, y + z) d\mu_1^x(y) d\mu_2^x(z) \right) d\mathcal{L}(x),$$

where $F : I \times \mathbb{T}^d \rightarrow \mathbb{C}$ is a continuous test-function with compact support. Equivalently, $(\mu * \nu)^x$ is the usual convolution product of μ^x and ν^x .

The following Lemma, whose proof is left to the reader, summarizes elementary properties of this fiberwise convolution product.

LEMMA 2.2. — *The following holds.*

- (1) $\forall (f, g) \in ((\mathbb{T}^d)^I)^2$, $\mathcal{D}_{f+g} = \mathcal{D}_f * \mathcal{D}_g$.
- (2) \mathcal{D}_0 is the neutral element of the commutative monoid $(\text{Prob}_{\mathcal{L}}, *)$, where $0 : I \rightarrow \mathbb{T}^d$ is the angle map almost everywhere zero.
- (3) The set of invertible elements for $*$ is \mathcal{G} , the set of graph measures.

Remark that, if $f : [0, 1] \rightarrow \mathbb{T}^1$, $f(x) = x \bmod 1$, then one can check by hand (or see e.g. Proposition 3.2) that \mathcal{D}_{nf} tends to λ as $n \rightarrow \pm\infty$, but

$$\mathcal{D}_{nf} * \mathcal{D}_{-nf} = \mathcal{D}_0 \neq \lambda * \lambda,$$

so the fiberwise convolution product is not continuous. However, we have:

LEMMA 2.3. — *For any $\nu \in \text{Prob}_{\mathcal{L}}$, the convolution map*

$$\begin{aligned} * \nu : \text{Prob}_{\mathcal{L}} &\longrightarrow \text{Prob}_{\mathcal{L}}, \\ \mu &\longmapsto \mu * \nu, \end{aligned}$$

is continuous.

Proof. — It is sufficient to check that the preimage by $*\nu$ of any set of the form $\mathcal{U}(F, \epsilon, \mu * \nu)$, for any F, ϵ, μ , contains a set of the form $\mathcal{U}(G, \delta, \mu)$ for some G and some $\delta > 0$. Let $\mathcal{U}(F, \epsilon, \mu * \nu)$ be such a neighborhood of $\mu * \nu$, and let δ such that $\delta(4\|F\|_{\infty} + 1) < 1$. As the map $x \mapsto \nu^x$ is measurable, again by Lusin's Theorem, it is continuous on a set J of measure $1 - \delta$. Define

$$H(x, y) = \int_{\mathbb{T}^d} F(x, y + z) d\nu^x(z).$$

It follows from the continuity of F and the continuity of $x \mapsto \nu^x$ that this is a continuous map on $J \times \mathbb{T}^d$, and moreover bounded by $\|F\|_{\infty}$. Thus it can be extended to a bounded continuous map, say G , on $I \times \mathbb{T}^d$, still

bounded by $\|F\|_\infty$. Notice that for any $\eta \in \text{Prob}_{\mathcal{L}}$,

$$\begin{aligned} \int_{J \times \mathbb{T}^d} Fd(\eta * \nu) &= \int_J \int_{\mathbb{T}^d} \left(\int_{\mathbb{T}^d} F(x, y + z) d\nu^x(z) \right) d\eta^x(y) d\mathcal{L}(x) \\ &= \int_{J \times \mathbb{T}^d} G d\eta. \end{aligned}$$

Let $\eta \in \mathcal{U}(G, \delta, \mu)$, then

$$\begin{aligned} \left| \int_{I \times \mathbb{T}^d} Fd(\eta * \nu) - \int_{I \times \mathbb{T}^d} Fd(\mu * \nu) \right| &\leq 2\delta \|F\|_\infty + \left| \int_{J \times \mathbb{T}^d} Fd(\eta * \nu) - \int_{J \times \mathbb{T}^d} Fd(\mu * \nu) \right| \\ &\leq 2\delta \|F\|_\infty + \left| \int_{J \times \mathbb{T}^d} G d\eta - \int_{J \times \mathbb{T}^d} G d\mu \right| \\ &\leq 4\delta \|F\|_\infty + \delta. \end{aligned}$$

By the choice of δ , this implies that $*\nu(\mathcal{U}(G, \delta, \mu)) \subset \mathcal{U}(F, \epsilon, \mu * \nu)$, as announced. \square

3. Asynchronous maps

In this section, we begin by characterizing asynchronous angle maps in terms of characters, then prove that for a sufficiently regular angle map f , the point $\lambda \in \text{Prob}_{\mathcal{L}}$ is attracting, using Fourier series. Then we prove Theorem 1.1, the main tool being a Cesaro-average of the sequence of square of Fourier coefficients of the measures \mathcal{D}_{nf} . Lastly, we prove the claims of the introduction about the existence of a set of integers of density one such that $\mathcal{D}_{nf}*$ attracts all points to λ for any sequence along that set.

3.1. Equivalent definitions of asynchronicity

Recall that a *character* on a torus \mathbb{T}^d is a continuous group morphism $\chi : (\mathbb{T}^d, +) \rightarrow (\mathbb{S}^1, \times)$; they are of the form $\chi(x) = e^{2i\pi \langle x, m \rangle}$, where $m \in \mathbb{Z}^d$. The group of characters is denoted by $\widehat{\mathbb{T}^d}$, and is isomorphic to \mathbb{Z}^d . The Fourier-Stieltjes coefficients of a measure μ on \mathbb{T}^d are the quantities $\widehat{\mu}(\chi) = \int_{\mathbb{T}^d} \chi(x) d\mu(x)$, where χ is a character.

LEMMA 3.1. — *Let $f : I \rightarrow \mathbb{T}^d$ be a measurable map. The following properties are equivalent:*

- (1) *The $f_*\mathcal{L}$ -measure of any translate of any proper closed subgroup of \mathbb{T}^d is zero.*
- (2) *For any non-trivial character $\chi \in \widehat{\mathbb{T}^d}$, $(\chi \circ f)_*\mathcal{L}$ is a measure without atoms on the circle \mathbb{S}^1 .*
- (3) *For $\mathcal{L} \otimes \mathcal{L}$ -almost every $(x, y) \in I^2$, and any non-trivial character $\chi \in \widehat{\mathbb{T}^d}$, $\chi(f(x) - f(y)) \neq 1$.*
- (4) *For $\mathcal{L} \otimes \mathcal{L}$ -almost every $(x, y) \in I^2$, $f(x) - f(y)$ generates a dense subgroup of \mathbb{T}^d .*

If one (and hence every) of these conditions is satisfied, the angle map f is said to be asynchronous.

Proof. — Assume f satisfies (1). Let $\chi : \mathbb{T}^d \rightarrow \mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$ be a non-trivial character, and $y \in \mathbb{S}^1$. Then

$$(\chi \circ f)_*\mathcal{L}(\{y\}) = f_*\mathcal{L}(\chi^{-1}\{y\}),$$

and $\chi^{-1}\{y\}$ is a coset of the proper subgroup $\ker \chi$, so has $f_*\mathcal{L}$ -measure zero. So (1) implies (2).

Assume now that f satisfies (2). Let χ be a non-trivial character. For any $x \in I$, the set

$$\{y \in I : \chi(f(x)) = \chi(f(y))\} = (\chi \circ f)^{-1}(\{\chi(f(x))\}),$$

has \mathcal{L} -measure zero by the assumption (2). So, by Fubini's Theorem, the set

$$E_\chi = \{(x, y) \in I^2 : \chi(f(x) - f(y)) \neq 1\},$$

has full measure for any $\chi \neq 1$. As $\widehat{\mathbb{T}^d} \simeq \mathbb{Z}^d$ is countable, $\bigcap_{\chi \neq 1} E_\chi$ still has full measure, proving that (2) implies (3).

We now assume (3). Let $(x, y) \in I^2$ be in the full measure set described by (3), that is for all $\chi \in \widehat{\mathbb{T}^d} - \{1\}$, $\chi(f(x)) \neq \chi(f(y))$. Let H be the closure of the subgroup of \mathbb{T}^d generated by $f(x) - f(y)$, we wish to prove that $H = \mathbb{T}^d$. If not, the quotient \mathbb{T}^d/H is a non-trivial compact, connected abelian Lie group, hence a torus. The composition of any non-trivial character of \mathbb{T}^d/H with the canonical map $\mathbb{T}^d \rightarrow \mathbb{T}^d/H$ is a non-trivial character χ of \mathbb{T}^d , for which $\chi(H) \equiv 1$. This contradicts $\chi(f(x) - f(y)) \neq 1$, so $H = \mathbb{T}^d$, proving that (3) implies (4).

Let us now assume that (1) is not satisfied, that is, for some closed, proper subgroup $H \subset \mathbb{T}^d$, and some $k \in \mathbb{T}^d$, $\mathcal{L}(f^{-1}(k + H)) > 0$. Let $(x, y) \in (f^{-1}(k + H))^2$ (this is a set of positive $\mathcal{L} \otimes \mathcal{L}$ -measure in I^2),

then $f(x) - f(y) \in H$, so the closure of the group generated by $f(x) - f(y)$ is a subgroup of H . This shows that (4) is not satisfied either. By contraposition, this finishes the proof. \square

3.2. A simple example

We claimed in the introduction that

$$(3.1) \quad \forall \mu \in \text{Prob}_{\mathcal{L}}, \lim_{n \rightarrow \pm\infty} \mathcal{D}_{nf} * \mu = \lambda,$$

is equivalent to

$$(3.2) \quad \lim_{n \rightarrow \pm\infty} \mathcal{D}_{nf} = \lambda.$$

Indeed, if one assumes (3.1), one gets (3.2) by choosing for μ the neutral element \mathcal{D}_0 . Conversely, if one assumes (3.2), continuity of the convolution with a fixed μ (Lemma 2.3) implies (3.1).

As stated in the introduction, if f has enough regularity properties, it turns out that λ is the limit point of the dynamic of \mathcal{D}_f* on $\text{Prob}_{\mathcal{L}}$. This result will not be used in the sequel.

PROPOSITION 3.2. — *Assume $f : [0, 1] \rightarrow \mathbb{R}/\mathbb{Z}$ is a C^2 -angle map, such that f' does not vanish. Then for all $\mu \in \text{Prob}_{\mathcal{L}}$, $\mathcal{D}_{kf}*\mu \rightarrow \lambda$ when $k \rightarrow \pm\infty$.*

Proof. — By continuity of $*\mu$, it is sufficient to check that \mathcal{D}_{kf} tends to λ as k tends to infinity, that is, for any continuous $F : [0, 1] \times \mathbb{T}^1 \rightarrow \mathbb{C}$,

$$(3.3) \quad \lim_{k \rightarrow \pm\infty} \int_{[0,1] \times \mathbb{T}^1} F \, d\mathcal{D}_{kf} = \int_{[0,1] \times \mathbb{T}^1} F(x, y) \, dx dy.$$

By Stone–Weierstrass’s Theorem, the set of trigonometric polynomials is dense in $C^0([0, 1] \times \mathbb{T}^1)$. Thus we only need to prove (3.3) for $F(x, y) = e_{n,m}(x, y) = e^{2i\pi(nx+my)}$, where n, m are integers. To do so, we compute the corresponding Fourier–Stieltjes coefficients

$$\widehat{\mathcal{D}_{kf}}(n, m) = \int_0^1 e^{2i\pi(nx+mkf(x))} dx.$$

If $m = 0$, this coefficient is 1 or 0, depending on whether $n = 0$ or not, so (3.3) is satisfied by $e_{n,m}$ in this case. If $m \neq 0$, we can write

$$\widehat{\mathcal{D}_{kf}}(n, m) = \int_0^1 \frac{e^{2i\pi nx}}{2i\pi mk f'(x)} \frac{\partial}{\partial x} \left(e^{2i\pi mkf(x)} \right) dx,$$

and integration by parts gives

$$k \widehat{\mathcal{D}}_{kf}(n, m) = \left[\frac{e^{2i\pi(nx+mkf(x))}}{2i\pi m f'(x)} \right]_0^1 - \int_0^1 \frac{(2i\pi n f'(x) - f''(x))}{2i\pi m (f'(x))^2} e^{2i\pi(nx+mkf(x))} dx.$$

Notice that the right-hand side is bounded independently of k when n, m are fixed. So $\widehat{\mathcal{D}}_{kf}(n, m) = O(\frac{1}{k})$, and (3.3) is satisfied by $e_{n,m}$ \square

3.3. Proof of Theorem 1.1, (1) \Leftrightarrow (2)

We now show the equivalence

$$f \text{ is asynchronous} \iff \lambda \in \overline{(D_{nf})_{n \in \mathbb{Z}}}.$$

We first show the implication \Leftarrow , and assume that there exists a sequence $(n_i)_{i \geq 0}$ such that $(\mathcal{D}_{n_i f})$ converges to λ . We will use characterisation (2) of asynchronicity given by Lemma 3.1. So let χ be any non-trivial character of \mathbb{T}^d , we wish to show that $(\chi \circ f)_* \mathcal{L}$ is atom-free. Assume by contradiction that $(\chi \circ f)_* \mathcal{L}$ has an atom. In this case, there would be a set $E \subset I$ of positive measure on which $\chi \circ f$ is a constant, say c . Thus $\chi \circ (n_i f)$ is a constant on E , namely c^{n_i} , and $(\chi \circ (n_i f))_* \mathcal{L}$ will have an atom of mass $\mathcal{L}(E)$. Note that

$$(\chi \circ (n_i f))_* \mathcal{L} = (\chi \circ \pi_{\mathbb{T}^d})_* \mathcal{D}_{n_i f}.$$

But $(\chi \circ \pi_{\mathbb{T}^d})_* \mathcal{D}_{n_i f}$ tends to $(\chi \circ \pi_{\mathbb{T}^d})_* \lambda$, namely the Lebesgue measure on \mathbb{S}^1 , which cannot be a limit of measures having an atom of fixed mass. This is a contradiction.

We now turn to the \Rightarrow implication, and assume that the angle map f is asynchronous. We may, and will, assume that $I = \mathbb{T}^1$ endowed with its Haar probability measure \mathcal{L} . The space $I \times \mathbb{T}^d$ is then a $(d+1)$ -dimensional torus. The main point will be that the square of the modulus of Fourier coefficients of the measures \mathcal{D}_{nf} converges to zero in Cesaro average, and that will be sufficient to find a n such that D_{nf} is close to λ .

The Haar measure λ of $I \times \mathbb{T}^d = \mathbb{T}^{d+1}$ is in the closure of $\{\mathcal{D}_{nf}\}_{n \in \mathbb{Z}}$ if and only if for any given finite family F_1, \dots, F_p of continuous functions on \mathbb{T}^{d+1} , and any $\epsilon > 0$, we can find some n such that for any $i \in \{1, \dots, p\}$,

$$(3.4) \quad \left| \int_{I \times \mathbb{T}^d} F_i d\mathcal{D}_{nf} - \int_{I \times \mathbb{T}^d} F_i d\lambda \right| < \epsilon.$$

By Stone–Weierstrass’s Theorem, each F_i can be uniformly approached by a linear combination of characters: there exists a finite family of characters $E \subset \widehat{\mathbb{T}^{d+1}}$ and coefficients $a_{i,\chi} \in \mathbb{C}$ such that for all $i \in \{1, \dots, p\}$,

$$\left\| F_i - \sum_{\chi \in E} a_{i,\chi} \chi \right\|_{\infty} < \epsilon/4.$$

Let $A = \max |a_{i,\chi}|$. To prove (3.4), it is thus sufficient to find a common n such that for any $\chi \in E$, we have

$$\left| \int_{I \times \mathbb{T}^d} \chi \, d\mathcal{D}_{nf} - \int_{I \times \mathbb{T}^d} \chi \, d\lambda \right| < \frac{\epsilon}{2A}.$$

When χ is the trivial character, the above inequality is true for all n ; for other characters $\chi \in E - \{1\}$, this condition means that the Fourier–Stieltjes coefficient is small:

$$\left| \widehat{\mathcal{D}_{nf}}(\chi) \right| < \frac{\epsilon}{2A}.$$

We first notice:

LEMMA 3.3. — *For any nontrivial character χ of $I \times \mathbb{T}^d = \mathbb{T}^{d+1}$, we have*

$$\lim_{N \rightarrow +\infty} \frac{1}{N} \sum_{n=0}^{N-1} \left| \widehat{\mathcal{D}_{nf}}(\chi) \right|^2 = 0.$$

Proof. — For an integer $k \in \mathbb{Z}$, let e_k be the character of \mathbb{T}^1 , $e_k(x) = e^{2i\pi kx}$. Any character χ of $I \times \mathbb{T}^d = \mathbb{T}^{d+1}$ can be written uniquely as a product $\chi(x, y) = e_k(x)\psi(y)$, for some $k \in \mathbb{Z}$, some $\psi \in \widehat{\mathbb{T}^d}$ and every $x \in I$, $y \in \mathbb{T}^d$. We have

$$\frac{1}{N} \sum_{n=0}^{N-1} \left| \widehat{\mathcal{D}_{nf}}(\chi) \right|^2 = \int_{I^2} e^{2i\pi k(x-x')} \left(\frac{1}{N} \sum_{n=0}^{N-1} \psi(f(x) - f(x'))^n \right) d\mathcal{L}^2(x, x').$$

If $\psi = 1$, then $k \neq 0$ since $\chi \neq 1$. In this case, we have

$$\frac{1}{N} \sum_{n=0}^{N-1} \left| \widehat{\mathcal{D}_{nf}}(\chi) \right|^2 = 0,$$

so the statement is trivial.

If $\psi \neq 1$, then since f is asynchronous, by Lemma 3.1 (3), $\psi(f(x) - f(x')) \neq 1$ for almost every (x, x') , so

$$\lim_{N \rightarrow +\infty} \frac{1}{N} \sum_{n=0}^{N-1} \psi(f(x) - f(x'))^n = 0.$$

Therefore, Lebesgue's dominated convergence Theorem applies and we obtain the desired result. \square

By the previous Lemma, since the set E is finite, we have

$$\lim_{N \rightarrow +\infty} \frac{1}{N} \sum_{n=0}^{N-1} \left(\sum_{\chi \in E - \{1\}} \left| \widehat{\mathcal{D}_{nf}}(\chi) \right|^2 \right) = 0,$$

so there exists $n \geq 0$ such that for all $\chi \in E - \{1\}$, $|\widehat{\mathcal{D}_{nf}}(\chi)| < \epsilon/2A$, meaning that \mathcal{D}_{nf} is close to λ .

3.4. Proof of Theorem 1.1, (1) \Leftrightarrow (3)

Our goal here is to prove the equivalence

$\lambda \in \overline{\{\mathcal{D}_{nf}\}_{n \in \mathbb{Z}}} \iff$ the action of \mathcal{D}_f^* is on $\text{Prob}_{\mathcal{L}}$ is uniquely ergodic.

Let us first check the implication \Rightarrow . Let $(n_i)_{i \geq 1}$ be a sequence such that $\mathcal{D}_{n_i f}$ converges weakly to λ when $i \rightarrow +\infty$. Let $\mu \in \text{Prob}_{\mathcal{L}}$, then by one-sided continuity of convolution,

$$\mathcal{D}_{n_i f} * \mu \xrightarrow{i \rightarrow +\infty} \lambda * \mu = \lambda.$$

Let m be any invariant measure on $\text{Prob}_{\mathcal{L}}$, $F : \text{Prob}_{\mathcal{L}} \rightarrow \mathbb{R}$ be a continuous function. Then

$$\int_{\text{Prob}_{\mathcal{L}}} F(\mu) dm(\mu) = \int_{\text{Prob}_{\mathcal{L}}} F(\mathcal{D}_{n_i f} * \mu) dm(\mu),$$

by invariance of m . The Lebesgue dominated convergence Theorem implies that as i goes to infinity:

$$\int_{\text{Prob}_{\mathcal{L}}} F(\mu) dm(\mu) \longrightarrow \int_{\text{Prob}_{\mathcal{L}}} F(\lambda) dm(\mu) = F(\lambda),$$

which means that m is the Dirac measure at λ , as required.

We now prove the converse implication. Assume that the convolution action of \mathcal{D}_f is uniquely ergodic. As λ is a fixed point, the invariant measure is necessarily δ_λ , and thus as the invariant measure is a Dirac mass, there exist a subsequence $n_i \rightarrow +\infty$ such that $\mathcal{D}_{n_i f}$ tends to λ as $i \rightarrow +\infty$.

3.5. Sets of natural density one

PROPOSITION 3.4. — Assume $f : I \rightarrow \mathbb{T}^d$ is asynchronous. Then there exists a set $E \subset \mathbb{Z}$ of full natural density such that for all $\mu \in \text{Prob}_{\mathcal{L}}$,

$$\lim_{n \rightarrow \pm\infty, n \in E} \mathcal{D}_{nf} * \mu = \lambda.$$

Proof. — We consider the measure on $\text{Prob}_{\mathcal{L}}$,

$$\nu_N = \frac{1}{2N+1} \sum_{|k| \leq N} \delta_{\mathcal{D}_{kf}}.$$

As any weak limit of ν_N is \mathcal{D}_f -invariant and $\text{Prob}_{\mathcal{L}}$ is compact, by unique ergodicity of $(\text{Prob}_{\mathcal{L}}, \mathcal{D}_f)$, ν_N converges to δ_λ when N goes to $+\infty$. This implies that for any neighborhood \mathcal{U} of λ , the proportion of $\{\mathcal{D}_{kf}\}_{|k| \leq N}$ outside \mathcal{U} goes to zero as $N \rightarrow +\infty$.

Let $(\mathcal{U}_m)_{m \geq 1}$ be a decreasing basis of neighborhood of λ , and

$$E_m = \{k \in \mathbb{Z} : \mathcal{D}_{kf} \in \mathcal{U}_m\}.$$

Let N_m be an integer such that for all $N \geq N_m$,

$$\mathbb{P}_N(E_m) \geq 1 - \frac{1}{m},$$

where \mathbb{P}_N is the uniform probability on $[-N, N]$. We can modify the sequence $(N_m)_{m \geq 0}$ to be strictly increasing, and choose $N_1 = -1$. Let E be the subset

$$E = \bigcup_{m \geq 1} E_m \cap \{k \in \mathbb{Z} : N_m < |k| \leq N_{m+1}\}.$$

Notice that since the sets E_m are decreasing with m , if $n \leq N_{m+1}$,

$$E \cap [-n, n] \subset E_m.$$

Thus, for n such that $N_m < n \leq N_{m+1}$, we have

$$\mathbb{P}_n(E) \geq 1 - \frac{1}{m}.$$

This proves that E is a set of natural density one. By construction, we have

$$\lim_{n \rightarrow \pm\infty, n \in E} \mathcal{D}_{nf} = \lambda.$$

By continuity of the convolution with μ , the latter limit holds for the sequence $\mathcal{D}_{nf} * \mu$ with the same set E . \square

4. On the smallest algebraic group containing return elements

We now change the setting completely and turn to the application in homogeneous dynamics. This section is devoted to the proof of the following Theorem, which will be a crucial ingredient of the proof of Theorem 1.2.

THEOREM 4.1. — *Let G_0 be the group of real points of an algebraic group \mathbb{G}_0 defined over \mathbb{Q} , without nontrivial \mathbb{Q} -characters, $\Gamma_0 = \mathbb{G}_0(\mathbb{Z})$ be its integer points, $a \in G_0$ be triangularizable with positive eigenvalues, and μ an a -invariant measure on G_0/Γ_0 . We assume that the measure μ is ergodic and non-concentrated. Let $I \subset G_0$ be a fundamental domain for Γ_0 , and denote by \mathcal{L} the lift to I of μ . Let $E \subset I$ be a subset of positive \mathcal{L} -measure. Define*

$$P_E = \left\{ \gamma \in \Gamma_0 : \mathcal{L} \left(\bigcup_{k \in \mathbb{Z}} a^k E \gamma \cap E \right) > 0 \right\},$$

the set of elements of Γ_0 associated to return times in E . Then the smallest algebraic subgroup of G_0 containing P_E is G_0 .

To prove this, let H be the smallest algebraic subgroup of G_0 containing P_E . Our aim is to show that $H = G_0$. This will be done in the following sequence of Lemmata. The strategy is to show the existence of a closed homogeneous subspace of the form $xH\Gamma_0$, invariant by a and which support the measure μ , and then conclude using the non-concentration hypothesis.

4.1. Closure of $H\Gamma_0$

LEMMA 4.2. — *The set $H\Gamma_0$ is closed*

Proof. — Notice that H is defined over \mathbb{Q} , since P_E consists of integer points. We claim that the non-trivial \mathbb{Q} -characters of H are of order 2. Indeed, if c is such a character defined over \mathbb{Q} , the image by c of the subgroup generated by $P_E \cap H^0 \subset \Gamma_0 = \mathbb{G}(\mathbb{Z})$ consists of rational with bounded denominators, and is a multiplicative subgroup, so $c(P_E) \subset \{-1, +1\}$. Therefore, P_E is contained in $\text{Ker}(c^2)$, an algebraic group defined over \mathbb{Q} . By definition of H , $H \subset \text{Ker}(c^2)$, so $H = \text{Ker}(c^2)$ as required. In particular, $H/(H \cap \Gamma_0)$ is of finite volume, by the Theorem of Borel and Harish-Chandra [3, Corollaire 13.2]. By [3, Proposition 8.1], this also implies that $H^0\Gamma_0$ is a closed subset of G_0/Γ_0 , where H^0 is the connected component of the identity of H , in the Zariski topology (a subgroup of finite index). This implies that $H\Gamma_0$ is closed. \square

4.2. Reduction step

LEMMA 4.3. — *To prove Theorem 4.1, we can (and will) assume that for all $k \in \mathbb{Z}$ and $\gamma \in \Gamma_0$ such that $a^k E \gamma \cap E \neq \emptyset$, then $\gamma \in P_E$.*

Proof. — Consider the subset

$$F = E - \bigcup_{(k,\gamma) \in \mathbb{Z} \times \Gamma_0 \text{ s.t. } \mathcal{L}(a^k E \gamma \cap E) = 0} a^k E \gamma.$$

Clearly, F is a subset of E of the same measure, and $P_F = P_E$. So it is sufficient to prove the statement of Theorem 4.1 for F instead of E , and F satisfies the above property. \square

4.3. Invariance of $xH\Gamma_0$

LEMMA 4.4. — *For \mathcal{L} -almost every $x \in E$, $a \in xHx^{-1}$.*

Proof. — By a Theorem of Chevalley [4, Theorem 5.1], there exists a finite dimensional representation α of \mathbb{G} such that H is the stabilizer of a line D , that is $H = \{g \in G_0 : \alpha(g)D = D\}$. By Poincaré recurrence Theorem, for \mathcal{L} -almost every $x \in E$, there exists a sequence $n_k \rightarrow +\infty$ and $\gamma_k \in \Gamma_0$ such that $a^{n_k} x \gamma_k \rightarrow x$ and $a^{n_k} x \gamma_k \in E$. Fix such an $x \in E$.

By Lemma 4.3, we know that $\gamma_k \in P_E \subset H$. It follows that $\alpha(\gamma_k)D = D$, so since $a^{n_k} x \gamma_k \rightarrow x$, we have

$$(4.1) \quad \lim_{k \rightarrow +\infty} \alpha(a)^{n_k} \alpha(x)D = \alpha(x)D.$$

By assumption, a is triangularizable with positive eigenvalues, so $\alpha(a)$ has only positive, real eigenvalues. We claim that (4.1) implies that $\alpha(x)D$ is contained in one of its eigenspaces.

Let

$$\alpha(a) = \delta + \eta,$$

be the Jordan–Chevalley decomposition of $\alpha(a)$, that is: δ and η commutes, η is nilpotent, δ diagonalizable (with positive, real eigenvalues). If p is the nilpotent index of η , for k such that $n_k > p$,

$$\alpha(a)^{n_k} = \sum_{i=0}^{p-1} \binom{n_k}{i} \delta^{n_k-i} \eta^i.$$

Let $v \in \alpha(x)D - \{0\}$, and $v = \sum_{\theta} v_{\theta}$ be its decomposition along the eigenspaces of δ corresponding to the eigenvalues $\{\theta\}$ of δ . Then

$$\alpha(a)^{n_k}(v) = \sum_{\theta} \left(\sum_{i=0}^{p-1} \binom{n_k}{i} \theta^{n_k-i} \eta^i(v_{\theta}) \right).$$

As a function of n_k , this is a combination of polynomials and powers of eigenvalues. If θ_0 is the highest eigenvalue θ for which $v_\theta \neq 0$, and i_0 is the largest i for which $\eta^i(v_{\theta_0}) \neq 0$, then we have the asymptotic as $k \rightarrow +\infty$,

$$\alpha(a)^{n_k}(v) \sim \binom{n_k}{i_0} \theta_0^{n_k - i_0} \eta^{i_0}(v_{\theta_0}).$$

However, we know that projectively, $\alpha(a)^{n_k} \alpha(x)D \rightarrow \alpha(x)D$, so v is colinear to $\eta^{i_0}(v_{\theta_0})$. As η preserves the eigenspace of δ associated to θ_0 ,

$$\alpha(a)\eta^{i_0}(v_{\theta_0}) = \theta_0 \eta^{i_0}(v_{\theta_0}) + \eta^{i_0+1}(v_{\theta_0}) = \theta_0 \eta^{i_0}(v_{\theta_0}),$$

because by definition of i_0 , $\eta^{i_0+1}(v_{\theta_0}) = 0$. This shows that $\eta^{i_0}(v_{\theta_0})$ is an eigenvector, and so is v .

We have proved that $\alpha(x)D$ is contained in an eigenspace of $\alpha(a)$. So D is stabilized by $\alpha(x^{-1}ax)$, meaning that $a \in xHx^{-1}$, as required. \square

Remark. — The positivity hypothesis on the eigenvalues of a can be weakened. The previous Lemma is the only place where positivity is used. If one assumes that a^2 is ergodic for μ instead but allow negative eigenvalues, one can adapt the previous proof, by noticing that one can choose the sequence n_k to have infinitely many odd and infinitely many even values.

4.4. Conclusion of the proof of Theorem 4.1

LEMMA 4.5. — We have $H = G_0$.

Proof. — By ergodicity of a with respect to μ , for μ -almost every $x\Gamma_0$, $a^{\mathbb{Z}}x\Gamma_0$ is dense in the support of μ . By the previous Lemma, we have also for almost every $x \in E$, $a \in xHx^{-1}$, so

$$a^{\mathbb{Z}}x\Gamma_0 \subset xH\Gamma_0.$$

Consider a typical $x \in E$ satisfying both of these properties. By Lemma 4.2, $xH\Gamma_0$ is a closed set. By density of the a -orbit of x in the support of μ , this implies that $\text{supp}(\mu) \subset xH\Gamma_0$, so

$$\mu(xH\Gamma_0) = \mu((xHx^{-1})x\Gamma_0) = 1.$$

By assumption, μ is non-concentrated, so $H = G_0$ necessarily. \square

5. Proof of Theorem 1.2

As in the introduction, let G_0 be the group of real points of a connected semisimple algebraic linear group defined over \mathbb{Q} , Γ_0 its integer points, ρ a irreducible representation over \mathbb{Q} on a space V of dimension $d > 1$, $G = G_0 \times_{\rho} V$ and $\Gamma = \Gamma_0 \times_{\rho} \mathbb{Z}^d$. Let $a \in G_0$ be triangularizable with positive eigenvalues, and μ an a -invariant, ergodic, non-concentrated probability measure on G_0/Γ_0 . Choose an eigenvector $\beta \in V$ for $\rho(a)$, a fundamental domain $I \subset G_0$ for Γ_0 , and denote by \mathcal{L} the lift of μ to I . We now define $f_{\beta} : I \rightarrow \mathbb{T}^d$ by

$$f_{\beta}(x) = \rho(x)^{-1}\beta \bmod \mathbb{Z}^d.$$

The reader should be warned that this map is not really canonical and depends on the chosen fundamental domain I . Under these hypotheses, let us recall the content of Theorem 1.2:

THEOREM. — *The map $f_{\beta} : I \rightarrow \mathbb{T}^d$ is asynchronous.*

The proof is by contradiction. We assume that f_{β} is not asynchronous.

DEFINITION. — *By translate of a \mathbb{Q} -subspace of $V = \mathbb{R}^d$, we mean a set $T \subset V$ of the form $T = v + W$, where $v \in V$, and $W \subset V$ a subspace of V defined over \mathbb{Q} . The \mathbb{Q} -subspace W is called the direction of T .*

In particular, there is no rationality assumption on v , and T itself does not have to be defined over \mathbb{Q} .

The proof of Theorem 2 goes as follows: we lift f_{β} to a map $I \rightarrow V$; not being asynchronous means that this lift, restricted to a set E of positive measure, takes values in a proper translate of a \mathbb{Q} -subspace (Lemma 5.1). We then check that the set of directions of this subspace is invariant under 'return elements' of Γ_0 for this set E (Lemma 5.2). Finally, using Theorem 4.1 and irreducibility of ρ , we conclude.

If non-empty, the intersection of two translates of \mathbb{Q} -subspaces is again a translate of \mathbb{Q} -subspace. This property allows us to define, for a set $E \subset I$ of positive \mathcal{L} -measure, the set T_E which is the smallest translate of \mathbb{Q} -subspace containing $\{\rho(x)^{-1}\beta\}_{x \in E}$. We denote by W_E its direction.

LEMMA 5.1. — *There exist $E \subset I$ of positive measure such that $W_E \neq V$.*

Proof. — Since f_{β} is not asynchronous, so for some non-trivial character $\chi \in \widehat{\mathbb{T}^d}$, $\chi \circ f_{\beta}$ is constant on a set F of positive \mathcal{L} -measure. There exists $\mathbf{n} \in \mathbb{Z}^d - \{0\}$, such that

$$\chi(\bar{v}) = e^{2i\pi \langle \mathbf{n}, v \rangle}.$$

So the set $\{\langle \mathbf{n}, \rho(x)^{-1}\beta \rangle\}_{x \in F}$ is contained in a countable set of the form $c + \mathbb{Z}$, for some $c \in \mathbb{R}$. This implies that at least one of the sets

$$F_m = \{x \in F : \langle \mathbf{n}, \rho(x)^{-1}\beta \rangle = c + m\},$$

for $m \in \mathbb{Z}$, has positive \mathcal{L} -measure. By construction, for such a $m \in \mathbb{Z}$,

$$T_{F_m} \subset \{v \in V : \langle \mathbf{n}, v \rangle = c + m\},$$

the right-hand side set being the translate of a proper \mathbb{Q} -subspace, $E = F_m$ satisfies the Lemma. \square

We now fix $E \subset I$ a set of positive measure, such that W_E is of minimal possible dimension (it exists). By Lemma 5.1, $W_E \neq V$. Like in Theorem 4.1, we define

$$P_E = \left\{ \gamma \in \Gamma_0 : \mathcal{L} \left(\bigcup_{k \in \mathbb{Z}} a^k E \gamma \cap E \right) > 0 \right\}.$$

LEMMA 5.2. — For all $\gamma \in P_E$, $\rho(\gamma)W_E = W_E$.

Proof. — Recall that β is an eigenvector for $\rho(a)$, denote by κ the corresponding eigenvalue. By definition of P_E , there exists $k \in \mathbb{Z}$ with $\mathcal{L}(a^k E \gamma \cap E) > 0$. Let $F = a^k E \gamma \cap E$, then for all $x \in F$, there exist $y \in E$ such that $x = a^k y \gamma$. We have:

$$\rho(x)^{-1}\beta = \rho(\gamma^{-1}y^{-1}a^{-k})\beta = \kappa^{-k}\rho(\gamma)^{-1}\rho(y)^{-1}\beta \in \kappa^{-k}\rho(\gamma)^{-1}T_E.$$

Note that since $\rho(\gamma)^{-1}$ is a matrix with integer coefficients, $\kappa^{-k}\rho(\gamma)^{-1}T_E$ is also the translate of a \mathbb{Q} -subspace, containing $\{\rho(x)^{-1}\beta\}_{x \in F}$. By definition of T_F , this means that

$$T_F \subset \kappa^{-k}\rho(\gamma)^{-1}T_E.$$

Since $F \subset E$, $T_F \subset T_E$, and because E was chosen such that T_E is of minimal possible dimension, we have $T_E = T_F$, so from the rank-nullity Theorem,

$$\kappa^{-k}\rho(\gamma)^{-1}T_E = T_E.$$

This implies equality for the directions, $\rho(\gamma)^{-1}W_E = W_E$, and multiplication by $\rho(\gamma)$ concludes the proof. \square

LEMMA 5.3. — We have $W_E = \{0\}$, that is, the map $x \mapsto \rho(x)^{-1}\beta$ is constant on E .

Proof. — The subgroup $\text{Stab}_{G_0}(W_E) = \{g \in G_0 : \rho(g)W_E = W_E\}$ is an algebraic subgroup containing P_E , by Lemma 5.2. By Theorem 4.1, this group is G_0 . Since ρ is irreducible over \mathbb{Q} and W_E is defined over \mathbb{Q} ,

$W_E = \{0\}$, or $W_E = V$. But the latter cannot happen, because of the choice of E . \square

From now on, we fix some $x_0 \in E$. By the previous Lemma, T_E is the point $\rho(x_0)^{-1}\beta$.

LEMMA 5.4. — *For any $\gamma \in P_E$, $\rho(\gamma) \in \text{Stab}(\mathbb{R}\rho(x_0)^{-1}\beta)$.*

Proof. — Let $\gamma \in P_E$. Thus there exist $k \in \mathbb{Z}$ such that $\mathcal{L}(a^k E \gamma \cap E) > 0$. In the proof of Lemma 5.2, we saw that

$$\kappa^{-k} \rho(\gamma)^{-1} T_E = T_E.$$

But since $T_E = \{\rho(x_0)^{-1}\beta\}$, this means that $\rho(\gamma)^{-1}$ stabilizes the line through $\rho(x_0)^{-1}\beta$. \square

The end of the proof of Theorem 1.2 is given by the following contradictory Claim.

LEMMA 5.5. — *The space V is one-dimensional.*

Proof. — The group

$$\{g \in G : \rho(g) \in \text{Stab}(\mathbb{R}\rho(x_0)^{-1}\beta)\},$$

is an algebraic group containing P_E . By Theorem 4.1, it follows that $\rho(x_0)^{-1}\beta$ is a common eigenvector for all elements of $\rho(G_0)$ (and so is β). Were the representation ρ irreducible over \mathbb{R} , this would be sufficient to conclude; however we assumed only \mathbb{Q} -irreducibility, and have no particular rationality assumption on β .

Since G_0 is semisimple and connected, the eigenvalue associated to $\rho(x_0)^{-1}\beta$ is 1 for every $g \in G_0$. Let $V_1(g)$ denote the eigenspace associated to the eigenvalue 1 for the operator $\rho(g)$. Let $Z = \bigcap_{\gamma \in P_E} V_1(\gamma)$. This subspace Z is defined over \mathbb{Q} , because P_E consists of integral points. The set of $g \in G_0$ acting trivially on Z is an algebraic subgroup containing P_E , so again is G_0 . Since $\rho(x_0)^{-1}\beta \in Z$, Z is of positive dimension. By \mathbb{Q} -irreducibility of ρ , $Z = V$ is an irreducible representation where G_0 acts trivially, so is one-dimensional. \square

6. Ergodicity and mixing

Let $\alpha \in V$ be arbitrary, we now consider the action of (a, α) on $(G/\Gamma, \lambda)$, which has the action of a on $(G_0/\Gamma_0, \mu)$ as a factor. We now prove Theorem 1.4, which claims that the action of (a, α) is ergodic with respect to λ , and, if a is not unipotent, then mixing properties for (a, α) with respect to

λ are inherited from the same mixing properties of a w.r.t. μ . By assumption, a is triangularizable with positive eigenvalues. We separate the proof in two cases.

Case 1: $\rho(a)$ is unipotent. — Let $\beta \in V - \{0\}$ be an eigenvector for $\rho(a)$, its eigenvalue is 1.

Notice that the actions of (e, β) and (a, α) commute: since $\rho(a)\beta = \beta$, we have

$$(a, \alpha)(e, \beta) = (a, \alpha + \rho(a)\beta) = (a, \rho(e)\alpha + \beta) = (e, \beta)(a, \alpha).$$

Consider the ergodic decomposition of λ with respect to the action of (a, α) : there exists a measure m on the set of ergodic, (a, α) -invariant measures on G/Γ , such that

$$\lambda = \int \nu \, dm(\nu).$$

If we apply the projection map $\pi : G/\Gamma \rightarrow G_0/\Gamma_0$ to this equality, we obtain

$$\mathcal{L} = \mu = \int (\pi)_* \nu \, dm(\nu),$$

where $(\pi)_* \nu$ are a -invariant. Since μ is a -ergodic, we have that $(\pi)_* \nu = \mu$, m -almost surely. Therefore, m is supported on Prob_μ . But since (e, β) commutes with (a, α) , $(e, \beta)_* m$ is the measure associated to the ergodic decomposition of $(e, \beta)_* \lambda = \lambda$. This implies that m is (e, β) -invariant, and by Corollary 1.3, m is the Dirac measure at λ . This concludes the proof of the ergodicity.

Case 2: $\rho(a)$ is not unipotent. — Thus $\rho(a)$ has some of its eigenvalues different from 1. Since G_0 is semisimple, $\det(\rho(a)) = 1$ so there exist at least one eigenvalue $\kappa < 1$. Let $\beta \in V$ be an eigenvector of $\rho(a)$ associated to κ .

Notice that, for $k \geq 0$,

$$(a, \alpha)^k (e, \beta) = (a^k, \rho(a)^k \beta) + \sum_{i=0}^{k-1} \rho(a)^i \alpha = (e, \kappa^k \beta)(a, \alpha)^k,$$

and $(e, \kappa^k \beta) \rightarrow (e, 0)$ when $k \rightarrow +\infty$. This implies that the distance (with respect to a distance $d_{G/\Gamma}$ on G/Γ induced by a right- G -invariant riemannian distance on G) between $(a, \alpha)^k(x, v)\Gamma$ and $(a, \alpha)^k(e, \beta)(x, v)\Gamma$ tends to zero as k tends to $+\infty$. In other words, we have proved:

LEMMA 6.1. — *The strong stable distribution for (a, α) , defined by*

$$W^{ss}((x, v)\Gamma) = \left\{ (y, w)\Gamma : \lim_{k \rightarrow +\infty} d_{G/\Gamma}((a, \alpha)^k(x, v)\Gamma, (a, \alpha)^k(y, w)\Gamma) = 0 \right\},$$
is invariant under the action of (e, β) .

We first prove the claims about ergodicity and weak-mixing.

Let $f \in L^2(G/\Gamma, \lambda) - \{0\}$ be an eigenvector for the Koopman operator of (a, α) , that is

$$f((a, \alpha)(x, v)\Gamma) = \omega f((x, v)\Gamma),$$

for λ -almost every $(x, v)\Gamma$, for some $\omega \in \mathbb{C}$ of modulus one. Recall that ergodicity states that any such eigenvector associated to $\omega = 1$ is constant almost everywhere, and weak-mixing that any such eigenvector is constant almost everywhere and moreover $\omega = 1$.

To prove ergodicity or weak-mixing, we may (and will) restrict to the case where f is bounded. Let us recall the conclusion of the Hopf argument, as theorized by Coudène [5]:

THEOREM ([5]). — *Let X be a metric space, equipped with a Borel measure on X , $T : X \rightarrow X$ a measure preserving conservative transformation of X . Assume that there exists a countable family of open sets of finite measure which covers almost all of X . Then any bounded T -eigenfunction u is W^{ss} -invariant, meaning: there exists a full measure set $\Omega \subset X$ such that for all $(x, y) \in \Omega^2$ such that $\lim_{n \rightarrow +\infty} d(T^n x, T^n y) = 0$, then $u(x) = u(y)$.*

This result applies here for the space G/Γ , the (finite, so conservative) measure λ , and the transformation by the left multiplication by (a, α) . Thus there exists a full measure set $\Omega \subset G/\Gamma$ such that for all $(x, v)\Gamma \in \Omega$, $(y, w)\Gamma \in \Omega$ such that $(y, w)\Gamma \in W^{ss}((x, v)\Gamma)$, $f((x, v)\Gamma) = f((y, w)\Gamma)$.

By Theorem 1.2, the angle map $x \mapsto \rho(x)^{-1}\beta$ is asynchronous. In particular, for every strict linear rational (closed) subtorus $\mathcal{T} \subset \mathbb{T}^d$, the set of x such that $\rho(x)^{-1}\beta$ does not belong to \mathcal{T} is of full measure. As the set of such subtorus is countable, this implies that for μ -almost every x , the translation on \mathbb{T}^d given by $\rho(x)^{-1}\beta$ is ergodic. Via a linear change of variable, this means that for μ -almost every $x\Gamma_0 \in G_0/\Gamma_0$, the translation by (e, β) on the fiber above $x\Gamma_0$ is ergodic.

As f is (e, β) -invariant, for μ -almost every $x\Gamma_0$, f is almost everywhere constant on the fiber above $x\Gamma_0$, and merely depends on $x\Gamma_0$. Write $F \in L^2(G_0/\Gamma_0, \mu)$ for its almost everywhere value, that is $f = F \circ \pi$, λ -almost everywhere, where $\pi : G/\Gamma \rightarrow G_0/\Gamma_0$ is the fiber bundle. We have then

$$F(ax\Gamma_0) = \omega F(x\Gamma_0).$$

If $\omega = 1$, then F is a -invariant, and by ergodicity of a , F is constant μ -a.e., and so f is constant λ -a.e.. This proves the ergodicity.

Assume now that the action of a is weakly mixing on $(G_0/\Gamma_0, \mu)$. Let $f \in L^2(G/\Gamma, \lambda)$ be like previously an eigenvector for the Koopman operator of (a, α) , $F \in L^2(G_0/\Gamma_0, \mu)$ its almost sure value depending on the fiber. By what we saw before, F is an eigenvector for the Koopman operator of a , so $\omega = 1$ by weak-mixing of a . By ergodicity, F is constant, as was to be proved.

We now assume that the action of a on $(G_0/\Gamma_0, \mu)$ is strongly mixing, and wish to prove that (a, α) is also strongly mixing. Recall that strong mixing of a is equivalent to the fact that for all $F \in L^2(G_0/\Gamma_0, \mu)$, $F \circ a^k$ converges weakly to a constant as $k \rightarrow +\infty$.

Let $f \in L^2(G/\Gamma, \lambda)$. Let $g \in L^2(G/\Gamma, \lambda)$ be any weak limit of $f \circ (a, \alpha)^k$ as $k \rightarrow +\infty$ along a subsequence. By another result of Coudène [6], generalizing a result of Babillot, g is W^{ss} -invariant. By ergodicity of (e, β) on almost every fiber, $g = G \circ \pi$ almost surely, where $G \in L^2(G_0/\Gamma_0, \mu)$. Define

$$F(x\Gamma_0) = \int_{\mathbb{R}^d/\rho(x)\mathbb{Z}^d} f((x, v)\Gamma) d\text{Haar}_{\mathbb{R}^d/\rho(x)\mathbb{Z}^d}(v),$$

the mean value of f on each fiber. Let $H \in L^2(G_0/\Gamma_0, \mu)$ be a test-function. Then

$$\int_{G/\Gamma} (f \circ (a, \alpha)^k)(H \circ \pi) d\lambda = \int_{G_0/\Gamma_0} (F \circ a^k) \cdot H d\mu.$$

Taking the limits in the left-hand side and right-hand side respectively along the subsequence, using the strong mixing property for a , gives:

$$\int_{G/\Gamma} (G \circ \pi)(H \circ \pi) d\lambda = \left(\int_{G_0/\Gamma_0} F d\mu \right) \left(\int_{G_0/\Gamma_0} H d\mu \right),$$

in other words,

$$\int_{G_0/\Gamma_0} G H d\mu = \left(\int_{G_0/\Gamma_0} F d\mu \right) \left(\int_{G_0/\Gamma_0} H d\mu \right),$$

which implies that G is μ -almost everywhere the constant $\int_{G_0/\Gamma_0} F d\mu = \int_{G/\Gamma} f d\lambda$. Therefore the only possible weak limit of $f \circ (a, \alpha)^k$ is the above constant. By weak compactness of the ball of radius $\|f\|_2$ in $L^2(G/\Gamma, \lambda)$, this proves that this sequence must converge weakly to $\int_{G/\Gamma} f d\lambda$, as required.

BIBLIOGRAPHY

- [1] J. AARONSON, *An introduction to infinite ergodic theory*, Mathematical Surveys and Monographs, vol. 50, American Mathematical Society, 1997, xii+284 pages.
- [2] P. ARNOUX & A. M. FISHER, “The scenery flow for geometric structures on the torus: the linear setting”, *Chin. Ann. Math., Ser. B* **22** (2001), no. 4, p. 427-470.
- [3] A. BOREL, *Introduction aux groupes arithmétiques*, Actualités Scientifiques et Industrielles, vol. 1341, Hermann, 1969, 125 pages.
- [4] ———, *Linear algebraic groups*, second ed., Graduate Texts in Mathematics, vol. 126, Springer, 1991, xii+288 pages.
- [5] Y. COUDÈNE, “The Hopf argument”, *J. Mod. Dyn.* **1** (2007), no. 1, p. 147-153.
- [6] Y. COUDÈNE, “On invariant distributions and mixing”, *Ergodic Theory Dyn. Syst.* **27** (2007), no. 1, p. 109-112.
- [7] N. D. ELKIES & C. T. MCMULLEN, “Gaps in \sqrt{n} mod 1 and ergodic theory”, *Duke Math. J.* **123** (2004), no. 1, p. 95-139.
- [8] H. FURSTENBERG, *Recurrence in ergodic theory and combinatorial number theory*, Princeton University Press, 1981, M. B. Porter Lectures, xi+203 pages.
- [9] D. KLEINBOCK, N. SHAH & A. STARKOV, “Dynamics of subgroup actions on homogeneous spaces of Lie groups and applications to number theory”, in *Handbook of dynamical systems, Vol. 1A*, North-Holland, 2002, p. 813-930.
- [10] J. MARKLOF & A. STRÖMBERGSSON, “The Boltzmann-Grad limit of the periodic Lorentz gas”, *Ann. Math. (2)* **174** (2011), no. 1, p. 225-298.
- [11] D. W. MORRIS, *Ratner’s theorems on unipotent flows*, Chicago Lectures in Mathematics, University of Chicago Press, 2005, xii+203 pages.
- [12] U. SHAPIRA, “A solution to a problem of Cassels and Diophantine properties of cubic numbers”, *Ann. Math. (2)* **173** (2011), no. 1, p. 543-557.
- [13] A. STRÖMBERGSSON, “An effective Ratner equidistribution result for $\mathbf{SL}(2, \mathbb{R}) \ltimes \mathbb{R}^2$ ”, *Duke Math. J.* **164** (2015), no. 5, p. 843-902.
- [14] D. THOMINE, “Keplerian shear in ergodic theory”, *Ann. Henri Lebesgue* **3** (2020), p. 649-676.

Manuscrit reçu le 19 décembre 2017,
révisé le 17 août 2023,
accepté le 4 avril 2024.

François MAUCOURANT
Université de Rennes, IRMAR
Campus de Beaulieu
35042 Rennes cedex (France)
francois.maucourant@univ-rennes.fr