

KOHUR GOWRISANKARAN

**Fatou-Naïm-Doob limit theorems in the
axiomatic system of Brelot**

Annales de l'institut Fourier, tome 16, n° 2 (1966), p. 455-467

http://www.numdam.org/item?id=AIF_1966__16_2_455_0

© Annales de l'institut Fourier, 1966, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

FATOU-NAIM-DOOB LIMIT THEOREMS IN THE AXIOMATIC SYSTEM OF BRELOT

by Kohur GOWRISANKARAN

1. Introduction.

Let Ω be a locally compact Hausdorff space which is connected and has a countable base. Let \mathfrak{H} be a class of real valued continuous functions, called harmonic functions, on open subsets of Ω such that for each open set $W \subset \Omega$, the set \mathfrak{H}_W , consisting of all functions in \mathfrak{H} defined on W , is a real vector space. Let this class \mathfrak{H} satisfy the axioms 1, 2 and 3 of M. Brelot [1]. Let, moreover, there exist a potential > 0 on Ω .

The classical Fatou-Naim-Doob limit theorems were extended to the axiomatic system of M. Brelot in [2]. But, besides the above mentioned axioms, we had assumed the validity of axioms D and \mathfrak{R}_n [2]. The object of this paper is to show that the Fatou-Naim-Doob limit theorems (cf. Theorem 8) hold good in the axiomatic set up without these supplementary axioms (viz. D and \mathfrak{R}_n). The method consists in proving first, the limit theorems for a special class of superharmonic functions (cf. Theorem 4), and using it systematically to prove the general result. A novel feature in our proof is the consideration of a modified Dirichlet problem. We shall mostly follow the notation of [1, 2, 3].

Notation.

S^+ : The set of all non-negative superharmonic functions on Ω .

H^+ : The set of all non-negative harmonic functions on Ω .

Λ : A compact base for S^+ (compact in the T-topology [3]).

Δ_1 : The set of minimal harmonic functions contained in Λ .

For any $E \subset \Omega$ and $\nu \in S^+$,

$$R_\nu^E = \text{Inf} \{ \omega : \omega \in S^+ \text{ and } \omega \geq \nu \text{ on } E \}.$$

For any $h \in \Delta_1$, $\mathcal{F}_h = \{ E \subset \Omega : R_h^E \not\equiv h \}$ [2]. A set E is thin at $h \in \Delta_1$ if $R_h^E \equiv h$ (i.e. if \mathcal{F}_h leaves no trace on E).

The limits of any function f following \mathcal{F}_h , for any $h \in \Delta_1$, are called the fine limits of $f(x)$, as x tends to h . To every harmonic function $\omega \in H^+$ corresponds a unique measure μ_ω on Δ_1 , called the canonical measure corresponding to ω , such that $\omega = \int h \mu_\omega(dh)$. For any regular domain $\delta \subset \Omega$, and $x \in \delta$, $d\rho_x^\delta$ is the measure on $\partial\delta$ which associates to a finite continuous function f on $\partial\delta$ the integral $H_f(x)$. For the considerations below, let us fix a $u \in H^+$ with $u > 0$. Let μ_u be the canonical measure on Δ_1 , corresponding to u . A function ν on Ω is said to be super- u -harmonic (respectively- u -harmonic) if $u\nu$ is superharmonic (resp. harmonic) on Ω .

2. Fine limits of bounded u -harmonic functions.

LEMMA 1. — *Let $V \subset \Omega$ be an open set. Then, for every $x \in \Omega$, the mapping $h \rightarrow R_h^V(x)$ of $H^+ \rightarrow \mathbf{R}^+$ is lower semi-continuous.*

Proof. — Let $h_n \in H^+$ converge to $h \in H^+$. Let $\nu_n = R_{h_n}^V$. Then, ν_n is a non-negative superharmonic function on Ω and $\nu_n = h_n$ on V . Let $\nu = \liminf_{n \rightarrow \infty} \nu_n$. Let ω be a regular domain of Ω . Then,

$$\nu_n(y) \geq \int \nu_n(z) \rho_y^\omega(dz) \quad \text{for all } y \in \omega.$$

Hence,

$$\begin{aligned} \nu(y) = \liminf_{n \rightarrow \infty} \nu_n(y) &\geq \liminf_{n \rightarrow \infty} \int \nu_n(z) \rho_y^\omega(dz) \\ &\geq \int \nu(z) \rho_y^\omega(dz) \quad (\text{Fatou's Lemma}). \end{aligned}$$

(Note here that ν is a ρ_x^ω -measurable function.) Since ν is also non-negative, it follows that ν is an $S_{\mathcal{B}}$ -function, where \mathcal{B} is the class of all regular domains of Ω [1]. Hence, $\hat{\nu}$, the lower semi-continuous regularisation of ν , is a superharmonic function. But $\nu(y) = h(y)$, for all $y \in V$, and hence $\hat{\nu} = h$ on V .

It follows that $\nu \geq \hat{\nu} \geq R_h^V$ on Ω . This gives the required lower semi-continuity.

COROLLARY. — *For any regular domain δ of Ω and all $x \in \delta$, the function $h \rightarrow \int R_h^V(z) \rho_x^{\delta}(dz)$ is lower semi-continuous on H^+ .*

The corollary follows from the lemma by the use of Fatou's lemma.

LEMMA 2. — *The set \mathcal{E}_V of points of Δ_1 , where an open set $V \subset \Omega$ is thin, is a borel subset of Δ_1 .*

Proof. — Let $\{\delta_n\}$ be a countable covering of Ω by regular domains. Let, for each n , $x_n \in \delta_n$. Define,

$$F'_n = \{h \in \Lambda \cap H^+ : \int R_h^V(y) \rho_{x_n}^{\delta_n}(dy) < h(x_n)\}.$$

In view of the above lemma and its corollary, F'_n is a borel subset of Λ (in fact, a K_σ — set). Hence, $F_n = F'_n \cap \Delta_1$ is a borel subset of Δ_1 . It can be proved as in [2], that $\bigcup_{n=1}^{\infty} F_n$ is precisely the set \mathcal{E}_V . The lemma is proved.

THEOREM 1. — *Let $V \subset \Omega$ be any open set. Then $R_u^V \equiv u$ if and only if $\mu_u(\mathcal{E}_V) = 0$.*

Proof. — Let $\mu_u(\mathcal{E}_V) = 0$. For any $x \in \Omega$, we have,

$$R_u^V(x) = \int R_h^V(x) \mu_u(dh) \quad (\text{Th. 22.3, [3]}).$$

Since $R_h^V(x) = h(x)$, for all $h \in \Delta_1 - \mathcal{E}_V$, and $\mu_u(\mathcal{E}_V) = 0$, we get,

$$R_u^V(x) = \int h(x) \mu_u(dh) = u(x).$$

This is true whatever be $x \in \Omega$.

Conversely, suppose that $R_u^V \equiv u$. Let $\{\delta_n\}$ be a sequence covering Ω , each δ_n being a regular domain, and consider the sets $F_n \subset \Delta_1$, as defined in the above lemma.

Let ν_k be the swept-out measure corresponding to the measure $d\rho_{x_k}^{\delta_k}$ relative to the sweeping out on V . (Th. 10. 1, [3]). (Note that $d\rho_{x_k}^{\delta_k}$ is with the compact support $\partial\delta_k$). The measure ν_k is such that, for any $\nu \in S^+$,

$$\int \nu(y) \nu_k(dy) = \int R_\nu^V(y) \rho_{x_k}^{\delta_k}(dy).$$

We have,

$$\begin{aligned} \int R_u^V(y) \rho_{x_k}^{\delta_k}(dy) &= \int u(z) \nu_k(dz) = \int \nu_k(dz) \int h(z) \mu_u(dh) \\ &= \int \mu_u(dh) \int h(z) \nu_k(dz) = \int \mu_u(dh) \int R_h^V(y) \rho_{x_k}^{\delta_k}(dy) \dots (1) \end{aligned}$$

(Lebesgue-Fubini Theorem).

Now,

$$\begin{aligned} \int h(x_k) \mu_u(dh) &= u(x_k) = \int R_u^V(y) \rho_{x_k}^{\delta_k}(dy) \quad (\text{hypothesis}) \\ &= \int \mu_u(dh) \int R_h^V(y) \rho_{x_k}^{\delta_k}(dy) \quad (\text{from (1)}). \end{aligned}$$

It follows that,

$$\int [h(x_k) - \int R_h^V(y) \rho_{x_k}^{\delta_k}(dy)] \mu_u(dh) = 0 \dots (2).$$

Since the integrand in the above equation is always ≥ 0 , we get, $h(x_k) = \int R_h^V(y) \rho_{x_k}^{\delta_k}(dy)$, for all $h \in \Delta_1$, except for a set of μ_u -measure zero. But the exceptional set where the inequality does not hold good is precisely F_k . Hence,

$$\mu_u(F_k) = 0.$$

It follows, from the above lemma, that $\mu_u(\mathcal{E}_V) = 0$. The theorem is proved.

COROLLARY. — *The greatest harmonic minorant of R_u^V is the function $\int h \mu_u^V(dh)$ where μ_u^V is the restriction of μ_u to $\Delta_1 - \mathcal{E}_V$. Hence, R_u^V is a potential if V is thin μ_u -almost everywhere on Δ_1 .*

The proof of the corollary is exactly as in (Cor. Th. II. 2, [2]).

THEOREM 2. — *Let $w > 0$ be a potential on Ω . Then $\frac{w}{u}$ has the fine limit zero, at μ_u -almost every element of Δ_1 .*

Proof. — It is enough to show that, for every rational number $r > 0$, the set $V_r = \left\{ x \in \Omega : \frac{w(x)}{u(x)} > r \right\}$ is thin μ_u -almost everywhere. But, since $\frac{w}{u}$ is a lower semi-continuous function, V_r is an open subset of Ω . Further, $R_{V_r}^V \leq \frac{w}{r}$. Hence $R_{V_r}^V$ is a potential and it follows (Cor. to Theorem 1) that V_r is

thin at μ_u -almost every element of Δ_1 . This is true for every $r > 0$. The proof is completed easily.

The following result is an important corollary of the above theorem.

THEOREM 3. — *Let ν and ω be two non-negative harmonic functions on Ω such that their canonical measures μ_ν and μ_ω on Δ_1 are singular relative to each other. If, $\nu > 0$ on Ω , then, $\frac{\omega}{\nu}$ has the fine limit zero, at μ_ν -almost every element of Δ_1 .*

Proof. — Let $\nu' = \text{Inf}(\nu, \omega)$. Then it is clear that ν' is a potential on Ω . By the above theorem, we can find a set $E \subset \Delta_1$ of μ_u -measure zero such that, for every $h \in \Delta_1 - E$, $\text{fine lim}_{x \rightarrow h} \frac{\nu'(x)}{\nu(x)} = 0$. From this we easily deduce that, the $\text{fine lim}_{x \rightarrow h} \frac{\omega(x)}{\nu(x)} = 0$, for every $h \in \Delta_1 - E$. This completes the proof.

THEOREM 4. — *Let ω be a bounded u -harmonic function on Ω . Then, ω has a fine limit at μ_u -almost every element of Δ_1 .*

Proof. — Define, for a μ_u -summable function f on Δ_1 ,

$$\sigma_f = \int f(h) \cdot \frac{h}{u} \mu_u(dh).$$

For the characteristic function χ_E of a μ_u -measurable set $E \subset \Delta_1$, let us denote by σ_E the function σ_{χ_E} and σ'_E the function $\sigma_{\chi_{E^c}}$. σ_f is a u -harmonic function, for every such f . Now, for a μ_u -measurable set $E \subset \Delta_1$, since $\sigma_E \leq 1$ on Ω ,

$$\text{fine lim sup}_{x \rightarrow h} \sigma_E(x) \leq 1 \quad \text{for all } h \in \Delta_1.$$

If either $\mu_u(E)$ or $\mu_u(\int E)$ is zero, then $\sigma'_E = 1$ (or respectively $\sigma_E = 1$), and the fine limits of σ_E and σ'_E , exist at all points of Δ_1 . On the other hand, suppose $\mu_u(E) \neq 0$ and also

$$\mu_u(\int E) \neq 0.$$

Then, $u\sigma_E$ and $u\sigma'_E$ are two harmonic functions > 0 on Ω and their canonical measures on Δ_1 (viz. μ_u restricted to E and $\int E$) are singular relative to each other. Hence, by the Theorem 3, σ_E/σ'_E has the fine limit zero at μ_u -almost every element of $\Delta_1 - E$. It follows then that,

$$\text{fine lim sup}_{x \rightarrow h} \sigma_E(x) \leq \text{fine lim}_{x \rightarrow h} \frac{\sigma_E(x)}{\sigma'_E(x)} = 0$$

for μ_u -almost every element of $\Delta_1 - E$, as $\sigma'_E(x) \leq 1$. Hence,

$$\text{fine lim sup}_{x \rightarrow h} \sigma_E(x) \leq \chi_E(h)$$

for μ_u -almost every $h \in \Delta_1 \dots$ (3).

In particular, the inequality (3) is valid for the complement of E and we deduce that,

$$\text{fine lim inf}_{x \rightarrow h} \sigma_E(x) \geq \chi_E(h) \quad \text{for } \mu_u\text{-almost every } h \in \Delta_1.$$

In any case we get, for the characteristic function χ_E of a μ_u -measurable set E contained in Δ_1 ,

$$\text{fine lim}_{x \rightarrow h} \sigma_E(x) = \chi_E(h) \quad \text{for } \mu_u\text{-almost every } h \in \Delta_1 \dots \text{ (4).}$$

Suppose, now, $f \geq 0$ is a μ_u -measurable function on Δ_1 . Then, there exists an increasing sequence of non-negative simple functions s_n such that $\lim_{n \rightarrow \infty} s_n = f$. We deduce easily from (4) that

$$\text{fine lim}_{x \rightarrow h} \sigma_{s_n}(x) = s_n(h) \quad \text{for } \mu_u\text{-almost every } h \in \Delta_1.$$

Hence, σ_f satisfies,

$$\text{fine lim inf}_{x \rightarrow h} \sigma_f(x) \geq s_n(h) \quad \text{for } \mu_u\text{-almost every } h \in \Delta_1.$$

Now, it is easily seen that,

$$\text{fine lim inf}_{x \rightarrow h} \sigma_f(x) \geq f(h) \quad \text{for } \mu_u\text{-almost every } h \in \Delta_1 \dots \text{ (5).}$$

Let us now consider a bounded μ_u -measurable function g on Δ_1 (say $|g| \leq M$). Then, applying the inequality (5) to the

two functions $\sigma_{(M \pm g)}$, and noting that, $\sigma_{M \pm g} = M \pm \sigma_g$, we get that

$$\text{fine lim}_{x \rightarrow h} \sigma_g(x) = g(h) \quad \text{for} \quad \mu_u\text{-almost every } h \in \Delta_1.$$

Now, the proof of the theorem is completed by noting that any bounded u -harmonic function ω is equal to $u\sigma_g$, for some bounded μ_u -measurable function g on Δ_1 ; this g is unique (depending on ω) upto a set of μ_u -measure zero.

Remark 1. — In the course of the proof of the theorem, we have shown that, for any $f \geq 0$, which is μ_u -measurable,

$$\text{fine lim inf}_{x \rightarrow h_0} \int f(h) \frac{h(x)}{u(x)} \mu_u(dh) \geq f(h_0),$$

for μ_u -almost every $h_0 \in \Delta_1$ (viz. the inequality (5)).

Remark 2. — For any bounded u -harmonic function ω on Ω , if $g(h) = \text{fine lim}_{x \rightarrow h} \omega(x)$, (the function g is defined upto a set of μ_u -measure zero), then g is μ_u -measurable and

$$\omega(x) = \int g(h) \frac{h(x)}{u(x)} \mu_u(dh).$$

In particular, if the fine limit is ≥ 0 for μ_u -almost every element of Δ_1 , then ω is non-negative.

Remark 3. — For any bounded super- u -harmonic function ν on Ω , the fine lim $\nu(x)$ exists for μ_u -almost every $h \in \Delta_1$.

THEOREM 5. — (*The Minimum Principle*). Let ν be a lower bounded super- u -harmonic function on Ω . Suppose that, for every $h \in \Delta_1 - E$, fine lim $\sup_{x \rightarrow h} \nu(x) \geq 0$, where E is a set with $\mu_u^*(E) = 0$. Then, ν is ≥ 0 on Ω .

Proof. — Let $\alpha > 0$ be such that $\nu \geq -\alpha$. Consider

$$\nu' = \text{Inf}(\nu, 1).$$

Then ν' is a super- u -harmonic function such that $\nu' \geq -\alpha$. The theorem would be proved if we show that $\nu' \geq 0$ on Ω .

Now, it is easily seen that $\text{fine lim sup}_{x \rightarrow h} \nu'(x) \geq 0$, for all $h \in \Delta_1 - E$. But, we know, (by the Remark 3 following the Theorem 4) that, the limit of ν' exists, following \mathcal{F}_h , for μ_u -almost every $h \in \Delta_1$; and this fine limit is precisely the fine limit of u_1 , where u_1 is the greatest u -harmonic minorant of ν' . Hence, we have that the fine limit of u' is ≥ 0 at μ_u -almost every element of Δ_1 . It follows that $u_1 \geq 0$ (from the Remark 2, Theorem 4). A fortiori, $\nu' \geq 0$. This completes the proof of the theorem.

3. A Dirichlet problem.

Let Σ be the set of all lower bounded super- u -harmonic functions on Ω . Corresponding to any extended real valued function f on Δ_1 , define,

$$\Sigma_f = \left\{ \begin{array}{l} \nu \in \Sigma : \exists \text{ a set } E_\nu \subset \Delta_1 \text{ of } \mu_u\text{-measure zero such that for} \\ \text{all } h \in \Delta_1 - E_\nu, \text{ fine lim inf}_{x \rightarrow h} \nu(x) \geq f(h) \end{array} \right\}$$

$$\tilde{\Sigma}_f = \left\{ \begin{array}{l} \nu \in \Sigma : \exists \text{ a set } F_\nu \subset \Delta_1 \text{ of } \mu_u\text{-measure zero such that for} \\ \text{all } h \in \Delta_1 - F_\nu, \text{ fine lim sup}_{x \rightarrow h} \nu(x) \geq f(h) \end{array} \right\}$$

DEFINITION. — Corresponding to any extended real valued function f on Δ_1 , define, for all $x \in \Omega$,

$$\overline{\mathcal{H}}_{f,u}(x) = \text{Inf} \{ \nu(x) : \nu \in \Sigma_f \}$$

$$\underline{\mathcal{H}}_{f,u}(x) = - \overline{\mathcal{H}}_{-f,u}(x)$$

and

$$\overline{\mathcal{D}}_{f,u}(x) = \text{Inf} \{ \nu(x) : \nu \in \tilde{\Sigma}_f \}.$$

It is easy to see that Σ_f is a saturated family of super- u -harmonic functions [1]. Hence $\overline{\mathcal{H}}_{f,u}$ is either identically $\pm \infty$ or it is a u -harmonic function. Moreover, from the minimum principle, we deduce that $\overline{\mathcal{H}}_{f,u} \geq \underline{\mathcal{H}}_{f,u}$ on Ω .

Also $\overline{\mathcal{H}}_{f,u} \geq \overline{\mathcal{D}}_{f,u}$.

DEFINITION 2. — Let $u(\mathcal{R})$ be the class of extended real valued functions f on Δ_1 such that, $\overline{\mathcal{H}}_{f,u} = \underline{\mathcal{H}}_{f,u}$ and this function u -harmonic on Ω . For functions $f \in u(\mathcal{R})$, we denote $\mathcal{H}_{f,u} = \overline{\mathcal{H}}_{f,u} = \underline{\mathcal{H}}_{f,u}$.

LEMMA 3. — Every bounded μ_u -measurable function f on Δ_1 belongs to $u(\mathcal{R})$ and moreover

$$\mathcal{H}_{f,u} = \int f(h) \frac{h}{u} \mu_u(dh).$$

Proof. — The u -harmonic function $\sigma_f = \int f(h) \frac{h}{u} \mu_u(dh)$ satisfies,

$$\text{fine lim}_{x \rightarrow h} \sigma_f(x) = f(h) \quad \text{for} \quad \mu_u\text{-almost every } h \in \Delta_1$$

(Theorem 4). Hence, $\overline{\mathcal{H}}_{f,u} \leq \sigma_f \leq \underline{\mathcal{H}}_{f,u}$. This completes the proof.

PROPOSITION 1. — Let $\{f_n\}$ be an increasing sequence of extended real functions such that $\overline{\mathcal{H}}_{f_n,u} > -\infty$. Then,

$$\lim \overline{\mathcal{H}}_{f_n,u} = \overline{\mathcal{H}}_{f,u}.$$

Proof. — Since $\overline{\mathcal{H}}_{f_n,u} \leq \overline{\mathcal{H}}_{f,u}$, for every n , it is enough to show that $\overline{\mathcal{H}}_{f,u} \leq \lim_{n \rightarrow \infty} \overline{\mathcal{H}}_{f_n,u}$, when the limit is not $+\infty$.

Let $x_0 \in \Omega$. Given $\varepsilon > 0$, choose for every n , an element $\nu_n \in \Sigma_{f_n}$ such that

$$\overline{\mathcal{H}}_{f_n,u}(x_0) \geq \nu_n(x_0) - \frac{\varepsilon}{2^n}.$$

Consider $\omega = \lim \overline{\mathcal{H}}_{f_n,u} + \sum_{n=1}^{\infty} (\nu_n - \overline{\mathcal{H}}_{f_n,u})$. It is easily seen that ω is a super- u -harmonic function. Moreover $\omega \geq \nu_n$, for every n . Hence ω is lower bounded on Ω . Also, if E_{ν_n} is the set contained in Δ_1 such that $\mu_u(E_{\nu_n}) = 0$ and for all $h \in \Delta_1 - E_{\nu_n}$, $\text{fine lim inf}_{x \rightarrow h} \nu_n(x) \geq f_n(h)$, then,

$$\text{fine lim inf}_{x \rightarrow h} \omega(x) \geq f(h),$$

for all $h \in \Delta_1 - \bigcup_{n=1}^{\infty} E_{\nu_n}$. It follows that $\omega \in \Sigma_f$. Hence $\omega \geq \overline{\mathcal{H}}_{f,u}$. But,

$$\overline{\mathcal{H}}_{f,u}(x_0) \leq \omega(x_0) \leq \lim \overline{\mathcal{H}}_{f_n,u}(x_0) + \varepsilon.$$

The proof is now completed easily.

The following proposition is proved easily.

PROPOSITION 2. — $u(\mathfrak{R})$ is a real vector space. Moreover, for $f, g \in u(\mathfrak{R})$, $\mathfrak{H}_{f,u} + \mathfrak{H}_{g,u} = \mathfrak{H}_{f+g,u}$.

LEMMA 4. — For any non-negative extended real valued function f on Δ_1 , $\overline{\mathfrak{H}}_{f,u} = 0$ is equivalent to the fact that $f = 0$ μ_u -almost everywhere.

Proof. — Suppose $f = 0$ except on a set of μ_u -measure zero. Let $\nu \in \Sigma_f$. Then clearly $\frac{1}{n} \nu \in \Sigma_f$, for all positive integers n . Hence $\overline{\mathfrak{H}}_{f,u} = 0$.

Conversely, suppose $\overline{\mathfrak{H}}_{f,u} = 0$. Let $A_n = \left\{ h : f(h) > \frac{1}{n} \right\}$. Then the characteristic function χ_n of $A_n \subset \Delta_1$ has the property that $\overline{\mathfrak{H}}_{\chi_n,u} = 0$. The lemma would be proved if we show that for any set $A \subset \Delta_1$, $\overline{\mathfrak{H}}_{\chi_A,u} = 0$ implies that $\mu_u^*(A) = 0$.

Let $\nu \in \Sigma_{\chi_A}$. That is, there exists a set E_ν of μ_u -measure zero such that $\liminf_{x \rightarrow h} \nu(x) \geq \chi_A(h)$, for all $h \in \Delta_1 - E_\nu$. Given $\varepsilon > 0$, let $V_\varepsilon = \{ x \in \Omega : \nu(x) > 1 - \varepsilon \}$. Then, V_ε is an open set and V_ε is not thin at any point of $h \in A - E_\nu$. Now,

$$\frac{u\nu}{1 - \varepsilon} \geq R_u^{V_\varepsilon} \geq \int h \chi_{A-E_\nu}(h) \mu_u(dh) = \int h \chi_A(h) \mu_u(dh).$$

This inequality is true for all $\varepsilon > 0$. Hence

$$\nu \geq \int \frac{h}{u} \chi_A(h) \mu_u(dh).$$

In turn, this inequality is true for all $\nu \in \Sigma_{\chi_A}$, and we deduce,

$$\overline{\mathfrak{H}}_{\chi_A,u} \geq \frac{1}{u} \int h \chi_A(h) \mu_u(dh).$$

Hence, if $\overline{\mathfrak{H}}_{\chi_A,u} = 0$, then $\int h \chi_A(h) \mu_u(dh) = 0$. Now, we deduce easily that $\mu_u^*(A) = 0$. This completes the proof.

THEOREM 6. — Every μ_u -summable function f on Δ_1 belongs to $u(\mathfrak{R})$ and moreover, $\mathfrak{H}_{f,u}(x) = \int f(h) \frac{h(x)}{u(x)} \mu_u(dh)$ on Ω .

Proof. — Suppose f is a non-negative μ_u -summable function on Δ_1 . For each positive integer n , if, $f_n = \inf(f, n)$, then

$f_n \in u(\mathcal{R})$ and $\mathcal{H}_{f_n, u} = \int f_n(h) \frac{h}{u} \lambda_u(dh)$. (Lemma 3). Hence, we have,

$$\begin{aligned} \overline{\mathcal{H}}_{f, u} &= \lim_{n \rightarrow \infty} \mathcal{H}_{f_n, u} \quad (\text{Proposition 1}) \\ &= \lim_{n \rightarrow \infty} \int f_n(h) \frac{h}{u} \mu_u(dh) \\ &= \int f(h) \frac{h}{u} \mu_u(dh). \end{aligned}$$

Also,

$$\int f(h) \frac{h}{u} \mu_u(dh) = \lim \mathcal{H}_{f_n, u} \leq \underline{\mathcal{H}}_{f, u}.$$

It follows that

$$f \in u(\mathcal{R}) \quad \text{and} \quad \mathcal{H}_{f, u} = \int f(h) \frac{h}{u} \mu_u(dh).$$

Now the proof is completed easily.

Remark. — It can be proved that any function $f \in u(\mathcal{R})$ is necessarily equal μ_u -almost everywhere to a μ_u -summable function and that $\mathcal{H}_{f, u}$ is precisely $\int f(h) \frac{h}{u} \mu_u(dh)$.

4. The Main Result.

THEOREM 7. — Let $f \geq 0$ be an extended real valued function on Δ_1 . Then, $\overline{\mathcal{D}}_{f, u} = \overline{\mathcal{H}}_{f, u}$.

Proof. — It is enough to show that $\overline{\mathcal{D}}_{f, u} \geq \overline{\mathcal{H}}_{f, u}$.

First of all consider a function $f \geq 0$ which is bounded, say $f \leq M$. Consider $\tilde{\Sigma}_f^M = \{\nu \in \tilde{\Sigma}_f : \nu \leq M\}$. We assert that $\overline{\mathcal{D}}_{f, u} = \text{Inf} \{\nu : \nu \in \tilde{\Sigma}_f^M\}$. For, suppose $\nu \in \tilde{\Sigma}_f$. Then $\nu_M = \text{inf}(\nu, M)$ is a super- u -harmonic function and satisfies

$$\text{fine lim sup}_{x \rightarrow h} \nu_M(x) \geq f(h),$$

for μ_u -almost every $h \in \Delta_1$. Hence, $\nu \geq \nu_M \geq \text{Inf} \{\nu : \nu \in \tilde{\Sigma}_f^M\}$. Hence $\overline{\mathcal{D}}_{f, u} \geq \text{Inf} \{\nu : \nu \in \tilde{\Sigma}_f^M\}$. The opposite inequality is obvious.

Now, let $\nu \in \tilde{\Sigma}_f^M$. Then, by Theorem 4, Remark 3, the

fine limit $\nu(x)$ exists for all $h \in \Delta_1 - E'_v$, where $\mu_u(E'_v) = 0$. But, by the defining property of $\nu \in \tilde{\Sigma}_f^M$, $\text{fine lim sup}_{x \succ h} \nu(x) \geq f(h)$ for all $h \in \Delta_1 - F_v$, where $\mu_u(F_v) = 0$. It follows that,

$$\text{fine lim inf}_{x \succ h} \nu(x) \geq f(h),$$

for all $h \in \Delta_1 - (E'_v \cup F_v)$. Hence, $\nu \geq \bar{\mathcal{H}}_{f,u}$. This is true for all $\nu \in \tilde{\Sigma}_f^M$ and we get that $\bar{\mathcal{D}}_{f,u} \geq \bar{\mathcal{H}}_{f,u}$.

Let us now consider any $f \geq 0$. Let, for every positive integer n , $f_n = \inf(f, n)$. Then, we have,

$$\bar{\mathcal{D}}_{f,u} \geq \lim \bar{\mathcal{D}}_{f_n,u} = \lim \bar{\mathcal{H}}_{f_n,u} = \bar{\mathcal{H}}_{f,u}.$$

This completes the proof of the theorem.

THEOREM 7. — *For every μ_u -summable function f on Δ_1 ,*

$$\text{fine lim}_{x \rightarrow h_0} \int f(h) \frac{h(x)}{u(x)} \mu_u(dh) = f(h_0),$$

for μ_u almost every $h_0 \in \Delta_1$.

Proof. — It is enough to prove the theorem assuming that $f \geq 0$. Define, for every $h_0 \in \Delta_1$,

$$\varphi'(h_0) = \text{fine lim sup}_{x \rightarrow h_0} \int f(h) \frac{h(x)}{u(x)} \mu_u(dh).$$

Let $\varphi = \sup(\varphi', f)$ and $\nu \in \tilde{\Sigma}_f$. Then, $\nu \geq \int f(h) \frac{h}{u} \mu_u(dh)$ and we see easily that the $\text{fine lim sup}_{x \rightarrow h} \nu(x) \geq \varphi(h)$, for μ_u -almost every $h \in \Delta_1$. It follows that $\nu \in \tilde{\Sigma}_\varphi$. This is true for all $\nu \in \tilde{\Sigma}_f$. Hence, $\bar{\mathcal{H}}_{f,u} \geq \bar{\mathcal{H}}_{\varphi,u}$. But $\bar{\mathcal{H}}_{f,u} = \mathcal{H}_{f,u} \leq \mathcal{H}_{\varphi,u}$. This implies that $\varphi \in u(\mathcal{R})$ and $\mathcal{H}_{\varphi,u} = \mathcal{H}_{f,u}$. Again, $\varphi - f \geq 0$ and $\mathcal{H}_{\varphi-f,u} = 0$. We get, from the Lemma 4, that, $\varphi = f$, μ_u -almost everywhere. Hence,

$$\text{fine lim sup}_{x \rightarrow h_0} \int f(h) \frac{h(x)}{u(x)} \mu_u(dh) \leq f(h_0)$$

for μ_u -almost every $h_0 \in \Delta_1$. But we have already proved that

the fine lim inf is $\geq f(h_0)$ for μ_u -almost every $h_0 \in \Delta_1$. Hence, we get,

$$\text{fine lim}_{x \rightarrow h_0} \int f(h) \frac{h(x)}{u(x)} \mu_u(dh) = f(h_0)$$

for μ_u -almost every $h_0 \in \Delta_1$, completing the proof of the theorem.

THEOREM 8. (Fatou-Naïm-Doob). *For any $\nu \in S^+$, $\frac{\nu}{u}$ has a finite limit at μ_u -almost every element of Δ_1 .*

Proof. — Let ν be the canonical measure on Δ_1 corresponding to the greatest harmonic minorant of ν . Let ν_1 (respectively ν_2) be the absolutely continuous (resp. singular) part of ν relative to μ_u . Let f be the Radon-Nikodym derivative of ν_1 relative to μ_u (f is defined upto a set of μ_u measure zero). Then

$$\nu = \nu_1 + \nu_2 + \nu_3$$

where ν_3 is a potential, $\nu_2 = \int h \nu_2(dh)$ and $\nu_1 = \int f(h) h \mu_u(dh)$.

Now, $\frac{\nu_1}{u}$ has the fine limit f (note that f is finite μ_u almost everywhere), for μ_u -almost every element of Δ_1 . Also, $\frac{\nu_2 + \nu_3}{u}$ has the fine limit zero at μ_u -almost every element of Δ_1 . This completes the proof of the theorem.

BIBLIOGRAPHY

- [1] M. BRELOT, *Lectures on Potential Theory*, Tata Institute of Fundamental Research, Bombay, (1960).
- [2] K. GOWRISANKARAN, *Ann. Inst. Fourier*, t. XIII, Fasc. 2, 307-356.
- [3] R. M. HERVÉ, *Ann. Inst. Fourier*, t. XII (1962), 415-571.

Manuscrit reçu le 21 juin 1966.

Kohur Gowrisankaran,
Tata Institute of Fundamental Research,
Bombay 5.