

ROBERT R. PHELPS

**Theorems of Krein Milman type for certain
convex sets of functions operators**

Annales de l'institut Fourier, tome 20, n° 2 (1970), p. 45-54

http://www.numdam.org/item?id=AIF_1970__20_2_45_0

© Annales de l'institut Fourier, 1970, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

THEOREMS OF KREIN-MILMAN TYPE FOR CERTAIN CONVEX SETS OF FUNCTIONS AND OPERATORS

by Robert R. PHELPS

Let X be a compact Hausdorff space and E a real [or complex] locally convex Hausdorff vector space. Denote by $C(X, E)$ the real [or complex] linear space of all continuous functions from X to E , provided with the topology of uniform convergence. (Thus, a typical neighborhood of 0 has the form

$$\{f \in C(X, E) : \sup_x p(f(x)) \leq 1\}$$

where p is a continuous seminorm on E .)

For any subset A of E we let

$$C(X, A) = \{f \in C(X, E) : f(X) \subset A\}.$$

It follows that if B is a bounded closed convex subset of E , then $C(X, B)$ is a bounded closed convex subset of $C(X, E)$. We denote by $\text{ext } S$ the set of extreme points of a given convex set S ; it is readily verified that

$$C(X, \text{ext } B) \subset \text{ext } C(X, B).$$

[It is known [2, p. 755] that this inclusion can be proper, even for four dimensional E . There are also examples where $\text{ext } C(X, B)$ is empty for every X ; for instance, if $E = c_0$ in the norm topology and B is its unit ball.] The main purpose of this note is to exhibit conditions under which the set $C(X, B)$ will be the closed convex hull $\overline{c_0} C(X, \text{ext } B)$ of this subset of extreme points.

[Note that, even for one dimensional B , the set $C(X, B)$ need not be compact, so the Krein-Milman theorem does not apply.] Our main result was proved in two special cases in [6] (Theorems 2.1 and 4.1), where applications were made to various convex sets of bounded (or of compact) linear operators from a Banach space into $C(X)$. The more general result of the present note may be applied to analogous sets of *weakly* compact operators. We give one such application, as well as two results which were overlooked in [6].

As in [6], the problem is handled in two steps. First, we consider a condition (D) (below) on a pair of spaces (X, A) , with X compact Hausdorff and A bounded in E , which implies that

$$C(X, \overline{\text{co}} A) = \overline{\text{co}} C(X, A).$$

[This formulation was first considered by G. Seever [7].] We then apply this to bounded closed convex subsets B of E such that (with $A = \text{ext } B$), the pair (X, A) satisfies condition (D) and $B = \overline{\text{co}} A$.

DEFINITION. — *A pair of Hausdorff spaces (X, A) is said to satisfy condition (D) if the following holds for each $n > 0$:*

Given nonempty open sets U_1, U_2, \dots, U_n in A and pairwise disjoint nonempty compact sets K_1, K_2, \dots, K_n in X , there exists $f \in C(X, A)$ such that $f(K_i) \subset U_i, i = 1, 2, \dots, n$.

Condition (D) is a sort of density property for the subspace $C(X, A)$ in the space A^X of all functions from X to A . Indeed, condition (D) implies that $C(X, A)$ is dense in the pointwise topology on the space A^X , while density of $C(X, A)$ in the compact-open topology implies condition (D). As noted in [6], if X is a totally disconnected compact space, then (X, A) satisfies condition (D) for any A . On the other hand, if A is arcwise connected (or even only « almost arcwise connected » [6]), then (X, A) satisfies (D) for any compact X .

THEOREM 1. — *Let E be a real or complex locally convex Hausdorff vector space, X a compact Hausdorff space and A a bounded subset of E . If (X, A) satisfies condition (D), then*

$$C(X, \overline{\text{co}} A) = \overline{\text{co}} C(X, A).$$

The proof of the theorem depends on the following two technical lemmas.

LEMMA 1. — Suppose that L is a continuous linear functional on $C(X, E)$. Then there exists a continuous seminorm p on E and a regular Borel positive measure μ on X such that $\mu(X) \leq 1$ and

$$|L(f)| \leq \int_X p(f(x)) d\mu(x) \quad (\text{for each } f \in C(X, E)).$$

LEMMA 2. — Suppose that X and A are as described in the statement of the theorem. Given a continuous seminorm p on E , $\varepsilon > 0$, a regular Borel probability measure μ on X and $f \in C(X, \overline{\text{co}} A)$, there exist $g \in \overline{\text{co}} C(X, A)$ and a compact subset $K \subset X$ such that

$$p(g(x) - f(x)) < \varepsilon \quad \text{for } x \in K \quad \text{and} \quad \mu(X \setminus K) < \varepsilon.$$

Assuming that these lemmas have been proved, the theorem follows readily. Indeed, since $C(X, A) \subset C(X, \overline{\text{co}} A)$ and since the latter is closed and convex, we have

$$\overline{\text{co}} C(X, A) \subset C(X, \overline{\text{co}} A).$$

To show equality, it suffices to show that for each $\varepsilon > 0$, each $L \in C(X, E)^*$ and each $f \in C(X, \overline{\text{co}} A)$, there exists $g \in \overline{\text{co}} C(X, A)$ with

$$\text{Re } L(g) > \text{Re } L(f) - \varepsilon.$$

Choose p and μ according to Lemma 1, and let $M = \sup\{p(a) : a \in A\}$. Choose $K \subset X$ and $g \in \overline{\text{co}} C(X, A)$ according to Lemma 2, with ε replaced by $\varepsilon/2(M + 1)$. It follows that

$$\text{Re } L(f) - \text{Re } L(g) \leq |L(f - g)| \leq \int_X p(f(x) - g(x)) d\mu(x).$$

The integral on the right is the sum of the integral over K and the integral over $X \setminus K$. From Lemma 2, the first summand is at most $\varepsilon/2(M + 1)$, while the second is at most $M\varepsilon/2(M + 1)$, hence the total is at most ε .

We now turn to the proof of Lemma 1. Since L is continuous on $C(X, E)$ it is bounded in absolute value by 1 on a

neighborhood of the form

$$\{f \in C(X, E) : p(f(x)) \leq 1, x \in X\},$$

where p is a continuous seminorm on E . Thus,

$$(*) \quad |L(f)| \leq \sup\{p(f(x)) : x \in X\}, \quad f \in C(X, E).$$

Let N denote the closed subspace $p^{-1}(0)$ and consider the space E/N , normed by the quotient norm $\|\cdots\|$ defined by p . Let φ denote the quotient map from E into $F = E/N$; the composition $f \rightarrow \varphi \circ f$ defines a linear mapping of $C(X, E)$ into $C(X, F)$ which satisfies

$$\|\varphi(f(x))\| = p(f(x))$$

for all $f \in C(X, E)$, $x \in X$. The space $C(X, F)$ has the norm

$$\|g\| = \sup\{\|g(x)\| : x \in X\}.$$

It follows from (*) that the formula $J(\varphi \circ f) = L(f)$ defines a continuous linear functional J of norm at most 1 on the subspace $\varphi \circ C(X, E)$ of $C(X, F)$, and we can extend J to a functional of norm at most 1 on all of $C(X, F)$. At this point we could apply known results, which represent $C(X, F)^*$ in terms of dominated vector valued measures [4, p. 387], but we prefer to use the following direct (and simple) proof which was kindly furnished us by Dr. Erik Thomas. Let us define, for $h \in C(X)$, $h \geq 0$,

$$(**) \quad \mu(h) = \sup \{ |J(g)| : g \in C(X, F), \\ \|g(x)\| \leq h(x) \text{ for } x \text{ in } X \}$$

It is straightforward to verify that $\mu(h) < \infty$, that $\mu(\lambda h) = \lambda \mu(h)$ for $\lambda > 0$, and that $\mu(h_1 + h_2) \geq \mu(h_1) + \mu(h_2)$ if $h_1, h_2 \geq 0$ are in $C(X)$. The reverse inequality follows easily once we have the following fact: If $h = h_1 + h_2$ ($h_i \geq 0$) and $\|g(x)\| \leq h(x)$ for all x in X , then there exists g_1, g_2 in $C(X, F)$ such that $g = g_1 + g_2$ and $\|g_i(x)\| \leq h_i(x)$, $i = 1, 2$ and $x \in X$. Indeed, let $V = \{x \in X : \|g(x)\| > 0\}$ and for x in V let

$$\alpha_1(x) = \min(1, h_1(x)/\|g(x)\|), \quad \alpha_2(x) = 1 - \alpha_1(x).$$

If we define $g_i(x) = \alpha_i(x)g(x)$ for $x \in V$, $= 0$ for $x \in X \setminus V$,

then $g = g_1 + g_2$, $\|g_i(x)\| \leq h_i(x)$ and $\|g_i(x)\| \leq \|g(x)\|$ ($x \in X$, $i = 1, 2$). (The last inequality shows that each g_i is continuous.) Thus, μ is additive, non negative and positive homogeneous on the positive cone in $C(X)$, hence can be considered as an integral with respect to a finite positive regular Borel measure, say μ , on X . Furthermore, from (**) it is obvious that $|J(g)| \leq \int \|g(x)\| d\mu(x)$ for all $g \in C(X, F)$ and that $\mu(1) = \|J\| \leq 1$. Finally, for $f \in C(X, E)$ we have $|L(f)| = |J(\varphi \circ f)| \leq \int \|\varphi(f(x))\| d\mu(x) = \int p(f(x)) d\mu(x)$, which completes the proof of Lemma 1.

We next give the proof of Lemma 2. For $x \in X$, let

$$V_x = \{y \in X : p(f(x) - f(y)) < \varepsilon/3\};$$

this is an open neighborhood of x , and we can choose x_1, \dots, x_n such that the collection $\{V_{x_1}, \dots, V_{x_n}\}$ covers X , and such that no proper subcollection covers X . An easy induction argument, using the regularity of μ , shows that we can find another cover $\{V_1, \dots, V_n\}$ of open sets V_i such that $V_i \subset V_{x_i}$ and such that $\mu(D) < \varepsilon$, where $D = \cup \{V_i \cap V_j : i, j = 1, 2, \dots, n; i \neq j\}$. Let

$$K_i = V_i \setminus \cup \{V_j : j \neq i\} = X \setminus \cup \{V_j : j \neq i\}, i = 1, 2, \dots, n.$$

Then each K_i is compact, nonempty and $K_i \cap K_j$ is empty if $i \neq j$. Furthermore, if $K = \cup K_i$, then K is compact and $X \setminus K \subset D$, hence $\mu(X \setminus K) < \varepsilon$. Now, for each $i = 1, 2, \dots, n$ we have $f(x_i) \in \overline{\text{co}} A$, hence we can find $u_i \in \text{co} A$, with $p[u_i - f(x_i)] < \varepsilon/3$, of the following form:

$$u_i = \sum_{k=1}^{m_i} \lambda_{ik} a_{ik}, \quad \{a_{ik}\}_{k=1}^{m_i} \subset A, \quad \lambda_{ik} > 0, \quad \sum_{i=1}^{m_i} \lambda_{ik} = 1$$

where each λ_{ik} is a rational number, $k = 1, 2, \dots, m_i$. We can assume that the numbers λ_{ik} have a common denominator $Q > 0$, so by allowing repetitions of the points a_{ik} and by relabelling, we have

$$u_i = Q^{-1} \sum_{k=1}^Q b_{ik}, \quad \{b_{ik}\}_{k=1}^Q \subset A, \quad i = 1, 2, \dots, n.$$

By property (D), for each $k = 1, 2, \dots, Q$, we can choose

$g_k \in C(X, A)$ such that

$$g_k(K_i) \subset \{\nu \in E : p(\nu - b_{ik}) < \varepsilon/3\}.$$

Let $g = Q^{-1} \sum_{k=1}^Q g_k$, so that $g \in \text{co } C(X, A)$.

Suppose that $x \in K$; then $x \in K_i$ for some i and

$$\begin{aligned} p[g(x) - u_i] &= p[Q^{-1} \sum g_k(x) - Q^{-1} \sum b_{ik}] \\ &\leq Q^{-1} \sum p[g_k(x) - b_{ik}] < \varepsilon/3. \end{aligned}$$

Since $K_i \subset V_i \subset V_{x_i}$, we have $p[f(x) - f(x_i)] < \varepsilon/3$. Thus,

$$\begin{aligned} p[g(x) - f(x)] &\leq p[g(x) - u_i] + p[u_i - f(x_i)] \\ &\quad + p[f(x_i) - f(x)] < \varepsilon, \end{aligned}$$

which completes Lemma 2.

COROLLARY 1. — *Suppose that B is a bounded closed convex subset of the locally convex space E , and that X is a compact Hausdorff space. Let $A \subset \text{ext } B$. If $B = \overline{\text{co}} A$ and if (X, A) satisfies condition (D), then*

$$\overline{\text{co}} C(X, A) = C(X, B);$$

in particular, the latter set is the closed convex hull of its extreme points.

The hypothesis in Corollary 1 that $B = \overline{\text{co}} A$ is obviously a necessary one for the conclusion; indeed, since

$$C(X, A) \subset C(X, \overline{\text{co}} A)$$

and since the latter is closed and convex, it contains $\overline{\text{co}} C(X, A)$. Thus, if $C(X, B) = \overline{\text{co}} C(X, A)$, then $C(X, B) \subset C(X, \overline{\text{co}} A)$, whence $B = \overline{\text{co}} A$.

In general, condition (D) is not a necessary one for the validity of the equality $C(X, \overline{\text{co}} A) = \overline{\text{co}} C(X, A)$. Consider, for instance, $X = [0, 1]$, $E = \mathbb{C}$ (complex plane) and

$$A = \{z : |z| < 1/4\} \cup \{z : 3/4 < |z| < 1\}.$$

Then $\overline{\text{co}} A = \{z : |z| \leq 1\}$ is compact, and the above equality holds, but it is easily seen that $C(X, A)$ is not even pointwise dense in A^X . If, however, A is the set of extreme points of $\overline{\text{co}} A$ — this is the situation we are mainly interested in — then there is a partial converse to Theorem 1.

THEOREM 2. — *If B is a compact convex subset of the locally convex space E and $A = \text{ext } B$ (so $B = \overline{\text{co}} A$), then the equality*

$$\overline{\text{co}} C(X, A) = C(X, \overline{\text{co}} A)$$

implies that $C(X, A)$ is pointwise dense in A^X .

We omit the proof, since it, closely parallels that of Theorem 3.1 of [6], in which E is a dual Banach space (in the weak* topology) and B is its unit ball. The same argument works in the general case, using the fact that each extreme point of B has a neighborhood base in B consisting of « slices » [3, p. 108].

We now consider some applications of the foregoing results to spaces of linear operators. Suppose that M is a real (resp. complex) Banach space and let $C(X)$ denote the real (resp. complex) continuous functions on the compact Hausdorff space X . The space $\mathcal{L}(M, C(X))$ (or simply \mathcal{L}) of all bounded linear operators from M into $C(X)$ is linearly isomorphic to the space $C(X, E)$, where $E = M_w^*$ is the space M^* in its weak* topology [5, p. 490]. The correspondence between an operator T in \mathcal{L} and a function f in $C(X, E)$ is defined by

$$(Tm)(x) = \langle m, f(x) \rangle, \quad (x \in X, m \in M).$$

Moreover, $\|T\| = \sup \{ \|f(x)\| : x \in X \} = \|f\|$.

Thus, the unit ball \mathcal{U} of \mathcal{L} may be identified with the subset $C(X, U^*)$ of $C(X, E)$, where U^* is the unit ball of M^* . This correspondence was used in [6] to obtain various corollaries to Theorem 1, which was proved there for this particular choice of E . Similarly, the subspace

$$\mathcal{L}_c = \mathcal{L}_c(M, C(X))$$

of all compact operators in \mathcal{L} can be identified with the subspace $C(X, M_n^*)$, of $C(X, E)$, where M_n^* is M^* in its norm topology [5], and Theorem 1 was also proved in [6] for this case. It is readily verified that the uniform topology on $C(X, E)$ carries over (under the correspondence indicated above) to the strong operator topology on \mathcal{L} , and that in $C(X, M_n^*)$ the uniform topology is the norm topology (norm defined as above) and this identifies on \mathcal{L}_c with the norm (or « uniform operator ») topology. The fact that Theorem 1

was proved for arbitrary E allows us to consider the case where $E = M_w^*$, the space M^* in its weak (i.e. $\sigma(M^*, M^{**})$) topology. Under the above correspondence, $C(X, M_w^*)$ is exactly the space $\mathcal{L}_{wc} = \mathcal{L}_{wc}(M, C(X))$ of all weakly compact operators from M into $C(X)$. The topology induced on \mathcal{L}_{wc} by the uniform topology on $C(X, M_w^*)$ is not one of the usual « operator » topologies, but is easily seen to be between the strong operator and norm topologies on \mathcal{L}_{wc} .

We will denote by \mathcal{U} , \mathcal{U}_c and \mathcal{U}_{wc} the unit ball of \mathcal{L} , \mathcal{L}_c and \mathcal{L}_{wc} respectively. These are, of course, the same as the sets $C(X, U^*)$, $C(X, U_n^*)$ and $C(X, U_w^*)$. An operator which corresponds to an element f of one of these sets such that $f(X) \subset \text{ext } U^*$ is called a *nice* (resp. nice compact, nice weakly compact) operator. They are of course, extreme points of the sets \mathcal{U} , \mathcal{U}_c and \mathcal{U}_{wc} respectively.

The next result is almost a direct application of Corollary 1 to the ball of weakly compact operators. The main point is to account for the difference between the two topologies involved.

PROPOSITION 1. — *Let M and $C(X)$ be as above, and let U^* be the unit ball of M^* . Suppose that there is a subset $A \subset \text{ext } U^*$ such that:*

- (i) *The pair (X, A_w) satisfies condition (D).*
- (ii) *U^* is the norm closed convex hull of A .*

Then the unit ball \mathcal{U}_{wc} of \mathcal{L}_{wc} is the strong operator closed convex hull of the nice weakly compact operators.

Proof. — Hypotheses (i) and (ii) allow us to apply Corollary 1 to obtain the equality $C(X, U_w^*) = \overline{\text{co}} C(X, A_w)$, where the closure is in the uniform topology of $C(X, M_w^*)$. Since $C(X, M_w^*) \subset C(X, M_w^{**})$, the uniform topology on the latter space induces a topology on $C(X, M_w^*)$ which is weaker than the original; we will call it the « strong » topology since it corresponds exactly to the strong operator topology on \mathcal{L}_{wc} . Thus, we want to show that $C(X, U_w^*)$ is the strong closed convex hull of $C(X, A_w)$, since the latter is clearly a subset of the nice weakly compact operators. But it is easily verified that (since U^* is weak* closed) $C(X, U_w^*)$ is strongly closed in $C(X, M_w^*)$, hence contains the strong closure of $\text{co } C(X, A_w)$, which in turn contains $\overline{\text{co}} C(X, A_w) = C(X, U_w^*)$.

The fact that in hypothesis (ii) above we used the norm closure instead of the weak closure (which Corollary 1 would have allowed) is no loss in generality, of course, since the set involved is convex.

Recall that a real or complex Banach space M is said to be *smooth* if for each point $x \in S(M) = \{x \in M : \|x\| = 1\}$ there exists a unique functional f_x in the unit sphere $S(M^*)$ of M^* such that $\operatorname{Re} f_x(x) = 1$. This is equivalent to Gateaux differentiability of the norm (at each nonzero point), and the functional f_x is the Gateaux differential of the norm at x .

THEOREM 3. — *Let M be a real or complex Banach space and X a compact Hausdorff space. In the real case, we assume that $\dim M > 1$.*

(a) *If M is smooth, then \mathcal{U} is the strong operator closed convex hull of the nice operators.*

(b) *If the norm in M is Fréchet differentiable at each nonzero point, then \mathcal{U}_{wc} [resp. \mathcal{U}_c] is the strong operator [resp. norm] closed convex hull of its nice operators.*

Proof. — (a) It is well known (and easily proved) that if M is smooth, then the map $x \rightarrow f_x$ defined above is continuous from $S(M)$ in its norm topology into $S(M^*)$ in its weak* topology. It is readily verified that U^* is the weak* closed convex hull of the image A of $S(M)$ under this map, and that $A \subset \operatorname{ext} U^*$. [In fact, A is known [1] to be norm dense in $S(M^*)$.] Since $S(M)$ is arcwise connected (in the real case this assertion obviously requires that $\dim M > 1$), the set A is arcwise connected in the weak* topology. Thus, (X, A) satisfies condition (D) so Corollary 1 yields the desired conclusion.

(b) The Fréchet differentiability of the norm in M implies that the derivative map $x \rightarrow f_x$ defined above is continuous from the norm topology on $S(M)$ into the norm topology on $S(M^*)$. With the same notation as in (a), the set A is norm arcwise connected and norm dense in $S(M^*)$, hence U^* is the norm closed convex hull of A and Proposition 1 [resp. Corollary 1] applies.

In the case when $M = C(X)$ for some compact Hausdorff space X , it is possible to obtain necessary and sufficient

conditions on X and Y that $\mathcal{U}_{wc} \subset \mathcal{L}_{wc}(C(X), C(Y))$ be the strong operator closed convex hull of the nice weakly compact operators. These conditions are the same as those in Theorem 4.6 of [6], and the methods for obtaining them are essentially the same. (We don't know, in this case, whether every extreme element of \mathcal{U}_{wc} is a nice operator.) Similar results hold in the real case for the set of positive normalized weakly compact operators.

The following problem arises in the context of Corollary 1: Suppose that $C(X, B) = \overline{\text{co}} \text{ ext } C(X, B)$. Must $\text{ext } B$ be nonempty?

[*Note added in proof:* J. Lindenstrauss (private communication) has answered this question in the negative by showing that there exists a normed linear space E , a nonempty convex closed and bounded subset $B \subset E$ and a nonempty compact Hausdorff space X such that $\text{ext } B$ is empty, but $C(X, B) = \overline{\text{co}} \text{ ext } C(X, B)$.]

BIBLIOGRAPHY

- [1] Erret BISHOP and R. R. PHELPS, The support functionals of a convex set, *Proc. Symp. Pure Math.* vol 7 (Convexity), A.M.S. (1963), p. 27-35.
- [2] R. M. BLUMENTHAL, J. LINDENSTRAUSS and R. R. PHELPS, Extreme operators into $C(K)$, *Pacific J. Math.* 15 (1965), p. 747-756.
- [3] N. BOURBAKI, *Espaces vectoriels topologiques*, Ch. 1 et 2, 2^e édition, Paris, 1966.
- [4] N. DINCULEANU, *Vector measures*, Berlin, 1967.
- [5] N. DUNFORD and J. T. SCHWARTZ, *Linear operators Part I*, (1958), Interscience.
- [6] P. D. MORRIS and R. R. PHELPS, Theorems of Krein-Milman type for certain convex sets of operators, *Trans. Amer. Math. Soc.* 150 (1970), 183-200.
- [7] G. SEEVER, Generalization of a theorem of Lindenstrauss (dittoed notes).

Manuscrit reçu le 6 mars 1970.

Robert R. PHELPS,
 Department of Mathematics,
 University of Washington
 Seattle, Wash, 98105
 (USA..)