

A. K. ROY

Closures of faces of compact convex sets

Annales de l'institut Fourier, tome 25, n° 2 (1975), p. 221-234

http://www.numdam.org/item?id=AIF_1975__25_2_221_0

© Annales de l'institut Fourier, 1975, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

CLOSURES OF FACES OF COMPACT CONVEX SETS

by **A.K. ROY**

1. Introduction.

It is well-known that one of the disconcerting facts in the theory of infinite-dimensional compact convex sets is that the closure of a face need not be a face. The main purpose of this paper is to determine necessary and sufficient conditions which ensure that this pathology does not occur for a given face. It should be emphasised that our results are purely individual in character. We do *not* characterise the class of compact convex sets which have the property that the closures of all their faces are again faces. (As a matter of fact, this appears to be a very difficult problem.) By way of applications, it is shown that several results scattered in the literature can be proved in a rather economical and uniform manner by our method.

We conclude by giving several characterizations of cases when face (C) is closed in a compact convex set K, for any closed convex subset C of K without core points. This generalises a recent result in [11]. Our method of proof is quite different.

It is a pleasure to thank Dr. A.J. Ellis for showing some interest in this investigation and for providing me with the example at the end of § 3.

2. Definitions & Notations.

We will work with a fixed compact convex set K in a locally convex Hausdorff topological vector space E defined over the reals

R. We assume throughout that K is "regularly embedded" in E in the sense defined in [1].

Following [1], we let $\partial_e K$ be the set of extreme points of K and let $C(K)$, $P(K)$ and $A(K)$ denote, respectively, the space of continuous functions, the cone of continuous convex functions and the space of continuous affine functions, on K . Let $M_1^+(K)$ denote the convex set of probability measures on K equipped with the weak* topology induced on it by $M(K)$, the dual of $C(K)$.

For each $x \in K$, we write

$$M_x = \{\mu \in M_1^+(K) : \mu(a) = a(x), \forall a \in A(K)\}$$

which is a non-empty weak* compact convex subset of $M_1^+(K)$. Let Z_x denote the set of *maximal or boundary measures* [1] in M_x .

If $f \in C(K)$, we define

$$\hat{f}(x) = \inf \{h(x) : h \in A(K), h \geq f\};$$

which is the least upper semicontinuous (u.s.c) concave majorant of f and, dually, we define \check{f} as the greatest convex minorant of f .

If C is a proper compact convex subset of K , we define for each $\alpha \geq 1$,

$$D_\alpha(C) = (\alpha C - (\alpha - 1)K) \cap K$$

and by face (C) we mean the σ -compact set $\bigcup_{n=1}^{\infty} D_n(C)$. We recall [2 : page 99] that face (C) is the smallest, not necessarily closed, face of K containing C .

If f is a function defined on K and S is a subset of K , we consistently employ the notation $f(S) \leq \alpha$ to mean $f(x) \leq \alpha$ for all $x \in S$. A similar meaning should be given to $f(S) = 0$.

3. Conditions for the closure of a face to be a face.

Let $F \subset K$ be a face and let $a \in A(K)$ be such that $a \leq 0$ on F , and hence on \bar{F} . The theorem in this section is motivated by the following simple observation :

$$\widehat{0 \vee a}(x) = 0 \text{ for all } x \in F.$$

This follows from the fact (see [1]) that

$$\widehat{0 \vee a}(x) = \sup \{ \mu(0 \vee a) : \mu \text{ discrete, } \mu \in M_x \}.$$

However, since $\widehat{0 \vee a}$ is u.s.c. we cannot, in general, assert that $\widehat{0 \vee a}(x) = 0$ for all $x \in \bar{F}$. But this is the case if and only if \bar{F} is also a face.

Let

$$F^* = \{a \in A(K) : a(F) \leq 0\}$$

and

$$(F^*)_* = \{x \in K : a(x) \leq 0 \forall a \in F^*\}.$$

Then we have the following.

LEMMA 3.1. $\bar{F} = (F^*)_*$

We omit the proof which is a simple application of the Hahn-Banach (separation) theorem. We will also need the following simple result.

LEMMA 3.2. — Let $f \in P(K)$ and let $\{f_n\}$ be a sequence of functions in $P(K)$ converging uniformly to f . Then $\{\hat{f}_n\}$ converges uniformly to f .

This is an obvious consequence of the fact that $f - \epsilon \leq g \leq f + \epsilon$ implies $\hat{f} - \epsilon \leq \hat{g} \leq \hat{f} + \epsilon$ for any $\epsilon > 0$.

Adopting the terminology of [6], we say that F^* is *perfect* if for any $a \in F^*$ and $\epsilon > 0$, there exists $a_\epsilon \in F^*$ such that $0, a \leq a_\epsilon + \epsilon$.

We can now state the main result of this section as follows :

THEOREM 3.3. — Let $F \subseteq K$ be a proper face. Then the following are equivalent :

- (1) \bar{F} is a face.
- (2) F^* is perfect.
- (3) $\widehat{0 \vee a}(\bar{F}) = 0 \forall a \in F^*$.

(4) $\widehat{0 \vee f}(\bar{F}) = 0 \ \forall f \in P(K)$ such that $f(F) \leq 0$.

(5) If $f - g, f \in P(K)$ with $f(F) \leq g(F)$, then $\widehat{f \vee g} = g$ on \bar{F} .

Comments 1. – If F is assumed to be closed, then the equivalence of (1) and a result similar to (2) has been proved in [4] by means of the “polar calculus”. However, our proof, which is an adaptation to this context of an argument in [8], and formulation are somewhat different.

2. – We should note that the statements (2) – (5) have obvious “duals” : for example, the dual of (2) is $0 \wedge a(\bar{F}) = 0 \ \forall a \in F^0$ where

$$F^0 = \{a \in A(K) : a(F) \geq 0\}.$$

Proof of Theorem 3.3

(1) \Rightarrow (2). Suppose F^* is not perfect. Then $\exists a_0 \in F^*$ and $\epsilon_0 > 0$ such that $\forall b \in F^*$,

either $a_0 \not\leq b + \epsilon_0$
 or $0 \not\leq b + \epsilon_0$ } (α)

If $A(K)^+$ denotes the positive cone in $A(K)$, define

$$U = \{a \in A(K) : \|a\| \leq \epsilon_0\}$$

and

$$H = \{(b - p, b - q) : b \in F^*, p, q \in A(K)^+\}.$$

Then (α) can be restated as

$$(a_0, 0) + (u_1, u_2) \notin H, \forall u_1, u_2 \in U.$$

This implies that $(a_0, 0) \notin \bar{H}$ and hence by the Hahn-Banach theorem, $\exists \varphi \in (A(K) \times A(K))^*$ such that

$$\sup \varphi(\bar{H}) < \varphi(a_0, 0). \quad (\beta)$$

H being a cone, (β) says that $\varphi \leq 0$ on H . Now, we can write

$$\varphi = \varphi_1 + \varphi_2 \quad \text{where } \varphi_i \in A(K)^* \ (i = 1, 2)$$

and

$$\varphi_1(a) = \varphi(a, 0), \varphi_2(b) = \varphi(0, b) \quad \text{for } a, b \in A(K).$$

If $c \in A(K)^+$ then $(-c, 0) \in H$ and hence $\varphi_1(-c) = \varphi(-c, 0) \leq 0$, showing that $\varphi_1 \geq 0$. Similarly, $\varphi_2 \geq 0$ and thus (by [13]),

$\varphi_i(a) = \lambda_i a(x_i) \forall a \in A(K)$, for some $\lambda_i \in R^+$ and $x_i \in K (i = 1, 2)$.

If $a \in F^*$, $(a, a) \in H$ and therefore

$$\begin{aligned} 0 &\geq \varphi(a, a) = \varphi_1(a) + \varphi_2(a) \\ &= \lambda_1 a(x_1) + \lambda_2 a(x_2) \\ &= a(\lambda_1 x_1 + \lambda_2 x_2), \end{aligned}$$

showing, by lemma 3.1, that $\frac{\lambda_1 x_1 + \lambda_2 x_2}{\lambda_1 + \lambda_2} \in \bar{F}$. But by (β) ,

$$0 < \varphi(a_0, 0) = \varphi_1(a_0, 0) = \lambda_1 a_0(x_1)$$

and this shows, again by lemma 3.1, that $x_1 \notin \bar{F}$, so \bar{F} is not a face.

(2) \Rightarrow (3). If $a \in F^*$ and $\epsilon > 0$ then by assumption there exists $a_\epsilon \in F^*$ such that

$$0, a \leq a_\epsilon + \epsilon$$

which implies that

$$\widehat{0 \vee a}(x) \leq a_\epsilon(x) + \epsilon \leq \epsilon, \forall x \in \bar{F}$$

and since ϵ is arbitrary, we can conclude that $\widehat{0 \vee a}(\bar{F}) = 0$.

(3) \Rightarrow (4). If $a_1, a_2 \in F^*$ then for each $x \in K$,

$$0 \vee a_1(x) \vee a_2(x) \leq 0 \vee a_1(x) + 0 \vee a_2(x)$$

and hence

$$\widehat{0 \vee a_1 \vee a_2} \leq \widehat{0 \vee a_1 + 0 \vee a_2} \leq \widehat{0 \vee a_1} + \widehat{0 \vee a_2}$$

by the subadditivity of the \wedge function and it follows that $0 \vee a_1 \vee a_2(\bar{F}) \leq 0$. By induction, this is true for any finite number of $a_i \in F^*$. If $f \in P(K)$ with $f(\bar{F}) \leq 0$, we know from [1] that f can be approximated uniformly by an increasing sequence of functions of the form $a_1^{(n)} \vee \dots \vee a_k^{(n)}$ where $a_i^{(n)} \in F^*$ for $i = 1, \dots, k$. Therefore, by lemma 3.2, $(0 \vee a_1^{(n)} \vee \dots \vee a_k^{(n)})^\wedge$ increases monotonically to $\widehat{0 \vee f}$ and it follows that $\widehat{0 \vee f}(\bar{F}) = 0$.

(4) \Rightarrow (5). If $f(F) \leq g(F)$ then $(f - g)(F) \leq 0$ and since $f - g \in P(K)$, by (4) $\widehat{0 \vee (f - g)} = 0$ on \bar{F} . But

$$f \vee g = 0 \vee (f - g) + g$$

and therefore

$$\widehat{f \vee g} = \widehat{0 \vee (f - g) + g} \leq \widehat{0 \vee (f - g)} + \hat{g} = g \text{ on } \bar{F}$$

and (5) follows.

(5) \Rightarrow (3). Obvious.

(3) \Rightarrow (1). Let $x \in \bar{F}$ and consider $\mu \in M_x$. Suppose, if possible, that $\text{supp. } (\mu) \setminus \bar{F} \neq \emptyset$, i.e. $\exists y \in \text{supp. } (\mu)$, $y \notin \bar{F}$. By the Hahn-Banach theorem, $\exists a \in A(K)$ such that

$$a(\bar{F}) \leq 0 < a(y).$$

So $a \in F^*$. By continuity, there exists a neighbourhood U of y with $U \cap \bar{F} = \emptyset$ and $a(U) \geq \alpha > 0$ for some α .

Now,

$$\begin{aligned} \widehat{0 \vee a}(x) &= \sup \{ \lambda(0 \vee a) : \lambda \in M_x \} \\ &\geq \int (0 \vee a) d\mu \geq \int_U (0 \vee a) d\mu \geq \alpha \mu(U \cap \text{supp.}(\mu)) > 0, \end{aligned}$$

which contradicts (3).

This completes the proof of theorem 3.3.

COROLLARY 3.4. — *If $F \subseteq K$ is a face and \bar{F} is also a face, then*

$$\bar{F} = \bigcap_{a \in F^*} (\widehat{0 \vee a})^{-1}(0)$$

Remark. — Suppose that \bar{F} is a proper face where $F \subseteq K$ is a face. If $(\bar{F})'$ denotes the *complementary set* of \bar{F} , i.e. the union of all faces disjoint from \bar{F} , then it is clear that $(\bar{F})' \subseteq F'$. It is natural to enquire whether this inclusion is an equality. That it is *not*, is shown by the following simple example :

$$K = M_1^+[0, 1], F = \{ \mu \in M_1^+[0, 1] : \mu[0, a] = \mu[b, 1] = 0 \}$$

where $0 < a < b < 1$. It is clear that F is a face as is

$$\bar{F} = \{\mu \in M_1^+[0, 1] : \text{supp. } (\mu) \subseteq [a, b]\}.$$

Now, $\epsilon_a \in F'$ but $\epsilon_a \notin (\bar{F})'$ as $\epsilon_a \in \bar{F}$, showing that $(\bar{F})' \subsetneq F'$.

If $\partial_e K$ is closed then the necessary and sufficient condition for the closure of a face $F \subseteq K$ to be a face is expressed below in a different way. This has the advantage that it gives a rather explicit description of \bar{F} .

Let $S = \bar{F} \cap \partial_e K$ and define

$$T = \{f \in C(\partial_e K) : 0 \leq f \leq 1, f(S) = 1\}.$$

Then we have the following

THEOREM 3.5. — *Assume $\partial_e K$ closed. Then \bar{F} is a face if and only if $\bigcap_{f \in T} [f = 1]$ is closed. When this condition is satisfied,*

$$\bar{F} = \bigcap_{f \in T} [f = 1].$$

This result has been extracted from [10] and since its proof is essentially the same as in [10], modulo some trivial details, we omit it.

An example. — There are of course several examples in the literature showing that the closure of a face may fail to be a face (see, for instance, [2]). However, to the best of our knowledge, all these examples deal with compact convex sets, usually Choquet simplexes, with non-closed boundaries. We now present an example to show that this pathology can occur even if the boundary is closed.

Let \mathcal{A} be the disc-algebra and let K be its state space. Following [7], we let $Z = \text{conv}(K \cup -iK)$. Then K is just the probability measures on the unit circle Γ , so that $\partial_e K$ and $\partial_e Z$ are both closed. Every point of $\partial_e Z$ is a split face of Z and so if $E \subseteq \partial_e Z$, the norm-closed convex hull of E (in $A(Z)^*$) is a norm-closed split face of Z (by the L -ideal theory of [3]). However, take $E = \gamma \cup -i\gamma$ where γ is a proper arc of Γ of length > 0 . Using the fact that K is a simplex, it is easy to check that $\text{conv}(E)$ is a face of Z . However, γ is not a peak set for \mathcal{A} since any function in \mathcal{A} which is constant

on γ is necessarily constant on Γ , so that $\overline{\text{conv}}(E)$ (w^* -closure) is not a face of Z (by Theorem 2 of [7]).

4. Applications.

We will now give some applications of the results established in the last section.

PROPOSITION 4.1. — *If K has the property that \hat{f} is continuous for all $f \in P(K)$, then the closure of every face F in K is again a face.*

Proof. — Let $a \in F^*$. As we have already remarked, $\widehat{0 \vee a}(F) = 0$. But $0 \vee a \in P(K)$ and therefore $\widehat{0 \vee a}$ is continuous by assumption. Thus $\widehat{0 \vee a}(\bar{F}) = 0$ and we can use (3) of Theorem 3.3 to conclude that \bar{F} is a face.

Remark. — This result was first proved in [10] by a more elaborate method.

DEFINITION 4.2 (after [12]). — *A compact convex set K has the strong equal support property (s.e.s.p. for short) if (i) $\partial_e K$ is closed and (ii) for any $x \in K$ and $\mu, \nu \in Z_x$ we have $\text{supp.}(\mu) = \text{supp.}(\nu)$.*

We now prove

PROPOSITION 4.3. — *If K has the strong equal support property, then the closure of every face F in K is again a face.*

Proof. — If $S = \bar{F} \cap \partial_e K$ then it is rather easy to show from the defining property of a face that $\bar{F} = \overline{\text{conv}}(S)$. Let $a \in F^*$. Then $\widehat{0 \vee a}(F) = 0$. If $x \in \bar{F}$, there exists a probability measure μ on S representing x (See [13]) : μ is obviously maximal since $S \subseteq \partial_e K$, hence all $\lambda \in Z_x$ have their supports in S by the s.e.s.p. But

$$\begin{aligned} \widehat{0 \vee a}(x) &= \lambda_1(0 \vee a) \text{ for some } \lambda_1 \in Z_x \\ &= 0 \end{aligned}$$

and hence \overline{F} is a face by Theorem 3.3.

PROPOSITION 4.4. — *Let K be a Choquet simplex. If $F \subseteq K$ is a face then \overline{F} is a face if and only if $\partial_e \overline{F} \subseteq \partial_e K$.*

Proof. — If \overline{F} is a face then by the Krein-Milman theorem, $\partial_e \overline{F}$ is non-empty and it is clear that $\partial_e \overline{F} \subseteq \partial_e K$.

On the other hand, suppose F is a face with the property that $\partial_e \overline{F} \subseteq \partial_e K$. Consider $a \in F^*$. If $x \in \partial_e \overline{F} \subseteq \partial_e K$, by Herve's criterion [1],

$$\widehat{0 \vee a}(x) = (0 \vee a)(x) = 0.$$

But K being a simplex, $\widehat{0 \vee a}$ is an u.s.c. affine function (by the Choquet-Meyer theorem), hence by the Bauer maximum principle, $\widehat{0 \vee a}(\overline{F}) = 0$ and we see that \overline{F} is a face.

COROLLARY 4.5. — *If K is a Choquet simplex and if S is a compact subset of $\partial_e K$ then $\text{conv}(S)$ is a face of K .*

Proof. — This is immediate from the preceding Proposition, once we observe that $F = \text{conv}(S)$ is a face in K and that $\partial_e \overline{F} = S \subseteq \partial_e K$.

Remarks 1. — It would be interesting to know whether Prop. 4.4 extends to compact convex sets with the equal support property [12].

2. — In [15], the following result of Mokobodzki is proved : *If K is a Choquet simplex and if B is a compact convex subset of K with $\partial_e B \subseteq \partial_e K$ then B is a face of K .*

This again follows immediately from the preceding discussion once we note that $F = \text{conv}(\partial_e B)$ is a face and that $\partial_e \overline{F} = \partial_e B \subseteq \partial_e K$.

5. Compactness of face (C).

Considerations of subsets of K of the form *face* (x) , $x \in K$, have proved useful in several contexts : for example, they are important

in the local version of Choquet's Uniqueness Theorem [9] and in Wil's proof of the existence and uniqueness of central measures for points of K (see [1]). Their usefulness is also suggested by the following simple result :

PROPOSITION 5.1. — *If K is metrisable then every closed face F of K has the form $\text{face}(x)$ for some $x \in K$.*

Proof. — The metrisability implies that K , and hence F , is separable. Let $\{k_n\}$ be a dense subset of F and define

$$x = \sum_{n=1}^{\infty} \frac{1}{2^n} k_n$$

It is clear that this series defines an element x of F , and that $k_n \in \text{face}(x) \subseteq F$; it follows that $\bar{F} = \text{face}(x)$.

In view of the preceding remarks, it is natural to look for conditions which ensure that $\text{face}(x)$ is closed. This was recently done in [11] where it is proved, among other things, that $\text{face}(x)$ is closed iff $\text{face}(x) = D_n(x)$ for some n (see § 2 for the definition of $D_n(x)$). We propose to generalize this result in the theorem below. It should be pointed out that the proof of the implication (1) \Rightarrow (2) in this theorem follows an argument used in [14] in a different situation. We denote by $P(K)^+$ the cone of non-negative continuous convex functions on K .

THEOREM 5.2. — *Let C be a proper compact convex subset of K without core points. Then the following are equivalent :*

- 1) *face(C) is closed.*
- 2) *face(C) = $D_n(C)$ for some n .*
- 3) *If $f_m \in P(K)^+$ and $\lim_{m \rightarrow \infty} f_m(u) = 0$ uniformly for $u \in C$ then $\lim_{m \rightarrow \infty} \hat{f}_m(y) = 0$ uniformly for $y \in \text{face}(C)$.*
- 4) *face(C) is norm-closed in the space $A(K)^*$.*

Moreover, all the above statements are implied by
 (*) : $\text{lin } M_c^+$ is a norm-closed (or weak*-closed) subspace of $M(K)$, where $M_c^+ = \{\mu \in M_1^+(K) : \text{resultant } (\mu) \in C\}$.

Proof. – (1) \Rightarrow (2). Suppose face (C) is closed and face (C) $\neq D_n(C)$ for all n . This means that given any $n \in \mathbb{N}^+$ (= set of positive integers), $\exists y_n \in \text{face (C)}$ such that $y_n \not\leq n2^n C$. (By this is meant that

$$y_n \not\leq n2^n u, \forall u \in C).$$

Define $y = \sum_{n=1}^{\infty} 2^{-n} y_n$. Now $y \in \text{face (C)}$ as this set is closed by assumption. But then $y \not\leq mC$ for all $m \in \mathbb{N}^+$ which is a contradiction as $\text{face (C)} = \bigcup_{m=1}^{\infty} D_m(C)$.

(2) \Rightarrow (3). By (2), if $y \in \text{face (C)}$ then $y \leq nu$ for some $u \in C$, hence $\hat{f}_m(y) \leq n \hat{f}_m(u)$ as \hat{f}_m is a concave function and thus $\lim_{m \rightarrow \infty} \hat{f}_m(y) = 0$ uniformly on face (C) if $\lim_{m \rightarrow \infty} \hat{f}_m(u) = 0$ uniformly on C.

(3) \Rightarrow (2). Suppose $\text{face (C)} \neq D_n(C) \forall n \in \mathbb{N}^+$. This means that given $n \in \mathbb{N}^+$, $\exists y_n \in \text{face (C)}$ such that $y_n \not\leq nC$ but $y_n \leq n^{m(n)}u_n$ for some $u_n \in C$ and for some sufficiently large $m(n) \in \mathbb{N}^+$. Since $(nC - y_n) \cap \tilde{K} = \emptyset$, where \tilde{K} is the (closed) cone generated by K, a standard Hahn-Banach argument shows that $\exists a_n \in A(K)^+$ such that

$$a_n(y_n) > n a_n(u) \forall u \in C$$

and

$$a_n(y_n) \leq n^{m(n)} a_n(u_n)$$

These inequalities imply that $\sup \{a_n(u) : u \in C\} > 0$

and

$$a_n(y_n) > n \sup \{a_n(u) : u \in C\}.$$

Let

$$b_n = a_n/n \sup \{a_n(u) : u \in C\}$$

Then

$$b_n \in A(K)^+ \text{ and if } u \in C,$$

$$b_n(u) = a_n(u)/n \sup \{a_n(u) : u \in C\} \leq \frac{1}{n}$$

showing that $\lim_{n \rightarrow \infty} b_n(u) = 0$ uniformly for $u \in C$: however, $b_n(y_n) > 1$ and so b_n does not tend to zero uniformly on face (C), contradicting (3).

(2) \Rightarrow (1). Obvious.

(2) \Rightarrow (4). Obvious.

(4) \Rightarrow (2). By the regular embedding of K in $A(K)^*$, we can regard each $x \in K$ as a member of the unit ball of $A(K)^*$. As a norm-closed subset of the complete metric space $A(K)^*$, face (C) is complete and hence by the Baire Category Theorem, some $D_{n_0}(C)$ must have non-empty relative interior, i.e. there exists some $y_0 \in D_{n_0}(C)$ such that for some neighbourhood of the origin,

$$U = \{u \in A(K)^* : \|u\| < \eta\}$$

$$(y_0 + U) \cap \text{face } (C) \subseteq D_{n_0}(C).$$

Let $y \in \text{face } (C)$ and define

$$z = \frac{\eta}{2 + \eta} y + \frac{2}{2 + \eta} y_0 \in \text{face } (C)$$

Then
$$\|z - y_0\| = \frac{\eta}{2 + \eta} \|y - y_0\| \leq \frac{2\eta}{2 + \eta} < \eta$$

and hence
$$z \in (y_0 + U) \cap \text{face } (C) \subseteq D_{n_0}(C).$$

Therefore, for some $c \in C$ and $k \in K$,

$$\frac{\eta}{2 + \eta} y + \frac{2}{2 + \eta} y_0 = n_0 c - (n_0 - 1) k$$

or,
$$c = \frac{1}{n_0} \cdot \frac{\eta}{2 + \eta} y + \left(1 - \frac{1}{n_0} \cdot \frac{\eta}{2 + \eta}\right) k'$$

where
$$k' = \frac{\left(1 - \frac{1}{n_0}\right) k + \frac{1}{n_0} \cdot \frac{2}{2 + \eta} y_0}{1 - \frac{1}{n_0} \cdot \frac{\eta}{2 + \eta}} \in K.$$

Thus,
$$y \in D_\alpha(C) \quad \text{where} \quad \alpha = \left(1 + \frac{2}{\eta}\right) n_0$$

and this implies (2).

This completes the proof of the equivalence of (1), (2), (3) and (4). As far as statement (*) is concerned, first note that since

M_c^+ is a w^* compact subset of $M_i^+(K)$, by a known result [1 : page 112] $\text{lin } M_c^+$ is w^* closed iff $\text{lin } M_c^+$ is norm-closed. Now the proof follows exactly the argument used in [11] to prove (vi) \Rightarrow (i) in Theorem 1.9 of that paper.

Remarks 1. – The use of the Baire Category Theorem above was suggested by the proof of a similar result in [5]. The argument proving (1) \Rightarrow (2) could also be used here.

2. – We have not been able to decide whether any of the first four statements in Theorem 5.2 implies (*).

BIBLIOGRAPHY

- [1] E.M. ALFSEN, Compact convex sets and boundary integrals, *Ergebnisse der Mathematik*, Springer-Verlag, Berlin, 1971.
- [2] E.M. ALFSEN, On the geometry of Choquet simplexes, *Math. Scand.*, 15 (1964), 97-110.
- [3] E.M. ALFSEN & E.G. EFFROS, Structure in real Banach spaces, Part I & II, *Annals of Math.*, 96, No. 1 (1972), 98-173.
- [4] L. ASIMOW, Exposed faces of dual cones and peak-set criteria for function spaces, *Journal of Function Analysis*, vol. 12, No. 4 (1973).
- [5] F. DEUTSCH & R.J. LINDAHL, Minimal extremal subsets of the unit sphere, *Math. Annalen*, 197 (1972).
- [6] A.J. ELLIS, On faces of compact convex sets and their annihilators, *Math. Annalen*, 184 (1969).
- [7] A.J. ELLIS, Split faces in function algebras, *Math Annalen*, 195 (1972).
- [8] G. JAMESON, Nearly directed subspaces of partially ordered linear spaces, *Proc. Edinburgh Math. Soc.*, (2) 16 (1968).
- [9] J. KOHN, Barycentres of unique maximal measures, *J. of Funct. Analysis*, 6 (1970).

- [10] A. LIMA, On continuous convex functions and split faces, *Proc. London Math. Soc.*, (3) 25 (1972).
- [11] A. LIMA, Closed faces with internal points, *Preprint series – Matematisk institutt*, Universiteteti Oslo (1972).
- [12] J.N. McDONALD, Compact convex sets with the equal support property, *Pac. J. of Math.*, vol. 37, No. 2 (1971).
- [13] R. PHELPS, Lectures on Choquet's Theorem, Van Nostrand, Princeton (1960).
- [14] M. RAJAGOPALAN & A.K. ROY, Maximal core representing measures and generalized polytopes, *Quart. J. of Math.*, Oxford, vol. 25, no. 99 (1974).
- [15] M. ROGALSKI, Etude du quotient d'un simplexe par une face fermée. . . relation d'équivalence, *Seminaire Brelot – Choquet – Deny* (Theorie du Potentiel), 1967/68, No. 2.

Manuscrit reçu le 11 avril 1974
accepté par G. Ghoquet.

A.K. ROY,
Tata Institute of Fundamental Research
Homi Bhabha Road
Bombay – 400005 (Inde).