

ANNALES DE L'INSTITUT FOURIER

MICHAEL DRMOTA

Irregularities of continuous distributions

Annales de l'institut Fourier, tome 39, n° 3 (1989), p. 501-527

http://www.numdam.org/item?id=AIF_1989__39_3_501_0

© Annales de l'institut Fourier, 1989, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

IRREGULARITIES OF CONTINUOUS DISTRIBUTIONS

by Michael DRMOTA

1. INTRODUCTION

Let $x : [0, 1] \rightarrow \mathbf{R}^K / \mathbf{Z}^K$ be a continuous function with finite arc length

$$(1) \quad s(x) = \int_0^1 |dx(t)|.$$

($\mathbf{R}^K / \mathbf{Z}^K$ denotes the K -dimensional unit torus. As usual it can be identified with the K -dimensional unit cube $[0, 1)^K$.) Such a function can be interpreted as a particle's movement on $\mathbf{R}^K / \mathbf{Z}^K$. It is of some interest to consider a measure for the irregularity of the distribution behaviour of this particle, the discrepancy

$$(2) \quad D^{(\mathcal{R})}(x) = \sup_R \left| \int_0^1 \chi_R(x(t)) dt - \lambda_K(R) \right|,$$

where the supremum is taken over all rectangles $R \subseteq \mathbf{R}^K / \mathbf{Z}^K$ of the form $R = [a_1, b_1] \times \cdots \times [a_K, b_K]$ with $\max_{1 \leq k \leq K} (b_k - a_k) \leq 1$. (χ_M denotes the characteristic function of a set M and λ_K the K -dimensional Lebesgue measure. For details see Section 2.1.) In other words, $D^{(\mathcal{R})}(x)$ measures the maximal difference between the time of the particles stay in a rectangle R

Key-words : Irregularities – Uniform distribution – Uniformly distributed functions – Discrepancy.

A.M.S Classification : 10K05 (11K05).

and the volume of R . R. J. Taschner [10] was the first who noticed that the discrepancy cannot be too small. He proved that for $K \geq 2$

$$(3) \quad D^{(\mathcal{R})}(x) > c_K s(x)^{-(1+1/(K-1))}$$

provided that $s(x) \geq 1$. (Another more general proof can be found in [5].)

It is interesting to see that the discrepancy must be essentially larger if the supremum in (2) is taken over all rectangles in arbitrary position. Here we have

$$(4) \quad D^{(\mathcal{R}')} (x) > c_K s(x)^{-(\frac{1}{2}+1/(K-1))}.$$

The same lower bound holds if we use balls instead of rectangles in arbitrary position or if we consider a particle's movement on the K -dimensional unit sphere and the discrepancy with respect to spherical caps. It should be stated that the problem to find bounds for the discrepancy for continuous functions on the sphere was proposed by R.F. Tichy [11].

In the first part of this paper (4) is proved by an application of J. Beck's Fourier transform method [2] and it is shown that the lower bounds (3) and (4) are optimal despite of logarithmic factors of $s(x)$. The next part deals with convex bodies and gives a continuous analogon to J. Beck's solution of a problem of K. F. Roth [1]. In the last section some problems and results concerning the behaviour of the discrepancy are discussed if one considers a function $x : [0, \infty) \rightarrow \mathbf{R}^K/\mathbf{Z}^K$.

It should be noted that the discrepancy of a continuous function, introduced by E. Hlawka [7], is an analogon to the discrepancy of sequences which measures the irregularity of a point distribution. An excellent survey of this subject including recent results with complete proofs can be found in J. Beck and W. Chen's book *Irregularities of Distribution* [2].

2. TORUS & SPHERE

2.1 Definitions and results.

Let $\mathcal{R}^K = \{\prod_{i=1}^K [a_i, b_i] \mid 0 < b_i - a_i \leq 1, i = 1, \dots, K\}$ be the set of all K -dimensional rectangles and $\overline{\mathcal{R}}^K = \mathcal{R}^K/\mathbf{Z}^K$ the set of rectangles contained in the unit torus $\mathbf{R}^K/\mathbf{Z}^K$. If $x : [0, 1] \rightarrow \mathbf{R}^K/\mathbf{Z}^K$ is a continuous

function with finite arclength $s(x) = \int_0^1 |dx(t)|$ the discrepancy of $x(t)$ with respect of $\overline{\mathcal{R}}^K$ is defined by

$$(5) \quad D^{(\mathcal{R})}(x) = \sup_{R \in \overline{\mathcal{R}}^K} \left| \int_0^1 \chi_R(x(t)) dt - \lambda_K(R) \right|,$$

where λ_K is the Lebesgue measure on $\mathbf{R}^K/\mathbf{Z}^K$ (i. e. the Haar measure on $\mathbf{R}^K/\mathbf{Z}^K$).

Set

$$(6) \quad \Delta_K^{(\mathcal{R})}(s) = \inf_{x \in C_s^K} D^{(\mathcal{R})}(x),$$

where C_s^K is the set of all continuous functions $x : [0, 1] \rightarrow \mathbf{R}^K/\mathbf{Z}^K$ with $s(x) = s$. Then we can formulate

THEOREM 1. — *If $s \geq 1$, then we have for $K \geq 2$*

$$(7) \quad \Delta_K^{(\mathcal{R})}(s) \gg s^{-(1+1/(K-1))}$$

and on the other hand

$$(8) \quad \Delta_2^{(\mathcal{R})} \ll s^{-2}$$

and for $K \geq 3$

$$(9) \quad \Delta_K^{(\mathcal{R})}(s) \ll s^{-(1+1/(K-1))} (\log(s+1))^{K-1},$$

where the constants implied by \ll or \gg are only depending on the dimension K .

Remark 1. — If $K = 1$ it is easy to see that $\Delta_1^{(\mathcal{R})}(s) = 0$ for $s \geq 1$ (cf. [3]).

Now let \mathcal{Q}^K denote the set of all cubes $Q \in \mathbf{R}^K$ in arbitrary position such that the length of the sides are $\leq K^{-1/2}$ and \mathcal{B}^K the set of balls $B \in \mathbf{R}^K$ with diameter ≤ 1 . Again we can define discrepancies

$$(10) \quad \begin{aligned} D^{(\mathcal{Q})}(x) &= \sup_{Q \in \mathcal{Q}^K} \left| \int_0^1 \chi_Q(x(t)) dt - \lambda_K(Q) \right| \\ D^{(\mathcal{B})}(x) &= \sup_{B \in \mathcal{B}^K} \left| \int_0^1 \chi_B(x(t)) dt - \lambda_K(B) \right| \end{aligned}$$

and

$$(11) \quad \begin{aligned} \Delta_K^{(\mathcal{Q})}(s) &= \inf_{x \in C_s^K} D^{(\mathcal{Q})}(x) \\ \Delta_K^{(\mathcal{B})}(s) &= \inf_{x \in C_s^K} D^{(\mathcal{B})}(x). \end{aligned}$$

In comparison to Theorem 1 there is a significant difference.

THEOREM 2. — *If $s \geq 1$, then we have for $K \geq 2$*

$$(12) \quad \left. \begin{array}{l} \Delta_K^{(\mathcal{Q})}(s) \\ \Delta_K^{(\mathcal{B})}(s) \end{array} \right\} \gg s^{-(\frac{1}{2}+1/(K-1))}$$

and on the other hand

$$(13) \quad \left. \begin{array}{l} \Delta_K^{(\mathcal{Q})}(s) \\ \Delta_K^{(\mathcal{B})}(s) \end{array} \right\} \ll s^{-(\frac{1}{2}+1/(K-1))} (\log(s+1))^{1/2},$$

where the constants implied by \ll or \gg are only depending on the dimension K .

Remark 2. — Since every cube is also a rectangle (12) implies (4).

Remark 3. — Since the torus $\mathbf{R}^K/\mathbf{Z}^K$ can be identified with the unit cube $U_0^K = [0, 1)^K$ it is also possible to consider the set of cubes \mathcal{Q}_0^K in arbitrary position contained in U_0^K and the set of balls \mathcal{B}_0 contained in U_0^K . Then it can be shown by a truncation technique that the corresponding $\Delta_K^{(\mathcal{Q}_0)}(s)$ and $\Delta_K^{(\mathcal{B}_0)}(s)$ satisfy

$$(14) \quad \left. \begin{array}{l} \Delta_K^{(\mathcal{Q}_0)}(s) \\ \Delta_K^{(\mathcal{B}_0)}(s) \end{array} \right\} \gg s^{-(\frac{1}{2}+1/(K-1)+\varepsilon)},$$

where $\varepsilon > 0$ is arbitrary but fixed and the constants implied by \gg are also depending on ε . (For details see section 3.)

Let $\mathbf{S}^K = \{x \in \mathbf{R}^{K+1} \mid |x| = 1\}$ denote the K -dimensional unit sphere and σ_K the normalized surface measure on \mathbf{S}^K , i. e. $\sigma_K(\mathbf{S}^K) = 1$. Here we consider the set \mathcal{C}^K of all spherical caps $C = \{x \in \mathbf{S}^K \mid |x - m| \leq r\}$ ($m \in \mathbf{S}^K, r > 0$) and the discrepancy of a continuous function $x : [0, 1] \rightarrow \mathbf{S}^K$

$$(15) \quad D^{(C)}(x) = \sup_{C \in \mathcal{C}^K} \left| \int_0^1 \chi_C(x(t)) dt - \sigma_K(C) \right|$$

If \mathcal{C}_s^K denotes the set of all continuous functions $x : [0, 1] \rightarrow \mathbf{S}^K$ with arclength $s(x) = s$ and

$$(16) \quad \Delta_K^{(C)}(s) = \inf_{x \in \mathcal{C}_s^K} D^{(C)}(x)$$

then we have similarly to Theorem 2.

THEOREM 3. — *If $s \geq 1$, then we have for $K \geq 2$*

$$(17) \quad \Delta_K^{(C)}(s) \gg s^{-(\frac{1}{2}+1/(K-1))}$$

and on the other hand

$$(18) \quad \Delta_K^{(C)}(s) \ll s^{-(\frac{1}{2}+1/(K-1))} (\log(s+1))^{1/2},$$

where the constants implied by \ll or \gg are only depending on the dimension K .

Remark 4. — The lower bounds of Theorem 2 and 3 sharpen the results of [4]:

$$(19) \quad \left. \begin{array}{l} \Delta_K^{(\mathcal{R}')} (s) \\ \Delta_K^{(\mathcal{B})} (s) \\ \Delta_K^{(C)} (s) \end{array} \right\} \gg \begin{array}{l} \text{for } K = 2, 3 \\ s^{(\frac{1}{2}+1/(K-1)+\varepsilon)} \text{ for } K \geq 2 \\ \text{for } K \geq 2 \end{array}$$

($\varepsilon > 0$ arbitrary but fixed.) The technique used in [4] is an application of W. Schmidt's integral equation method [9].

2.2. Lower bounds.

The lower bound (7) of Theorem 1 is due to R. J. Taschner [10] (see also [5]).

For the proof of the lower bounds (12) and (17) of Theorem 2 and 3 we will use J. Beck's Fourier transform technique. If $f \in L^1(\mathbf{R}^K)$ then the Fourier transform \hat{f} of f is defined by

$$(20) \quad \hat{f}(t) = \frac{1}{(2\pi)^{K/2}} \int_{\mathbf{R}^K} e^{-i\langle x,t \rangle} f(x) dx,$$

where $\langle x, t \rangle$ denotes the usual inner produkt in \mathbf{R}^K . Let $x : [0, 1] \rightarrow \mathbf{R}^K/\mathbf{Z}^K$ be a continuous function with $s(x) = s$, then we can introduce a measure z on \mathbf{R}^K by defining it on measurable subsets $A \in \mathbf{R}^K$ that are contained in some translated unit cube $y + U_0^K$ by

$$(21) \quad z(A) = \int_0^1 \chi_{A/\mathbf{Z}^K}(x(t)) dt.$$

Let τ denote a proper orthogonal transformation (i. e. a rotation) then $\chi_{r,\tau}$ shall denote the characteristic function of the rotated cube $\tau[-r, r]^K = \{\tau x \mid x \in [-r, r]^K\}$ ($r \leq K^{-1/2}/2$). (For simplicity we will write $Q(x)$ for the cube $[-x, x]^K$ in the sequel.) Let $M > 0$ be a parameter to be fixed later, and set

$$(22) \quad \mu_M(A) = \lambda_K(A \cap Q(M)) \quad \text{and} \quad z_M(A) = z(A \cap Q(M)).$$

Now consider the function

$$(23) \quad F_{r,\tau} = \chi_{r,\tau} * (dz_M - d\mu_M) ,$$

where $*$ denotes the convolution operation. More precisely we have

$$(24) \quad \begin{aligned} F_{r,\tau}(x) &= \int_{\mathbf{R}^K} \chi_{r,\tau}(x - y)(dz_M(y) - d\mu_M(y)) \\ &= \int_0^1 \chi_{(\tau Q(r)+x) \cap Q(M)/\mathbf{Z}^K}(x(t))dt - \lambda_K((\tau Q(r) + x) \cap Q(M)) . \end{aligned}$$

Let \mathbf{T} be the group of proper orthogonal transformations in \mathbf{R}^K and let $d\tau$ be the volume element of the normalized Haar measure on \mathbf{T} . Introducing the notation

$$(25) \quad \Phi(q) = \frac{1}{q} \int_q^{2q} \int_{\mathbf{T}} \int_{\mathbf{R}^K} |F_{r,\tau}(x)|^2 dx d\tau dr$$

we derive by using the Parseval—Plancherel identity and the convolution relation $\widehat{f * g} = \widehat{f} \cdot \widehat{g}$ that

$$(26) \quad \Phi(q) = \int_{\mathbf{R}^K} \left(\frac{1}{q} \int_q^{2q} \int_{\mathbf{T}} |\widehat{\chi}_{r,\tau}(t)|^2 d\tau dr \right) \left| (dz_M - \widehat{d\mu_M})(t) \right|^2 dt .$$

For brevity set

$$(27) \quad \varphi_q(t) = \frac{1}{q} \int_q^{2q} \int_{\mathbf{T}} |\widehat{\chi}_{r,\tau}(t)|^2 d\tau dr$$

and

$$(28) \quad \psi(t) = (dz_M - \widehat{d\mu_M})(t) = \frac{1}{(2\pi)^{K/2}} \int_{\mathbf{R}^K} e^{-i(x,t)}(dz_M - d\mu_M)(x) .$$

We will need two Lemmata.

LEMMA 1 ([2] P. 134). — *If $0 < q < p$, then*

$$(29) \quad \frac{\varphi_p(t)}{\varphi_q(t)} \gg \left(\frac{p}{q}\right)^{K-1}$$

uniformly in all $t \in \mathbf{R}^K$.

LEMMA 2. — *If $s(x) = s \geq 1$, then*

$$(30) \quad \Phi \left(\frac{c_K}{s^{1/(K-1)}} \right) \gg s^{-2K/(K-1)} M^K,$$

where c_K is a proper constant only depending on the dimension K .

Proof. — Identify the unit torus $\mathbf{R}^K/\mathbf{Z}^K$ with the unit cube $U_0^K = [0, 1)^K$. Now subdivide U_0^K into N^K cubes $Q_{(m_i)} = \prod_{i=1}^K [m_i/N, (m_i + 1)/N)$ ($0 \leq m_i < N, i = 1, \dots, K$). Furthermore subdivide the interval $[0, 1]$ into $[s(N + 1)]$ intervals $[t_j, t_{j+1})$ ($j = 0, \dots, [s(N + 1)] - 1$), $0 = t_0 < t_1 < \dots < t_{[s(N+1)]} = 1$, such that the arclength $\int_{t_j}^{t_{j+1}} |dx(t)| < 1/N$ ($j = 0, \dots, [s(N + 1)] - 1$). Trivially, the number of cubes $Q_{(m_i)}$ such that $x([t_j, t_{j+1})) \cap Q_{(m_i)}$ is non-void is less or equal 2^K . Therefore there are at most $2^K [s(N + 1)]$ cubes $Q_{(m_i)}$ such that $x([0, 1]) \cap Q_{(m_i)}$ is non void. Now choose the minimal N such that $2^K s(N + 1) \leq N^K/2$. Hence there are at least $N^K/2$ cubes $Q_{(m_i)}$ such that $x([t_j, t_{j+1})) \cap Q_{(m_i)}$ is void. Now consider the subcube $C_{(m_i)} = \prod_{i=1}^K [(m_i + 1/4)/N, (m_i + 3/4)/N)$ of such a cube $Q_{(m_i)}$. If $r < (4NK^{1/2})^{-1}$ and $x \in C_{(m_i)}$ we have

$$(31) \quad F_{r,\tau}(x) = -(2r)^K$$

and therefore

$$(32) \quad \Phi(q) \gg M^K q^{2K},$$

if $q < (8NK^{1/2})^{-1}$. Since $N \gg\gg s^{1/(K-1)}$, (32) implies (30). Thus Lemma 2 is proved.

Now set $q = c_K s^{-1/(K-1)}$ and $p = (5K^{1/2})^{-1}$, then we can use (29) and (28) to conclude

$$(33) \quad \begin{aligned} \Phi(p) &= \int_{\mathbf{R}^K} \varphi_p(t) |\psi(t)|^2 dt \gg s \Phi(q) \\ &\gg s^{1-2K/(K-1)} M^K = s^{-(1+2/(K-1))} M^K. \end{aligned}$$

Since $F_{r,\tau}(x) = 0$ for $x \notin Q(M + 1)$, (33) implies that either

$$(34) \quad \frac{1}{p} \int_p^{2p} \int_{\mathbf{T}} \int_{Q(M+1) \setminus Q(M-1)} |F_{r,\tau}(x)|^2 dx d\tau dr \gg s^{-(1+2/(K-1))} M^K$$

or

$$(35) \quad \frac{1}{p} \int_p^{2p} \int_{\mathbf{T}} \int_{Q(M-1)} |F_{r,\tau}(x)|^2 dx d\tau dr \gg s^{-(1+2/(K-1))} M^K.$$

Since $F_{r,\tau}(x)$ is absolutely bounded by 1, (34) cannot be true if M is large enough, e. g. $M = \bar{c}_K s^{1+2/(K-1)}$ for some proper chosen constant \bar{c}_K . Therefore (35) implies that there is an $r \in [p, 2p]$, a $\tau \in \mathbf{T}$, and an $x \in Q(M - 1)$ with

$$(36) \quad |F_{r,\tau}(x)| = \left| \int_0^1 \chi_{(\tau Q(r)+x)/\mathbf{Z}^K}(x(t)) dt - \lambda_K(\tau Q(r) + x) \right| \gg s^{-(\frac{1}{2}+1/(K-1))}.$$

Thus

$$(37) \quad \Delta_K^{(Q)}(s) \gg s^{-(\frac{1}{2}+1/(K-1))}.$$

The second part of (12), $\Delta_K^{(B)}(s) \gg s^{-(1/2+1/(K-1))}$, can be proved similarly. Instead of (22) set

$$(38) \quad \bar{F}_r = \bar{\chi}_r * (dz_M - d\mu_M),$$

where $\bar{\chi}_r$ denotes the characteristic function of the ball $B(r) = \{x \in \mathbf{R}^K \mid |x| \leq r\}$. Now consider

$$(39) \quad \begin{aligned} \bar{\Phi}(q) &= \frac{1}{q} \int_q^{2q} \int_{\mathbf{R}^K} |\bar{F}_r(x)|^2 dx dr \\ &= \int_{\mathbf{R}^K} \frac{1}{q} \int_q^{2q} |\hat{\bar{\chi}}_r(t)|^2 dr |(dz_M - \widehat{d\mu_M})(t)|^2 dt \\ &= \int_{\mathbf{R}^K} \bar{\varphi}_q(t) |\psi(t)|^2 dt. \end{aligned}$$

Again we have

LEMMA 3. — If $0 < q < p$, then

$$(40) \quad \frac{\bar{\varphi}_p(t)}{\bar{\varphi}_q(t)} \gg \left(\frac{p}{q}\right)^{K-1}$$

uniformly in all $t \in \mathbf{R}^K$.

Proof. — (40) is an immediate consequence of formula (201)

$$(41) \quad \frac{1}{y} \int_y^{2y} |\hat{\bar{\chi}}_r(t)|^2 dr \gg \ll \frac{y^{K-1}}{|t|^{K+1}} \quad \text{for } y|t| > c_K$$

and formula (203)

$$(42) \quad \frac{1}{y} \int_y^{2y} |\widehat{\chi}_r(t)|^2 dr \gg \ll y^{2K} \quad \text{for } y|t| \leq c_K$$

of [2], p. 228.

LEMMA 4. — *If $s(x) = s \geq 1$, then*

$$(43) \quad \overline{\Phi} \left(\frac{c_K}{s^{1/(K-1)}} \right) \gg s^{-2K/(k-1)} M^K ,$$

where c_K is a proper constant only depending on the dimension K .

The proof of Lemma 4 is verbally the same as the proof of Lemma 2. Hence we can deduce the bound $\Delta_K^{(B)}(s) \gg s^{-(1/2+1/(K-1))}$ as before. Therefore the proof of (12) in Theorem 2 is finished.

Now let $x : [0, 1] \rightarrow \mathbf{S}^K$ be a continuous function with $s(x) = s \geq 1$. Define two measures on \mathbf{R}^{K+1} , one by $\sigma_0(A) = \sigma_K(A \cap \mathbf{S}^K)$ and a second by

$$(44) \quad z_0(A) = \int_0^1 \chi_{A \cap \mathbf{S}^K}(x(t)) dt .$$

Set

$$(45) \quad G_r = \overline{\chi}_r * (dz_0 - d\sigma_0)$$

and

$$(46) \quad \begin{aligned} \Psi(q) &= \frac{1}{q} \int_q^{2q} \int_{\mathbf{R}^{K+1}} |G_r(x)|^2 dx dr \\ &= \int_{\mathbf{R}^{K+1}} \overline{\varphi}_q(t)(t) \left| (dz_0 - d\sigma_0)(t) \right|^2 dt . \end{aligned}$$

Then we can prove

LEMMA 5. — *If $s(x) = s \geq 1$, then*

$$(47) \quad \Psi \left(\frac{c_K}{s^{1/(K-1)}} \right) \gg s^{-\frac{2K+1}{K-1}} ,$$

where c_K is a proper chosen constant only depending on the dimension K .

Proof. — It is well known that \mathbf{S}^K can be represented by $Y : (0, \infty) \times [0, \pi]^{K-1} \times [0, 2\pi) \rightarrow \mathbf{R}^{K+1}$:

$$\begin{aligned}
 y_1 &= \rho \sin \varphi_1, \sin \varphi_2 \cdots \sin \varphi_{K-2}, \sin \varphi_{K-1}, \sin \varphi_K \\
 y_2 &= \rho, \sin \varphi_1, \sin \varphi_2 \cdots \sin \varphi_{K-2}, \sin \varphi_{K-1}, \cos \varphi_K \\
 y_3 &= \rho, \sin \varphi_1, \sin \varphi_2 \cdots \sin \varphi_{K-2}, \cos \varphi_{K-1} \\
 &\vdots \\
 y_K &= \rho, \sin \varphi_1, \cos \varphi_2
 \end{aligned}$$

where ρ is set equal to 1. Now consider the images $R_{(m_i)} = Y(\{1\} \times B_{(m_i)})$ of the $4N^K$ cubes

$$\begin{aligned}
 (49) \quad B_{(m_i)} &= \prod_{j=1}^{K-1} \left[\frac{\pi}{4} \left(1 + 2 \frac{m_j}{N} \right), \frac{\pi}{4} \left(1 + 2 \frac{m_j + 1}{N} \right) \right) \times \left[\pi \frac{m_K}{2N}, \pi \frac{m_K + 1}{2N} \right) \\
 &0 \leq m_1, \dots, m_{K-1} < N, \quad 0 \leq m_K < 4N.
 \end{aligned}$$

As in Lemma 2 it is possible to choose $N \gg\gg s^{1/(K-1)}$ such that for at least $2N^K$ sets $R_{(m_i)}$ the intersection $x([0, 1] \cap R_{(m_i)})$ is void. For these $R_{(m_i)}$ consider the images $\bar{R}_{(m_i)} = Y(C_{(m_i)})$ of the rectangles

$$\begin{aligned}
 (50) \quad C_{(m_i)} &= \left[1, 1 + \frac{1}{8N} \right] \times \prod_{j=1}^{K-1} \left[\frac{\pi}{4} \left(1 + 2 \frac{m_j + \frac{1}{4}}{N} \right), \frac{\pi}{4} \left(1 + 2 \frac{m_j + \frac{3}{4}}{N} \right) \right) \\
 &\times \left[\frac{\pi(m_K + 1/4)}{2N}, \frac{\pi(m_K + 3/4)}{2N} \right) .
 \end{aligned}$$

If $\rho \in \left[\frac{\pi}{16N}, \frac{\pi}{8N} \right]$ and $y \in C_{(m_i)}$ (for such $R_{(m_i)}$) we have similarly to (31)

$$(51) \quad |G_r(x)| \gg N^{-K} .$$

This implies (47). Now it is easy to verify the lower bound (17) in Theorem 3. By Lemma 3 we have

$$(52) \quad \frac{\bar{\varphi}_1(t)}{\bar{\varphi}_q(t)} \gg s^{K/(K-1)} ,$$

where $q = c_K s^{-1/(K-1)}$. (Note that we are working in \mathbf{R}^{K+1} .) Thus

$$(53) \quad \Psi(1) \gg s^{K/(K-1)} \psi(q) \gg s^{-(\frac{1}{2} + 1/(K-1))} .$$

2.3. Upper bounds.

The essential ideas of this section can be stated in two simple Lemmata. (We postpone the proofs at the end of this section.)

LEMMA 6. — *Let X be a convex body in \mathbf{R}^K and x_1, \dots, x_N N points in X . Then there is a permutation π of $\{1, \dots, N\}$ such that*

$$(54) \quad \sum_{i=1}^{N-1} |x_{\pi(i+1)} - x_{\pi(i)}| \ll N^{1-1/K} .$$

If x_1, \dots, x_N are on the boundary ∂X of X , then there is a permutation ρ of $\{1, \dots, N\}$ such that

$$(55) \quad \sum_{i=1}^{N-1} \delta(x_{\rho(i+1)}, x_{\rho(i)}) \ll N^{1-1/(K-1)} ,$$

where $\delta(\cdot, \cdot)$ denotes the geodesic distance on ∂X . The constants implied by \ll in (54) and (55) are only depending on the diameter $\text{diam}(X)$ of X .

Remark 5. — It should be noted that the estimates (54) and (55) are best possible. For every $N \geq 1$ there are N points x_1, \dots, x_N in X such that for every permutation π of $\{1, \dots, N\}$

$$(56) \quad \sum_{i=1}^{N-1} |x_{\pi(i+1)} - x_{\pi(i)}| \gg N^{1-1/K} .$$

(For example take an orthogonal lattice in X .) Similarly it can be shown that (55) is optimal, too.

LEMMA 7. — *Let X be a convex body in \mathbf{R}^K with $\lambda_K(X) = 1$ and let σ denote the normalized surface measure on ∂X .*

If x_1, \dots, x_N are N points in X , then for every $\varepsilon > 0$ there is a continuous function $x : [0, 1] \rightarrow X$ with $s(x) \ll N^{1-1/K}$ such that

$$(57) \quad \left| \int_0^1 \chi_A(x(t)) dt - \lambda_K(A) \right| \leq \left| \frac{1}{N} \sum_{n=1}^N \chi_A(x_n) - \lambda_K(A) \right| + \varepsilon$$

holds uniformly in all measurable subsets $A \subseteq X$.

If x_1, \dots, x_N are on ∂X , then for every $\varepsilon > 0$ there is a continuous function $x : [0, 1] \rightarrow \partial X$ with $s(x) \ll N^{1-1/(K-1)}$ such that

$$(58) \quad \left| \int_0^1 \chi_C(x(t)) dt - \sigma(C) \right| \leq \left| \frac{1}{N} \sum_{n=1}^N \chi_C(x_n) - \sigma(C) \right| + \varepsilon$$

holds uniformly in all measurable subsets $C \subseteq \partial X$.

COROLLARY. — Let \mathcal{D} be a system of measurable subsets of a convex body $X \subseteq \mathbf{R}^K$ with $\lambda_K(X) = 1$. If we set

$$(59) \quad \Omega^{(\mathcal{D})}(N) = \inf_{\{x_1, \dots, x_N\} \subseteq X} \sup_{D \in \mathcal{D}} \left| \frac{1}{N} \sum_{n=1}^N \chi_D(x_n) - \lambda_K(D) \right|$$

and

$$(60) \quad \Delta^{(\mathcal{D})}(s) = \inf_{x \in C_s} \sup_{D \in \mathcal{D}} \left| \int_0^1 \chi_D(x(t)) dt - \lambda_K(D) \right|,$$

where C_s is the set of all continuous functions $x : [0, 1] \rightarrow X$ with $s(x) = s$, then there is a constant $c > 0$ such that

$$(61) \quad \Delta^{(\mathcal{D})}(s) \leq \Omega^{(\mathcal{D})}(N)$$

for $s = cN^{1-1/K}$. A similar statement is true for the boundary ∂X of X .

Remark 6. — This Corollary has two interesting consequences. The first one, that will be used here, is that an upper bound for $\Omega(N)$, the smallest discrepancy of sequences, yields an upper bound for $\Delta(s)$, the smallest discrepancy of continuous functions. On the other hand, a lower bound for functions in terms of the arc length s gives a lower bound for sequences in terms of the number N . For example, consider N points on x_1, \dots, x_n on the sphere \mathbf{S}^K . Then Theorem 3 and the above Corollary imply that there is a spherical cap $C \subseteq \mathbf{S}^K$ such that

$$(62) \quad \left| \frac{1}{N} \sum_{n=1}^N \chi_C(x_n) - \sigma_K(C) \right| \gg N^{-(\frac{1}{2}+1/(2K))}.$$

(Compare with [2], Theorem 24C.) Therefore the lower bounds for the discrepancy of functions proved here are in some sense more general than the corresponding results for sequences [2].

Now consider the K -dimensional torus $\mathbf{R}^K/\mathbf{Z}^K$ which can be identified with $[0, 1)^K$. Every subset of $\mathbf{R}^K/\mathbf{Z}^K$ can be considered as a subset of $[0, 1)^K$, and a continuous function $x : [0, 1] \rightarrow [0, 1)^K$ can be interpreted as a continuous function $x : [0, 1] \rightarrow \mathbf{R}^K/\mathbf{Z}^K$. Hence we can use Lemma 7 to construct well distributed continuous functions $x : [0, 1] \rightarrow \mathbf{R}^K/\mathbf{Z}^K$, too.

It is well known (cf.[2]) that

$$\Omega^{(\mathcal{R})}(N) \ll \frac{(\log N)^{K-1}}{N}$$

$$(63) \quad \left. \begin{matrix} \Omega^{(Q)}(N) \\ \Omega^{(B)}(N) \\ \Omega^{(C)}(N) \end{matrix} \right\} \ll N^{-(\frac{1}{2}+1/(2K))}(\log N)^{1/2}$$

Thus the above Corollary implies the upper bounds (9), (13), and (18) for the discrete values $s = cN^{1-1/K}$. But $\Delta_K^{(\cdot)}(s)$ is a decreasing function in s , since $C_{s_1}^K \subseteq C_{s_2}^K$ for $s_1 \leq s_2$. Therefore (9), (13), and (18) are satisfied for all $s \geq 1$ for proper chosen constants. A proof for the upper bound (8) can be found in [3] (or cf. Remark 7 in section 4).

Proof of Lemma 6. — First assume that X is a ball which center is the origin of \mathbf{R}^K . For $K = 1$ (54) is trivial. Now suppose that $K \geq 2$ and let $X_1 = X \cap \{x_1 = 0\}$ be the intersection with the hyperplane $\{x_1 = 0\}$. Set $M = [N^{1/K}]$ and let $Z \subseteq X_1$ be the set of all points $z \in X_1$ such that $M \cdot z$ is an integer point. Trivially $|Z| \ll M^{K-1} \ll N^{1-1/K}$. For every $i = 1, \dots, N$ let $z(x_i)$ be a point $z \in Z$ such that the distance between x_i and the line $l(z) = \{y \in X \mid \langle y - z, x \rangle = 0 \text{ for all } x \in \{x_1 = 0\}\}$ is minimal and let $x_i^0 \in l(z(x_i))$ be the corresponding point. (This means that $|x_i, x_i^0|$ is minimal on $l(z(x_i))$.) Let $z_1, \dots, z_{|Z|}$ be a numeration of Z , k_j be the number of points x_i^0 on $l(z_j)$, and I_j the set of indices $i \in \{1, \dots, N\}$ with $x_i^0 \in l(z_j)$. If I_j is non-void, consider the points $x_{i_1}^0, \dots, x_{i_{k_j}}^0 \in l(z_j)$ in a way that the first coordinates $x_{i_1,1}^0 = x_{i_1,1}$ are ordered, i. e. $x_{i_1,1}^0 \leq x_{i_2,1}^0 \leq \dots \leq x_{i_{k_j},1}^0$. Now define a permutation π by $\pi \left(\sum_{l=1}^{j-1} k_l + m \right) = i_m$ ($m = 1, \dots, k_j$). Thus we have

$$(64) \quad \begin{aligned} \sum_{i=1}^{N-1} |x_{\pi(i+1)} - x_{\pi(i)}| &= \sum_{j=1}^{|Z|} \sum_{i=\sum_{l=1}^{j-1} k_l + 1}^{\sum_{l=1}^j k_l} |x_{\pi(i+1)} - x_{\pi(i)}| \\ &\leq 2|Z|\text{diam}(X) + \sum_{j=1}^{|Z|} k_j \frac{\sqrt{K-1}}{M} \\ &\ll N^{1-1/K} . \end{aligned}$$

Now consider N points on the surface ∂X . Since the case $K = 2$ is trivial we can assume $K > 2$. It is well known that for every positive integer L there are L points z_1, \dots, z_L on $X_1 \cap \partial X$ such that the maximal geodesic distance to the next point $\delta = \max_{1 \leq i \leq L} \min_{j \neq i} \delta(x_i, x_j) \ll L^{-1/(K-2)}$. If $L = [N^{1-1/(K-1)}]$ then $\delta \ll N^{-1/(K-1)}$. The set $\partial X \cap \{x_2 = \dots = x_K = 0\}$ consists of two points \underline{z} and \bar{z} which can be considered as the poles of the sphere ∂X . Let $l(z_j)$ ($j = 1, \dots, L$) be the meridian joining \underline{z} and \bar{z} and

containing z_j . Now the construction of a permutation ρ can be managed as before to get the estimate

$$(65) \quad \sum_{i=1}^{N-1} \delta(x_{\rho(i+1)}, x_{\rho(i)}) \ll L + N \cdot N^{-1/(K-1)} \\ \ll N^{1-1/(K-1)} .$$

It should be noted that the constants implied by \ll in (64) and (65) only depend on the diameter $\text{diam}(X)$.

If X is a general convex body let Y denote the smallest closed ball containing X . Trivially $\text{diam}(X) \gg \ll \text{diam}(Y)$. Therefore (64) implies (54). For the proof of (55) we need a surjective non-expansive function $p : \partial Y \rightarrow \partial X$. Then, if x_1, \dots, x_N are points on ∂X , we can choose points y_1, \dots, y_N on ∂Y with $p(y_i) = x_i$. Therefore $\delta(x_i, x_j) \leq \delta(y_i, y_j)$ and (65) imply (55). Such a function p can be constructed in the following way. Let $B(r) = \{x \in \mathbf{R}^K \mid |x| \leq r\}$ be the closed ball with radius r . Then the boundary of $X_r = \frac{R}{r+R} \cdot (X + B(r))$ ($R = \text{diam}(Y)/2$) converges uniformly to ∂Y as $r \rightarrow \infty$. Furthermore there is a canonical surjective non-expansive function $p_r : \partial X_r \rightarrow \partial X$ for every $r > 0$. Now let $p(y) = x$ if for every $\varepsilon > 0$ there is a $r > 0$ and a point $y_r \in \partial X_r \cap (y + B(\varepsilon))$ with $p_r(y_r) = x$. This completes the proof of Lemma 6.

Proof of Lemma 7. — It is no loss of generality to assume that $\varepsilon < 1$ and (by Lemma 6) that

$$(66) \quad s = \sum_{n=1}^{N-1} |x_{n+1} - x_n| \ll N^{1-1/K} .$$

Now set

$$(67) \quad x(t) = \begin{cases} x_i & \text{for } \frac{i-1}{N} \leq t \leq \frac{i-\varepsilon}{N}, \quad 1 \leq i < N \\ x_i + (x_{i+1} - x_i) \frac{t - \frac{i-\varepsilon}{N}}{\varepsilon/N} & \text{for } \frac{i-\varepsilon}{N} \leq t \leq \frac{i}{N}, \quad 1 \leq i < N \\ x_N & \text{for } 1 - \frac{1}{N} \leq t \leq 1 . \end{cases}$$

Trivially

$$(68) \quad \left| \frac{1}{N} \sum_{n=1}^N \chi_A(x_n) - \int_0^1 \chi_A(x(t)) dt \right| \leq \varepsilon .$$

Thus (68) implies (57). The construction for (58) can be managed similarly.

3. CONVEX BODIES

3.1. Definitions and results.

Let $X \subseteq \mathbf{R}^K$ ($K \geq 2$) be a convex body with $\lambda_K(X) = 1$ and let $x : [0, 1] \rightarrow X$ be a continuous function with arclength $s(x) = s$. Furthermore let \mathcal{Q} denote the set of all cubes $Q \subseteq \mathbf{R}^K$ in arbitrary position and \mathcal{Q}_0 the cubes contained in X , respectively let \mathcal{B} denote the set of all balls and \mathcal{B}_0 the set of all balls contained in X . Now set

$$(69) \quad \Delta^{(\mathcal{Q})}(s) = \inf_{x \in C_s} \sup_{Q \in \mathcal{Q}} \left| \int_0^1 \chi_Q(x(t)) dt - \lambda_K(Q \cap X) \right|$$

and

$$(70) \quad \Delta^{(\mathcal{Q}_0)}(s) = \inf_{x \in C_s} \sup_{Q \in \mathcal{Q}_0} \left| \int_0^1 \chi_Q(x(t)) dt - \lambda_K(Q) \right|,$$

where C_s denotes the set of all continuous functions $x : [0, 1] \rightarrow X$ with $s(x) = s$. $\Delta^{(\mathcal{B})}(s)$ and $\Delta^{(\mathcal{B}_0)}(s)$ can be defined similarly. Here we can formulate

THEOREM 4. — *Let $X \subseteq \mathbf{R}^K$ ($K \geq 2$) be a convex body with $\lambda_K(X) = 1$. If $s \geq 1$, then we have*

$$(71) \quad \left. \begin{array}{l} \Delta^{(\mathcal{Q})}(s) \\ \Delta^{(\mathcal{B})}(s) \end{array} \right\} \gg s^{-(\frac{1}{2}+1/(K-1))}$$

and

$$(72) \quad \left. \begin{array}{l} \Delta^{(\mathcal{Q})}(s) \\ \Delta^{(\mathcal{B})}(s) \end{array} \right\} \ll s^{-(\frac{1}{2}+1/(K-1))} (\log(s+1))^{1/2},$$

where the constants implied by \ll and \gg are only depending on the convex body $X \subseteq \mathbf{R}^K$.

For \mathcal{Q}_0 and \mathcal{B}_0 we must use a truncation method to get

THEOREM 5. — *Let $X \subseteq \mathbf{R}^K$ ($K \geq 2$) be a convex body with $\lambda_K(X) = 1$. If $s \geq 1$, then we have for arbitrary $\epsilon > 0$*

$$(73) \quad \left. \begin{array}{l} \Delta^{(\mathcal{Q}_0)}(s) \\ \Delta^{(\mathcal{B}_0)}(s) \end{array} \right\} \gg s^{-(\frac{1}{2}+1/(K-1)+\epsilon)}$$

and

$$(74) \quad \left. \begin{array}{l} \Delta^{(Q_0)}(s) \\ \Delta^{(B_0)}(s) \end{array} \right\} \ll s^{-(\frac{1}{2}+1/(K-1))} (\log(s+1))^{1/2} ,$$

where the constants implied by \ll and \gg are only depending on the convex body $X \subseteq \mathbf{R}^K$ and on $\varepsilon > 0$ in (73).

It is also interesting to consider segments instead of cubes or balls. Let \mathcal{S} denote the set of all half-spaces $\{y = (y_1, \dots, y_K) \in \mathbf{R}^K \mid a_1 y_1 + \dots + a_K y_K \geq b\}$ ($a_1, \dots, a_K, b \in \mathbf{R}$) and

$$(75) \quad \Delta^{(\mathcal{S})}(s) = \inf_{x \in C_s} \sup_{S \in \mathcal{S}} \left| \int_0^1 \chi_S(x(t)) dt - \lambda_K(S \cap X) \right| ,$$

then one gets

THEOREM 6. — *Let $X \subseteq \mathbf{R}^K$ ($K \geq 2$) be a convex body with $\lambda_K(X) = 1$. If $s \geq 1$, then we have*

$$(76) \quad \Delta^{(\mathcal{S})}(s) \gg s^{-(\frac{1}{2}+1/(K-1))} (\log(s+1))^{(1-K^2)/4} \frac{1}{\phi(s)} ,$$

where $\phi(s)$ is an arbitrary positive monotonic function with $\lim_{s \rightarrow \infty} \phi(s) = \infty$. Furthermore

$$(77) \quad \Delta^{(\mathcal{S})}(s) \ll s^{-(\frac{1}{2}+1/(K-1))} (\log(s+1))^{1/2} .$$

The constants implied by \ll and \gg are only depending on the convex body $X \subseteq \mathbf{R}^K$ and on $\phi(s)$ in (76).

Theorem 6 is a continuous analogon to J. Beck's [1] solution of a problem of K. F. Roth.

3.2. Lower bounds.

The proof of (71) is quite similar to the proofs of (12) and (17). Let $x : [0, 1] \rightarrow X$ be a continuous function with $s(x) = s \geq 1$. Define two measures

$$(78) \quad \mu(A) = \lambda_K(A \cap X) \quad \text{and} \quad z(A) = \int_0^1 \chi_A(x(t)) dt$$

and set

$$(79) \quad F_{r,\tau} = \chi_{r,\tau} * (dz - d\mu)$$

as in (22). Again we can conclude for

$$\begin{aligned}
 \Phi(q) &= \frac{1}{q} \int_q^{2q} \int_{\mathbf{T}} \int_{\mathbf{R}^K} |F_{r,\tau}(x)|^2 dx d\tau dr \\
 (80) \quad &= \int_{\mathbf{R}^K} \left(\frac{1}{q} \int_q^{2q} \int_{\mathbf{T}} |\hat{\chi}_{r,\tau}(t)|^2 d\tau dr \right) |(dz - d\mu)(t)|^2 dt \\
 &= \int_{\mathbf{R}^K} \varphi_q(t) |\psi(t)|^2 dt
 \end{aligned}$$

that there is a constant c with

$$(81) \quad \Phi\left(\frac{c}{s^{1/(K-1)}}\right) \gg s^{-2K/(K-1)}.$$

Therefore Lemma 1 yields

$$(82) \quad \Phi(1) \gg s^{1-2K/(K-1)} = s^{-(1+2/(K-1))},$$

which implies $\Delta^{(\mathcal{Q})}(s) \gg s^{-(1/2+1/(K-1))}$. The second part of (71) can be shown similarly.

We omit the proof of (73) since the major ideas are also included in the simpler proof of (76). In addition we generalize [1] to higher dimensions.

Let $x(t)$ be a continuous function as above. Set $n = c s^{1/(K-1)}$, where the constant c will be fixed in the sequel. Now $y(t) = n x(t/T)$ with $T = n^K$ is a continuous function $y : [0, T] \rightarrow Y$ with $Y = nX$. The arclength of $y(t)$ can be evaluated by $s(y) = n s = s'$. Now (76) can be restated in the following way: There is a halfspace S with

$$(83) \quad \left| \int_0^T \chi_S(y(t)) dt - \lambda_K(S \cap Y) \right| \gg \frac{T^{\frac{1}{2}-1/(2K)}}{(\log T)^{(K^2-1)/4} \phi(T)}.$$

Set

$$(84) \quad z(A) = \int_0^T \chi_A(y(t)) dt, \quad \mu(A) = \lambda_K(A \cap Y),$$

(85)

$$R = \{x = (x_1, \dots, x_K) \in \mathbf{R}^K \mid |x_1| \leq m, \dots, |x_{K-1}| \leq m, |x_K| \leq \frac{1}{100}\},$$

where $m = n(\log n)^{K/2} \phi(n)^{2K-1}$,

$$(86) \quad E(x) = e^{-|x|^2/r^2},$$

where $r = n(\log n)^{(K-1)/2} \phi(n)^{2K}$, and

$$(87) \quad F_\tau = E \cdot (\chi_{\tau R} * (dz - d\mu)) ,$$

where τ is the class of all proper orthogonal transformations τ' that create the same image $\tau'R$ of R , i. e. $\tau'R$ is independent of the choice $\tau' \in \tau$. Let \mathbf{T} denote the homogeneous space consisting of all these classes. Since $R \cap Y$ is the set-theoretic difference of two parallel segments $S_1 \cap Y, S_2 \cap Y$, we have

$$(88) \quad \int_{\mathbf{R}^K} F_\tau(x)^2 dx = \int_{\mathbf{R}^K} E(x)^2 dx \cdot \left| \int_0^1 \chi_{\tau R+x}(y(t)) dt - \mu(\tau R + x) \right|^2 + T^2 \int_{|x| \geq m/2} E(x)^2 dx \ll r^K \Delta^2 + T^2 r^K e^{-\left(\frac{m/2}{r}\right)^2} ,$$

where

$$(89) \quad \Delta = \sup_{S \in \mathcal{S}} |z(S) - \mu(S)| .$$

Thus

$$(90) \quad \int_{\mathbf{T}} \int_{\mathbf{R}^K} F_\tau(x)^2 dx d\tau \ll r^K \Delta^2 + O(1) ,$$

where $d\tau$ is the volume element of the homogeneous space \mathbf{T} induced by the Haar measure of the compact group of all proper orthogonal transformations. Therefore (83) follows from

$$(91) \quad \int_{\mathbf{T}} \int_{\mathbf{R}^K} |\hat{F}_\tau(t)|^2 dt d\tau \gg \frac{n^{2K-1} r^{K-1}}{m^{K-1}} .$$

As in [1] it can be shown that for a proper constant c_0

$$(92) \quad \int_{T(c_0)} |\psi(t)|^2 dt \leq \frac{1}{2} \int_{Q(100)} |\psi(t)|^2 dt ,$$

where

$$(93) \quad \psi(t) = \widehat{(dz - d\mu)}(t) = \frac{1}{(2\pi)^{K/2}} \int_{\mathbf{R}^K} e^{-i\langle x, t \rangle} (dz - d\mu)(x)$$

and

$$(94) \quad T(c_0) = \left\{ t \in Q(100) \mid \text{there are indices } l_1, \dots, l_K \text{ with } \begin{aligned} & l_i \geq 0, \quad i = 1, \dots, K, \text{ and } l_1 + \dots + l_K \geq 1 \text{ such that} \\ & \left| \frac{\partial^{l_1 + \dots + l_K}}{\partial t_1^{l_1} \dots \partial t_K^{l_K}} \psi(t) \right| \geq c_0 (2n)^{k_1 + \dots + l_K} |\psi(t)| \end{aligned} \right\} .$$

But we need following additional observations for the proof of (92). Set

$$g(x) = \prod_{i=1}^K \left(\frac{2}{\pi} \left(\frac{\sin(50 x_i)}{x_i} \right)^2 \right) ,$$

$$(95) \quad G(x) = \int_{\mathbf{R}^K} g(x - y) (dz - d\mu)(y) ,$$

and

$$v(x) = \int_0^T \chi_{Q+x}(y(t)) dt ,$$

where $Q = Q(\frac{1}{100})$. Instead of $|G(x)| \geq v(x)$ (inequality (19) of [1] suffices to prove

$$(96) \quad \int_{\mathbf{R}^K} G(x)^2 dx \gg \int_{\mathbf{R}^K} v(x)^2 dx .$$

We will need

LEMMA 8. — *If $s(y) = s' = n s$ and $n = c s^{1/(K-1)}$, then*

$$(97) \quad \int_{\mathbf{R}^K} v(x)^2 dx \gg c^{K-1} n^K .$$

Proof. — Let t_i ($i = 0, \dots, [100 s']$) be defined by

$$(98) \quad \int_0^{t_i} |dy(t)| = \frac{i}{100} .$$

Therefore $s, t \in [t_i, t_{i+1}]$ implies $|y(t) - y(s)| \leq \frac{1}{100}$. Since

$$(99) \quad v(x)^2 = \int_0^T \int_0^T \chi_{Q+x}(y(t)) \chi_{Q+x}(y(s)) ds dt ,$$

Cauchy—Schwarz's inequality yields

$$\begin{aligned}
 \int_{\mathbf{R}^K} v(x)^2 dx &= \int_0^T \int_0^T \int_{\mathbf{R}^K} \chi_{Q+x}(y(t)) \chi_{Q+x}(y(s)) dx ds dt \\
 &\geq \int_0^T \int_0^T \int_{|y(t)-y(s)| \leq \frac{1}{100}} 50^{-K} ds dt \\
 &\geq 50^{-K} \sum_{i=1}^{[100s']} (t_i - t_{i-2})^2 + 50^{-K} (T - t_{[100s']})^2 \\
 &\geq 50^{-K} \frac{T^2}{100s' + 1} \geq \frac{T^2}{ns} \gg c^{K-1} n^K.
 \end{aligned}
 \tag{100}$$

If we set

$$I = \int_{\mathbf{R}^K} g(x) dx = 100^K,$$

we get

$$\underline{g} = \inf_{x \in Q} g(x) = \left(\frac{2}{\pi}\right)^K \left(100 \sin \frac{1}{2}\right)^{2K} \geq 10I.$$

Define $c_1 = I/(10I - 1)$. Then $\frac{1}{10} < c_1 < \frac{1}{9}$. Let R_1, R_2 denote the sets $R_1 = \{x \in \mathbf{R}^K \mid v(x) \geq c_1\}$ and $R_2 = \mathbf{R}^K \setminus R_1$, then we have for $x \in R_1$

$$\begin{aligned}
 G(x) &= \int_0^T g(x - y(t)) dt - \int_Y g(x - y) dy \\
 &\geq \int_0^T \underline{g} \chi_{Q+x}(y(t)) dt - I = \underline{g} v(x) - I \\
 &\geq 10I v(x) - I \geq v(x) + (10I - 1)c_1 - I \geq v(x).
 \end{aligned}
 \tag{103}$$

Since $\int_{R_2} v(x)^2 dx \ll n^K$, Lemma 8 implies that c can be chosen in a way that

$$\int_{R_2} v(x)^2 dx \leq \frac{1}{2} \int_{\mathbf{R}^K} v(x)^2 dx.$$

Thus (100), (104), and (97) imply

$$\int_{\mathbf{R}^K} G(x)^2 dx \geq \int_{R_1} G(x)^2 dx \geq \int_{R_1} v(x)^2 dx \gg \int_{\mathbf{R}^K} v(x)^2 dx,$$

if c is chosen in this proper way.

If $t \in \mathbf{R}^K \setminus \{0\}$ let $\alpha(t) \in \mathbf{T}$ be the class of all proper orthogonal transformations that can be represented by a proper orthogonal matrix A with columns a_1, \dots, a_K such that $\langle a_1, t \rangle = \dots = \langle a_{K-1}, t \rangle = 0$. If $\tau \in \mathbf{T}$, let $e_K(\tau) = \pm a_K$ be the last column of a corresponding orthogonal matrix. Set $B(x) = (\sin x)/x$ and $\beta(t, \tau) = B(\frac{1}{100} \langle t, e_K(\tau) \rangle)$. For $|\Delta\tau| = \text{dist}(\tau, \alpha(t))$ we have (similarly to [1])

$$(106) \quad \hat{\chi}_{\tau R}(t+u) = \hat{\chi}_{\tau R}(\beta(t, \tau) + O(|u|) + O(m^K |\Delta\tau|))$$

and for t not contained in $T(c_0)$

$$(107) \quad |\psi(t+u) - \psi(t)| \leq c_0 |\psi(t)| (e^{2Kn|u|} - 1).$$

Using

$$(108) \quad \left| \int_{\mathbf{R}^K} \hat{E}(-u) \hat{\chi}_{\tau R} \beta(t, \tau) \psi(t) du \right| \gg |\psi(t)| r^{K-1}$$

and some elementary estimates we can derive

$$(109) \quad \begin{aligned} |\hat{F}_{\tau}(t)| &\geq c' |\psi(t)| r^{K-1} \left(1 - O\left(\frac{m^{K-1}n}{r}\right) - O\left(\frac{m^K}{r^{K-1}}|\Delta\tau|\right) \right) \\ &\gg |\psi(t)| r^{K-1} \end{aligned}$$

for t not contained in $T(c_0)$ and $|\Delta\tau| \leq n/(mr)$. Thus we get

$$(110) \quad \begin{aligned} \int_{\mathbf{T}} \int_{\mathbf{R}^K} |\hat{F}_{\tau}(t)|^2 dt d\tau &\geq \int_{Q(100) \setminus T(c_0)} \int_{|\Delta\tau| \leq \frac{n}{mr}} |\hat{F}_{\tau}(t)|^2 d\tau dt \\ &\gg \left(\frac{n}{mr}\right)^{K-1} r^{2K-2} \int_{Q(100) \setminus T(c_0)} |\psi(t)|^2 dt \\ &\gg \left(\frac{n}{mr}\right)^{K-1} r^{2K-2} n^K = \frac{n^{2K-1} r^{K-1}}{m^{K-1}} \end{aligned}$$

since we have by (92), (95), (97), and (105)

$$(111) \quad \begin{aligned} \int_{Q(100) \setminus T(c_0)} |\psi(t)|^2 dt &\gg \int_{Q(100)} |\psi(t)|^2 dt \\ &\gg \int_{\mathbf{R}^K} |\hat{g}(t) \psi(t)|^2 dt = \int_{\mathbf{R}^K} G(x)^2 dx \\ &\gg \int_{\mathbf{R}^K} v(x)^2 dx \gg n^K. \end{aligned}$$

This completes the proof of (76).

3.3 Upper bounds.

A combination of the results for sequences [2] and the Corollary of Lemma 7 immediately implies the upper bounds (72), (74), and (77).

4. FURTHER PROBLEMS & RESULTS

It is also interesting to consider continuous functions $x : [0, \infty) \rightarrow \mathbf{R}^K / \mathbf{Z}^K$, where we can define the discrepancy

$$(112) \quad D_T^{(\mathcal{R})} = \sup_{R \in \mathcal{R}} \left| \frac{1}{T} \int_0^T \chi_R(x(t)) dt - \lambda_K(R) \right|$$

as a function of $T > 0$. (The same can be done with balls, squares, or with caps on the sphere \mathbf{S}^K .)

If the arc length

$$(113) \quad s_T(x) = \int_0^T |dx(t)|$$

is finite for all $T > 0$, we can apply Theorem 1 and 2 (for $K \geq 2$) to get

$$(114) \quad \liminf_{T \rightarrow \infty} D_T^{(\mathcal{R})}(x) s_T(x)^{1+1/(K-1)} > c_K > 0$$

and

$$(115) \quad \liminf_{T \rightarrow \infty} D_T^{(\mathcal{B})}(x) s_T(x)^{\frac{1}{2}+1/(K-1)} > c_K > 0.$$

It is surprising that the exponents $1+1/(K-1)$ ($K \geq 2$) and $1/2+1/(K-1)$ ($K = 2, 3$) are not optimal if \liminf is replaced by \limsup .

THEOREM 7 ([3]). — *If $\phi(T)$ is an arbitrary monotonically increasing function with*

$$(116) \quad \int_0^\infty \frac{1}{\phi(T)} dT < \infty$$

and $x : [0, \infty) \rightarrow \mathbf{R}^K / \mathbf{Z}^K$ (or $\rightarrow \mathbf{S}^K$) ($K \geq 1$) is a continuous function with $\lim_{T \rightarrow \infty} s_T(x) = \infty$, then

$$(117) \quad \limsup_{T \rightarrow \infty} D_T^{(\mathcal{A})}(x) \phi(s_T(x)) > 0,$$

where A is one of the sets $\mathcal{R}, \mathcal{Q}, \mathcal{B}$, (or \mathcal{C}).

For example we have

$$(118) \quad \limsup_{T \rightarrow \infty} D_T^{(\mathcal{R})}(x) s_T(x) (\log s_T(x))^{1+\varepsilon} > 0 .$$

It is worth remarking that Theorem 7 is best possible for $K = 1$ (cf. [3]).

For every increasing function $\phi(T)$ with

$$(119) \quad \int_0^T \frac{1}{\phi(T)} dT = \infty$$

there exists a function $x : [0, \infty) \rightarrow \mathbf{R}^K / \mathbf{Z}^K$ with $\lim_{T \rightarrow \infty} s_T(x) = \infty$ and

$$(120) \quad \limsup_{T \rightarrow \infty} D_T^{(\mathcal{R})}(x) \phi(s_T(x)) = 0 .$$

It seems that (117) is not optimal for $K \geq 2$. For example, one may conjecture that

$$(121) \quad \limsup_{T \rightarrow \infty} D_T^{(\mathcal{R})}(x) s_T(x) (\log s_T(x))^{2-K+\varepsilon} > 0 .$$

For $K = 2$ this would be best possible, since we can prove

THEOREM 8. — *Let $0 < \alpha < 1$ be an irrational number of finite approximation type $\eta < 2$. Then $x(t) = (t, \alpha t)$ satisfies*

$$(122) \quad \limsup_{T \rightarrow \infty} D_T^{(\mathcal{R})}(x) s_T(x) = \sqrt{1 + \alpha^2} \limsup_{T \rightarrow \infty} T D_T^{(\mathcal{R})}(x) < \infty .$$

Proof. — We will use the inequality of Erdős—Turan—Koksma for functions

$$(123) \quad D_T^{(\mathcal{R})}(t, \alpha t) \leq C \left(\frac{1}{H} + \sum_{0 < \max(|h_1|, |h_2|) \leq H} \frac{1}{R(h_1, h_2)} \left| \frac{1}{T} \int_0^T e^{2\pi i(h_1 t + h_2 \alpha t)} dt \right| \right)$$

to estimate the discrepancy. ($R(h_1, h_2) = \max(|h_1|, 1) \max(|h_2|, 1)$) Since

$$(124) \quad \left| \int_0^T e^{2\pi i(h_1 + \alpha h_2)t} dt \right| \leq \frac{1}{\pi |h_1 + h_2 \alpha|} ,$$

we have to estimate

$$(125) \quad 2 \sum_{h_1=1}^H \frac{1}{h_1^2} + 2 \sum_{h_2=1}^H \frac{1}{\alpha h_2^2} + 2 \sum_{h_1, h_2=1}^H \frac{1}{h_1 h_2 (h_1 + \alpha h_2)} + 2 \sum_{h_1, h_2=1}^H \frac{1}{h_1 h_2 |h_1 - \alpha h_2|} = S_1 + S_2 + S_3 + S_4 .$$

Trivially $S_1 = O(1)$ and $S_2 = O(1)$. Since $h_1 + \alpha h_2 \geq 2\sqrt{\alpha h_1 h_2}$, $S_3 = O(1)$, too. In order to estimate S_4 we will split up the sum into three parts

$$(126) \quad \sum_{h_1, h_2=1}^H = \sum_{h_1 \leq \alpha h_2 - \frac{1}{2}} + \sum_{|h_1 - \alpha h_2| < \frac{1}{2}} + \sum_{h_1 \geq \alpha h_2 + \frac{1}{2}} = R_1 + R_2 + R_3$$

and use the fact that

$$(127) \quad \frac{1}{h_1(h_1 - \alpha h_2)} = \frac{1}{\alpha h_2(h_1 - \alpha h_2)} - \frac{1}{\alpha h_1 h_2}.$$

Now it is easy to show that

$$(128) \quad \sum_{h_1 \geq \alpha h_2 + \frac{1}{2}} \frac{1}{\alpha h_2^2(h_1 - \alpha h_2)} = \frac{\pi^2}{6\alpha} \log H + O(1),$$

$$\sum_{h_1 \geq \alpha h_2 + \frac{1}{2}} \frac{1}{\alpha h_1 h_2^2} = \frac{\pi^2}{6\alpha} \log H + O(1),$$

and

$$\sum_{h_1 \leq \alpha h_2 - \frac{1}{2}} \left(\frac{1}{\alpha h_1 h_2^2} + \frac{1}{\alpha h_2^2(\alpha h_2 - h_1)} \right) = O(1).$$

Thus $R_1 + R_3 = O(1)$. Now consider R_2 . For every $h_2 > 1/(2\alpha)$ there is a unique h_1 satisfying $|h_1 - \alpha h_2| < 1/2$. These numbers satisfy $h_1 \gg h_2$ and $|h_1 - \alpha h_2| = \|\alpha h_2\|$ ($\|x\| = \min_{n \in \mathbf{Z}} |x - n|$). α is of approximation type $\eta < 2$. This means that there is a constant c such that $\|\alpha h_2\| \geq c h_2^{-\eta}$ for all $h_2 \geq 1$. It is easy to show that $|\|\alpha p\| - \|\alpha q\|| \geq c(2r)^{-\eta}$ if $0 \leq p < q \leq r$. Thus

$$(129) \quad \sum_{h=1}^r \frac{1}{\|\alpha h\|} \leq \frac{1}{c} \sum_{h=1}^r \frac{(2r)^\eta}{h} = O(r^\eta \log r)$$

implies by partial summation that

$$(130) \quad R_2 = \sum_{|h_1 - \alpha h_2| < \frac{1}{2}} \frac{1}{h_1 h_2 \|\alpha h_2\|} \ll \sum_{1/(2\alpha) < h_2 \leq H} \frac{1}{h_2^2 \|\alpha h_2\|}$$

$$= O\left(\frac{H^\eta \log H}{H^2}\right) + O\left(\sum_{h_2=1}^H \frac{1}{h_2^3} h_2^\eta \log h_2\right)$$

$$= O(1).$$

Therefore we can tend $H \rightarrow \infty$ in (123) to prove (122), i.e. $D_T^{(R)}(t, \alpha t) = O(1/T)$.

Remark 7. — Consider $x(t) = (pt, qt)$ ($0 \leq t \leq 1$), where $p < q$ are distinct prime numbers. Then we can use the inequality of Erdős—Turan—Koksma to get ($H \rightarrow \infty$)

$$(131) \quad D^{(\mathcal{R})}(x) \leq c \left(\frac{1}{H} + 2 \sum_{1 \leq k \leq H/q} \frac{1}{pqk^2} \right) = O \left(\frac{1}{pq} \right).$$

In fact it can be shown that $D^{(\mathcal{R})}(pt, qt) = 1/(4pq)$. By using $p = p_n$ and $q = p_{n+1}$ (p_n is the n -th prime number satisfying $p_n \sim n \log n$) we can deduce (10)

$$(132) \quad \Delta_2^{(\mathcal{R})}(s) \leq \left(\frac{1}{2} + o(1) \right) s^{-2} \quad (s \rightarrow \infty).$$

It is a pity that this method fails to reach the lower bound (9) in higher dimensions.

Remark 8. — The example $x(t) = (t, \alpha t)$ of Theorem 7 seems to be optimal in another sense, too. The author conjectures that there is no continuous function $x : [0, \infty) \rightarrow \mathbf{R}^2/\mathbf{Z}^2$ with finite arc length $s_T(x)$ (for all $T > 0$) such that

$$(133) \quad \lim_{T \rightarrow \infty} T D_T^{(\mathcal{R})} = 0.$$

Only in the one dimensional case it is possible to find functions $x(t)$ satisfying (133). But this is the only example known. On the other hand it is easy to construct functions $x(t)$ (for a very general setting of the discrepancy on compact spaces) such that $\limsup_{T \rightarrow \infty} T D_T^{(\cdot)}(x)$ is arbitrarily small but not 0. (The construction is similar to that of [6].)

The higher dimensional case ($K \geq 3$) seems to be more difficult. Until now the author only knows

THEOREM 9. — *If $K \geq 2$, then there exist continuous functions $x_K : [0, \infty) \rightarrow \mathbf{R}^K/\mathbf{Z}^K$ with $\lim_{T \rightarrow \infty} s_T(x_K) = \infty$ and*

$$(134) \quad \limsup_{T \rightarrow \infty} D_T^{(\mathcal{R})}(x_K) s_T(x) (\log s_T(x))^{1-K} < \infty.$$

Proof. — It is well known that there are sequences $(x_n)_{n=1}^\infty$, on the torus $\mathbf{R}^K/\mathbf{Z}^K$, with

$$(135) \quad D^{(\mathcal{R})}((x_n)_{n=1}^N) \leq c_k \frac{(\log N)^K}{N} \quad (N > 1).$$

Set $l_n = \lfloor \log n \rfloor^K$, $L_N = \sum_{n=1}^N l_n$, and $X_n = \{x_i \mid L_{n-1} \leq i < L_n\}$. Thus $|X_n| = l_n$ and Lemma 6 imply that there is a sequence $(y_n)_{n=1}^\infty$

with $Y_n = \{y_i \mid L_{n-1} \leq i < L_n\} = X_n$ and

$$(136) \quad \sum_{i=L_{n-1}}^{L_n-1} |y_{i+1} - y_i| \leq C(\log n)^{K-1} + \sqrt{K}.$$

Since $X_n = Y_n$ and $|Y_n| \leq (\log n)^K$ we have

$$(137) \quad D^{(R)}((y_n)_{n=1}^N) \leq (c_K + 1) \frac{(\log N)^K}{N}.$$

Therefore the continuous function

$$(138) \quad x_K(t) = \begin{cases} y_n & \text{for } t \in [n-1, n-2^{-n}] \\ y_n + 2^n(t - n + 2^{-n})(y_{n+1} - y_n) & \text{for } t \in [n-2^{-n}, n] \end{cases}$$

satisfies

$$(139) \quad D_T^{(R)}(x_K) \leq D^{(R)}((y_n)_{n=1}^{[T]}) + \frac{2}{T} = O\left(\frac{(\log s_T(x_K))^{K-1}}{s_T(x_K)}\right),$$

since

$$(140) \quad s_T(x_K) \leq \sum_{L_{n-1} \leq T} (C(\log n)^{K-1} + \sqrt{K}) = O\left(\frac{T}{\log T}\right).$$

Remark 9. — It is apparent from the proof on Theorem 9 that lim sup-bounds for $D_T^{(R)}$ imply lim sup-bounds for the discrepancy of sequences. If one could prove $(\alpha < 0)$

$$(141) \quad \limsup_{T \rightarrow \infty} D_T^{(R)}(x) s_T(x) (\log s_T(x))^\alpha > c_K > 0$$

for functions $x : [0, \infty) \rightarrow \mathbf{R}^K / \mathbf{Z}^K$, then (141) would imply

$$(142) \quad \limsup_{T \rightarrow \infty} D^{(R)}((x_n)_{n=1}^N) N (\log N)^\alpha \frac{K}{K-1} > c'_K > 0.$$

Maybe one can improve Roth's bound [8] by this method.

BIBLIOGRAPHIE

- [1] J. BECK, On a problem of K.F. Roth concerning irregularities of point distribution, *Invent. Math.*, 74 (1980), 477-487.
- [2] J. BECK and W. CHEN, *Irregularities of distribution*, Cambridge University Press, Cambridge-New York, 1987.
- [3] M. DRMOTA, An optimal lower bound for the discrepancy of C -uniformly distributed functions modulo 1, *Indag. Math.* (1), 50 (1988), 21-28.

- [4] M. DRMOTA, Untere Schranken für die C -Diskrepanz, Österr. Akad. Wiss., Math. Naturw. Kl. SB II, 196 (1987), 107-117.
- [5] M. DRMOTA and R.F. TICHY, C -uniform distribution on compact metric spaces, J. Math. Analysis & Appl. (1), 129 (1988), 284-292.
- [6] M. DRMOTA and R.F. TICHY, Distribution properties of continuous curves, in : Théorie des nombres, Comptes Rendus de la Conférence internationale de Théorie des nombres tenue à l' Université Laval en 1987 (J.M. De Koninck, C. Levesque eds.), de Gruyter, Berlin - New York, 1989, 117-127.
- [7] E. HLAWKA, Über C -Gleichverteilung, Ann. Mat. Pura Appl. (IV), 49 (1960), 311-366.
- [8] K.F. ROTH, On irregularities of distribution, Mathematika, 1 (1954), 73-79.
- [9] W.M. SCHMIDT, Irregularities of distribution IV, Invent. Math., 7 (1969), 55-82.
- [10] R.J. TASCHNER, The discrepancy of C -uniformly distributed multidimensional functions, J. Math. Analysis and Appl. (2), 78 (1980), 400-404.
- [11] R.F. TICHY, Konvexe Mengen auf speziellen Mannigfaltigkeiten, Proc. 3. Kolloquium über Diskrete Geometrie (1985), 263-275.

Manuscrit reçu le 15 décembre 1988.

Michael DRMOTA,
Technical University of Vienna
Department of Discrete Mathematics
Wiedner Hauptstrasse 8-10/118
A - 1040 Vienna (Austria).