

FINNUR LÁRUSSON

**Compact quotients of large domains in
complex projective space**

Annales de l'institut Fourier, tome 48, n° 1 (1998), p. 223-246

http://www.numdam.org/item?id=AIF_1998__48_1_223_0

© Annales de l'institut Fourier, 1998, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

COMPACT QUOTIENTS OF LARGE DOMAINS IN COMPLEX PROJECTIVE SPACE

by Finnur LÁRUSSON

Introduction.

It has been known for a long time that if a bounded domain Ω in \mathbb{C}^n is a Galois covering space of a compact manifold M , then Ω is a domain of holomorphy and M is projective, meaning that M is isomorphic to a subvariety of some complex projective space \mathbb{P}^N . In fact, the canonical bundle of M is ample. Bounded domains in \mathbb{C}^n can be viewed as domains in \mathbb{P}^n with a large complement: the complement is so large that it contains a hyperplane in its interior. In this paper, we study the other end of the spectrum, following suggestions of Nori [Nor] and Yau [Yau] that this might lead to new and interesting compact complex manifolds, outside the well-known and much-studied classes of manifolds that are algebraic or in some sense close to being algebraic.

We will consider compact complex manifolds M covered by a domain Ω in \mathbb{P}^n whose complement $E = \mathbb{P}^n \setminus \Omega$ is non-empty and small in the sense that the $(2n - 2)$ -dimensional Hausdorff measure $\Lambda_{2n-2}(E)$ vanishes. This condition is just strong enough to exclude hypersurfaces in E . Little work seems to have been done on this subject. Among the few relevant papers in the literature are [Kat1], [Kat2], [Kat3], [Kat4], [Nor], and [Yam].

This work was supported in part by the Natural Sciences and Engineering Research Council of Canada, and by a VPRSC grant from the University of Western Ontario.

Key words: Complex projective space – Covering space – Generalized Kleinian group – Schottky covering – Blanchard manifold.

Math. classification: 32J17 – 32J18 – 32M99.

In Section 1, we establish basic properties of manifolds M of this kind. These include:

- (1) The Kodaira dimension of M is $-\infty$.
- (2) The covering group $\text{Aut}_M \Omega$ is a subgroup of the automorphism group of \mathbb{P}^n , so $\pi_1(M)$ is in fact a generalized Kleinian group.
- (3) There is a lower bound on the size of E , which implies that no 2-dimensional examples exist.
- (4) M is rationally chain connected. The limit set E can be described in terms of rational curves in M that respect the unique projective structure on M .
- (5) M is not of class \mathcal{C} . In particular, M is neither Kähler nor Moishezon.

In Sections 2 and 3, we study in some detail two classes of 3-dimensional examples: the generalized Schottky coverings constructed by Nori, and Blanchard manifolds, for which E is a line, which is the smallest it can be. We determine their fields of meromorphic functions, and describe the surfaces they contain. In Section 4, we make some final remarks on the general 3-dimensional case.

Let us clarify a few terms. By a curve in a complex manifold, we shall mean a (closed analytic) subvariety of pure dimension 1. A surface is a subvariety of pure dimension 2, and a hypersurface is a subvariety of pure codimension 1. When we speak of a manifold, we assume that it is connected.

Acknowledgements. I would like to thank Donu Arapura and Frédéric Campana for helpful discussions, and Sergei Ivashkovich and Masahide Kato for valuable comments on a draft of this paper.

1. Properties of the quotient manifolds.

Let M be an n -dimensional compact complex manifold, $n \geq 2$, covered by a domain Ω in complex projective space \mathbb{P}^n such that the $(2n - 2)$ -dimensional Hausdorff measure $\Lambda_{2n-2}(E)$ of the complement $E = \mathbb{P}^n \setminus \Omega$ is zero. Then Ω is simply connected, so it is the universal covering space of M . Let $\pi : \Omega \rightarrow M$ be the covering map, and $\Gamma \cong \pi_1(M)$ be the covering group. We assume that $\Omega \neq \mathbb{P}^n$, so Γ is infinite.

Let us note that if U is a domain in \mathbb{P}^n , then $U \cap \Omega$ is connected. For this, it actually suffices to have $\Lambda_{2n-1}(E) = 0$. Hence, the compactification \mathbb{P}^n of Ω is finer than the end compactification of Ω . Indeed, the connected components of E correspond bijectively to the ends of Ω , which in turn correspond bijectively to the ends of Γ since M is compact. In particular, E is connected if and only if Γ has only one end.

We will make much use of the following extension theorem, due to Shiffman [Shi1], [Shi2]. See also [HP].

1.1. THEOREM (Shiffman). — *Let E be a closed subset of an n -dimensional complex manifold X . If $\Lambda_{2n-2}(E) = 0$, then holomorphic, meromorphic, and plurisubharmonic functions extend from $X \setminus E$ to X . If $\Lambda_{2n-3}(E) = 0$, then the closure of a hypersurface in $X \setminus E$ is a hypersurface in X .*

The theorem implies that M inherits many properties from \mathbb{P}^n . We see that Ω has no non-constant holomorphic or plurisubharmonic functions, and no non-zero holomorphic p -forms for $p \geq 1$, so

$$H^{p,0}(M) = 0, \quad p \geq 1,$$

and M has a trivial Albanese. Also, no positive power of the canonical bundle of M has any non-zero holomorphic sections, so M has Kodaira dimension $-\infty$.

1.2. PROPOSITION. — *Let $\varphi : \Omega \rightarrow \mathbb{P}^n$ be a holomorphic map.*

(1) *φ extends to a rational map $\mathbb{P}^n \rightarrow \mathbb{P}^n$.*

(2) *If φ is an immersion, then φ extends to an automorphism of \mathbb{P}^n . In particular, every automorphism of Ω is the restriction of a unique automorphism of \mathbb{P}^n , so*

$$\Gamma \subset \text{Aut } \mathbb{P}^n = \text{PGL}(n + 1, \mathbb{C}).$$

By Selberg’s theorem [Sel], the proposition implies that Γ has a normal torsion-free subgroup of finite index. It also implies that $E \subset \mathbb{P}^n$ is a biholomorphic invariant of M , modulo automorphisms of \mathbb{P}^n .

Proof. — The meromorphic function $(z_i/z_0) \circ \varphi$ on Ω extends to a meromorphic function ψ_i on \mathbb{P}^n , and $\psi = [1, \psi_1, \dots, \psi_n]$ is a rational map $\mathbb{P}^n \rightarrow \mathbb{P}^n$ extending φ .

Write $\psi = [q_0, \dots, q_n]$, where q_0, \dots, q_n are homogeneous polynomials in z_0, \dots, z_n of the same degree d , and let $\Psi = (q_0, \dots, q_n) : \mathbb{C}^{n+1} \rightarrow \mathbb{C}^{n+1}$.

Let $p : \mathbb{C}^{n+1} \setminus \{0\} \rightarrow \mathbb{P}^n$ be the canonical projection. Suppose φ is an immersion. Then $\Psi|_{p^{-1}(\Omega)}$ is an immersion. The zero set of the Jacobian determinant $J = \det[\partial q_i / \partial z_j]$ of Ψ is either empty or a hypersurface in $p^{-1}(E) \cup \{0\}$. Since $p^{-1}(E)$ cannot contain a hypersurface, J is a non-zero constant. Also, J is homogeneous of degree $(d - 1)^{n+1}$. Hence $d = 1$, so q_0, \dots, q_n are linear, and ψ is an automorphism of \mathbb{P}^n . \square

Part 2 of the proposition can also be deduced from [Iva1], Theorem 1.

We remark that Ω is maximal among domains in \mathbb{P}^n on which Γ acts with a Hausdorff quotient. Indeed, if Ω' is a domain containing Ω on which Γ acts with a Hausdorff quotient $M' = \Omega' / \Gamma$, then $M \subset M'$ is both open and compact, and hence closed, so since M' is connected, $M = M'$. Therefore, $\Omega' \subset \Gamma\Omega = \Omega$, so $\Omega' = \Omega$.

We now show that there is a lower bound on the size of E .

1.3. PROPOSITION. — *If n is even, then $\Lambda_n(E) > 0$. If n is odd, then $\Lambda_{n-1}(E) > 0$.*

Proof. — Suppose $\Lambda_{2n-2k}(E) = 0$ for an integer k in $[0, n]$. Then Ω contains a k -dimensional complex linear subspace S . Find a sequence $\gamma_i \rightarrow \infty$ in Γ . Then $\gamma_i(S)$ converge to a k -dimensional linear subspace in E , so $\Lambda_{2k}(E) > 0$. Hence, $2k < 2n - 2k$, so $k < n/2$. This shows that if $k \geq n/2$, then $\Lambda_{2n-2k}(E) > 0$, and the proposition follows. \square

Since $\Lambda_{2n-2}(E) = 0$, the proof shows that E contains a line.

1.4. COROLLARY. — *If a domain Ω in \mathbb{P}^2 covers a compact complex manifold, then $\Omega = \mathbb{P}^2$ or $\Lambda_2(\mathbb{P}^2 \setminus \Omega) > 0$.*

The proposition is sharp in the sense that we may have $\Lambda_{n+\epsilon}(E) = 0$ when n is even and $\Lambda_{n-1+\epsilon}(E) = 0$ when n is odd for all $\epsilon > 0$. To see this, let k be $n/2$ if n is even and $(n - 1)/2$ if n is odd, and consider the automorphism φ given by the formula

$$\varphi[z_0, \dots, z_k, z_{k+1}, \dots, z_n] = [2z_0, \dots, 2z_k, z_{k+1}, \dots, z_n].$$

The group Γ of iterates of φ acts freely and properly on $\Omega = \mathbb{P}^n \setminus E$, where E is the union of the two linear subspaces $\{z_0, \dots, z_k = 0\}$ and $\{z_{k+1}, \dots, z_n = 0\}$, and the quotient manifold $M = \Omega / \Gamma$ is compact.

Next we show that M contains many rational curves.

1.5. PROPOSITION. — *If L is a line in Ω , then $\pi(L)$ is a rational curve in M . Hence, M is rationally chain connected.*

For a very general point $p \in M$ and every v in a dense set of tangent vectors at p , there is a smooth rational curve through p which is tangent to v .

For very general points p_1, p_2 in M , there is a connected curve containing p_1 and p_2 which is the union of two smooth rational curves.

Recall that a general point is a point outside a finite union of proper subvarieties, and a very general point is a point outside a countable union of proper subvarieties.

For the proof, we need the following lemma.

1.6. LEMMA. — *For an automorphism φ of \mathbb{P}^n , $n \geq 3$, the following are equivalent:*

- (1) $L \cap \varphi(L) \neq \emptyset$ for all lines L in \mathbb{P}^n .
- (2) φ has a hyperplane of fixed points.

Proof. — (2) \Rightarrow (1) is clear. For the converse, represent φ by a matrix A in $\text{GL}(n + 1, \mathbb{C})$. First let $n = 3$. In suitable coordinates z_0, z_1, z_2, z_3 in \mathbb{C}^4 , A has the Jordan form

$$A = \begin{bmatrix} a_0 & 0 & 0 & 0 \\ \epsilon_1 & a_1 & 0 & 0 \\ 0 & \epsilon_2 & a_2 & 0 \\ 0 & 0 & \epsilon_3 & a_3 \end{bmatrix},$$

where $\epsilon_1, \epsilon_2, \epsilon_3 \in \{0, 1\}$. Suppose $S \cap AS \neq 0$ for all 2-dimensional subspaces S in \mathbb{C}^4 . We need to show that A has a 3-dimensional eigenspace.

Suppose first that A is diagonal. Let $S = \{z_{i_0} = z_{i_1}, z_{i_2} = z_{i_3}\}$ with i_0, i_1, i_2, i_3 mutually distinct. Since $S \cap AS \neq 0$, we get $a_{i_0} = a_{i_1}$ or $a_{i_2} = a_{i_3}$. This means that three of the diagonal entries must be the same, so A has a 3-dimensional eigenspace.

Now suppose A is not diagonal; say $\epsilon_3 = 1$, so $a_2 = a_3$. Let $S = \{z_0 = z_3, z_1 = 0\}$. Since $S \cap AS \neq 0$, there are $x, y \in \mathbb{C}$, not both zero, such that

$$A \begin{bmatrix} x \\ 0 \\ y \\ x \end{bmatrix} = \begin{bmatrix} a_0x \\ \epsilon_1x \\ a_2y \\ y + a_3x \end{bmatrix} \in S,$$

so $\epsilon_1 x = 0$ and $a_0 x = y + a_3 x$. This implies that $\epsilon_1 = 0$. Taking $S = \{z_0 = z_2, z_3 = 0\}$, we get $\epsilon_2 = 0$. Taking $S = \{z_0 = z_1, z_3 = 0\}$, we get $a_0 = a_1$. Finally, taking $S = \{z_0 = z_2, z_1 = z_3\}$, we get $a_0 = a_2$, so A has only one eigenvalue and a 3-dimensional eigenspace corresponding to it.

To disprove (1) in general, it suffices to find a 4-dimensional A -invariant subspace in \mathbb{C}^{n+1} which does not contain a 3-dimensional eigenspace. By examining Jordan forms, it is easy to see that such a subspace exists precisely when A does not have an n -dimensional eigenspace, i.e., when (2) fails. \square

Proof of Proposition 1.5. — Let L be a line in Ω . Then $\pi(L)$ is an irreducible curve in M . Now $y \in L$ is in a fibre of $\pi|L$ with more than one element if and only if $y \in L \cap \gamma L$ for some $\gamma \in \Gamma$, $\gamma \neq \text{id}$. Since Γ acts properly on Ω , there are at most finitely many $\gamma \in \Gamma$ with $L \cap \gamma L \neq \emptyset$. Also, if $\gamma L = L$, then γ has a fixed point in L , so $\gamma = \text{id}$. Hence, $\pi|L$ is injective outside a finite set, so $\pi(L)$ is rational.

For $\gamma \in \Gamma$, let $Y(\gamma)$ be the set of $y \in \mathbb{P}^n$ such that $L \cap \gamma L \neq \emptyset$ for all lines L through y . Then $Y(\gamma)$ is a subvariety of \mathbb{P}^n . We have $\beta Y(\gamma) = Y(\beta\gamma\beta^{-1})$ for $\beta \in \Gamma$. Also, for a compact subset K of Ω , we have $Y(\gamma) \cap K \neq \emptyset$ for only finitely many $\gamma \in \Gamma$ since Γ acts properly on Ω . This implies that $X(\gamma) = \pi(Y(\gamma) \cap \Omega)$ is a subvariety of M . Now $n \geq 3$ by Corollary 1.4, so if $\gamma \neq \text{id}$, then $X(\gamma) \neq M$ by Lemma 1.6. Let $X = \bigcup_{\gamma \neq \text{id}} X(\gamma)$.

Let $p \in M \setminus X$ (so p is a very general point) and $q \in \pi^{-1}(p)$. For $\gamma \in \Gamma$, $\gamma \neq \text{id}$, let \mathcal{L}_γ be the set of lines L in Ω through q such that $L \cap \gamma L = \emptyset$. Then \mathcal{L}_γ is open and dense in the $(n-1)$ -dimensional projective space of lines through q in \mathbb{P}^n . By the Baire category theorem, the intersection $\bigcap \mathcal{L}_\gamma$ is dense. If L is in the intersection, then $\pi|L$ is injective, so $\pi(L)$ is a smooth rational curve through p .

Now let $p_1, p_2 \in M \setminus X$ and $q_k \in \pi^{-1}(p_k)$, $k = 1, 2$. For $\gamma \in \Gamma$, $\gamma \neq \text{id}$, and $k = 1, 2$, let $S(\gamma, k)$ be the union of lines L in Ω through q_k such that $L \cap \gamma L = \emptyset$. Then $S(\gamma, k)$ is open and dense in $\Omega \setminus \{q_k\}$, so the intersection $\bigcap_{\gamma, k} S(\gamma, k)$ is dense in Ω . Hence there are intersecting lines L_1, L_2 in Ω through q_1, q_2 respectively, such that π is injective on both L_1 and L_2 . Then $\pi(L_1) \cup \pi(L_2)$ is a connected union of two smooth rational curves in M containing both p_1 and p_2 . \square

The proposition implies that if $n = 3$ and the algebraic dimension a of M is 0 or 1 (examples of which will be given in sections 2 and 3), then there is no holomorphic surjection f from M onto a 2-dimensional complex manifold. Namely, if $a = 0$, then there are only finitely many surfaces in M . If $a = 1$, then there is a complex manifold X , a compact Riemann surface Y (the algebraic reduction of M), a proper modification $g : X \rightarrow M$, and a holomorphic surjection $h : X \rightarrow Y$, such that with only finitely many exceptions, an irreducible surface in M is an irreducible component of $g(h^{-1}(y))$ for some $y \in Y$. See [FF]. In either case, by the proposition, there is a smooth rational curve C in M which is not contained in any surface in M . But C is contained in the surface $f^{-1}(f(C))$, which is absurd.

Let us recall that an atlas of holomorphic charts on M is called *projective* if the charts map to open sets of \mathbb{P}^n and the coordinate changes are restrictions of automorphisms of \mathbb{P}^n . A projective atlas on M defines an element of $H^1(M, \text{PGL}(n + 1, \mathbb{C}))$, called a *projective structure* on M . Equivalently, a projective structure on M is given by a conjugacy class of group homomorphisms $\pi_1(M) \rightarrow \text{PGL}(n + 1, \mathbb{C})$. Clearly, M has a projective structure. For more information and references, see [Sim]. A projective structure on M yields a developing map, which is a holomorphic immersion from the universal covering space Ω of M to \mathbb{P}^n . By Proposition 1.2, any such map is an automorphism of \mathbb{P}^n , so the projective structure on M is unique.

The projective structure on M defines a germ \mathcal{F}_p of a holomorphic foliation at each point p in M , obtained by pulling back a pencil of lines by a projective chart. If $p \in M$ and $q \in \pi^{-1}(p)$, then the leaf space D_p of \mathcal{F}_p is naturally identified with the space \mathbb{P}^{n-1} of lines through q , so we have a linear projection of $\mathbb{P}^n \setminus \{q\}$ onto D_p . Since E is Γ -invariant, its image in D_p is well defined, regardless of the choice of q .

We say that a curve in M respects the projective structure on M if it appears as a union of straight lines in each projective chart, i.e., if its germ at every point p (or merely at some point in each of its irreducible components) is a union of germs in D_p . Note that the rational curves constructed in the proof of Proposition 1.5 are of this kind.

Our next result relates E to rational curves in M that respect the projective structure.

1.7. PROPOSITION. — *A germ in D_p extends to a rational curve in*

M if and only if it does not lie in the image of E in D_p .

A germ in the image of E in D_p may or may not extend to a curve in M . This can be verified by explicit computations for the example of a Blanchard manifold of type A given in [Kat2] (see Section 3). There, some germs extend to a torus in M , and others do not extend to a curve at all.

Proof. — Let L be the line through q corresponding to a germ in D_p outside the image of E , so $L \subset \Omega$. Then the curve $\pi(L)$ in M is rational by Proposition 1.5.

Conversely, suppose L is a line through q that intersects E , and that the corresponding germ in D_p extends to a rational curve C in M . Then there is a non-constant map $\mathbb{P}^1 \rightarrow C \subset M$, which lifts by π to a map $\mathbb{P}^1 \rightarrow \Omega$, whose image lies in $\pi^{-1}(C) \subset \Gamma(L \setminus E)$. Hence, the image lies in a connected component of $\gamma(L \setminus E)$ for some $\gamma \in \Gamma$, but such a component is isomorphic a domain in \mathbb{C} , which is absurd. \square

It remains to be seen if new information about E can be obtained from this result. It does, however, say something about rational curves in M through a given point that respect the projective structure.

1.8. COROLLARY. — *The set of germs in D_p that do not extend to a rational curve in M is closed, nowhere dense, non-empty, and, when $n = 3$, connected.*

Proof. — Only the last statement needs to be proved. Let L be a line in Ω . Let e be an end of Γ and (γ_n) be a sequence in Γ converging to e . Then the lines $\gamma_n(L)$ converge to the connected component E_0 of E corresponding to e , so E_0 contains a line. This shows that if E_0 is a connected component of E , then the image of E_0 in $D_p \cong \mathbb{P}^{n-1}$ contains a line. If $n = 3$, then two such lines must intersect, so the image of E in D_p is connected. \square

We conclude this section by showing that M is far from being projective.

1.9. PROPOSITION. — *M does not carry a Kähler metric.*

It is easy to see that a domain in \mathbb{P}^n that contains a complex line does not admit a Shafarevich map (also known as a $\tilde{\Gamma}$ -reduction). Since universal covering spaces of compact Kähler manifolds have Shafarevich

maps [Cam2], this implies Proposition 1.9. Also, a domain in a complex manifold is locally Stein if it covers a compact Kähler manifold [Iva2]. Since Ω is not locally Stein, Proposition 1.9 follows. We will give a detailed proof using much simpler means.

The Fubini-Study metric on \mathbb{P}^n is Kähler and invariant under unitary transformations, so the following corollary is immediate.

1.10. COROLLARY. — $\Gamma \not\subset \text{PU}(n + 1)$.

The following lemma is well known. We supply a proof for the convenience of the reader.

1.11. LEMMA. — *If M is a compact Kähler manifold and $H^{2,0}(M) = 0$, then M is projective.*

Proof. — Let ω be a Kähler form on M . We can find $\alpha \in H^2(M, \mathbb{Q})$ so close to $[\omega] \in H^2(M, \mathbb{R})$ that α is positive (but a priori not necessarily of type $(1, 1)$ any more). For some integer $k > 0$ we have $k\alpha \in H^2(M, \mathbb{Z})$. Since $H^{2,0}(M) = 0$, the long exact sequence obtained from the exponential sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathcal{O} \rightarrow \mathcal{O}^\times \rightarrow 0$ shows that $k\alpha$ is the Chern class of a line bundle L on M . Then L is positive, so M is projective. \square

Proof of Proposition 1.9. — Suppose M is Kähler. By the lemma, M is projective, and hence Moishezon, but this contradicts the following proposition. \square

1.12. PROPOSITION. — *M is not Moishezon.*

Proof. — By the extension theorem 1.1, the field $\mathcal{M}(M)$ of meromorphic functions on M can be identified with the field of Γ -invariant meromorphic functions on \mathbb{P}^n . Suppose M is Moishezon, so $\mathcal{M}(M)$ has transcendence degree n over \mathbb{C} . Now $\mathcal{M}(\mathbb{P}^n) = \mathbb{C}(X_1, \dots, X_n)$ also has transcendence degree n over \mathbb{C} , so $\mathcal{M}(\mathbb{P}^n)$ is algebraic over $\mathcal{M}(M)$.

Let $f \in \mathcal{M}(\mathbb{P}^n)$. Then there are $g_1, \dots, g_k \in \mathcal{M}(M)$ such that

$$f^k + g_1 f^{k-1} + \dots + g_k = 0.$$

Say $p \in \Omega$ and g_1, \dots, g_k are all finite at p . Then $f(\gamma p)$, $\gamma \in \Gamma$, are roots of the same polynomial, so the set $f(\Gamma p)$ is finite. Taking $f = z_i/z_0$, $i = 1, \dots, n$, we see that Γ has a finite orbit in Ω , which is absurd. \square

When $n = 3$, by a result of Kato [Kat1], page 53, if the algebraic dimension of M is non-zero, then there is a plane P in \mathbb{P}^3 which is invariant under a subgroup Γ_0 of finite index in Γ . Then we have an embedding of Γ_0 into $\text{Aut } P \cong \text{PGL}(3, \mathbb{C})$ by $\gamma \mapsto \gamma|P$. One might say, therefore, that the “truly” 3-dimensional examples have algebraic dimension zero. In the following sections, we will see 3-dimensional examples with algebraic dimensions 0, 1, and 2.

Let us recall that a compact reduced complex space X belongs to the class \mathcal{C} (as defined by Fujiki) if there is a compact Kähler manifold Y and a holomorphic surjection $Y \rightarrow X$. Equivalently, X is bimeromorphically equivalent to a compact Kähler manifold. All reduced Moishezon spaces are contained in \mathcal{C} . For more information, see [CP] and the references therein.

We need the following property of the class \mathcal{C} .

1.13. THEOREM (Campana [Cam1]). — *Let X be an irreducible compact complex space of class \mathcal{C} . Then a very general point in X is contained in a largest irreducible Moishezon subvariety.*

The following result combines and strengthens Propositions 1.9 and 1.12.

1.14. THEOREM. — *M is not of class \mathcal{C} .*

Proof. — Let p be a point in M . If there is a largest irreducible Moishezon subvariety Y through p , then Y must be M itself by Proposition 1.5. But M is not Moishezon by Proposition 1.12, so M is not of class \mathcal{C} by Campana’s theorem. \square

2. Schottky coverings.

Nori [Nor] has constructed higher-dimensional analogues of the classical Schottky coverings in the following way. Let $n = 2k + 1$, $k \geq 1$, and $g \geq 1$. Choose $2g$ mutually disjoint linear subspaces L_1, \dots, L_{2g} of dimension k in \mathbb{P}^n . Fix an integer i with $1 \leq i \leq g$ and choose a basis so that

$$L_i = \{z_0, \dots, z_k = 0\}, \quad L_{g+i} = \{z_{k+1}, \dots, z_n = 0\}.$$

Define $\varphi_i : \mathbb{P}^n \rightarrow \mathbb{R}$ by the formula

$$\varphi_i[z_0, \dots, z_n] = \frac{|z_0|^2 + \dots + |z_k|^2}{|z_0|^2 + \dots + |z_n|^2},$$

and define open neighbourhoods

$$V_i = \{x \in \mathbb{P}^n : \varphi_i(x) < \alpha\}, \quad V_{g+i} = \{x \in \mathbb{P}^n : \varphi_i(x) > 1 - \alpha\},$$

of L_i, L_{g+i} respectively, where $0 < \alpha < \frac{1}{2}$. Define an automorphism γ_i of \mathbb{P}^n by the formula

$$\gamma_i[z_0, \dots, z_n] = [\lambda z_0, \dots, \lambda z_k, z_{k+1}, \dots, z_n],$$

where $\lambda \in \mathbb{C}$ and $|\lambda| = \frac{1}{\alpha} - 1$. Then $\gamma_i(V_i) = \mathbb{P}^n \setminus \overline{V}_{g+i}$. Let Γ be the subgroup of $\text{PGL}(n + 1, \mathbb{C})$ generated by $\gamma_1, \dots, \gamma_g$. Let A be the complement of $V_1 \cup \dots \cup V_{2g}$, and let $\Omega = \bigcup_{\gamma \in \Gamma} \gamma A$.

Suppose α is so small that the closures of the sets V_1, \dots, V_{2g} are mutually disjoint. Any positive power of γ_i maps $\mathbb{P}^n \setminus \overline{V}_i$ into V_{g+i} , and any negative power of γ_i maps $\mathbb{P}^n \setminus \overline{V}_{g+i}$ into V_i . Hence, any non-trivial word in $\gamma_1, \dots, \gamma_g$ maps the interior of A into its complement, so it is not the identity, and Γ is free on the generators $\gamma_1, \dots, \gamma_g$. Also, Ω is a domain on which Γ acts freely and properly with compact quotient M . Let $\pi : \Omega \rightarrow M$ be the covering map. The compact manifold M is precisely the quotient space of A obtained by identifying the disjoint subsets ∂V_i and ∂V_{g+i} of A by the transformation γ_i for $i = 1, \dots, g$.

The complement E of Ω in \mathbb{P}^n is the closure of the Γ -orbit of $L_1 \cup \dots \cup L_{2g}$. Its connected components are k -dimensional linear subspaces. When $g \geq 2$, E is a ‘‘Cantor set of k -dimensional linear subspaces’’. Given $\epsilon > 0$, E has $(2k + \epsilon)$ -dimensional Hausdorff measure zero if α is small enough. Suppose α is so small that $\Lambda_{2n-2}(E) = 0$. Then Ω is simply connected, so $\pi_1(M) = \Gamma$ is free on g generators.

In this section, M will denote a manifold constructed as above. We will call M a *Schottky manifold*.

Note that if $g = 1$, then the functions

$$\frac{z_1}{z_0}, \dots, \frac{z_k}{z_0}, \frac{z_{k+2}}{z_{k+1}}, \dots, \frac{z_n}{z_{k+1}}$$

descend to algebraically independent meromorphic functions on M , so the algebraic dimension $a(M)$ of M is at least $n - 1$. Also, $E = L_1 \cup L_2$, so $\Lambda_{2n-2}(E) = 0$, and $a(M) = n - 1$ by Proposition 1.12.

Now let M be a 3-dimensional Schottky manifold with $g \geq 2$. The remainder of this section will be concerned with determining the function field of M and the surfaces contained in M for small values of α .

Let Y be an irreducible surface in \mathbb{P}^3 which is invariant under a subgroup of finite index in Γ . Let $1 \leq i \leq g$. Choose a basis so that L_i and L_{g+i} are given as above. Now Y is invariant under γ_i^m for some natural number $m \geq 1$, so if $[z_0, z_1, z_2, z_3] \in Y \setminus (L_i \cup L_{g+i})$, then $[\lambda^{mj} z_0, \lambda^{mj} z_1, z_2, z_3] \in Y$ for all $j \in \mathbb{Z}$. Hence, Y has infinitely many points in common with the line L through the points $[0, 0, z_2, z_3] \in L_i$ and $[z_0, z_1, 0, 0] \in L_{g+i}$, so $L \subset Y$. This shows that Y is a union of lines intersecting both L_i and L_{g+i} , along with either L_i or L_{g+i} . If Y contains only one of the lines L_i, L_{g+i} , then Y is a plane containing that line. The same will hold for other values of i , but that is absurd, since two lines in a plane intersect. Hence, Y contains all the lines L_1, \dots, L_{2g} , and for each i with $1 \leq i \leq g$, Y is a union of lines intersecting both L_i and L_{g+i} .

If Z is another irreducible surface in \mathbb{P}^3 which is invariant under a subgroup of finite index in Γ , then both Y and Z are invariant under the same subgroup Γ' of finite index in Γ . Then $Y \cap Z$ contains the Γ' -orbit of $L_1 \cup \dots \cup L_{2g}$, which is a union of an infinite number of mutually disjoint lines, so $Y = Z$. This shows that there is at most one surface in \mathbb{P}^3 invariant under a subgroup of finite index in Γ .

If f is a meromorphic function on M and $\Lambda_4(E) = 0$, then $f \circ \pi$ extends to a meromorphic function h on \mathbb{P}^3 . Applying the above to the level sets of h gives the following result.

2.1. PROPOSITION. — *Let M be a 3-dimensional Schottky manifold with $g \geq 2$. If $\Lambda_4(E) = 0$, which is the case if α is small enough, then M has no non-constant meromorphic functions.*

In the remainder of the section, we assume that $\Lambda_3(E) = 0$.

Let S be a smooth surface in M . The closure Y of $\pi^{-1}(S)$ is a Γ -invariant surface in \mathbb{P}^3 . If Y is not smooth with singular locus Z , then $Z \subset E$ and Z is Γ -invariant. Since a group with infinitely many ends acts on its space of ends with dense orbits [Kul], this contradicts Z having only a finite number of connected components. Hence, Y is smooth and irreducible. Since Y is covered by rational curves, its degree is 1, 2, or 3. Since Y contains disjoint lines, it cannot be a plane. Since Y contains more than 27 lines, it cannot be a cubic. Hence, Y is a quadric. In suitable projective coordinates, Y is the image of $\mathbb{P}^1 \times \mathbb{P}^1$ under the Segre embedding, and Y has two rulings by lines, which are the only lines in Y . Two disjoint lines in Y must belong to one of the rulings. In particular, the lines L_1, \dots, L_{2g} lie in a quadric.

Conversely, suppose L_1, \dots, L_{2g} lie in a smooth quadric Q in \mathbb{P}^3 . A line in \mathbb{P}^3 is Γ -invariant if and only if it intersects all the lines L_1, \dots, L_{2g} . Since Q is ruled by such lines, it is Γ -invariant. Also, E is a union of lines in one of the rulings. This ruling consists of the fibres of a projection $p : Q \cong \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$. There is an induced action of Γ on $\mathbb{P}^1 \setminus p(E)$, and the quotient map $\mathbb{P}^1 \setminus p(E) \rightarrow C$ is a 1-dimensional Schottky covering onto a compact Riemann surface of genus g . The image of $Q \setminus E$ in M is a smooth surface, ruled over C .

We have proved the following result.

2.2. PROPOSITION. — *Let M be a 3-dimensional Schottky manifold with $g \geq 2$. Suppose $\Lambda_3(E) = 0$, which is the case if α is small enough. Then M contains at most one surface. Also, M contains a smooth surface S if and only if the lines L_1, \dots, L_{2g} lie in a smooth quadric. Then S is a ruled surface of genus g .*

It is easy to see that four lines in \mathbb{P}^3 in general position do not lie in a quadric.

Finally, suppose S is a non-smooth surface in M . Then the singular locus Z of Y is Γ -invariant and not contained in E . For each point $x \in Z \setminus E$ and each i with $1 \leq i \leq g$, there is a line in Z through x intersecting both L_i and L_{g+i} . The singular locus of Z is finite, but also Γ -invariant, so it is empty and Z is smooth. Hence, through every point in $Z \setminus E$ there is a line in Z intersecting all the lines L_1, \dots, L_{2g} , and Z is a disjoint union of a finite number of such lines.

We see that if M contains a surface, then there is a Γ -invariant line in \mathbb{P}^3 . If $g \geq 3$ and L_1, \dots, L_{2g} are in general position, then no such line exists.

3. Blanchard manifolds.

As before, we let M be a compact complex manifold whose universal covering space is a domain Ω in \mathbb{P}^n , such that the complement $E = \mathbb{P}^n \setminus \Omega$ is non-empty with $\Lambda_{2n-2}(E) = 0$. The covering map is $\pi : \Omega \rightarrow M$, and the covering group $\Gamma \cong \pi_1(M)$ can be considered as a subgroup of $\text{PGL}(n + 1, \mathbb{C})$.

3.1. PROPOSITION. — *Suppose E is a subvariety of \mathbb{P}^n . If there is a Γ -invariant curve X in \mathbb{P}^n , not contained in E , then Γ is a finite extension*

of \mathbb{Z} or \mathbb{Z}^2 . Moreover, the normalization of each irreducible component of $\pi(X \setminus E)$ is a torus.

Proof. — Now $X \setminus E$ is Γ -invariant and has finitely many irreducible components. The elements of Γ that leave each component invariant form a normal subgroup Γ' of finite index. Let C be an irreducible component of the curve $\pi(X \setminus E)$ in M , and let Y be an irreducible component of $\pi^{-1}(C)$. Then Y is an irreducible component of $X \setminus E$. Also, Y is a non-compact covering space over C whose covering group Γ'' contains Γ' .

Let Z be the irreducible component of X containing Y . Then $Z \setminus Y$ is finite and non-empty. The normalization \hat{Y} of Y is naturally identified with the complement of a non-empty finite set in the normalization of Z . Each element of Γ'' lifts to a unique automorphism of \hat{Y} . These are the deck transformations of the induced covering map $\hat{Y} \rightarrow \hat{C}$. Since \hat{Y} is a compact Riemann surface with a finite non-zero number of points removed, and \hat{Y} has an infinite automorphism group, \hat{Y} is either the complex plane or the punctured plane. In the first case, $\Gamma'' = \mathbb{Z}^2$, and in the second case, $\Gamma'' = \mathbb{Z}$. In both cases, \hat{C} is a torus. \square

3.2. COROLLARY. — Suppose E is a curve (so $n = 3$). If M contains two distinct irreducible surfaces S_1 and S_2 which intersect, then Γ is a finite extension of \mathbb{Z} or \mathbb{Z}^2 . The intersection $S_1 \cap S_2$ is a curve, and the normalization of each of its irreducible components is a torus.

Proof. — The closure Y_k of $\pi^{-1}(S_k)$ is a surface in \mathbb{P}^3 , and $Y_1 \cap Y_2 \not\subset E$ is a Γ -invariant curve. \square

Now we restrict ourselves to the lowest dimensional case, so we let $n = 3$. We have seen that the smallest E can be is a line. If E is a line, then M is called a *Blanchard manifold*. The first example of such a manifold was given in [Bla]; see also [Kat1]. Kato [Kat2] has shown that if M is a Blanchard manifold, then Γ is torsion-free and contains a subgroup Γ_0 of finite index which is isomorphic to \mathbb{Z}^4 . Furthermore, in a suitable system of homogeneous coordinates z_0, z_1, z_2, z_3 , which we will henceforth adopt, we have $E = \{z_2, z_3 = 0\}$, and either $\Gamma_0 = \Gamma \cap G$ or $\Gamma_0 = \Gamma \cap H$, where

$$G = \left\{ \begin{bmatrix} 1 & a & b & c \\ 0 & 1 & a & b \\ 0 & 0 & 1 & a \\ 0 & 0 & 0 & 1 \end{bmatrix} : a, b, c \in \mathbb{C} \right\}$$

and

$$H = \left\{ \begin{bmatrix} 1 & 0 & a & b \\ 0 & 1 & c & d \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} : a, b, c, d \in \mathbb{C} \right\},$$

considered as subgroups of $\text{PGL}(4, \mathbb{C})$. Note that the groups G and H are abelian. A Blanchard manifold M is said to be of type A if $\pi_1(M)$ contains an abelian subgroup of finite index which is conjugate to a subgroup of G , but $\pi_1(M)$ does not contain an abelian subgroup of finite index which is conjugate to a subgroup of H . If M is not of type A, then M is said to be of type B. Blanchard manifolds of different types are non-isomorphic. There are examples of both types [Kat2].

The Blanchard manifold M constructed in [Bla] is of type B. It is diffeomorphic to the product of the 2-sphere and the real 4-torus [Kat1]. Hence, the underlying smooth manifold of M carries a projective algebraic complex structure, in addition to the complex structure being considered here, which by Theorem 1.14 is not of class C . In particular, M does not fail to be Kähler due to any topological obstruction. Note that there are no 2-dimensional examples of this kind, due to the fact that a compact complex surface carries a Kähler metric if and only if its first Betti number is even.

From now on, we let M be a Blanchard manifold.

Consider the planes in \mathbb{P}^3 containing the line E . They foliate Ω , and the leaf space is \mathbb{P}^1 . The covering group Γ permutes these planes, so there is an induced holomorphic foliation \mathcal{F} of M . Its leaves are the images of the planes. The leaves are smooth, but not necessarily closed in M .

3.3. PROPOSITION. — *A plane P in \mathbb{P}^3 containing E is invariant under a subgroup of finite index in Γ if and only if $T = \pi(P \setminus E)$ is closed in M . Then T is a smooth connected surface in M , and T is either a torus or a hyperelliptic surface. If $\Gamma_0 = \Gamma$, then T is a torus.*

Proof. — First of all, T is closed in M if and only if $\Gamma(P \setminus E) = \pi^{-1}(T)$ is closed in Ω if and only if the Γ -orbit of P consists of finitely many planes if and only if P is invariant under a subgroup of finite index in Γ .

Suppose the subgroup $\Gamma' = \{\gamma \in \Gamma : \gamma P = P\}$ is of finite index in Γ . If $p \in T$ and $x, y \in \pi^{-1}(p)$, then $y = \gamma x$ for some $\gamma \in \Gamma$, so $\gamma P \cap P \cap \Omega \neq \emptyset$,

and $\gamma P = P$. This shows that $P \setminus E \cong \mathbb{C}^2$ is a Galois covering space of T with covering group Γ' .

Now P is given by an equation of the form $sz_2 + tz_3 = 0$ with $s, t \in \mathbb{C}$. Simple computations show that the zero-free holomorphic 2-form $dz_0 \wedge dz_1$ on $P \setminus E$ is Γ_0 -invariant, so it descends to $T_0 = (P \setminus E)/(\Gamma' \cap \Gamma_0)$. Hence, the canonical bundle of T_0 is trivial, and the Kodaira dimension of T_0 is zero. Also, $\Gamma' \cap \Gamma_0$ is of finite index in Γ_0 , which is isomorphic to \mathbb{Z}^4 , so the first Betti number of T_0 is 4. Hence, by the Enriques-Kodaira classification [BPV], T_0 is a torus.

This shows that T has a torus as a finite unbranched covering. Hence, T is Kähler, the Kodaira dimension of T is zero, and the fundamental group of T is infinite. By the Enriques-Kodaira classification, T is either a torus or a hyperelliptic surface. □

We recall that every hyperelliptic surface X is the quotient of a torus (in fact, a product of two smooth elliptic curves) by a finite group acting freely. In particular, X is projective, so any torus covering X must be an abelian variety. Furthermore, a torus is never homeomorphic to a hyperelliptic surface, e.g. since they have different first Betti numbers (4 and 2 respectively).

If M is of type B, then all the planes containing E are invariant under Γ_0 , so by the proposition, M is foliated by smooth surfaces, and each leaf is either a torus or a hyperelliptic surface. On the other hand, if M is of type A, then only the plane $\{z_3 = 0\}$ is invariant under a subgroup of finite index in Γ , so by the proposition, \mathcal{F} has only one closed leaf, and this leaf is a torus or a hyperelliptic surface.

Now let $T = \pi(P \setminus E)$, where $P = \{z_3 = 0\}$. We know that T is a smooth connected surface in M . Suppose $S \neq T$ is a surface in M . The closure X of $\pi^{-1}(S)$ is a Γ -invariant surface in \mathbb{P}^3 which intersects P in a curve. By Proposition 3.1 (applied with Γ replaced by Γ_0), the intersection must be E . Also by Proposition 3.1, $X \setminus E$ is smooth, so S is smooth. Choose an irreducible component Y of X . It is the zero locus of an irreducible homogeneous polynomial g . Since $Y \cap P = E$, we may take $g = z_2^m + z_3h$, where $m \geq 1$ is the degree of g , and h is a polynomial. Now Y is invariant under a subgroup of finite index in Γ , so for every γ in a subgroup of finite index in Γ_0 there is a constant $c \neq 0$ such that

$$cg = \gamma^*g = (z_2 + az_3)^m + z_3\gamma^*h = z_2^m + z_3k,$$

where $a \in \mathbb{C}$ and k is a polynomial, so $c = 1$ and g is γ -invariant. Hence, the non-constant meromorphic function $f = g/z_3^m$ on \mathbb{P}^3 is invariant under

a normal subgroup Γ_1 of finite index in Γ . The indeterminacy locus of f is E . The restriction $f|_\Omega : \Omega \rightarrow \mathbb{P}^1$ is a Γ_1 -invariant holomorphic map with smooth fibres, and it descends to a holomorphic map $f_1 : M_1 = \Omega/\Gamma_1 \rightarrow \mathbb{P}^1$. Let $f_2 : M_1 \rightarrow C_1$ be the Stein factorization of f_1 . The 1-dimensional complex space C_1 is normal, and hence smooth, so it is a compact Riemann surface. The fibres of f_2 are the connected components of the fibres of f_1 .

The finite group Γ/Γ_1 acts on M_1 with quotient M . By Corollary 3.2, the fibres of f_2 are the only irreducible surfaces in M_1 , so they are permuted by Γ/Γ_1 . Hence, there is an induced action of Γ/Γ_1 on C_1 that makes f_2 equivariant. Passing to quotients, we obtain a non-constant holomorphic map $\psi : M \rightarrow C$, where C is a normal complex space, and hence a compact Riemann surface. The fibres of ψ are smooth, and they are the images of the fibres of f_2 , so they are connected. The map ψ lifts to a non-constant holomorphic map from Ω to the universal covering \tilde{C} of C , but Ω carries no non-constant holomorphic functions, so \tilde{C} must be \mathbb{P}^1 , and $C = \mathbb{P}^1$.

So far we have proved the following.

3.4. THEOREM. — *If M is a Blanchard manifold, then one of the following holds.*

(1) *M contains only one surface, which is a torus or a hyperelliptic surface. Hence, M has no non-constant meromorphic functions.*

(2) *There is a holomorphic map $\psi : M \rightarrow \mathbb{P}^1$ with smooth connected fibres, which are the only irreducible surfaces in M . Hence, $\mathcal{M}(M) = \mathcal{M}(\mathbb{P}^1) \circ \psi$, and the algebraic dimension of M is 1.*

If M is of type B, then (2) holds, and the fibres of ψ are the leaves of \mathcal{F} .

We now wish to understand the dichotomy in the theorem when M is of type A. First we consider the automorphism group of M . Let N_Γ be the normalizer of Γ in $\text{PGL}(4, \mathbb{C})$. If $\nu \in N_\Gamma$, then $\Gamma\nu E = \nu\Gamma E = \nu E$, so the line νE is Γ -invariant. Now $\nu E \not\subset \Omega$, so νE must intersect E in the unique Γ -fixed point $[1, 0, 0, 0]$ of E . Computations show that E is the only Γ -invariant line through $[1, 0, 0, 0]$, so $\nu E = E$. This shows that Ω is N_Γ -invariant. Therefore, every $\nu \in N_\Gamma$ induces an automorphism of M , which is the identity if and only if $\nu \in \Gamma$. Conversely, every automorphism of M lifts to an element of N_Γ , so

$$\text{Aut } M = N_\Gamma/\Gamma.$$

Let $P = \{z_3 = 0\}$ be the unique plane in \mathbb{P}^3 containing E , which is

invariant under a subgroup of finite index in Γ . Since P is Γ_0 -invariant, νP is invariant under the group $\nu\Gamma_0\nu^{-1}$, which is of finite index in Γ , so $\nu P = P$. This shows that the surface $T = (P \setminus E)/\Gamma$ in M is $\text{Aut } M$ -invariant.

Assume now that $\Gamma \subset G$. This amounts to replacing M by the finite unbranched covering space $M_0 = \Omega/\Gamma_0$. Clearly, (1) in Theorem 3.4 holds for M if and only if it holds for M_0 .

Since G is abelian, $G \subset N_\Gamma$. This shows that the dimension of the complex Lie group $\text{Aut } M$ is at least 3. It is easy to see that G acts transitively on $\Omega \setminus P$ (with trivial stabilizers) and on $P \setminus E$, so G acts transitively on $M \setminus T$ and on T . Hence, $\text{Aut } M$ has precisely two orbits, namely $M \setminus T$ and T . In particular, M is almost homogeneous.

Let

$$N = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

The commuting matrices N , N^2 , and N^3 form a basis for the Lie algebra \mathfrak{g} of G , and the exponential map $\exp : \mathfrak{g} \rightarrow G$,

$$rN + sN^2 + tN^3 \mapsto I + rN + \left(\frac{1}{2}r^2 + s\right)N^2 + \left(t + \frac{1}{6}r^3 + rs\right)N^3,$$

is an algebraic group isomorphism with inverse $\log : G \rightarrow \mathfrak{g}$,

$$I + aN + bN^2 + cN^3 \mapsto aN + \left(b - \frac{1}{2}a^2\right)N^2 + \left(c - ab + \frac{1}{3}a^3\right)N^3.$$

Suppose now that T is not the only surface in M . By Theorem 3.4, there is a holomorphic map $\psi : M \rightarrow \mathbb{P}^1$ with smooth connected fibres, which are the only irreducible surfaces in M . Say $T = \psi^{-1}(\infty)$. Then G permutes the fibres of ψ , and we get an induced action of G on \mathbb{P}^1 which is transitive on \mathbb{C} and fixes ∞ . The stabilizer of $0 \in \mathbb{C}$ is a proper algebraic subgroup H of G , and $\Gamma \subset H$. In particular, Γ is not Zariski-dense in G . We have $H = \exp \mathfrak{h}$, where \mathfrak{h} is a 2-dimensional subspace of \mathfrak{g} . If \mathfrak{h} is defined by an equation of the form $ar + bs = 0$ with $a, b \in \mathbb{C}$, then the polynomial function

$$f[z, w, 1, 0] = bz - \frac{1}{2}bw^2 + aw$$

on $P \setminus E$ is H -invariant, and hence Γ -invariant, so f descends to a non-constant holomorphic function on T , but this is absurd since T is compact. Hence, \mathfrak{h} is defined by an equation of the form $t = ar + bs$ with $a, b \in \mathbb{C}$.

For $r \in \mathbb{C}$, let w_r be a solution of the quadratic equation

$$w^2 + (r + b)w + \frac{1}{3}r^2 + \frac{1}{2}br - a = 0,$$

and let

$$s_r = -r \left(\frac{1}{2}r + w_r \right).$$

Then

$$h_r = \exp(\tau N + s_r N^2 + (ar + bs_r)N^3) \in H$$

maps $p_r = [0, 0, w_r, 1]$ to $[0, 0, w_r + r, 1]$. Since $w_r \rightarrow \infty$ and $w_r + r \rightarrow \infty$ as $r \rightarrow \infty$, both p_r and $h_r p_r$ converge to the point $[0, 0, 1, 0]$ in Ω . This shows that H does not act properly on Ω .

Now $\Gamma = \exp \Lambda$, where Λ is a lattice in \mathfrak{h} . There is a constant $c_0 > 0$ such that for every $v \in \mathfrak{h}$ there is $u \in \Lambda$ with $|u - v| < c_0$. Hence, there is a constant $c > 0$ such that for every $h \in H$ there is $\gamma \in \Gamma$ such that all the off-diagonal entries of γh^{-1} have absolute value less than c . Let $\gamma_r \in \Gamma$ be associated to h_r in this way. Then $\gamma_r \rightarrow \infty$ as $r \rightarrow \infty$. Let U be the neighbourhood of the point $[0, 0, 1, 0]$ defined by the inequality

$$|z_0| + |z_1| + |z_3| < |z_2|.$$

Then U is relatively compact in Ω . There is $\rho > 0$, depending only on c , such that if $g \in G$ and all the off-diagonal entries of g have absolute value less than c , then gU lies in the compact subset K of Ω defined by the inequality

$$|z_0| + |z_1| \leq \rho(|z_2| + |z_3|).$$

For $r \in \mathbb{C}$ sufficiently large, we have $h_r p_r \in U$, so $\gamma_r p_r = (\gamma_r h_r^{-1}) h_r p_r \in K$. This shows that Γ does not act properly on Ω , contrary to hypothesis.

We have proved the following.

3.5. THEOREM. — *If M is a Blanchard manifold of type A, then M contains only one surface, and Γ is Zariski-dense in G .*

This shows that the classification of Blanchard manifolds by type is the same as their classification by algebraic dimension.

Note that Γ is Zariski-dense in G if and only if $\log(\Gamma \cap G)$ spans \mathfrak{g} as a \mathbb{C} -vector space.

It would be interesting to have a characterization of those discrete Zariski-dense subgroups Γ of G of rank 4 that act freely and properly on Ω with compact quotient, so that Ω/Γ is a Blanchard manifold of type A.

Proposition 3.3 and Theorems 3.4 and 3.5 imply the following.

3.6. COROLLARY. — *An irreducible surface in a Blanchard manifold is a torus or a hyperelliptic surface. If $\Gamma_0 = \Gamma$, then it is a torus.*

If M is of type B, then the map $\psi : M \rightarrow \mathbb{P}^1$ in Theorem 3.4 is a proper submersion, and hence a smooth fibre bundle by Ehresmann's fibration theorem [BJ], over the complement of its finite set of critical values. Hence, all but finitely many of the surfaces in M are mutually diffeomorphic, so either all but finitely many surfaces in M are tori, or all but finitely many surfaces in M are hyperelliptic.

3.7. Example. — We will now show that hyperelliptic surfaces can occur in Blanchard manifolds of type A.

In [Kat2], Kato constructs a Blanchard manifold $M_0 = \Omega/\Gamma_0$ of type A, such that Γ_0 is the subgroup of G generated by $I + N$, $I + iN$, $I + N^2$, and $I + iN^2$.

Let

$$\varphi = \begin{bmatrix} 1 & 1 & i/2 & 0 \\ 0 & -1 & -1 & -i/2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & -1 \end{bmatrix} \in \text{PGL}(4, \mathbb{C}).$$

Then

$$\varphi N = -N\varphi,$$

so φ commutes with $I + N^2$ and $I + iN^2$. Also,

$$\varphi(I + N)\varphi^{-1} = I - N \quad \text{and} \quad \varphi(I + iN)\varphi^{-1} = I - iN$$

are in Γ_0 , since their logarithms lie in the \mathbb{Z} -span of the logarithms of the given generators of Γ_0 . Hence, Γ_0 is a normal subgroup of the subgroup Γ of $\text{PGL}(4, \mathbb{C})$ spanned by Γ_0 and φ . Moreover, $\varphi^2 = I + (i - 1)N^2 \in \Gamma_0$, so Γ_0 is of index 2 in Γ .

This shows that φ induces a holomorphic involution of M_0 . It has a fixed point in M_0 if and only if there are $\gamma \in \Gamma_0$ and $x \in \Omega$ such that $\varphi x = \gamma x$. Then

$$\varphi^2 x = \varphi \gamma x = (\varphi \gamma \varphi^{-1})(\varphi x) = \varphi \gamma \varphi^{-1} \gamma x,$$

so $\varphi^2 = \varphi \gamma \varphi^{-1} \gamma$ since Γ_0 acts freely on Ω . Hence, $\varphi \gamma^{-1} \in \Gamma$ is of order 2. We will show that this cannot happen.

For $\gamma = I + aN + bN^2 + cN^3 \in \Gamma_0$, we have

$$(\varphi\gamma)^2 = \begin{bmatrix} 1 & 0 & 2b - a^2 - 1 + i & * \\ 0 & 1 & 0 & * \\ 0 & 0 & 1 & * \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

so if $(\varphi\gamma)^2 = \text{id}$, then $a^2 + 1 = 2b + i$. Now $\log \gamma$ is a \mathbb{Z} -linear combination of the logarithms of the given generators of Γ_0 , so

$$a = n_1 + in_2, \quad b = \frac{1}{2}(n_1 + in_2)^2 + \frac{1}{2}(n_2 - n_1) + n_3 + in_4,$$

where $n_1, \dots, n_4 \in \mathbb{Z}$. Hence, the imaginary part of $a^2 + 1$ is an even integer, whereas the imaginary part of $2b + i$ is an odd integer. This shows that Γ has no elements of order 2, so φ has no fixed points in M_0 , and $M = M_0/\varphi = \Omega/\Gamma$ is a Blanchard manifold of type A with fundamental group Γ .

The unique surface $T = (P \setminus E)/\Gamma$ in M has the torus $(P \setminus E)/\Gamma_0$ in M_0 as a 2-sheeted unbranched covering space, and its fundamental group is Γ . Let $\alpha : \Gamma \rightarrow \Gamma' = \Gamma/[\Gamma, \Gamma]$ be the abelianization map. Then $\alpha(\Gamma_0)$ has finite index (1 or 2) in Γ' . Also, Γ_0 contains non-trivial commutators, such as $\varphi(I - N)\varphi^{-1}(I - N)^{-1}$, so the rank of Γ' , i.e., the first Betti number of T , is less than 4. Hence, T is hyperelliptic.

It has been pointed out that certain Blanchard manifolds of type B are twistor spaces, as are certain 3-dimensional Schottky manifolds with $g = 1$. See [Bes], Chapter 13.D,E and [Hit]. On the other hand, we have the following result.

3.8. PROPOSITION. — *A Blanchard manifold of type A is not a twistor space.*

Proof. — Suppose a Blanchard manifold of type A is a twistor space. The twistor structure pulls up to the universal covering Ω , so Ω is smoothly fibred by smooth rational curves, and Ω has a Γ -equivariant fixed-point-free anti-holomorphic involution τ preserving the fibres, which extends to a Γ -equivariant anti-holomorphic involution of \mathbb{P}^3 . The normal bundle N of a fibre C is $H_1 \oplus H_1$, where H_k denotes the hyperplane bundle on \mathbb{P}^k .

By the short exact sequence

$$0 \rightarrow TC \rightarrow T\mathbb{P}^3|_C \rightarrow N \rightarrow 0$$

on C , we have

$$4c_1(H_3|C) = c_1(T\mathbb{P}^3|C) = c_1(TC) + c_1(N) = 2c_1(H_1) + 2c_1(H_1)$$

in $H^2(C, \mathbb{Z}) = \mathbb{Z}$, so $\deg C = \deg(H_3|C) = 1$, and C is a line.

Let $P = \{z_3 = 0\}$ be the unique Γ_0 -invariant plane in \mathbb{P}^3 containing E . Now $\tau(z) = \varphi(\bar{z})$ for some $\varphi \in \text{Aut } \mathbb{P}^3$, so τP is a plane in \mathbb{P}^3 containing $\tau E = E$, invariant under $\tau\Gamma_0\tau^{-1} = \Gamma_0$, so $\tau P = P$.

Let C be a fibre in Ω . Clearly, $C \not\subset P$, so C intersects P in a single point x . Since C and P are τ -invariant, $\tau x = x$, which is absurd. \square

4. The 3-dimensional case.

In conclusion, we will make some remarks on the general 3-dimensional case. Let us make the following definition, motivated by [Kat2]. A 3-dimensional compact complex manifold M is called a *Kato manifold* if

- (1) the universal covering space of M is a domain Ω in \mathbb{P}^3 such that every connected component of the complement $\mathbb{P}^3 \setminus \Omega$ is a line,
- (2) the fundamental group of M has a torsion-free subgroup of finite index, and
- (3) M contains a domain which is biholomorphic to a neighbourhood of a line in \mathbb{P}^3 .

The following main result of [Kat2] classifies Kato manifolds and describes their fundamental groups. For terms left undefined here, we refer the reader to [Kat2].

4.1. THEOREM (Kato). — *Let M be a Kato manifold and let Γ_0 be a torsion-free subgroup of finite index in $\pi_1(M)$. Then Γ_0 is isomorphic to the free product*

$$\Gamma_1 * \dots * \Gamma_r * \Gamma_{r+1} * \dots * \Gamma_s,$$

where $0 \leq r \leq s$, $\Gamma_1, \dots, \Gamma_r$ are infinite cyclic, and $\Gamma_{r+1}, \dots, \Gamma_s$ contain \mathbb{Z}^4 as a subgroup of finite index. Also, the finite covering of M with fundamental group Γ_0 is a Klein combination of r primary L -Hopf manifolds and $s - r$ Blanchard manifolds.

Now we let M be a compact complex manifold whose universal covering space is a domain Ω in \mathbb{P}^3 , such that the complement $E = \mathbb{P}^3 \setminus \Omega$ is non-empty with $\Lambda_4(E) = 0$. The covering map is $\pi : \Omega \rightarrow M$.

By Proposition 1.2 and Selberg's Theorem, M satisfies (2). As for (3), by the proof of Proposition 1.5, there is a line L in Ω such that $\pi|L$

is injective. If π is not injective on any neighbourhood of L , then there are $p_n \rightarrow p \in L$ and $q_n \rightarrow q \in L$ with $p_n \neq q_n$ such that $\pi(p_n) = \pi(q_n)$, so $\pi(p) = \pi(q)$ and $p = q$, but this is absurd because π is locally injective. Hence, M satisfies (3).

This shows that M is a Kato manifold if and only if all the connected components of E are lines. Recall that in the proof of Corollary 1.8, we observed that every connected component of E contains a line, so roughly speaking, M is a Kato manifold if the connected components of E are as small as they can possibly be.

If M is Blanchard or Schottky, then M is Kato. However, M need not be Kato. Kato [Kat3] has constructed examples where E is a smooth submanifold of \mathbb{P}^3 of real dimension 3 (in fact, E is a circle of projective lines), and Γ is the fundamental group of a hyperbolic compact Riemann surface. These examples are twistor spaces of algebraic dimension zero.

All the examples of compact manifolds covered by “large” domains in projective space that I know of have been mentioned in this paper. There are clearly very few of them, especially in dimensions greater than 3, and they do not seem to give any clues to a possible classification. In particular, we have seen that although the class of these manifolds intersects the classes of twistor spaces and Kato manifolds, it is not contained in either of them. The work that has been done to date gives a glimpse of a rich and varied theory, but it is only a beginning.

BIBLIOGRAPHY

- [BPV] W. BARTH, C. PETERS, A. VAN DE VEN, *Compact Complex Surfaces*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, 3. Folge, Band 4, Springer-Verlag, 1984.
- [Bes] A. L. BESSE, *Einstein Manifolds*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, 3. Folge, Band 10, Springer-Verlag, 1987.
- [Bla] A. BLANCHARD, *Sur les variétés analytiques complexes*, *Ann. Sci. Ecole Norm. Sup.*, 73 (1956), 157–202.
- [BJ] T. BRÖCKER, K. JÄNICH, *Introduction to differential topology*, Cambridge University Press, 1982.
- [Cam1] F. CAMPANA, *Coréduction algébrique d’un espace analytique faiblement kählérien compact*, *Invent. Math.*, 63 (1981), 187–223.
- [Cam2] F. CAMPANA, *Remarques sur le revêtement universel des variétés kählériennes compactes*, *Bull. Soc. Math. France*, 122 (1994), 255–284.
- [CP] F. CAMPANA, Th. PETERNELL, *Cycle spaces*, *Several Complex Variables VII*, *Encyclopaedia of Mathematical Sciences*, vol. 74, Springer-Verlag, 1994.

- [FF] G. FISCHER, O. FORSTER, Ein Endlichkeitssatz für Hyperflächen auf kompakten komplexen Räumen, *J. reine angew. Math.*, 306 (1979), 88–93.
- [HP] R. HARVEY, J. POLKING, Extending analytic objects, *Comm. Pure Appl. Math.*, 28 (1975), 701–727.
- [Hit] N. J. HITCHIN, Kählerian twistor spaces, *Proc. London Math. Soc.*, (3) 43 (1981), 133–150.
- [Iva1] S. M. IVASHKOVICH, Extension of locally biholomorphic mappings of domains into complex projective space, *Math. USSR Izvestiya*, 22 (1984), 181–189.
- [Iva2] S. M. IVASHKOVICH, The Hartogs-type extension theorem for meromorphic maps into compact Kähler manifolds, *Invent. Math.*, 109 (1992), 47–54.
- [Kat1] M. KATO, On compact complex 3-folds with lines, *Japan. J. Math.*, 11 (1985), 1–58.
- [Kat2] M. KATO, Factorization of compact complex 3-folds which admit certain projective structures, *Tôhoku Math. J.*, (2) 41 (1989), 359–397.
- [Kat3] M. KATO, A non-Kähler structure on an S^2 -bundle over a ruled surface (1992), unpublished preprint.
- [Kat4] M. KATO, Compact quotient manifolds of domains in a complex 3-dimensional projective space and the Lebesgue measure of limit sets, *Tokyo J. Math.*, 19 (1996), 99–119.
- [Kul] R. S. KULKARNI, Some topological aspects of Kleinian groups, *Amer. J. Math.*, 100 (1978), 897–911.
- [Nor] M. V. NORI, The Schottky groups in higher dimensions, *Contemporary Mathematics*, 58, Part I (1986), 195–197.
- [Sel] A. SELBERG, On discontinuous groups in higher-dimensional symmetric spaces, *Contributions to Function Theory*, Tata Institute of Fundamental Research, (1960), 147–164.
- [Shi1] B. SHIFFMAN, On the removal of singularities of analytic sets, *Michigan Math. J.*, 15 (1968) 111–120.
- [Shi2] B. SHIFFMAN, Extension of positive line bundles and meromorphic maps, *Invent. Math.*, 15 (1972), 332–347.
- [Sim] R. R. SIMHA, The monodromy representation of projective structures, *Arch. Math.*, 52 (1989), 413–416.
- [Yam] A. YAMADA, Small deformations of certain compact manifolds of class L . II, *Tokyo J. Math.*, 10 (1987), 189–202.
- [Yau] S.-T. YAU, A review of complex differential geometry, *Proc. Symp. Pure Math.*, 52, Part 2 (1991), 619–625.

Manuscrit reçu le 1er septembre 1997,
accepté le 23 septembre 1997.

Finnur LÁRUSSON,
University of Western Ontario
Department of Mathematics
London, Ont. N6A 5B7 (Canada).
larusson@uwo.ca