

# ANNALES DE L'INSTITUT FOURIER

F. Y. WU

C. KING

W. T. LU

## **On the rooted Tutte polynomial**

*Annales de l'institut Fourier*, tome 49, n° 3 (1999), p. 1103-1114

[http://www.numdam.org/item?id=AIF\\_1999\\_\\_49\\_3\\_1103\\_0](http://www.numdam.org/item?id=AIF_1999__49_3_1103_0)

© Annales de l'institut Fourier, 1999, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## ON THE ROOTED TUTTE POLYNOMIAL (\*)

by F.Y. WU, C. KING and W.T. LU

---

### 1. The Tutte polynomial.

Consider a finite graph  $G$  with vertex set  $V$  and edge set  $E$ . A *spanning subgraph*  $G'(S) \subseteq G$  is a subgraph of  $G$  containing all members of  $V$  and an edge set  $S \subseteq E$ . Let  $C$  be a set of  $q$  distinct colors. A  $q$ -*coloring* of  $G$  is a coloring of the vertices in  $V$  such that two vertices connected by an edge bear different colors. It is well-known that the number of  $q$ -colorings of  $G$  is given by the chromatic polynomial (see [1])

$$(1) \quad P(G; q) = \sum_{S \subseteq E} q^{p(S)} (-1)^{|S|},$$

where  $p(S)$  is the number of components in the spanning subgraph  $G'(S)$ . Alternately, we can regard (1) as generating colorings of components of spanning subgraphs of  $G$  with  $q$  colors with an edge weight  $-1$ .

As an extension of the chromatic polynomial, Tutte [2], [3], [4] introduced what is now known as the *Tutte polynomial*

$$(2) \quad Q(G; t, v) = \sum_{S \subseteq E} t^{p(S)} v^{|S| - |V| + p(S)}.$$

Indeed, one has the relation

$$(3) \quad P(G; q) = (-1)^{|V|} Q(G; -q, -1).$$

---

(\*) This work is supported in part by the National Science Foundation grants DMR-9614170 (FYW and WTL) and DMS-9705779 (CK).

Keywords: Graph colorings – Rooted Tutte polynomial – Planar partition – Duality relation.

Math. classification: 05C30 – 05C15.

In view of (3), it is useful to write (2) as

$$(4) \quad Q(G; t, v) = v^{-|V|} \sum_{S \subseteq E} (vt)^{p(S)} v^{|S|},$$

so that for  $vt = q =$  positive integers, the Tutte polynomial (4) generates colorings of components of spanning subgraphs of  $G$  with  $q$  colors and edge weights  $v$ , instead of  $v = -1$ .

For planar  $G$  with dual graph  $G_D$ , it is well-known that the Tutte polynomial possesses the duality relation

$$(5) \quad v Q(G; t, v) = t Q(G_D; v, t),$$

a relation first observed by Whitney [5].

## 2. The rooted Tutte polynomial.

We extend the definition (4) to a rooted Tutte polynomial.

A vertex is rooted, or is a root, if it is colored with a prescribed (fixed) color. A graph is rooted if it contains rooted vertices.

Let  $R$  denote a set of  $n$  roots located at vertices  $\{r_1, r_2, \dots, r_n\}$ . A *color configuration* is a map  $x: R \mapsto C$ , and as a convenient shorthand we write  $x(r_i) = x_i$  for  $i = 1, 2, \dots, n$ . A component of a spanning subgraph is *exterior* if it contains one or more roots, and is *interior* otherwise. An exterior component is *proper* if all roots in the component are of the same color. A spanning subgraph  $G'(S)$  is proper if all its exterior components are proper. An edge set  $S_x \subseteq E$  is proper if the spanning subgraph  $G'(S_x)$  it generates is proper.

For a prescribed color configuration  $\{x_1, x_2, \dots, x_n\}$  of the  $n$  roots, we introduce in analogy to (4) the *rooted* Tutte polynomial<sup>(1)</sup>

$$(6) \quad Q_{x_1 x_2 \dots x_n}(G; t, v) = v^{-|V|} \sum_{S_x \subseteq E} (vt)^{p_{\text{in}}(S_x)} v^{|S_x|},$$

where the summation is taken over all proper edge sets  $S_x$ , and  $p_{\text{in}}(S_x)$  is the number of interior components of  $G'(S_x)$ . Thus, as in (3), we have for

<sup>(1)</sup> Strictly speaking, it is the expression  $v^{|V|} Q_{x_1 x_2 \dots x_n}(G; t, v)$  which is a polynomial in  $v$  and  $t$ .

positive integral  $q$  the relation

$$(7) \quad (-1)^{|V|} Q_{x_1 x_2 \dots x_n}(G; -q, -1) \\ = \text{the number of } q\text{-colorings of } G \\ \text{with color configuration } \{x_1, x_2, \dots, x_n\}.$$

Clearly, the expression (6) depends on how the  $n$  roots are partitioned into subsets of different colors, and the actual colors do not enter the picture.

The coloring configuration  $\{x_1, x_2, \dots, x_n\}$  induces a partition  $X$  of  $R$  into blocks (subsets) such that all roots in one block are of one color, and colors of different blocks are different. Namely, two elements  $r_i, r_j \in R$  belong to the same block of  $X$  if and only if they have the same prescribed color  $x_i = x_j$ .

Consider now the summation in (6). Let  $G'(S)$  be any (not necessarily proper) spanning subgraph of  $G$ . The connected components of  $G'(S)$  induce a partition on the set of vertices  $V$  of  $G$ . We get hence also a partition  $\pi(S)$  on the set of rooted vertices  $R$  by restricting this partition to  $R$ . Clearly, the spanning subgraph  $G'(S_x)$  is proper if and only if the partition  $\pi(S_x)$  is a refinement of the partition  $X$ . It follows that we can rewrite (6) as

$$(8) \quad Q_X(G; t, v) = \sum_{X' \preceq X} F_{X'}(G; t, v),$$

where

$$(9) \quad F_{X'}(G; t, v) \equiv v^{-|V|} \sum_{\substack{S_x \subseteq E \\ \pi(S_x) = X'}} (vt)^{p_{\text{in}}(S_x)} v^{|S_x|}.$$

Here, we have abbreviated  $Q_{x_1 x_2 \dots x_n}$  by  $Q_X$ , which is permitted since the actual colors do not enter the picture at this point. Also it is understood that  $G$  is now a rooted graph, with root set  $R$ .

The expression (8) assumes the form of a transformation of a partially ordered set. Its inverse is given by the Möbius inversion

$$(10) \quad F_X(G; t, v) = \sum_{X'} \mu(X', X) Q_{X'}(G; t, v),$$

where (see [6])

$$(11) \quad \mu(X', X) = \begin{cases} (-1)^{|X'| - |X|} \prod_{\text{blocks} \in X} (n_b(X') - 1)!, & \text{if } X' \preceq X, \\ 0, & \text{otherwise,} \end{cases}$$

$n_b(X')$  being the number of blocks of  $X'$  that are contained in the block  $b$  of  $X$ . Note that for  $n = 1$  we have  $|X| = |X'| = 1$ ,  $p_{\text{in}}(S_x) = p(S_x) - 1$ ,

and all edge sets  $S \subseteq E$  are proper. Hence we have

$$(12) \quad F_X(G;t, v) = Q_X(G;t, v) = (vt)^{-1}Q(G;t, v), \quad n = 1.$$

This completes the definition and general description of the rooted Tutte polynomial for any graph  $G$ .

### 3. Planar graphs.

From here on we consider  $G$  being planar with the  $n$  roots residing around a single face of  $G$ . Without the loss of generality, we can choose the face to be the infinite face and order the roots in the sequence  $\{r_1, r_2, \dots, r_n, r_1\}$  as shown in Fig. 1.




Figure 1. A planar graph  $G$  with  $n$  roots. The graph is denoted by the shaded region and the  $n$  roots by the black circles.

A partition  $X$  of the  $n$  roots is *non-planar* if two roots of one block separate two roots of another block in the cyclic sequence. Otherwise  $X$  is *planar*. For a given  $n$ , there are  $b_n$  partitions, where (see [9])

$$(13) \quad b_n = \sum_{m_\nu=0}^{\infty} \left[ n! / \prod_{\nu=1}^{\infty} (\nu!)^{m_\nu} m_\nu! \right], \quad \sum_{\nu=1}^{\infty} \nu m_\nu = n,$$

and of the  $c_n$  partitions

$$(14) \quad c_n = \frac{(2n)!}{n!(n+1)!}$$

are planar (see [7], [8]). We shall adopt the convention of writing

$$X = \{ij, kl, \dots, \dots\}$$

for colors  $\{x_i = x_j, x_k = x_l = \dots, \dots\}$ , with  $\{ij\}, \{kl, \dots\}, \dots$  each in order (see [8]). For example, two partitions for  $n = 5$  are

$$(15) \quad \begin{aligned} X_1 &= \{123, 4, 5\}, & |X_1| &= 3, & \text{planar,} \\ X_2 &= \{24, 351\}, & |X_2| &= 2, & \text{non-planar.} \end{aligned}$$

Now if  $G$  is planar and  $X'$  is non-planar then by definition the summand in (9) is empty and one has  $F_{X'}(G;t, v) = 0$ . Thus we have

PROPOSITION 1. — *For planar  $G$*

$$(16) \quad F_X(G; t, v) = 0, \quad \text{if } X \text{ is non-planar.}$$

This proposition was first established in [9] for the Potts model correlation function (see section 6) by considering its graphical expansion similar to the consideration given in the above.

As a consequence of Proposition 1 and the use of (10), we now have:

COROLLARY 1. — *Rooted Tutte polynomials associated with non-planar partitions can be written as linear combinations of the rooted Tutte polynomials associated with (refined) planar partitions.*

Corollary 1 leads to the sum-rule identities reported in [9] for the Potts correlation function. In the case of  $n = 4$ , for example, the identity

$$F_{\{13,24\}}(G; t, v) = 0$$

leads to the sum rule (see [9])

$$(17) \quad Q_{\{13,24\}}(G; t, v) = Q_{\{13,2,4\}}(G; t, v) + Q_{\{1,3,24\}}(G; t, v) \\ - Q_{\{1,2,3,4\}}(G; t, v).$$

From here on we shall restrict our considerations to rooted Tutte polynomials associated with the  $c_n$  planar partitions only.

#### 4. The graph $G^*$ .

The rooted Tutte polynomial (6) possesses a duality relation for planar graphs, which relates the rooted Tutte polynomial on a graph  $G$  to that of a related graph  $G^*$ . Here we define  $G^*$ . Starting from a planar  $G$ , place an extra vertex  $f$  in the infinite face and connect it to each root of  $G$  by an edge. This gives a new graph  $G''$ , which has one more vertex than  $G$  and  $n$  additional edges. The dual graph of  $G''$  is also planar, and it has a face  $F$  containing the extra vertex  $f$ . Now remove the  $n$  edges on the boundary of  $F$ , and the resulting graph is  $G^*$ .

It is readily seen that the graph  $G^*$  has

$$(18) \quad |V^*| = |V_D| + n - 1$$

vertices where  $|V_D|$  is the number of vertices of  $G_D$ , the dual of  $G$ , and there is a one-one correspondence between the edges of  $G$  and  $G^*$ . We denote the set of  $n$  vertices  $\{r_1^*, r_2^*, \dots, r_n^*\}$  of  $G^*$  surrounding the face  $F$  by  $R^*$ , with  $r_i^*$  residing between the two edges  $\langle f, r_{i-1} \rangle$  and  $\langle f, r_i \rangle$  of  $G''$ , where  $r_0 = r_n$ .

An example of a  $G$  and the related  $G^*$  for  $n = 4$  is shown in Fig. 2.




Figure 2. A graph  $G$  (solid lines) and the related graph  $G^*$  (broken lines) for  $n = 4$ . Black circles denote roots of  $G$  and crosses denote vertices of  $G^*$ .

Clearly, the relation of  $G$  to  $G^*$  is reciprocal, namely, we have

$$(G^*)^* = G.$$

Now each planar partition  $X$  of  $R$  induces a partition  $X^*$  of the set  $R^*$  (see [10]). In order to define  $X^*$ , for each block  $b$  of  $X$  we choose a point in the infinite face of  $G$ , and connect all roots  $r_i$  in  $b$  to this point by drawing new edges. Because  $X$  is planar, the points for the blocks can be chosen so that the edges of different blocks do not cross, and the resulting extended graph is still planar. So this process divides the infinite face into regions. The induced partition  $X^*$  is then described by the condition that all roots of  $R^*$  in one region are regarded as belonging to one block of  $X^*$ . Alternately, for another way of defining  $X^*$ , let  $P_{ij}$ ,  $1 \leq i < j \leq n$ , be a partition of  $R$  into two blocks, the sets  $\{r_i, \dots, r_{j-1}\}$  and  $\{r_j, \dots, r_{i-1}\}$ , where all numbers are modulo  $n$ . Then, the partition  $X^*$  induced by  $X$  is defined by the condition that the two roots  $r_i^*$  and  $r_j^*$  belong to the same block in  $X^*$  if and only if  $X$  is a refinement of  $P_{ij}$ .

If  $X$  induces  $X^*$ , we write

$$X \longrightarrow X^*.$$

Clearly,  $X^*$  is planar, and we have

$$(19) \quad |X| + |X^*| = n + 1.$$

For the planar partition  $X_1$  in (15), for example, we have

$$(20) \quad X_1 = \{123, 4, 5\} \longrightarrow X_1^* = \{2, 3, 451\}, \quad |X_1| = |X_1^*| = 3.$$

However, as a result of our labelling convention, the color configuration of the partition  $(X^*)^*$  further induced by  $X^*$  is a cyclic shift of that of  $X$ , namely,

$$(21) \quad \{x_1, x_2, \dots, x_n\} \longrightarrow X^* \longrightarrow \{x_n, x_1, \dots, x_{n-1}\}.$$

In the example above, for instance, we have

$$(22) \quad \{123, 4, 5\} \longrightarrow \{2, 3, 451\} \longrightarrow \{234, 5, 1\}.$$

Finally, there is a one-one correspondence between the edge sets  $E$  of  $G$  and  $E^*$  of  $G^*$ , an edge set  $S \subseteq E$  defines a “complement” edge set  $S^* \subseteq E^*$  by the condition that an edge is included in  $S^*$  if and only if its corresponding edge is not included in  $S$ . Clearly, we have

$$(S^*)^* = S.$$

## 5. The duality relation.

The rooted Tutte polynomial arises in statistical physics as the correlation function of the Potts model (see next section). In a recent paper [10] we have established a duality relation for the Potts correlation function for planar  $G$ . However, the proof of the duality relation given in [10] is cumbersome and not easily deciphered in graphical terms. Here we re-state the results as two propositions in the context of the rooted Tutte polynomial, and present direct graph-theoretical proofs of the propositions.

**PROPOSITION 2.** — *For planar  $G$  and  $G^*$  and the associated planar partitions  $X \rightarrow X^*$ , we have*

$$(23) \quad v^{|X|} F_X(G; t, v) = t^{|X^*|} F_{X^*}(G^*; v, t).$$



*Proof.* — Let  $S_x$  be a proper edge set on  $G$ . We have the Euler relation

$$(24) \quad |S_x| + |S_x^*| = |V| + |V_D| - 2$$

and, after eliminating  $n$  and  $|V_D|$  using (18), (19) and (24), the identity

$$(25) \quad |S_x| + |X| - |V| = |S_x^*| + |X^*| - |V^*|,$$

which holds for any proper edge set  $S_x$ . Note that we have also the fact

$$(26) \quad \pi(S_x) = X \quad \text{if and only if} \quad \pi(S_x^*) = X^*.$$

Let  $c(S_x^*)$  the number of independent circuits in the spanning subgraph  $G'(S_x^*)$ . Then we have

$$(27) \quad p(S_x) = c(S_x^*) + |X|.$$

Also, starting from the  $|V^*|$  isolated vertices on  $G^*$ , one constructs  $G'(S_x^*)$  by drawing edges of  $S_x^*$  on  $G^*$  one at a time. Since each edge reduces the number of components by one except when the adding of an edge completes an independent circuit, one has also

$$(28) \quad p(S_x^*) = |V^*| - |S_x^*| + c(S_x^*).$$

Eliminating  $c(S_x^*)$  using (27) and (28) and making use of the relations

$$(29) \quad \begin{cases} p(S_x) = p_{\text{in}}(S_x) + |X|, \\ p(S_x^*) = p_{\text{in}}(S_x^*) + |X^*|, \end{cases}$$

one obtains

$$(30) \quad p_{\text{in}}(S_x^*) = p_{\text{in}}(S_x) + |V^*| - |S_x^*| - |X^*|.$$

Proposition 2 now follows from the substitution of (30) into the right-hand side of (23) where, explicitly,

$$(31) \quad F_{X^*}(G^*; v, t) = t^{-|V^*|} \sum_{\substack{S_x^* \subseteq E^* \\ \pi(S_x^*) = X^*}} (vt)^{p_{\text{in}}(S_x^*)} t^{|S_x^*|},$$

and the use of the identities (25) and (26). This completes the proof of Proposition 2.

Proposition 2 was first conjectured in [9] and established later in [10] in the context of Potts correlation functions (see next section) without the explicit reference to the polynomial form (9).

*Remark.* — For  $n = 1$ , the duality relation (23) for the rooted Tutte polynomial becomes the duality relation (5) for the Tutte polynomial. This is a consequence of (12).

PROPOSITION 3.

1) *The rooted Tutte polynomials associated with the  $c_n$  planar partitions for  $G$  and  $G^*$  are related by the duality transformation*

$$(32) \quad Q_X(G; t, v) = \sum_Y \mathbf{T}_n(X, Y) Q_Y(G^*; v, t),$$

where  $\mathbf{T}_n$  is a  $c_n \times c_n$  matrix with elements

$$(33) \quad \mathbf{T}_n(X, Y) = t^{n+1} \sum_{X' \preceq X} (vt)^{-|X'|} \mu(Y, Y'), \quad X' \rightarrow Y'.$$

2) *The matrix  $\mathbf{T}_n$  satisfies the identity*

$$(34) \quad [\mathbf{T}_n]^2(X, X') = \delta(x_1, x'_2) \delta(x_2, x'_3) \cdots \delta(x_n, x'_1).$$

*Proof.* — The transformation (32) follows by combining (8) and (10) with Proposition 2, and its uniqueness is ensured by the uniqueness of the Möbius inversion. The property (34) is a consequence of (21). □

Proposition 3 was first given in [10] in the context of the Potts correlation function (see next section). Explicit expression of  $\mathbf{T}_n$  for  $n = 2, 3, 4$  can be found in [10] and [11].

### 6. The Potts and the random cluster models.

It is well-known in statistical physics that the Tutte polynomial gives rise to the partition function of the Potts model (see [12]). In view of the prominent role played by the Potts model in many fields in physics, it is useful to review this equivalence and the further equivalence of the rooted Tutte polynomial with the Potts correlation function.

The  $q$ -state Potts model [13] is a spin model defined on a graph  $G$ . The spin model consists of  $|V|$  spins placed at the vertices of  $G$  with each spin taking on  $q$  different states and interacting with spins connected by edges. Without going into details of the physics [12] which lead to the Potts model, it suffices for our purposes to define the Potts partition function

$$(35) \quad Z(G; q, v) \equiv \sum_{S \subseteq E} q^{p(S)} v^{|S|},$$

the  $n$ -point partial partition function

$$(36) \quad Z_X(G; q, v) \equiv \sum_{S_x \subseteq E} q^{p_{\text{in}}(S_x)} v^{|S_x|},$$

and the  $n$ -point correlation function

$$(37) \quad P_n(G; x_1, x_2, \dots, x_n) = P_n(G; X) \equiv Z_X(G; q, v) / Z(G; q, v),$$

where again, in analogy to notation in Sections 1 and 2, we have denoted the color configuration  $\{x_1, x_2, \dots, x_n\}$  by the associated partition  $X$ . More generally, for any real or complex  $q$ , the partition function (35) defines the random cluster model of Fortuin and Kasteleyn [14], which coincides with the Potts model for integral  $q$ .

Relating this to the Tutte polynomial, we now have

$$(38) \quad \begin{cases} Z(G; q, v) = v^{|V|} Q(G; t, v), \\ Z_X(G; q, v) = v^{|V|} Q_X(G; t, v), \\ P_n(G; X) = Q_X(G; t, v) / Q(G; t, v), \end{cases}$$

for  $q = vt$ . The duality relation (5) for the Tutte polynomial then implies the following duality relation for the Potts partition function [10], [13], [15]

$$(39) \quad v^{1-|V|} Z(G; q, v) = (v^*)^{1-|V_D|} Z(G_D; q, v^*),$$

where

$$(40) \quad vv^* = q.$$

One further defines the dual correlation function

$$(41) \quad P_n^*(G^*; X^*) \equiv q Z_{X^*}(G^*; q, v^*) / Z(G_D; q, v^*),$$

and also the functions  $A_X$  and  $B_{X^*}$  by

$$(42) \quad P_n(G; X) = \sum_{X' \preceq X} A_{X'}(G; q, v)$$

and

$$(43) \quad P_n^*(G^*; X^*) = \sum_{X^{*'} \preceq X^*} B_{X^{*'}}(G^*; q, v^*).$$

Then, Proposition 2 leads to the relation

$$(44) \quad A_X(G; q, v) = q^{-|X|} B_{X^*}(G^*; q, v^*), \quad X \rightarrow X^*.$$

which is the main result of [10].

### 7. Summary and discussions.

We have introduced the rooted Tutte polynomial (6) as a two-variable polynomial associated with a rooted graph and deduced a number of pertinent results.

Our first result is that the rooted Tutte polynomial assumes the form (8) of a partially order set for which the inverse can be uniquely determined. For planar graphs and all roots residing surrounding a single face, we showed that (Proposition 1) the inverse function vanishes for non-planar partitions of the roots. We further showed that the inverse function satisfies the duality relation (23) (Proposition 2) which, in turn, leads to the duality (32) for the rooted Tutte polynomial (Proposition 3). We also reviewed the connection of the Tutte and rooted Tutte polynomials with the Potts model in statistical physics.

Finally, we remark that results reported here have previously been obtained in [9] and [10] in the context of the Potts correlation function. Here, the results are reformulated as properties of the rooted Tutte polynomial and thereby permitting graph-theoretical proofs.

*Noted added.* — The polynomial  $Q(G; x, y)$  is now commonly referred to as the *dichromatic polynomial*, and the dichromate

$$\chi(G; x, y) = (x - 1)^{-1} Q(G; x - 1, y - 1)$$

is now commonly known as the *Tutte polynomial*. We are indebted to Professor W.T. Tutte for this remark.

*Acknowledgments.* — We are grateful to the referee for providing an independent proof of Proposition 1 and numerous suggested improvements on an earlier version of this paper.

## BIBLIOGRAPHY

- [1] G.D. BIRKHOFF, A determinant formula for the number of ways of coloring of a map, *Ann. Math.*, 14 (1912), 42–46.
- [2] W.T. TUTTE, A contribution to the theory of chromatic polynomials, *Can. J. Math.*, 6 (1954), 80–91.
- [3] W.T. TUTTE, On dichromatic polynomials, *J. Comb. Theory*, 2 (1967), 301–320.
- [4] W.T. TUTTE, Graph Theory, in *Encyclopedia of Mathematics and Its Applications*, Vol. 21, Addison–Wesley, Reading, Massachusetts, 1984, Chap. 9.
- [5] H. WHITNEY, The coloring of graphs, *Ann. Math.*, 33 (1932), 688–718.
- [6] See, for example, L.J. VAN LINT and R.M. WILSON, *A course in combinatorics*, Cambridge University Press, Cambridge, 1992, p. 301.
- [7] H.N.V. TEMPERLEY and E.H. LIEB, Relations between the percolation and colouring problem and other graph–theoretical problems associated with regular planar lattice: some exact results for the percolation problem, *Proc. Royal Soc. London A*, 322 (1971), 251–280.
- [8] W.T. TUTTE, The matrix of chromatic joins, *J. Comb. Theory B*, 57 (1993), 269–288.
- [9] F.Y. WU and H.Y. HUANG, Sum rule identities and the duality relation for the Potts  $n$ -point boundary correlation function, *Phys. Rev. Lett.*, 79 (1997), 4954–4957.
- [10] W.T. LU and F.Y. WU, On the duality relation for correlation functions of the Potts model, *J. Phys. A: Math. Gen.*, 31 (1998), 2823–2836.
- [11] F.Y. WU, Duality relations for Potts correlation functions, *Phys. Letters A*, 228 (1997), 43–47.
- [12] See, for example, F.Y. WU, The Potts Model, *Rev. Mod. Phys.*, 54 (1982), 235–268.
- [13] R.B. POTTS, Some generalized order–disorder transformations, *Proc. Camb. Philos. Soc.*, 48 (1954), 106–109.
- [14] C.M. FORTUIN and P.W. KASTELEYN, On the random–cluster model I. Introduction and relation to other models, *Physica*, 57 (1972), 536–564.
- [15] F.Y. WU and Y.K. WANG, Duality transformation in a many–component spin model, *J. Math. Phys.*, 17 (1976), 439–440.

F.Y. WU & W.T. LU,  
 Northeastern University  
 Department of Physics  
 Boston, Massachusetts 02115 (USA).  
 fywu@neu.edu  
 wtlu@albert.physics.neu.edu  
 &  
 C. KING,  
 Northeastern University  
 Department of Mathematics  
 Boston, Massachusetts 02115 (USA).  
 king@neu.edu