



ANNALES

DE

L'INSTITUT FOURIER

Carolyn S. GORDON, Pierre GUERINI,
Thomas KAPPELER & David L. WEBB

Inverse spectral results on even dimensional tori

Tome 58, n° 7 (2008), p. 2445-2501.

http://aif.cedram.org/item?id=AIF_2008__58_7_2445_0

© Association des Annales de l'institut Fourier, 2008, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

INVERSE SPECTRAL RESULTS ON EVEN DIMENSIONAL TORI

by Carolyn S. GORDON, Pierre GUERINI,
Thomas KAPPELER & David L. WEBB (*)

ABSTRACT. — Given a Hermitian line bundle L over a flat torus M , a connection ∇ on L , and a function Q on M , one associates a Schrödinger operator acting on sections of L ; its spectrum is denoted $\text{Spec}(Q; L, \nabla)$. Motivated by work of V. Guillemin in dimension two, we consider line bundles over tori of arbitrary even dimension with “translation invariant” connections ∇ , and we address the extent to which the spectrum $\text{Spec}(Q; L, \nabla)$ determines the potential Q . With a genericity condition, we show that if the connection is invariant under the isometry of M defined by the map $x \rightarrow -x$, then the spectrum determines the even part of the potential. In dimension two, we also obtain information about the odd part of the potential. We obtain counterexamples showing that the genericity condition is needed even in the case of two-dimensional tori. Examples also show that the spectrum of the Laplacian defined by a connection on a line bundle over a flat torus determines neither the isometry class of the torus nor the Chern class of the line bundle.

In arbitrary dimensions, we show that the collection of all the spectra $\text{Spec}(Q; L, \nabla)$, as ∇ ranges over the translation invariant connections, uniquely determines the potential. This collection of spectra is a natural generalization to line bundles of the classical Bloch spectrum of the torus.

RÉSUMÉ. — À un fibré en droites hermitien sur un tore plat M , une connexion ∇ sur L et une fonction Q sur M , on associe un opérateur de Schrödinger agissant sur les sections de L ; on note $\text{Spec}(Q; L, \nabla)$ son spectre. À la suite du travail de V. Guillemin en dimension deux, on considère des fibrés en droites complexes au-dessus de tores de dimension paire quelconque ainsi qu’une connexion « invariante par translation » fixée et on se demande dans quelle mesure $\text{Spec}(Q; L, \nabla)$ détermine le potentiel Q . Sous une condition générique, on montre que le spectre détermine la partie paire du potentiel, à condition que la connexion soit invariante par l’isométrie du tore définie par l’application $x \mapsto -x$. En dimension deux, on obtient également des informations sur sa partie impaire. On obtient des contre-exemples qui montrent que la condition générique utilisée est nécessaire même dans le cas des tores de dimension deux. Ces exemples montrent aussi que le spectre du laplacien

Keywords: Schrödinger operator, spectrum, line bundles over tori.

Math. classification: 58J50, 58J53.

(*) Gordon and Webb were supported in part by NSF grants DMS-0072534 and DMS-0306752; Guerini and Kappeler were supported in part by the Swiss National Science Foundation and by the European Research Training Network HPRN-CT-1999-00118.

défini par une connexion sur un fibré en droites sur un tore plat ne détermine ni la classe d'isométrie du tore ni la classe de Chern du fibré.

En dimension quelconque, on montre que la collection de tous les spectres $Spec(Q; L, \nabla)$, lorsque ∇ parcourt l'ensemble des connexions invariantes, détermine le potentiel de manière unique. Cette collection de spectres est une généralisation naturelle aux fibrés en droites du spectre classique de Bloch sur le tore.

CONTENTS

1.	Introduction	2446
2.	Line bundles over tori: construction and classification	2450
3.	Negative inverse spectral results on rectangular tori	2464
4.	Positive inverse spectral results on tori	2472
	Appendices	2478
	Appendix A. Wave kernel on \mathbb{R}^n	2478
	Appendix B. Wave kernel acting on sections of a line bundle	2485
	Appendix C. Expansion of the wave trace at singularities	2486
	Appendix D. Proof of Proposition C.2	2495
	Bibliography	2500

1. Introduction

Let M be a flat n -dimensional torus $\mathcal{L} \backslash \mathbb{R}^n$. To each smooth function Q on M , viewed as an \mathcal{L} -periodic function $Q : \mathbb{R}^n \rightarrow \mathbb{R}$, one associates a Schrödinger operator $\Delta + Q$. In their pioneering work [4], G. Eskin, J. Ralston, and E. Trubowitz address the extent to which the potential Q is determined by its periodic spectrum $Spec(Q)$, i.e., by the spectrum of the Schrödinger operator acting on \mathcal{L} -periodic functions, and also by the more extensive data of the Bloch spectrum. The latter associates to each linear functional α on \mathbb{R}^n the spectrum $Spec_\alpha(Q)$ of the Schrödinger operator acting on smooth functions on \mathbb{R}^n satisfying the condition $f(x+l) = e^{2\pi i \alpha(l)} f(x)$ for all $l \in \mathcal{L}$. In what follows, we will refer to the Bloch spectrum as the “classical Bloch spectrum.” As explained in Section 2, the classical Bloch spectrum can be viewed as the spectrum of the bundle Laplacian determined by a flat connection on the trivial complex line bundle over M .

Given a Hermitian line bundle L over M and a connection ∇ on L , one associates a Laplace operator acting on smooth sections of L . A smooth function Q on M then gives rise to a Schrödinger operator acting on sections of the bundle. The spectrum $Spec(Q; L, \nabla)$ of this operator depends both on the connection and on the potential. In case L is the trivial bundle and ∇ the trivial connection, then $Spec(Q; L, \nabla)$ coincides with the periodic spectrum of Q . The torus $M = \mathcal{L} \backslash \mathbb{R}^n$ is said to have *nondegenerate length spectrum* if for each $l \in \mathcal{L}$, the only elements of \mathcal{L} of length $|l|$ are $\pm l$. G. Eskin [3] addressed the problem of determining both the connection

and potential from $Spec(Q; L, \nabla)$ in the case of an arbitrary connection and potential on trivial line bundles L over two-dimensional tori with nondegenerate length spectrum. Our work is motivated by that of V. Guillemin [10] who addressed the same question for line bundles of Chern number one. For 2-tori with nondegenerate length spectrum and for the class of connections and potentials that are invariant under the isometry of M defined by the map $x \rightarrow -x$, he first proved under curvature bounds on the connection that $Spec(Q; L, \nabla)$ determines the connection. He then proved that for a given connection, $Spec(Q; L, \nabla)$ determines the potential.

In this article, we consider Schrödinger operators on Hermitian line bundles L over higher-dimensional tori. In general dimensions, it seems unlikely that one could recover the connection from the spectrum of the Schrödinger operator. Thus we fix either a single connection or a family of specified connections on a line bundle L and ask whether the potential function can be determined from the associated spectral data. Associated to each connection on L is a curvature 2-form on M . By an abuse of language, we will refer to a connection as translation-invariant if the associated curvature 2-form is translation-invariant, i.e., invariant under the group of isometries of the torus given by the translations. For a given potential function Q , the mapping $\nabla \mapsto Spec(Q; L, \nabla)$ as ∇ ranges over the translation-invariant connections is a natural analog of the classical Bloch spectrum (as explained in Remark 2.23) and thus will be referred to as the *L-Bloch spectrum* of Q . Among the translation invariant connections on L , there is a unique connection (up to gauge equivalence) that is also \mathbb{Z}_2 -invariant, i.e., invariant under the map $x \rightarrow -x$ of \mathbb{R}^n , and exactly 2^n connections that are *weakly* \mathbb{Z}_2 -invariant in the sense that they are gauge equivalent to their pullbacks with respect to the map $x \rightarrow -x$. We address two questions in this article:

- Given a weakly \mathbb{Z}_2 -invariant, translation invariant connection ∇ on L , does $Spec(Q; L, \nabla)$ determine Q ?
- Does the *L-Bloch spectrum* of Q determine Q ?

A line bundle over a torus is uniquely determined up to equivalence by its Chern class. We consider bundles whose Chern class is represented by a nondegenerate two-form. This condition forces the dimension of M to be even, say $n = 2m$. As will be explained in Section 2, associated to the Chern class is an ordered m -tuple (r_1, \dots, r_m) of positive integers such that $r_i | r_{i+1}$ for $i = 1, \dots, m - 1$; we refer to the integers r_j as the *invariant factors* of the Chern class. The m -tuple of invariant factors is a topological invariant of the line bundle but not a complete invariant (although in the case $m = 1$, r_1 is, up to sign, the Chern number of the bundle).

For our positive results, we restrict attention to the case in which all the invariant factors of the Chern class equal 1, the analog of the setting studied by Guillemin in dimension 2. In this setting, we give an affirmative answer to the second question even without assuming nondegeneracy of the length spectrum. For the first question, we obtain some positive results in the setting of tori with nondegenerate length spectrum. We also construct counterexamples illustrating that, both in the two-dimensional case studied by Guillemin and in the higher dimensional case studied here, the hypothesis of nondegeneracy of the length spectrum cannot be dropped.

We first state the negative results. Two potentials Q_1 and Q_2 on a torus M are said to be *congruent* if there exists an isometry σ of M such that $Q_2 = Q_1 \circ \sigma$.

THEOREM 1.1. — *Let M be a rectangular nonsquare 2-torus for which the ratio of the side lengths is rational. Then for every line bundle L over M with nondegenerate Chern class and for every translation invariant connection ∇ on L , there exist pairs of noncongruent potentials Q_1 and Q_2 on M such that $\text{Spec}(Q_1; L, \nabla) = \text{Spec}(Q_2; L, \nabla)$. The potentials may be chosen to be real analytic.*

A slightly weaker result holds in arbitrary even dimensions, as explained in Section 3.

According to Theorem 4.2 in [6] Part I, the potentials we construct do not have the same periodic spectrum. This example is contrary to the general belief that “spectral rigidity” in the case of a non trivial line bundle is more pronounced than in the case of a trivial line bundle.

The method used to construct the examples in Theorem 1.1 also yields results concerning the spectrum of the Laplace operator itself. Letting $\text{Spec}(L, \nabla)$ denote the spectrum of the Laplacian defined by ∇ acting on sections of L , we show (in contrast to the case of the Schrödinger operator with nontrivial potential) that within the class of translation invariant connections ∇ on L , the spectrum $\text{Spec}(L, \nabla)$ is independent of the choice of ∇ . We refer to this common spectrum as the *spectrum* of L . We show:

- For every even integer $n \geq 4$, there exists an n -dimensional flat rectangular torus M and a pair of topologically distinct, isospectral line bundles over M . The Chern classes of these line bundles have different invariant factors.
- For every even integer $n = 2m$, there exist pairs of nonisometric n -dimensional flat tori M and M' such that for every choice of $r = (r_1, \dots, r_m)$ as above, there is a pair of isospectral line bundles

$L_r \rightarrow M$ and $L'_r \rightarrow M'$. Thus the spectrum does not determine the isometry class of the torus.

We next address the positive results. First note that the spectrum of the Schrödinger operator associated with a fixed connection on a line bundle L may sometimes distinguish between congruent potentials. In particular, given a smooth potential Q on M , let \check{Q} denote the congruent potential given by $\check{Q}(x) = Q(-x)$. For the bundles L that we consider, $Spec(\check{Q}; L, \nabla)$ will typically differ from $Spec(Q; L, \nabla)$ when ∇ is not weakly \mathbb{Z}_2 -invariant. In fact, a consequence of Theorem 1.3 below is that Q and \check{Q} are distinguished by their L -Bloch spectra when all the Chern invariant factors of L equal one. On the other hand, we will see that $Spec(\check{Q}; L, \nabla)$ and $Spec(Q; L, \nabla)$ do agree when ∇ is weakly \mathbb{Z}_2 -invariant.

Our first theorem below generalizes to tori M of arbitrary even dimension $n = 2m$ a result of Guillemin [10] in dimension two. Both theorems below are based on wave invariants constructed in the appendix.

Given a connection ∇ on L and a potential Q , we let $Iso(Q; L, \nabla)$ denote the set of all potentials P such that $Spec(P; L, \nabla) = Spec(Q; L, \nabla)$.

THEOREM 1.2. — *Let M^{2m} be an even-dimensional torus with nondegenerate length spectrum, and let $L \rightarrow M$ be a line bundle with Chern invariant factors $r_1 = \cdots r_m = 1$. Let Q be a smooth potential on M . Then the following statements hold for every translation invariant, weakly \mathbb{Z}_2 -invariant connection ∇ on L :*

(a) *For any $P \in Iso(Q; L, \nabla)$,*

$$\frac{1}{2}(P + \check{P}) = \frac{1}{2}(Q + \check{Q}),$$

i.e. the even part $Q^+ := \frac{1}{2}(Q + \check{Q})$ of Q is uniquely determined by $Spec(Q; L, \nabla)$.

(b) *If Q is assumed to be even, then $Iso(Q; L, \nabla) = \{Q\}$.*

We will show in Section 4 that when $m = 1$, the odd part of Q has strong spectral rigidity properties as well (see Theorem 4.9) and its corollaries.

Theorem 1.1 shows that the nondegeneracy hypothesis cannot be dropped in Theorem 1.2, in contrast to the following result.

THEOREM 1.3. — *Let M^{2m} be an even-dimensional torus and let $L \rightarrow M$ be a line bundle with Chern invariant factors $r_1 = \cdots r_m = 1$. Then every smooth potential Q on M is uniquely determined by its L -Bloch spectrum.*

We emphasize the generality of Theorem 1.3; in particular, we do not assume nondegeneracy of the length spectrum. In the special case in which

the length spectrum is nondegenerate, then Theorem 1.3 can be improved; see Remark 4.5.

Since, as noted above, $\text{Spec}(Q; L, \nabla) = \text{Spec}(\tilde{Q}; L, \nabla)$ when ∇ is translation invariant and weakly \mathbb{Z}_2 -invariant, Theorem 1.3 shows that $\text{Spec}(Q; L, \nabla)$ does *not* suffice to determine the full L -Bloch spectrum of Q even in the nondegenerate case.

In contrast, when the bundle L in Theorem 1.3 is replaced by a trivial bundle, then the conclusion of Theorem 1.3 fails. In fact, it can be easily verified that Q and \tilde{Q} have the same classical Bloch spectrum. Moreover, G. Eskin, J. Ralston, and E. Trubowitz [4] showed that for real analytic potentials Q , the periodic spectrum of Q determines the classical Bloch spectrum of Q . See Remark 4.6 for further discussion of this contrasting behavior as well as possible similarities.

The paper is organized as follows. In section 2 we introduce the line bundles, connections and spectra as well as the notation that are used throughout the paper. The reader interested only in wave invariants and positive results need only read the first two subsections of Section 2. The third subsection gives the background used in the construction of the counterexamples. In section 3 we prove Theorem 1.1 and other negative inverse spectral results for rectangular tori. In section 4 we prove Theorem 1.2, Theorem 1.3 and additional positive results. Recall that a quantity $I(Q, L, \nabla)$ is a spectral invariant of (Q, L, ∇) if $I(Q', L', \nabla') = I(Q, L, \nabla)$ whenever $\text{Spec}(Q'; L', \nabla') = \text{Spec}(Q; L, \nabla)$. The proofs of the positive results use wave invariants, which are spectral invariants arising from the asymptotics of the wave trace. In the appendices, we construct the necessary wave invariants by adapting to our setting constructions developed in [4] (cf also [3] and [10]) to the case of non-trivial line bundles.

Acknowledgment. The authors wish to thank Y. Kurylev for pointing out that an earlier version of Theorem 1.3 could be strengthened. In particular, the earlier version assumed that the length spectrum of the torus was nondegenerate. We also thank Dorothee Schueth for bringing to our attention the distinction between the notions of \mathbb{Z}_2 -invariance and weak \mathbb{Z}_2 -invariance of connections (see Definition 2.12). Finally, we thank Thomas Haller and the referee for suggesting various improvements to the text.

2. Line bundles over tori: construction and classification

The results in this section are standard; we include them for completeness and for setting up notation.

2A. Invariant factors of 2-forms

Let $n \geq 2$, let \mathcal{L} be a lattice of maximal rank in \mathbb{R}^n , and let g_0 be an inner product on \mathbb{R}^n . Then $M := (\mathcal{L} \backslash \mathbb{R}^n, g_0)$ is a flat Riemannian torus of dimension n . We review the classification and construction of complex line bundles over M .

A complex line bundle L over M is classified up to equivalence by its Chern class $c_1(L)$ in $H^2(M; \mathbb{Z})$. Like every real or integer cohomology class on the torus M , the Chern class may be uniquely represented by a translation invariant 2-form Ω on \mathbb{R}^n ; this is the harmonic representative of the class. This form Ω may be viewed as an antisymmetric bilinear map on $\mathbb{R}^n \times \mathbb{R}^n$ that takes integer values on $\mathcal{L} \times \mathcal{L}$. We will consider only bundles with nondegenerate Chern class; i.e., we require that Ω be a nondegenerate bilinear map. In order that such a nondegenerate antisymmetric bilinear form exist, the dimension n of M must be even.

The following result is standard. See, for example, [8], page 304.

LEMMA 2.1. — *Let $n = 2m$ be an even integer. Given a nondegenerate antisymmetric bilinear map Ω on $\mathbb{R}^n \times \mathbb{R}^n$ that takes integer values on $\mathcal{L} \times \mathcal{L}$, there exists a unique m -tuple (r_1, \dots, r_m) of positive integers with the following properties:*

- $r_1 | r_2 | \dots | r_m$;
- *there exists a lattice basis $\mathcal{B} = \{U_1, \dots, U_m, V_1, \dots, V_m\}$ of \mathcal{L} such that $\Omega(U_j, V_j) = r_j$ for $j = 1, \dots, m$ and Ω vanishes on all other pairs of vectors from \mathcal{B} .*

Equivalently, letting $(u, v) := (u_1, \dots, u_m, v_1, \dots, v_m)$ be the coordinates on \mathbb{R}^n associated with the basis \mathcal{B} and viewing Ω as a 2-form on \mathbb{R}^n , we have

$$\Omega = \sum_{j=1}^m r_j du_j \wedge dv_j.$$

Remark 2.2. — The nondegenerate form $\Omega : \mathcal{L} \times \mathcal{L} \rightarrow \mathbb{Z}$ induces an injective map $\mathcal{L} \rightarrow \mathcal{L}^* := \text{Hom}_{\mathbb{Z}}(\mathcal{L}, \mathbb{Z})$, i.e., an injection $\mathbb{Z}^n \rightarrow \mathbb{Z}^n$. The sequence $r_1, r_1, r_2, r_2, \dots, r_m, r_m$ is just the sequence of invariant factors associated with this injection, i.e., the entries of its Smith normal form.

DEFINITION 2.3. — *In view of Remark 2.2, in the notation of Lemma 2.1, we will refer to the integers (r_1, \dots, r_m) as the invariant factors of the cohomology class $[\Omega] \in H^2(M; \mathbb{Z})$.*

Notation 2.4.

- (i) Throughout the rest of this section, we fix $r = (r_1, \dots, r_m)$ (all nonzero) with $r_1 | r_2 | \dots | r_m$, and we fix a lattice basis

$$\mathcal{B} = \{U_1, \dots, U_m, V_1, \dots, V_m\},$$

thus defining coordinates $(u_1, \dots, u_m, v_1, \dots, v_m)$. For $x \in \mathbb{R}^n$, we will express x in these coordinates as

$$x = (u(x), v(x)) = \sum_{j=1}^m u_j(x)U_j + v_j(x)V_j.$$

Define a 2-form Ω by $\Omega = \sum_{j=1}^m r_j du_j \wedge dv_j$.

- (ii) Given x, y in \mathbb{R}^n , define

$$(2.1) \quad e_x(y) = \sum_{j=1}^m r_j u_j(x) v_j(y).$$

In particular, viewing Ω as a bilinear form on \mathbb{R}^n , we have $\Omega(x, y) = e_x(y) - e_y(x)$.

Remark 2.5. — In the notation of 2.4, $[\Omega]$ depends on the choice of \mathcal{B} as well as on (r_1, \dots, r_m) . However, if the torus M is two-dimensional, the cohomology class depends only on the invariant factor $r (= r_1)$ and the orientation of the basis \mathcal{B} . In this case, if L is a line bundle with Chern class $c_1(L) = [\Omega]$, then the Chern number of L is $\pm r$, with the sign depending on the orientation of the basis \mathcal{B} . While we will normally assume that the integers r_i in 2.4 are positive (i.e., that they are the invariant factors as defined in 2.3), the constructions in Subsections 2B and 2C below never use the positivity of the integers r_i . In considering two-dimensional tori, it is convenient to allow r to be an arbitrary nonzero integer so that as r varies (with the basis \mathcal{B} remaining fixed), the corresponding 2-forms Ω range over all nontrivial cohomology classes in $H^2(M; \mathbb{Z})$, which parametrize nontrivial complex line bundles over a two-dimensional torus M .

2B. Line bundles, connections, and spectra.

Given the data in Notation 2.4(i), we now construct a complex Hermitian line bundle L over the torus M and a connection on L . By computing the curvature of this connection, we will see below that the line bundle has Chern class $[\Omega]$.

DEFINITION 2.6. — *Define an action of the group \mathcal{L} on the total space $\mathbb{R}^n \times \mathbb{C}$ of the trivial complex line bundle over \mathbb{R}^n as follows: for $l \in \mathcal{L}$,*

$x \in \mathbb{R}^n$, and $z \in \mathbb{C}$,

$$(2.2) \quad l.(x, z) = (l + x, e^{2\pi i e_l(x)} z),$$

where $e_l(x)$ is given by Notation 2.4(ii). We define L to be the orbit space of the \mathcal{L} -action on $\mathbb{R}^n \times \mathbb{C}$.

Denote elements of L by $[(x, z)]$. The bundle projection $\mathbb{R}^n \times \mathbb{C} \rightarrow \mathbb{R}^n$ is \mathcal{L} -equivariant, so induces $\pi : L \rightarrow M$ by $\pi([(x, z)]) = \bar{x} \in \mathcal{L} \backslash \mathbb{R}^n = M$. Then L is a complex line bundle over M . Moreover, L pulls back to our original trivial line bundle $\mathbb{R}^n \times \mathbb{C}$ over \mathbb{R}^n . Many of our considerations – sections, connections, curvature, etc. – will be simplified by expressing data associated with the bundle L via corresponding “pullback” data associated with the trivial complex line bundle over \mathbb{R}^n .

(Aside: We can endow L with local trivializations in such a way that its structure group is the unitary group $U(1)$. To describe the local trivializations and transition functions explicitly, let $p : \mathbb{R}^n \rightarrow M = \mathcal{L} \backslash \mathbb{R}^n$ be the projection. Given $\bar{x} \in M$, let \bar{U} be an evenly covered neighborhood of \bar{x} in M , so $p^{-1}(\bar{U})$ is a union of disjoint open sheets each carried diffeomorphically by p to \bar{U} . Let U be one of these sheets. For each $\bar{y} \in \bar{U}$, let $y \in U$ be the unique lift of \bar{y} in U . Define $t_U : L|_{\bar{U}} \rightarrow \bar{U} \times \mathbb{C}$ by $t_U([(y, z)]) = (\bar{y}, z)$. If U' is another of the sheets of $p^{-1}(\bar{U})$, then there exists $l \in \mathcal{L}$ such that $U' = l + U = \{l + y : y \in U\}$. The transition function $t_{U'} \circ (t_U)^{-1} : \bar{U} \times \mathbb{C} \rightarrow \bar{U} \times \mathbb{C}$ is given by $(\bar{y}, z) \mapsto (\bar{y}, \sigma_{U'U}(\bar{y})z)$, where $\sigma_{U'U} : \bar{U} \cap \bar{U}' \rightarrow U(1)$ is given by $\bar{y} \mapsto e^{2\pi i e_l(y)}$. We emphasize that $e^{2\pi i e_l(y)}$ is independent of the choice of lift y of \bar{y} in \mathbb{R}^n since $e_l(\mathcal{L}) \subset \mathbb{Z}$.)

The Hermitian structure on the trivial bundle $\mathbb{R}^n \times \mathbb{C}$ given by the standard Hermitian inner product on \mathbb{C} induces a Hermitian structure on L .

Given any bundle B , we denote by $\mathcal{E}(B)$ the space of smooth sections of B . An element of $\mathcal{E}(\mathbb{R}^n \times \mathbb{C})$ is a map $s : \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{C}$ of the form $s(x) = (x, h(x))$ for some smooth complex valued function h on \mathbb{R}^n . The group \mathcal{L} acts on $\mathcal{E}(\mathbb{R}^n \times \mathbb{C})$ on the left via the action

$$(2.3) \quad (l \lrcorner s)(x) = l.(s(x - l)) = (x, e^{2\pi i e_l(x)} h(x - l)).$$

Sections of L pull back to sections $s : \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{C}$ of the trivial bundle that are \mathcal{L} -invariant for this action, i.e., to sections that satisfy

$$(2.4) \quad s(l + x) = l.s(x)$$

for all $l \in \mathcal{L}$ and $x \in \mathbb{R}^n$. Such a section has the form

$$(2.5) \quad s(x) = (x, h(x))$$

with h a complex-valued function on \mathbb{R}^n satisfying the functional equation

$$(2.6) \quad h(l+x) = e^{2\pi i e_l(x)} h(x)$$

for $x \in \mathbb{R}^n$ and $l \in \mathcal{L}$. Given such a function h , the corresponding section \bar{h} of L is given by

$$(2.7) \quad \bar{h}(\bar{x}) = [(x, h(x))].$$

DEFINITION 2.7. — Given a complex-valued 1-form A on \mathbb{R}^n , we obtain a connection ∇ on the trivial bundle $\mathbb{R}^n \times \mathbb{C}$ over \mathbb{R}^n by

$$\nabla = d + A.$$

Thus for a section s given by $s(x) = (x, h(x))$ and a tangent vector $X \in T_x \mathbb{R}^n$,

$$(\nabla_X s)(x) = (x, dh|_x(X) + A|_x(X)h(x)).$$

The connection ∇ descends to a connection on L if and only if it carries \mathcal{L} -invariant sections to \mathcal{L} -invariant sections. More precisely, the condition that ∇ descend to a connection on L is the following: if s is a section of the trivial complex line bundle over \mathbb{R}^n satisfying (2.4), then for every $X \in T_x \mathbb{R}^n$ and $l \in \mathcal{L}$, we have that

$$(2.8) \quad (\nabla_{T_{l*}(X)} s)(l+x) = l. ((\nabla_X s)(x))$$

where $T_l : \mathbb{R}^n \rightarrow \mathbb{R}^n$ denotes translation by l . In this case we will use the same notation ∇ for the connection thus obtained on L .

One checks that the condition on A that guarantees that the connection on the trivial line bundle $\mathbb{R}^n \times \mathbb{C} \rightarrow \mathbb{R}^n$ descends in this way to a connection on the line bundle L over the torus is that, for any $l \in \mathcal{L}$,

$$(2.9) \quad T_l^* A = A - 2\pi i e_l$$

(one uses the fact that e_l can be viewed as either a function or a 1-form, and for a tangent vector $X \in T_x \mathbb{R}^n$, $X(e_l) = e_l|_x(X)$).

A necessary and sufficient condition that the connection be compatible with the Hermitian structure on the bundle is that the 1-form A be purely imaginary; for convenience, we will write $A = 2\pi i \varphi$ for some real-valued 1-form φ on \mathbb{R}^n . We will assume in what follows that *all connections are Hermitian*.

Remark 2.8. — For $l \in \mathcal{L}$, view the linear functional e_l as a harmonic (equivalently, translation-invariant) 1-form on \mathbb{R}^n . Then by Equation (2.9), a connection $\nabla = d + 2\pi i \varphi$ satisfies Equation (2.8), and thus defines a connection on L , if and only if $T_l^* \varphi = \varphi - e_l$ for all $l \in \mathcal{L}$. By identifying

each tangent space $T_x\mathbb{R}^n$ with \mathbb{R}^n and thus viewing the covectors $\varphi|_x$ and $\varphi|_{x+l}$ as linear functionals on \mathbb{R}^n , this equation may be rewritten as

$$(2.10) \quad \varphi|_{x+l} = \varphi|_x - e_l.$$

DEFINITION 2.9. — Let φ^D be the 1-form on \mathbb{R}^n given by $\varphi^D = -\sum_{j=1}^m r_j u_j dv_j$, and set $\nabla^D = d + 2\pi i \varphi^D$. Note that under the identification of the tangent space $T_x\mathbb{R}^n$ with \mathbb{R}^n , we have $\varphi^D|_x = -e_x$ for $x \in \mathbb{R}^n$. Thus φ^D trivially satisfies Equation (2.10) and hence defines a connection on L . This connection will be called the distinguished connection on L .

We will see in Remark 2.19(iii) that this connection is distinguished by its invariance properties.

PROPOSITION 2.10. — The Chern class of the line bundle L constructed in 2.6 is represented by the 2-form $\Omega = \sum_{j=1}^m r_j du_j \wedge dv_j$ on $\mathcal{L}\mathbb{R}^n$.

Proof. — The Chern class is represented by $-\frac{1}{2\pi i} \kappa$, where κ is the curvature form defined by a choice of connection on L . (The cohomology class of this form is independent of the choice of connection.) We use the distinguished connection ∇^D . Pulling back to the trivial bundle over \mathbb{R}^n , the curvature form is given by $d(2\pi i \varphi^D) = -2\pi i \sum_{j=1}^m r_j du_j \wedge dv_j$. This form, being translation-invariant, descends to a 2-form on M which is the curvature form of the distinguished connection ∇^D on L . The proposition now follows. □

Thus for every 2-form Ω of the type given by Notation 2.4(i), we have constructed a line bundle with Chern class $[\Omega]$. Hence every complex line bundle over the torus with Chern class represented by a nondegenerate harmonic form arises from the construction above.

DEFINITION 2.11. — Two Hermitian connections ∇_1 and ∇_2 on L are said to be gauge equivalent if there exists a Hermitian bundle automorphism $F : L \rightarrow L$ that intertwines the two connections: $F \circ \nabla_2 = \nabla_1 \circ F$.

DEFINITION 2.12.

- (i) Denote by $\bar{\rho}$ both the involutive isometry $\bar{\rho} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by $x \mapsto -x$ and the induced isometry of M . This isometry of M is covered by the map ρ of L given by $[(x, z)] \mapsto [(-x, z)]$. We obtain an involution $\rho^* : \mathcal{E}(L) \rightarrow \mathcal{E}(L)$ given by $\rho^*(s) = \rho \circ s \circ \bar{\rho}$; in particular, in the notation of Equation (2.7), we have $\rho^*(\bar{h})(\bar{x}) = [(x, h(-x))]$. We will say that a connection ∇ on L is \mathbb{Z}_2 -invariant if it is invariant under ρ , i.e., $\nabla_{\bar{\rho}_*(X)} \circ \rho^* = \rho^* \circ \nabla_X$ for all $X \in TM$. Expressing the connection (pulled back to the trivial bundle over

\mathbb{R}^n) as in Definition 2.7, we see that it is \mathbb{Z}_2 -invariant if and only if the 1-form φ is invariant under the involution $\bar{\rho}$. We will also say that φ is \mathbb{Z}_2 -invariant in this case. Under the identification of each tangent space $T_x(\mathbb{R}^n)$ with \mathbb{R}^n , the \mathbb{Z}_2 -invariance of φ says that $\varphi|_{-x}(-X) = \varphi|_x(X)$; i.e., $\varphi|_{-x} = -\varphi|_x$.

- (ii) We will say that a connection ∇ is weakly \mathbb{Z}_2 -invariant if the pullback of ∇ by ρ is gauge equivalent to ∇ ; i.e., if there exists a Hermitian bundle automorphism F of L such that $\rho^{*-1} \circ \nabla_{\bar{\rho}_*(X)} \circ \rho^* = F^{-1} \circ \nabla_X \circ F$ for all $X \in TM$.
- (iii) By an abuse of language, we will say that a connection ∇ on a line bundle L over M is a translation-invariant connection if its curvature is a translation-invariant 2-form. (The pullback of such a connection to the bundle $\mathbb{R}^n \times \mathbb{C} \rightarrow \mathbb{R}^n$ is translation-invariant in the sense that each translation of \mathbb{R}^n is covered by a connection preserving map from $\mathbb{R}^n \times \mathbb{C}$ to itself. However, in general the analogous statement does not hold for the bundle $L \rightarrow M$.) We remark that the translation-invariant forms on the torus are precisely the harmonic forms.

Remark 2.13. — Since the curvature form of any connection is cohomologous to that of ∇^D , and since the translation-invariant forms are precisely the harmonic forms, it follows by Hodge theory that the translation-invariant connections all have the same curvature form.

Remark 2.14. — Every Hermitian connection on L is of the form

$$\nabla = \nabla^D + 2\pi i\nu = d + 2\pi i(\varphi^D + \nu)$$

where ν is a real-valued 1-form on M . By the Hodge decomposition, we may write

$$(2.11) \quad \nu = \alpha + dg + d^*\mu$$

where α is a harmonic 1-form on M , $g \in C^\infty(M)$, and μ is a smooth 2-form on M . The harmonic form α may be viewed as a linear functional on \mathbb{R}^n .

Notation 2.15. — Let $\mathcal{L}' \subset (\mathbb{R}^n)'$ denote the dual lattice to \mathcal{L} ; \mathcal{L}' consists of linear functionals on \mathbb{R}^n taking integer values on \mathcal{L} .

PROPOSITION 2.16. — We use the notation of Definitions 2.6, 2.7, 2.9, and 2.12 and Remark 2.14. Let $\nabla = \nabla^D + 2\pi i\nu$ be a Hermitian connection on L with ν given as in Equation (2.11).

- (i) The gauge equivalence class of ∇ is given by $\{\nabla + 2\pi i(\beta + dh) : \beta \in \mathcal{L}', h \in C^\infty(M)\}$. In particular, the gauge equivalence class of

∇ is independent of g in Equation (2.11) and depends on α only through the coset of α in $\mathcal{L}' \setminus (\mathbb{R}^n)'$.

- (ii) ∇ is translation-invariant if and only if $d^*\mu = 0$.
- (iii) ∇ is both translation-invariant and \mathbb{Z}_2 -invariant if and only if $\alpha = 0 = d^*\mu$ and g is \mathbb{Z}_2 -invariant. (Thus $\nabla = \nabla^D + dg$.)
- (iv) ∇ is both translation-invariant and weakly \mathbb{Z}_2 -invariant if and only if $d^*\mu = 0$ and $\alpha(\mathcal{L}) \subset \frac{1}{2}\mathbb{Z}$.

Proof.

- (i) Any Hermitian bundle automorphism $F : L \rightarrow L$ is of the form $[(x, z)] \mapsto [(x, e^{2\pi i f(x)} z)]$, where $f \in C^\infty(\mathbb{R}^n)$ satisfies $f(x + l) \equiv f(x) \pmod{\mathbb{Z}}$ for all $x \in \mathbb{R}^n$ and $l \in \mathcal{L}$. The function f may be written in the form $f = h + \beta$ where h is periodic with respect to \mathcal{L} and $\beta \in \mathcal{L}'$. We then have $F^{-1} \circ \nabla \circ F = \nabla + 2\pi i(\beta + dh)$.
- (ii) The curvature forms of ∇ and ∇^D differ by $2\pi i d\nu = 2\pi i dd^*\mu$. Thus, if $d^*\mu = 0$, then ∇ is translation-invariant, since ∇^D is. Conversely, suppose that ∇ is translation-invariant. Then $dd^*\mu$ is translation-invariant, i.e., harmonic. By Hodge theory, this means that $dd^*\mu = 0$, i.e., $d^*\mu$ is closed. But being both closed and coclosed, $d^*\mu$ is harmonic; in view of the Hodge decomposition (2.11), it must be zero.
- (iii) One easily checks that ∇^D is \mathbb{Z}_2 -invariant. The connection $\nabla = \nabla^D + 2\pi i\nu$ is thus \mathbb{Z}_2 -invariant if and only if $\bar{\rho}^*\nu = \nu$. By (ii) and Equation (2.11), we may write $\nu = \alpha + dg$ and compute $\bar{\rho}^*\nu = -\alpha + d(\bar{\rho}^*g)$. Hence $\alpha = 0$ and g is even.
- (iv) By (i) and the proof of (iii), we see that the translation invariant connection ∇ is weakly \mathbb{Z}_2 -invariant if and only if $\alpha \equiv -\alpha \pmod{\mathcal{L}'}$, i.e., $2\alpha \in \mathcal{L}'$, so $\alpha(\mathcal{L}) \subset \frac{1}{2}\mathbb{Z}$. □

DEFINITION 2.17. — *The torus $\mathcal{L} \setminus \mathbb{R}^n$ has a flat Riemannian metric defined by the inner product g_0 on \mathbb{R}^n . Given a connection $\nabla : \mathcal{E}(L) \rightarrow \mathcal{E}(T^*(M) \otimes L)$ on L , we can thus define an associated Laplacian $\Delta : \mathcal{E}(L) \rightarrow \mathcal{E}(L)$ by*

$$(2.12) \quad \Delta = - \sum_{j=1}^n (\nabla_{X_j} \circ \nabla_{X_j} - \nabla_{\nabla_{X_j}^{LC}(X_j)})$$

where ∇^{LC} is the Levi-Civita connection on the tangent bundle to M and X_1, \dots, X_n is an orthonormal frame field on M . The expression (2.12) is independent of the choice of the orthonormal frame field. Since (M, g) is a flat torus, we may choose the vector fields X_j to be translation-invariant

vector fields (equivalently, coordinate vector fields), in which case the second term in each of the summands vanishes.

As ∇ is assumed to be Hermitian, Δ is essentially self-adjoint with respect to the L^2 -inner product on $\mathcal{E}(L)$ induced by the metric g on M and the Hermitian structure on L .

Notation 2.18. — For $\nabla = d + 2\pi i\varphi$ as in Definition 2.7, we will denote the associated Laplacian by Δ_φ . We will use the shorthand notation Δ^D for Δ_{φ^D} .

Remarks 2.19.

- (i) The connection ∇ , viewed as a connection on the trivial bundle $\mathbb{R}^n \times \mathbb{C}$ over \mathbb{R}^n , gives rise to a Laplace operator $\tilde{\Delta}$ on $\mathcal{E}(\mathbb{R}^n \times \mathbb{C})$ defined by the same expression (2.12), where X_1, \dots, X_n is any orthonormal frame field on \mathbb{R}^n . If each X_i is chosen to be a translation-invariant vector field on \mathbb{R}^n , then Equation (2.8) shows that ∇_{X_i} commutes with the action of \mathcal{L} on $\mathcal{E}(\mathbb{R}^n \times \mathbb{C})$ defined in Equation (2.3). It follows that $\tilde{\Delta}$ commutes with the action and thus induces an operator on $\mathcal{E}(L)$, namely the Laplacian defined in Definition 2.17.
- (ii) If ∇ and ∇' are gauge equivalent connections, say $\nabla' = F^{-1} \circ \nabla \circ F$ for some Hermitian bundle automorphism F , then F intertwines the associated Laplacians, i.e., $\Delta' = F^{-1} \circ \Delta \circ F$. Moreover, for any $Q \in C^\infty(M)$, we have $\Delta' + Q = F^{-1} \circ (\Delta + Q) \circ F$. Thus the Schrödinger operators $\Delta + Q$ and $\Delta' + Q$ are isospectral. For this reason, we will in general only be concerned with the gauge equivalence classes of the connections. Hence in the notation of Equation (2.11) and Proposition 2.16, we will *always* assume $g = 0$.
- (iii) The notion of weak \mathbb{Z}_2 -invariance of a connection is invariant under gauge transformations, but the notion of \mathbb{Z}_2 -invariance is not. As seen in Proposition 2.16, the only translation-invariant \mathbb{Z}_2 -invariant connection (with our convention that $g = 0$) is ∇^D . The one setting in which we will need \mathbb{Z}_2 -invariance rather than just weak \mathbb{Z}_2 -invariance is in Theorem 4.9; in that case we will work specifically with ∇^D .
- (iv) If the connection ∇ is \mathbb{Z}_2 -invariant as in Definition 2.12, then the associated Laplacian Δ commutes with the involution ρ^* of $\mathcal{E}(L)$. Equivalently, under the identification of sections of L with complex-valued functions on \mathbb{R}^n satisfying Equation (2.6), we have $\Delta(h \circ \bar{\rho})(x) = (\Delta h)(-x)$.

- (v) More generally, if ∇ is weakly \mathbb{Z}_2 -invariant, let F be a bundle automorphism satisfying $\rho^{*-1} \circ \nabla_{\tilde{\rho}^*(X)} \circ \rho^* = F^{-1} \circ \nabla_X \circ F$ as in Definition 2.12, and let $T = \rho^* \circ F^{-1}$. Then T commutes with the Laplacian. Moreover, if $Q \in C^\infty(M)$, then $T \circ (\Delta + Q) = (\Delta + \check{Q}) \circ T$, where $\check{Q}(x) = Q(-x)$, so the Schrödinger operators $\Delta + Q$ and $\Delta + \check{Q}$ are isospectral.

PROPOSITION 2.20. — *Let L be a complex line bundle over the flat torus (M, g) . In the notation of Definition 2.17, we have*

$$\Delta_\varphi = -\operatorname{div} \circ \operatorname{grad} - 4\pi i \varphi \circ \operatorname{grad} + 4\pi^2 |\varphi|^2 - 2\pi i \operatorname{div}(\varphi^\#)$$

where the “index-raising” map $\# : T^*M \rightarrow TM$ is the inverse of the duality isomorphism $\flat : TM \rightarrow T^*M$ arising from the Riemannian metric g_0 on \mathbb{R}^n .

Proof. — For h a smooth, complex-valued function on \mathbb{R}^n satisfying Equation (2.6) and for X a tangent vector to \mathbb{R}^n , we have $\nabla_X h = dh(X) + 2\pi i \varphi(X)h = (X + 2\pi i \varphi(X))h$. Thus

(2.13)

$$\nabla_X \circ \nabla_X(h) = X^2 h - 4\pi^2 \varphi(X) \varphi(X)h + 2\pi i X(\varphi(X))h + 4\pi i \varphi(X)Xh.$$

Let $\{X_1, \dots, X_n\}$ be an orthonormal, translation-invariant frame field on M . We have $\sum_{j=1}^n X_j^2(h) = \operatorname{div} \operatorname{grad} h$, $\sum_{j=1}^n \varphi(X_j) \varphi(X_j) = |\varphi|^2$, $\sum_{j=1}^n X_j(\varphi(X_j)) = \operatorname{div}(\varphi^\#)$, and $\sum_{j=1}^n \varphi(X_j) X_j(h) = \varphi(\operatorname{grad} h)$. Thus the proposition follows from Equation (2.13). □

Notation 2.21. — Given $[\alpha] \in \mathcal{L}' \setminus (\mathbb{R}^n)'$ and $Q \in C^\infty(M)$ (viewed as a periodic function on \mathbb{R}^n), let $\operatorname{Spec}_\alpha(Q; L)$ denote the spectrum of the Schrödinger operator $\Delta_{\varphi^D + \alpha} + Q$ acting on smooth sections of L . Since $\Delta_{\varphi^D + \alpha} + Q$ is essentially self-adjoint with respect to the L^2 -inner product on $\mathcal{E}(L)$ and M is closed, $\operatorname{Spec}_\alpha(Q; L)$ is real and discrete. The correspondence that associates to each $[\alpha] \in \mathcal{L}' \setminus (\mathbb{R}^n)'$ the spectrum $\operatorname{Spec}_\alpha(Q; L)$ will be called the L -Bloch spectrum of Q . Usually, the line bundle L will be understood and we will use “Bloch spectrum” to mean L -Bloch spectrum.

PROPOSITION 2.22. — *In the notation of 2.21, $\operatorname{Spec}_\alpha(Q; L)$ coincides with the spectrum of the operator $\Delta^D + Q$ acting on the space of all smooth functions on \mathbb{R}^n satisfying*

$$(2.14) \quad h(x + l) = e^{2\pi i \alpha(l)} e^{2\pi i e_l(x)} h(x)$$

for all $x \in M, l \in \mathcal{L}$, where e_l is given as in Notation 2.4.

Proof. — Let L_α be the bundle over M given by the quotient of $\mathbb{R}^n \times \mathbb{C}$ by the equivalence relation $(x, z) \equiv (x + l, e^{2\pi i \alpha(l)} e^{2\pi i e_l(x)} z)$ for $l \in \mathcal{L}$.

Sections of L_α may be viewed as functions on \mathbb{R}^n satisfying Equation (2.14). The automorphism of $\mathbb{R}^n \times \mathbb{C}$ given by $(x, z) \mapsto (x, e^{2\pi i\alpha(x)}z)$ intertwines $\nabla^D + 2\pi i\alpha$ with ∇^D and induces an isomorphism from L to L_α . Thus it intertwines $\Delta_{\varphi^D+\alpha}$ with the operator Δ^D acting on the space of all smooth functions on \mathbb{R}^n satisfying Equation (2.14). □

Remark 2.23. — In the case of the trivial line bundle $M \times \mathbb{C}$, the distinguished connection is given by $\nabla = d$, (i.e., $\varphi^D = 0$), the associated Laplacian Δ is the Euclidean Laplacian Δ defined by the Euclidean metric, and $e_l = 0$ for every $l \in \mathcal{L}$. Thus $Spec_\alpha(Q)$ is the spectrum of $\Delta + Q$ acting on the space of smooth functions on \mathbb{R}^n satisfying $f(x + l) = e^{2\pi i\alpha(l)}f(x)$ for all $x \in \mathbb{R}^n$ and $l \in \mathcal{L}$. Hence the notion of Bloch spectrum in Definition 2.21 in the case of a trivial bundle agrees with the classical Bloch (or Floquet) spectrum in the literature.

2C. Nilmanifold structure on the principal circle bundle.

We continue to use the conventions of Notation 2.4.

Let $\pi : L \rightarrow M$ be a smooth Hermitian complex line bundle over M . The unit sphere bundle $P \rightarrow M$ of $L \rightarrow M$ whose fiber P_x , above $x \in M$ is given by $P_x = \{\xi \in L_x \mid \langle \xi, \xi \rangle = 1\}$, is a principal S^1 -bundle. Conversely, given the principal circle bundle P , one recovers the original line bundle L as the associated vector bundle given as follows: The circle $S^1 = U(1)$ acts on \mathbb{C} by unitary transformations ($t \in S^1$ acts as multiplication by $e^{2\pi it}$). We thus obtain a left action of S^1 on $P \times \mathbb{C}$ given by $g \cdot (p, z) = (p \cdot g^{-1}, g \cdot z)$ for $g \in S^1$, $(p, z) \in P \times \mathbb{C}$. We let L be the orbit space of this action,

$$(2.15) \quad L = P \times_{S^1} \mathbb{C} = S^1 \backslash (P \times \mathbb{C}).$$

Then L is a complex line bundle over $\mathcal{L} \backslash \mathbb{R}^n$.

The space $\mathcal{E}(L)$ of smooth sections of L may be identified with the space of S^1 -equivariant smooth maps from P to the representation space \mathbb{C} :

$$(2.16) \quad C^\infty(P, \mathbb{C})^{S^1} = \{f \in C^\infty(P, \mathbb{C}) \mid f(pg^{-1}) = gf(p) \ \forall p \in P, \forall g \in U(1)\}.$$

(Given $f \in C^\infty(P, \mathbb{C})^{S^1}$, define its associated section $\sigma \in \mathcal{E}(L)$ by $\sigma(x) = [p, f(p)]$ where p is any element of P for which $\pi(p) = x$ and where $[p, z]$ denotes the equivalence class of (p, z) in $L' = S^1 \backslash (P \times \mathbb{C})$.)

R. S. Palais and T. E. Stewart [14] proved (by explicit construction) that every principal torus bundle over a torus has the structure of a two-step nilmanifold, i.e., a quotient of a two-step nilpotent Lie group by a cocompact discrete subgroup. In what follows, we will first describe the nilmanifold structure in the case of the circle bundles discussed above. We

will then construct an associated bundle L' and show that it coincides with the line bundle L constructed in Subsection 2B.

2C.1. *Principal bundles.*

Define a two-step nilpotent Lie group structure on \mathbb{R}^{n+1} by setting (2.17)

$$(x, t)(y, s) = \left(x + y, s + t + \sum_{j=1}^m r_j u_j(x) v_j(y) \right) = (x + y, s + t + e_x(y))$$

for all $x, y \in \mathbb{R}^n$ and $t, s \in \mathbb{R}$. (See Notation 2.4.) Let N denote \mathbb{R}^{n+1} with this Lie group structure. The derived group (i.e., the subgroup generated by all $\eta_1 \eta_2 \eta_1^{-1} \eta_2^{-1}$ with $\eta_1, \eta_2 \in N$) is given by $\{(0, t) : t \in \mathbb{R}\}$ and is central in N . We will denote the derived group by $(0, \mathbb{R})$. It is easily checked that the center consists of elements (z, u) such that z is in the kernel of the bilinear form Ω . In this paper, we are interested only in the case in which r_1, \dots, r_m are all nonzero, i.e., Ω is nondegenerate, in which case the derived group coincides with the center. Thus we will refer to the derived group as the center in what follows.

The lattice \mathcal{L} in \mathbb{R}^n extends to a cocompact discrete subgroup Γ of N given by

$$\Gamma = \{(l, k) : l \in \mathcal{L}, k \in \mathbb{Z}\}.$$

Letting $\bar{N} = (0, \mathbb{Z}) \backslash N$, where $(0, \mathbb{Z})$ is the integer lattice in $(0, \mathbb{R})$, then \bar{N} is a nilpotent Lie group with center isomorphic to S^1 . Elements in \bar{N} will be denoted by (x, \bar{z}) . Set $\bar{\Gamma} = (0, \mathbb{Z}) \backslash \Gamma \cong \mathcal{L}$ and

$$P' = \Gamma \backslash N = \bar{\Gamma} \backslash \bar{N}.$$

Denote elements of P' by $\overline{(x, t)}$, with $(x, t) \in N$. The surjective group homomorphism $N \rightarrow \mathbb{R}^n$ given by $(x, t) \mapsto x$ induces a submersion

$$\pi : P' \rightarrow \mathcal{L} \backslash \mathbb{R}^n = M,$$

with fiber $S^1 = (0, \mathbb{Z}) \backslash (0, \mathbb{R})$:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{Z} & \longrightarrow & \Gamma & \longrightarrow & \mathcal{L} \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \mathbb{R} & \longrightarrow & N & \longrightarrow & \mathbb{R}^n \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \mathbb{Z} \backslash \mathbb{R} & \longrightarrow & \Gamma \backslash N & \longrightarrow & \mathcal{L} \backslash \mathbb{R}^n \\ & & \parallel & & \parallel & & \parallel \\ & & S^1 & & P' & & M \end{array}$$

The action by translation of the center $(0, \mathbb{R})$ of N on N gives rise to a right (and left) action of S^1 on P' given by

$$\overline{(x, t)} \cdot e^{2\pi is} = \overline{(x, t + s)}.$$

This action gives P' the structure of a principal circle bundle over M .

Let L' be the line bundle over M associated with P' as in Equation (2.15). Elements of L' will be written $[(x, t), z]$. We have $[(x, t + s), z] = [(x, t), e^{2\pi is} z]$.


Notation 2.24. — By Equation (2.16), sections of L' may be viewed as smooth functions $f : P' \rightarrow \mathbb{C}$ satisfying $f(\overline{(x, t - s)}) = e^{2\pi is} f(\overline{(x, t)})$. Such a function pulls back to a \mathbb{C} -valued function, which we again denote by f , on \overline{N} satisfying the S^1 -equivariance condition

$$f(x, \overline{t - s}) = e^{2\pi is} f(x, \overline{t});$$

the latter in turn pulls back to a Γ -invariant function (still denoted f) on N satisfying

$$(2.18) \quad f(x, t - s) = e^{2\pi is} f(x, t)$$

for all $s, t \in \mathbb{R}, x \in \mathbb{R}^n$:



Note that the Γ -invariance of $f : N \rightarrow \mathbb{C}$ together with Equation (2.18) says that

$$(2.19) \quad f(x, t) = f((l, k)(x, t)) = f(l+x, k+t+e_l(x)) = e^{-2\pi i e_l(x)} f(l+x, t).$$

In order to compare the bundle L' with the bundle L constructed in Subsection 2B, we consider the pullbacks of P' and L' to bundles over \mathbb{R}^n . The former is simply \overline{N} , with S^1 action given by $(x, \overline{t}) \cdot e^{2\pi is} = (x, \overline{t + s})$. The latter is the bundle $\tilde{L}' = \overline{N} \times_{S^1} \mathbb{C}$ over \mathbb{R}^n given by equivalence classes $[(x, \overline{t}), z]$ with $[(x, \overline{t}), z] = [(x, \overline{t - s}), e^{2\pi is} z]$. In particular, $[(x, \overline{t}), z] = [(x, \overline{0}), e^{2\pi it} z]$.

PROPOSITION 2.25. — *In the notation above, define $T : \tilde{L}' \rightarrow \mathbb{R}^n \times \mathbb{C}$ by $T([(x, \overline{t}), z]) = (x, e^{2\pi it} z)$. Then T induces an equivalence, also denoted T , between the line bundle L' over M constructed above and the line bundle*

L constructed in Definition 2.6. Under this equivalence, a section f of L' , viewed as a function on N satisfying Equation (2.18), corresponds to the section h of L given by $h(x) = f(x, 0)$, where h is as in Equation (2.5).

Proof. — The group $\bar{\Gamma} \cong \mathcal{L}$ acts on the line bundle \tilde{L}' by

$$(l, \bar{0}) \cdot [(x, \bar{t}), z] = [(l, \bar{0})(x, \bar{t}), z] = [(l+x, \bar{t} + \overline{e_l(x)}), z] = [(l+x, \bar{t}), e^{2\pi i e_l(x)} z].$$

The bundle L' is the quotient of \tilde{L}' by this group action. The map T carries orbits of the action of $\bar{\Gamma}$ on \tilde{L}' to equivalence classes of $\mathbb{R}^n \times \mathbb{C}$ with respect to the equivalence relation in Equation (2.2). The proposition follows. \square

2C.2. *Connections on the principal bundle.*

The coordinate vector field $\frac{\partial}{\partial t}$ on \mathbb{R}^{n+1} with respect to the global coordinates (x, t) is invariant under left translations by elements of Γ and thus induces a vector field, which we still denote by $\frac{\partial}{\partial t}$, on P' . This vector field is the infinitesimal generator of the S^1 action on P' . A connection on P' is specified by a real-valued 1-form ω on P' which is invariant under the S^1 action and satisfies $\omega(\frac{\partial}{\partial t}) = 1$. (Here we are using the canonical identification of the Lie algebra of the Lie group S^1 with \mathbb{R} .) The kernel \mathcal{H} of ω is referred to as the horizontal distribution defined by the connection. We have

$$(2.20) \quad \mathcal{H} = \left\{ X - \omega(X) \frac{\partial}{\partial t} : X \in TP' \right\}$$

and π_{*p} carries \mathcal{H}_p isomorphically to $T_{\pi(p)}(M)$ for all $p \in P'$. The horizontal distribution \mathcal{H} is invariant under the action of S^1 . Conversely, any S^1 -invariant n -plane distribution on P' , complementary at each point to $\frac{\partial}{\partial t}$, defines a connection on P' .

A connection specified by a 1-form ω on P' defines a connection ∇ on the associated line bundle L' as follows. Given $x \in M$ and $X \in T_x(M)$, choose $p \in \pi^{-1}(x)$ and let \tilde{X} denote the unique horizontal lift of X in $T_p(P')$; i.e., $\tilde{X} \in \mathcal{H}_p$ and $\pi_*(\tilde{X}) = X$. Given a section of L' , viewed as an element of $C^\infty(P', \mathbb{C})^{S^1}$ (i.e., as an S^1 -equivariant smooth map $P' \rightarrow \mathbb{C}$), we then define $\nabla_X(f) = \tilde{X}(f)$. (The fact that \mathcal{H} is S^1 -invariant guarantees that this definition is independent of the choice of p .)

Example 2.26. — Let e_1, \dots, e_n denote the standard basis vectors of \mathbb{R}^n . The left-invariant vector field \tilde{X}_j on N whose value at the identity is given by $\frac{\partial}{\partial x_j}$ is given at (x, t) by

$$(2.21) \quad \tilde{X}_j = \left. \frac{d}{ds} \right|_{s=0} (x, t)(se_j, 0) = \left. \frac{d}{ds} \right|_{s=0} (x + se_j, t + se_x(e_j)) = \frac{\partial}{\partial x_j} + e_x(e_j) \frac{\partial}{\partial t}.$$

A left-invariant vector field \tilde{X} on N induces a vector field, also denoted \tilde{X} , on P' . We will abuse terminology and refer to \tilde{X} as a left-invariant vector field on P' . Let \mathcal{H} be the distribution on P' spanned by the left-invariant vector fields $\tilde{X}_1, \dots, \tilde{X}_n$ given in Equation (2.21). Viewing sections of L' as Γ -invariant functions f on N satisfying Equation (2.18), then $\frac{\partial f}{\partial t} = -2\pi i f$. Thus comparing Equation (2.21) and Definition 2.9, we see that the connection on L' defined by \mathcal{H} corresponds to the distinguished connection on L under the equivalence between L and L' given in Proposition 2.25.

More generally, for $\alpha = \sum_{j=1}^n a_j dx_j$ a harmonic 1-form on M , the translation-invariant connection $\nabla = d + 2\pi i(\varphi^D + \alpha)$ on L corresponds to the connection on L' for which the horizontal distribution (on P') is spanned by the left-invariant vector fields $\tilde{X}_j - a_j \frac{\partial}{\partial t}$, $j = 1, \dots, n$. (Note that $\frac{\partial}{\partial t}$ is a left-invariant vector field.) Thus by Proposition 2.16, a connection on L' corresponds to a translation-invariant connection on L if and only if its associated horizontal distribution \mathcal{H} is spanned by left-invariant vector fields, or equivalently, if the one-form ω on P' lifts to a left-invariant one-form on N .

Suppose that ∇ is a translation-invariant connection so that \mathcal{H} is spanned by left-invariant vector fields. The Riemannian inner product on M gives rise to an inner product on \mathcal{H} . Let $\tilde{X}_1, \dots, \tilde{X}_n$ denote an orthonormal basis of \mathcal{H} . Then the Laplacian on L' associated to the connection ∇ is given by

$$(2.22) \quad \Delta(f) = - \sum_{j=1}^n \tilde{X}_j^2(f)$$

where, as usual, we view f as a Γ -invariant function on N satisfying equation (2.18).

3. Negative inverse spectral results on rectangular tori

Notation 3.1.

- (i) Let $n = 2m$ be an even integer and let $(u, v) = (u_1, \dots, u_m, v_1, \dots, v_m)$ be the standard coordinates on \mathbb{R}^n . Given m -tuples $a = (a_1, \dots, a_m)$ and $b = (b_1, \dots, b_m)$ of positive real numbers, give \mathbb{R}^n the inner product for which the standard ordered basis elements are orthogonal and have lengths $a_1, \dots, a_m, b_1, \dots, b_m$, respectively. Let $M = M_{a,b}$ be the rectangular torus $\mathbb{Z}^n \backslash \mathbb{R}^n$ with the metric defined by this inner product.

(ii) Given an m -tuple $r = (r_1, \dots, r_m)$ of positive integers such that $r_1|r_2|\dots|r_m$, let $\Omega_r = \sum_{j=1}^m r_j du_j \wedge dv_j$. The construction in Section 2 gives us a line bundle over M with Chern class represented by Ω_r ; we denote this bundle by L_r . We will work in the setting of Subsection 2C, viewing the principal circle bundle given by the unit vectors in L_r as a nilmanifold. We will write elements of the nilpotent group N in the form (u, v, t) , with $u, v \in \mathbb{R}^m, t \in \mathbb{R}$. Comparing with the notation in Subsection 2C, an element $(x, t) \in N$ in the notation of Subsection 2C will be written as $(u(x), v(x), t)$, where $u(x)$ and $v(x)$ are defined as in Notation 2.4(i). The group multiplication in N is given by

$$(3.1) \quad (u, v, t)(u', v', t') = \left(u + u', v + v', t + t' + \sum_{j=1}^m r_j u_j v'_j \right).$$

The discrete group Γ is given by

$$\Gamma = \{(u, v, t) \in N : u, v \in \mathbb{Z}^m, t \in \mathbb{Z}\}.$$

Thus under the identifications discussed in Subsection 2C, sections of L_r will be viewed as Γ -invariant functions on N satisfying $f(u, v, t + s) = e^{-2\pi i s} f(u, v, t)$ (see Equation (2.18)).

(iii) Each harmonic 1-form α on M (equivalently, linear functional on \mathbb{R}^{2m}) will be identified with a pair of elements $\mu, \nu \in \mathbb{R}^m$ by writing $\alpha(u, v) = \sum_{j=1}^m (\mu_j u_j + \nu_j v_j)$.

Remark 3.2. — Recall Remark 2.5. Since in Notation 3.1, we fixed the basis used to construct Ω_r and the associated line bundle L_r , the bundles L_r do not represent all possible line bundles over M . However, in the case of two-dimensional tori (so $m = 1$ and we drop the subscripts on u, v), the bundles L_r , as r varies over the positive integers, range over all equivalence classes of line bundles with positive Chern number. As in Subsections 2B and 2C, all the results below continue to make sense if we allow r to be negative, so as to obtain all nontrivial line bundles over M in the two-dimensional case.

Notation 3.3.

(i) Given m -tuples a, b , and r as in Notation 3.1, define an operator $D_{r,a,b}$ on $C^\infty(\mathbb{R}^m)$ by

$$D_{r,a,b} = \sum_{j=1}^m \left(\left(2\pi \frac{r_j}{b_j} s_j \right)^2 - \frac{1}{a_j^2} \frac{\partial^2}{\partial s_j^2} \right)$$

where s_1, \dots, s_m are the standard coordinates on \mathbb{R}^m .

Under the change of coordinates on \mathbb{R}^m given by setting $w_j = \sqrt{\frac{a_j}{b_j}} s_j$, the operator $D_{r,a,b}$ transforms to the operator

$$\tilde{D}_{r,a,b} = \sum_{j=1}^m \frac{1}{a_j b_j} \left((2\pi r_j w_j)^2 - \frac{\partial^2}{\partial w_j^2} \right).$$

In particular, the operator depends only on the products $a_j b_j$, not on a_j and b_j individually. Note that $\tilde{D}_{r,a,b}$ represents m uncoupled harmonic oscillators.

- (ii) Suppose $q \in C^\infty(\mathbb{R}^m)$ is periodic with respect to \mathbb{Z}^m . For $c = (c_1, \dots, c_m) \in \mathbb{Z}^m$ and for $\mu = (\mu_1, \dots, \mu_m) \in \mathbb{R}^n$, define $q_{r,c,\mu} \in C^\infty(\mathbb{R}^m)$ by

$$q_{r,c,\mu}(s) = q \left(s_1 + \mu_1 - \frac{c_1}{r_1}, \dots, s_m + \mu_m - \frac{c_m}{r_m} \right).$$

Since q is periodic, $q_{r,c,\mu}$ depends only on the coset of c in $(r_1\mathbb{Z} \times \dots \times r_m\mathbb{Z}) \backslash \mathbb{Z}^m$. As q is real-valued, $D_{r,a,b} + q_{r,c,\mu}$ is essentially self-adjoint with respect to the standard L^2 -inner product. Its spectrum is real and discrete. Let $S(q, r, a, b, \mu)$ denote the join of the spectra of the operators $D_{r,a,b} + q_{r,c,\mu}$ as c varies over the right coset space $(r_1\mathbb{Z} \times \dots \times r_m\mathbb{Z}) \backslash \mathbb{Z}^m$.

THEOREM 3.4. — *Let a, b, r, α, μ and ν be as in Notation 3.1, and let $M = M_{a,b}$ and L_r be the corresponding rectangular torus and line bundle as in Notation 3.1. Let $q \in C^\infty(\mathbb{R}^m)$ be periodic with respect to \mathbb{Z}^m .*

- (1) *Define $Q \in C^\infty(M)$ by $Q(u, v) = q(u)$. Then the spectrum $Spec_\alpha(Q; L_r)$ coincides with $S(q, r, a, b, \nu)$ as defined in Notation 3.3.*
- (2) *Define $Q \in C^\infty(M)$ by $Q(u, v) = q(v)$. Then $Spec_\alpha(Q; L_r)$ coincides with $S(q, r, b, a, \mu)$.*

Before proving the theorem, we give several applications.

COROLLARY 3.5. — *Let M be a rectangular 2-dimensional torus for which the side lengths a and b are distinct positive integers. Let L be any line bundle over M with nondegenerate Chern class. Then for each translation invariant connection ∇ on L , there exist pairs of noncongruent potentials Q_1 and Q_2 on M such that $Spec(Q_1; L, \nabla) = Spec(Q_2; L, \nabla)$. The potentials may be chosen to be real analytic.*

Proof of Corollary 3.5. — By Remark 3.2, it suffices to consider the line bundles L_r given in Notation 3.1, where we allow r to be an arbitrary

nonzero integer. Fix r and α . Let $q \in C^\infty(\mathbb{R})$ be periodic with period $\frac{1}{ar}$, and let $q_1(s) = q(s - \nu)$ and $q_2(s) = \tilde{q}(s - \mu)$ where $\tilde{q}(s) = q(\frac{b}{a}s)$. Define $Q_1(u, v) = q_1(u)$ and $Q_2(u, v) = q_2(v)$. We show that $Spec_\alpha(Q_1; L_r) = Spec_\alpha(Q_2; L_r)$.

In the notation of 3.3(ii), we have that $(q_1)_{r,c,\nu} = q$ and $(q_2)_{r,c,\mu} = \tilde{q}$ for all $c \in \mathbb{Z}$ due to the periodicity condition on q . By Theorem 3.4, the spectrum of $\Delta + Q_1$ is given by $S(q, r, a, b, \nu)$, which is equal to the join of r copies of the spectrum of $D_{r,a,b} + q$. By the change of variables in Notation 3.3(ii), the spectrum of $\Delta + Q_1$ is thus equal to the join of r copies of the spectrum of $\tilde{D}_{r,a,b} + g$ where $g(s) = q\left(\sqrt{\frac{b}{a}}s\right)$. Again by Theorem 3.4, the spectrum of $\Delta + Q_2$ is equal to $S(\tilde{q}, r, b, a, \mu)$, which is in turn equal to the join of r copies of the spectrum of $D_{r,b,a} + \tilde{q}$. By the change of variables $w = \sqrt{\frac{b}{a}}s$, the operator $D_{r,b,a} + \tilde{q}$ is carried to $\tilde{D}_{r,a,b} + g$, since $\tilde{q}\left(\sqrt{\frac{a}{b}}w\right) = q\left(\frac{b}{a}\sqrt{\frac{a}{b}}w\right) = q\left(\sqrt{\frac{b}{a}}w\right) = g(w)$. This proves the isospectrality statement. Since $a \neq b$, it is clear that the potentials Q_1 and Q_2 are not congruent. The corollary follows. \square

Remark 3.6. — For $2m$ -dimensional rectangular tori given as in Notation 3.1 with $\frac{a_j}{b_j}$ rational and not equal to one for all j , a similar proof shows that for each $r = (r_1, \dots, r_m)$ and each translation invariant connection ∇ on L_r , there exist pairs of (analytic) noncongruent potentials on M such that $Spec(Q_1; L, \nabla) = Spec(Q_2; L, \nabla)$.

COROLLARY 3.7. — *In the setting of Theorem 3.4, suppose that Q is the zero potential. Then the spectrum $Spec_\alpha(Q; L)$ is independent of α . We will refer to this common spectrum as the spectrum of the line bundle L .*

The corollary is immediate from Theorem 3.4 and the fact that $S(q, r, a, b, \mu)$ depends on μ only through q .

COROLLARY 3.8. — *For every even integer $n \geq 4$, there exist an n -dimensional rectangular flat torus M and a pair of isospectral (in the sense of Corollary 3.7), topologically distinct, line bundles over M . The Chern classes of these bundles have different invariant factors.*

Proof of Corollary 3.8. — For an example in dimension $n = 2m$, choose a so that $a_1 = a_2 = \dots = a_m$ and let $b = (1, 2, 1, \dots, 1)$. Set $r = (1, 4, 1, \dots, 1)$ and $r' = (2, 2, 1, \dots, 1)$. Since $\frac{r_1}{b_1} = \frac{r'_1}{b_2}$ and $\frac{r_2}{b_2} = \frac{r'_1}{b_1}$, we see from Theorem 3.4 (with $Q = 0$) and from the definition of $D_{r,a,b}$ in Notation 3.3 that the line bundles L_r and $L_{r'}$ over M are isospectral. \square

COROLLARY 3.9. — *In every even dimension $2m$, $m \geq 1$, there exist pairs of nonisometric rectangular tori M and M' such that for every choice of $r = (r_1, \dots, r_m)$ as in Notation 3.1, the corresponding line bundles L_r and L'_r over M and M' , respectively, have the same spectrum in the sense of Corollary 3.7. In the case of two-dimensional rectangular tori M , the spectrum of a nontrivial line bundle over M depends only on the area of the torus and the Chern number of the bundle.*

Proof of Corollary 3.9. — We use the notation of 3.1. Choose m -tuples a, b, a' and b' of positive real numbers in such a way that $a_j b_j = a'_j b'_j$ for every $j = 1, \dots, m$. Let $M = M_{a,b}$ and $M' = M_{a',b'}$. Choose r arbitrarily subject only to the condition $r_1 | \dots | r_m$ and let L_r and L'_r be the corresponding line bundles over M and M' respectively. From the expression $\tilde{D}_{r,a,b}$ in Notation 3.3(ii) and by Theorem 3.4, with $Q = 0$, we see that L_r and L'_r have the same spectrum. In particular, in the two-dimensional case, (writing $a = a_1$ and $b = b_1$), the spectrum of the line bundle L_r over M depends only on ab and r . Since ab is the area of the torus and r the Chern number of the line bundle, the final statement of the corollary follows. (Here we allow r to take on negative as well as positive values; see Remark 3.2.) □

We now turn to the proof of Theorem 3.4.

Proof. —

(1) We use the notation of 3.1. As in Notation 2.24, smooth sections of the line bundle L_r may be identified with elements of the space of Γ -periodic complex-valued smooth functions on N satisfying

$$(3.2) \quad f(u, v, t) = e^{-2\pi i t} f(u, v, 0).$$

Let U_j , respectively V_j , denote the left-invariant vector field on N whose value at the identity is $\frac{\partial}{\partial u_j}$, respectively, $\frac{\partial}{\partial v_j}$, and let $Z = \frac{\partial}{\partial t}$. (Note that Z is also left-invariant.) By Example 2.26 and Equation (2.22), the Laplacian $\Delta_{\varphi^D + \alpha}$ with respect to the connection $\nabla^D + 2\pi i \alpha$ on L_r is given by

$$\Delta_{\varphi^D + \alpha} = - \sum_{j=1}^m \left(\frac{1}{a_j^2} (U_j - \mu_j Z)^2 + \frac{1}{b_j^2} (V_j - \nu_j Z)^2 \right).$$

The integral curve of U_j (respectively, V_j) through the identity element of N is given by $\exp(sU_j) = (se_j, 0, 0)$ (respectively, $\exp(sV_j) = (0, se_j, 0)$), where e_j is the j th standard basis vector of \mathbb{R}^m .

Left-invariant vector fields are defined as derivatives of right translations (i.e., $Xf(p) = \frac{d}{dt} \Big|_{t=0} f(p \exp(tX))$ for $p \in N$), so we can understand the action of the Laplacian on the smooth sections of L_r by analyzing the

right action of N on the larger space \mathcal{K} of square integrable sections. As will be explained below, it is well known that this right action is unitarily equivalent to the direct sum of $r_1 \cdots r_m$ copies of the action τ of N on $L^2(\mathbb{R}^m)$ given by

$$(3.3) \quad (\tau(u, v, t)g)(w) = e^{-2\pi i(t + \sum_{j=1}^m r_j w_j v_j)} g(w + u).$$

This action of N on $L^2(\mathbb{R}^m)$ gives rise to an infinitesimal action of the Lie algebra \mathfrak{n} of N ; this action satisfies $\tau_*(U_j)g = \frac{\partial g}{\partial w_j}$, $\tau_*(V_j)g = -2\pi i r_j w_j g$, and $\tau_*(Z)g = -2\pi i g$. Thus, under this equivalence, the Laplacian associated with the distinguished connection ∇^D is carried to the operator $D_{r,a,b}$ defined in Notation 3.3. (Later in the proof, we will also construct an intertwining map between $D_{r,a,b}$ and $\Delta_{\varphi^D + \alpha}$ for each α .)

We will make the equivalence between the action of N on \mathcal{K} and the action τ of N on $L^2(\mathbb{R}^m)$ precise in order to see its effect on the Schrödinger operator $\Delta + Q$. The presentation below of this equivalence was shown to the first author by E. N. Wilson during preparation of the article [7].

In order to avoid cumbersome notation, we will carry out the proof in the case $m = 1$. For the general case, the reader may simply replace rkv by $\sum_{j=1}^m r_j k_j v_j$, replace $c + rk$ by $(c_1 + r_1 k_1, \dots, c_m + r_m k_m)$, replace \mathbb{Z} by \mathbb{Z}^m , etc.

Let $f \in \mathcal{K}$. By Equation (3.2) and Γ -periodicity, we have

$$(3.4) \quad f(u, v, t) = f(u + k, v + l, t + rk v) = e^{-2\pi i rk v} f(u + k, v + l, t)$$

for all $k, l \in \mathbb{Z}$. In particular, f is periodic with respect to \mathbb{Z} in v and we may perform a Fourier decomposition in the v variable:

$$(3.5) \quad f(u, v, t) = \sum_{c \in \mathbb{Z}} \hat{f}_c(u) e^{-2\pi i c v} e^{-2\pi i t}.$$

(We are writing \hat{f}_c rather than \hat{f}_{-c} here for notational convenience.)

By Equation (3.4), we see that

$$\hat{f}_{c+rk}(u) = \hat{f}_c(u + k)$$

for all $k \in \mathbb{Z}$. Thus choosing a representative c of each coset \bar{c} of $r\mathbb{Z}$ in \mathbb{Z} , we may write

$$f(u, v, t) = \sum_{\bar{c} \in r\mathbb{Z} \backslash \mathbb{Z}} f_{\bar{c}}(u, v, t)$$

where

$$f_{\bar{c}}(u, v, t) = e^{-2\pi i t} \sum_{k \in \mathbb{Z}} \hat{f}_c(u + k) e^{-2\pi i (c+rk)v}.$$

Each of the functions $f_{\bar{c}}$ on N is Γ -invariant. Thus we obtain a decomposition

$$(3.6) \quad \mathcal{K} = \bigoplus_{\bar{c} \in r\mathbb{Z} \backslash \mathbb{Z}} \mathcal{K}_{\bar{c}}$$

where $\mathcal{K}_{\bar{c}} = \{f_{\bar{c}} : f \in \mathcal{K}\}$. (In the case of a $2m$ -dimensional torus, $r\mathbb{Z} \backslash \mathbb{Z}$ is replaced by $(r_1\mathbb{Z} \times \dots \times r_m\mathbb{Z}) \backslash \mathbb{Z}^m$. In particular, the index set in the right hand side of Equation (3.6) has order $r_1 \dots r_m$.)

For $f \in \mathcal{K}$, the square of the L^2 norm of f as a section of the bundle L_r is given by

$$ab \sum_{c \in \mathbb{Z}} \int_0^1 |\hat{f}_c|^2(x) dx = ab \sum_{\bar{c} \in r\mathbb{Z} \backslash \mathbb{Z}} \int_{\mathbb{R}} |\hat{f}_{\bar{c}}|^2(x) dx.$$

Thus for each coset \bar{c} , we obtain a unitary isomorphism $T_{\bar{c}} : \mathcal{K}_{\bar{c}} \rightarrow L^2(\mathbb{R})$ given by

$$T_{\bar{c}}(f) = \sqrt{ab} \hat{f}_c$$

where c is an arbitrary but fixed choice of coset representative.

Let ρ denote the right action of N on \mathcal{K} . We have

$$(\rho(u', v', t')f)(u, v, t) = f(u + u', v + v', t + t' + ruv')$$

and thus the Fourier coefficients of $\rho(u', v', t')f$ are given by

$$(\rho(u', v', t')f)_{\hat{c}} = e^{-2\pi i[t' + (ru+c)v']} \hat{f}_c(u + u').$$

In particular, the right action leaves each subspace $\mathcal{K}_{\bar{c}}$ -invariant. Moreover, $T_{\bar{c}}$ intertwines the right action ρ on $\mathcal{K}_{\bar{c}}$ with the action η of N on $L^2(\mathbb{R})$ given by $\eta(u', v', t')g(w) = e^{-2\pi i[t' + (rw+c)v']}g(w + u')$. For each c as above, define $S_{\bar{c}} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ by

$$S_{\bar{c}}(g)(w) = g_{r,c}(w) = g\left(w - \frac{c}{r}\right).$$

Then $S_{\bar{c}} \circ T_{\bar{c}}$ intertwines the right action ρ on $\mathcal{K}_{\bar{c}}$ with the action τ given in Equation (3.3) and carries the distinguished Laplacian Δ on $\mathcal{K}_{\bar{c}}$ to the operator $D_{r,a,b}$ on $L^2(\mathbb{R})$.

We next construct an intertwining operator between the actions of $\Delta_{\varphi^D + \alpha}$ acting on $\mathcal{K}_{\bar{c}}$ and $D_{r,a,b}$ acting on $L^2(\mathbb{R})$. Define $P_{\alpha} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ by

$$(3.7) \quad P_{\alpha}(g)(w) = e^{2\pi i\mu w} g(w + \nu).$$

(In case $m > 1$, then μw is replaced by $\mu \cdot w$.) Then we have

$$(3.8) \quad P_{\alpha} \circ \tau(u, v, t) = \sigma(u, v, t) \circ P_{\alpha}$$

where

$$(3.9) \quad \sigma(u, v, t)g(w) = e^{-2\pi i(t+\mu u+\nu v+rvv)}g(w + u).$$

Thus $P_\alpha \circ S_{\bar{c}} \circ T_{\bar{c}}$ intertwines the right action ρ of N on $\mathcal{K}_{\bar{c}}$ with the action σ of N on $L^2(\mathbb{R})$. An elementary computation shows that $\sigma_*(\Delta_{\varphi^{D+\alpha}}) = D_{r,a,b}$; i.e., $P_\alpha \circ S_{\bar{c}} \circ T_{\bar{c}}$ intertwines $\Delta_{\varphi^{D+\alpha}}$ with $D_{r,a,b}$.

Note that multiplication by the potential function Q (given as in part (i) of the theorem by $Q(u, v) = q(u)$) carries \mathcal{K} to itself, leaves each $\mathcal{K}_{\bar{c}}$ invariant, and satisfies, for $f \in \mathcal{K}$, $(Qf)\widehat{c}(u) = q(u)\widehat{f}_c(u)$. Hence $T_{\bar{c}}(Qf) = qT_{\bar{c}}f$. We thus have $P_\alpha \circ S_{\bar{c}} \circ T_{\bar{c}}(Qf) = q_{r,c,\nu}P_\alpha \circ S_{\bar{c}} \circ T_{\bar{c}}(f)$. It follows that $P_\alpha \circ S_{\bar{c}} \circ T_{\bar{c}}$ intertwines the Schrödinger operator $\Delta_{\varphi^{D+\alpha}} + Q$ on $\mathcal{K}_{\bar{c}}$ with the operator $D_{r,a,b} + q_{r,c,\nu}$ on $L^2(\mathbb{R})$. This completes the proof of (1).

(2) The asymmetry between the roles of u and v above is due to our choice of coordinates on N . We have been denoting elements of N as (u, v, t) . To avoid confusion as we introduce new coordinates, we will temporarily write $(u, v, t)_1$ for (u, v, t) . Define new coordinates by setting $(u, v, t)_2 = (u, v, -t + ruv)_1$. An elementary computation using Equation (3.1) shows that

$$(u, v, t)_2(u', v', t')_2 = (u + u', v + v', t + t' + rvu')_2.$$

Observe that the discrete subgroup Γ of N given by all elements with integer coordinates $(u, v, t)_1$ still consists precisely of the points whose coordinates $(u, v, t)_2$ are integers. However, in the new coordinates, we have for $k, l \in \mathbb{Z}$,

$$(k, 0, 0)_2(u, v, t)_2 = (u + k, v, t)_2$$

and

$$(0, l, 0)_2(u, v, t)_2 = (u, v + l, t + rlu)_2.$$

Thus the left Γ -invariant functions are periodic in u rather than in v . The integral curves of U and V through the identity element of N are given by exactly the same expression in the new coordinates as in the old.

The function space \mathcal{K} is given in our new coordinates by those left Γ -invariant functions on N satisfying $f((u, v, t)_2) = e^{2\pi it}f((u, v, 0)_2)$. Write $\bar{\mathcal{K}} = \{\bar{f} : f \in \mathcal{K}\}$, where \bar{f} denotes the complex conjugate, and define $B : \mathcal{K} \rightarrow \bar{\mathcal{K}}$ by $f \mapsto \bar{f}$. Note that the actions of the Schrödinger operator $\Delta_{\varphi^{D+\alpha}} + Q = -(\frac{1}{a^2}(U - \mu Z)^2 + \frac{1}{b^2}(V - \nu Z)^2) + Q$ on \mathcal{K} and $\bar{\mathcal{K}}$ are intertwined by B since Q is real-valued and $\Delta_{\varphi^{D+\alpha}}$ is a real operator. Thus we may consider $\bar{\mathcal{K}}$ rather than \mathcal{K} . Since $\bar{\mathcal{K}}$ consists of Γ -invariant functions satisfying $f((u, v, t)_2) = e^{-2\pi it}f((u, v, 0)_2)$, the computation of the spectrum is now identical to that carried out in part (i) except that the roles of u and v are switched. Thus (ii) follows. □

4. Positive inverse spectral results on tori

In this section we prove positive inverse spectral results, including Theorem 1.2, Theorem 1.3, and additional results concerning the odd part of the potential.

Let L be a line bundle over the torus $M = \mathcal{L} \backslash \mathbb{R}^{2m}$ for which all the invariant factors r_1, \dots, r_m of the Chern class are equal to one. Let Ω be the translation-invariant 2-form representing the Chern class of L . We may view Ω as a bilinear form on \mathbb{R}^{2m} . Recall that $\Omega(x, y) = e_x(y) - e_y(x)$ (see Notation 2.4). Let \mathcal{L}' be the lattice in \mathbb{R}^{2m} dual to \mathcal{L} , i.e., $\mathcal{L}' = \{\lambda \in \mathbb{R}^{2m} : \lambda \cdot l \in \mathbb{Z} \text{ for all } l \in \mathcal{L}\}$. Define a linear isomorphism $S : \mathbb{R}^{2m} \rightarrow \mathbb{R}^{2m}$ by $S(u) \cdot x = \Omega(u, x)$ for all $x \in \mathbb{R}^{2m}$. Since Ω takes integer values on $\mathcal{L} \times \mathcal{L}$, the map S carries \mathcal{L} (injectively) into \mathcal{L}' . The fact that the invariant factors of Ω are all equal to one implies that $S(\mathcal{L}) = \mathcal{L}'$, thus S gives an isomorphism between the lattices \mathcal{L} and \mathcal{L}' .

The proof of Theorem 1.2 relies on the following result, due to Guillemin [10] in the case $m = 1$ and $\alpha = 0$. For a proof with m arbitrary, see Appendix C.

PROPOSITION 4.1. — *Let $L \rightarrow M$ be a line bundle over $M = \mathcal{L} \backslash \mathbb{R}^{2m}$ with Chern invariant factors $r_1 = \dots = r_m = 1$ and assume that \mathcal{L} has nondegenerate length spectrum. Let $\alpha \in \mathbb{R}^{2m}$. For $l \in \mathcal{L} \setminus \{0\}$, set*

$$W_l^\alpha := e^{-2\pi i \alpha \cdot l} \int_{\mathcal{F}} dx e^{2\pi i \Omega(x, l)} \int_0^{|l|} d\tau Q \left(x - \tau \frac{l}{|l|} \right)$$

(Here \mathcal{F} is a fundamental domain for $\mathcal{L} \backslash \mathbb{R}^{2m}$.) Then $W_l^\alpha + W_{-l}^\alpha$ is an invariant of $\text{Spec}_\alpha(Q; L)$ for every $l \in \mathcal{L} \setminus \{0\}$. (See Notation 2.21.)

COROLLARY 4.2. — *Express a potential Q in its Fourier series*

$$(4.1) \quad Q(x) = \sum_{\gamma \in \mathcal{L}'} c_\gamma e^{2\pi i \gamma \cdot x}.$$

Under the hypothesis of Proposition 4.1, for each nonzero $\lambda \in \mathcal{L}'$, the expression

$$e^{-2\pi i \alpha \cdot l} c_\lambda + e^{2\pi i \alpha \cdot l} c_{-\lambda},$$

where $l = S^{-1}(\lambda)$, is an invariant of $\text{Spec}_\alpha(Q; L)$.

Proof of Corollary 4.2. — Let $\lambda \in \mathcal{L}'$ and let $l = S^{-1}(\lambda)$, so that $\Omega(l, x) = \lambda \cdot x$ for all $x \in \mathbb{R}^{2m}$. Then

$$W_l^\alpha = e^{-2\pi i \alpha \cdot l} \int_{\mathcal{F}} dx e^{-2\pi i \lambda \cdot x} \int_0^{|l|} d\tau \left(\sum_{\gamma \in \mathcal{L}'} c_\gamma e^{2\pi i \gamma \cdot x} e^{-2\pi i \tau \gamma \cdot \frac{l}{|l|}} \right)$$

$$= e^{-2\pi i \alpha \cdot l} \sum_{\gamma \in \mathcal{L}'} c_\gamma \int_{\mathcal{F}} dx e^{-2\pi i (\lambda - \gamma) \cdot x} \int_0^{|\lambda|} d\tau \left(e^{-2\pi i \tau \gamma \cdot \frac{1}{|\lambda|}} \right)$$

Since the first integral in each term of the latter sum vanishes when $\gamma \neq \lambda$ and since $\lambda \cdot l = \Omega(l, l) = 0$, we thus have $W_l^\alpha = |\lambda| \text{vol}(\mathcal{F}) e^{-2\pi i \alpha \cdot l} c_\lambda$ and thus

$$W_l^\alpha + W_{-l}^\alpha = |\lambda| \text{vol}(\mathcal{F}) \left(e^{-2\pi i \alpha \cdot l} c_\lambda + e^{2\pi i \alpha \cdot l} c_{-\lambda} \right).$$

The corollary follows. □

PROPOSITION 4.3. — [5] *Let L be a line bundle over the torus M , let ∇ be an arbitrary connection on L , and let Δ be the associated Laplacian. Then $\int_M Q(x) dx$ and $\int_M Q^2(x) dx$ are spectral invariants of the Schrödinger operator $\Delta + Q$.*

In particular, the spectrum determines the coefficient c_0 in the Fourier expansion (4.1) of Q .

Proof of Proposition 4.3. — This result follows immediately from the heat asymptotics computed by P. Gilkey in [5]. Indeed, Gilkey considered an arbitrary second order operator D on a vector bundle V over a Riemannian manifold M such that the leading symbol of D is the metric tensor. In his notation, $D = D_\nabla - \mathcal{E}$, where D_∇ is the Laplacian associated with the unique connection on V for which $D_\nabla - D$ is a zeroth order operator. In our case, D is the Schrödinger operator $\Delta + Q$, $D_\nabla = \Delta$, and $\mathcal{E} = -Q$. Consider the asymptotics of the trace of the heat kernel e_Q of the Schrödinger operator $\Delta + Q$ as $t \rightarrow 0^+$:

$$e_Q(t, x, x) = \frac{1}{(4\pi t)^{n/2}} (1 + tu_1(x) + t^2 u_2(x) + o(t^2)).$$

The integral $\int_M u_i(x) dx$ is a spectral invariant of $\Delta + Q$ for each i . By Theorem 4.3 in [5] and the fact that M is flat, we have $u_1 = \mathcal{E} = -Q$; hence $\int_M Q(x) dx$ is a spectral invariant of $\Delta + Q$. Next $u_2 = \frac{1}{2} Q^2 - \frac{1}{6} \Delta Q$ plus terms depending only on the curvature of the connection. Thus by the Divergence Theorem, the L^2 norm of Q is a spectral invariant. □

Proof of Theorem 1.2. —

- (a) By hypothesis, the connection ∇ is translation-invariant and weakly \mathbb{Z}_2 -invariant. By Proposition 2.16 and our standing hypothesis that $g = 0$ (cf Remark 2.19(ii)), $\nabla = \nabla^D + 2\pi i \alpha$ with $\alpha(\mathcal{L}) \subset \frac{1}{2}\mathbb{Z}$. Hence for $l \in \mathcal{L}$, we have $e^{2\pi i \alpha \cdot l} = e^{-2\pi i \alpha \cdot l}$, and thus by Corollary 4.2, $c_\gamma + c_{-\gamma}$ is a spectral invariant of $\Delta_{\varphi^D + \alpha} + Q$ for all $0 \neq \gamma \in \mathcal{L}'$. So is c_0 by Proposition 4.3. Thus the even part Q^+ of Q is spectrally determined by $\Delta_{\varphi^D + \alpha} + Q$.

- (b) By (a) and Proposition 4.3, the spectrum of $\Delta_{\varphi^{D+\alpha}} + Q$ determines the L^2 norms of both Q and of the even part of Q , and hence it determines whether Q is even. Thus (b) follows from (a). □

Remark 4.4. — Theorem 1.2 together with Proposition 4.3 show that, under the hypotheses of Theorem 1.2, the L^2 norm of the odd part $Q^- = \frac{1}{2}(Q - \tilde{Q})$ of Q is a spectral invariant of the Schrödinger operator $\Delta + Q$.

Proof of Theorem 1.3. — When one drops the assumption that \mathcal{L} has nondegenerate length spectrum, then the spectral invariant $W_l^\alpha + W_{-l}^\alpha$ in Proposition 4.1 is replaced by

$$\sum_{|l|=d} W_l^\alpha$$

for each length d appearing in $|\mathcal{L}|$. Thus from the computation in Corollary 4.2, we obtain the following invariant of $Spec_\alpha(Q; L)$ for each such d :

$$(4.2) \quad V_d^\alpha := \sum_{l \in \mathcal{L}, |l|=d} e^{-2\pi i \alpha \cdot l} c_{S(l)}.$$

Since the mapping $S : \mathcal{L} \rightarrow \mathcal{L}'$ is invertible, we need only show that for each d , we can recover the individual Fourier coefficients $c_{S(l)}$ from the expressions V_d^α by varying the choices of α .

Given d , let $\pm l_1, \dots, \pm l_k$ be all the lattice vectors of norm d . We need to choose $\alpha_1, \dots, \alpha_{2k}$ so that the matrix

$$\begin{pmatrix} e^{-2\pi i \alpha_1 \cdot l_1} & e^{2\pi i \alpha_1 \cdot l_1} & \dots & e^{-2\pi i \alpha_1 \cdot l_k} & e^{2\pi i \alpha_1 \cdot l_k} \\ \vdots & \vdots & & \vdots & \vdots \\ e^{-2\pi i \alpha_{2k} \cdot l_1} & e^{2\pi i \alpha_{2k} \cdot l_1} & \dots & e^{-2\pi i \alpha_{2k} \cdot l_k} & e^{2\pi i \alpha_{2k} \cdot l_k} \end{pmatrix}$$

is nonsingular. Choose $\alpha_1 = 0$ and choose α_2 subject only to the genericity condition that the entries in the second row of the matrix are all distinct. Then by choosing $\alpha_j = (j - 1)\alpha_2$ for $j = 2, \dots, 2k$, the determinant of the matrix above is a nonzero Vandermonde determinant. The theorem follows. □

Remark 4.5. — In the special case that the length spectrum is nondegenerate, then one needs only two Bloch spectra, $Spec_\alpha(Q; L)$ and $Spec_\beta(Q; L)$ to determine Q , where α and β are chosen subject only to the condition that $(\alpha - \beta) \cdot l$ is irrational for all $l \in \mathcal{L}$. Indeed, in that case the matrix

$$\begin{bmatrix} e^{-2\pi i \alpha \cdot l} & e^{2\pi i \alpha \cdot l} \\ e^{-2\pi i \beta \cdot l} & e^{2\pi i \beta \cdot l} \end{bmatrix}$$

is invertible for every l , so the spectral invariants $W_l^\alpha + W_{-l}^\alpha$ and $W_l^\beta + W_{-l}^\beta$ determine $S(\pm l)$ for each l .

This result fits nicely with results of the third author [11] concerning a discrete version of the Schrödinger equation on a 2-dimensional torus. There it was shown that, generically, the periodic spectrum does not determine the Bloch spectrum of Q but, again generically, two spectra $Spec_\alpha(Q)$ and $Spec_\beta(Q)$ do.

Remark 4.6. — Theorem 1.3 contrasts with spectral results for the Schrödinger operator on trivial line bundles over tori. (See Remark 2.23 for notation.) In the case of a trivial bundle, we have the following: (i) Given any smooth real potential Q , then Q and \check{Q} necessarily have the same classical Bloch spectrum. (Indeed, given a smooth function $f : \mathbb{R}^n \rightarrow \mathbb{C}$ satisfying $(\Delta + Q)f = \mu f$ and the Bloch condition $f(x + l) = e^{-2\pi i \alpha \cdot l} f(x)$ for all $l \in \mathcal{L}$, let $g := \bar{f}$ be the complex conjugate of f . Then g satisfies $(\Delta + \check{Q})g = \mu g$ and $g(x + l) = e^{-2\pi i \alpha \cdot l} g(x)$ for all $l \in \mathcal{L}$.) Moreover, (ii) Eskin, Ralston and Trubowitz [4], Theorem 6.2, proved that if Q is a real analytic potential on a torus with nondegenerate length spectrum, then the spectrum of the Schrödinger operator $\Delta + Q$ acting on periodic functions determines the classical Bloch spectrum of Q . See also G. Eskin [3] and Gordon Kappeler [6] for related results.

Theorem 1.3 shows that (i) always fails in the setting of line bundles L with Chern invariant factors $r_1 = \dots = r_m = 1$. Also by Remark 2.19(v), Q and \check{Q} are isospectral with respect to the distinguished connection (and in fact, with respect to any weakly \mathbb{Z}_2 -invariant connection). Thus (ii) also fails. However, we do have that $Spec_\alpha(Q) = Spec_{-\alpha}(\check{Q})$ for all α ; indeed the map $g \rightarrow \check{g}$ of $C^\infty(\mathbb{R}^n)$ intertwines the Schrödinger operators $\Delta_{\varphi^D + \alpha} + Q$ and $\Delta_{\varphi^D - \alpha} + \check{Q}$. Thus the result (ii) above of Eskin, Ralston, and Trubowitz suggests the following question: Under the hypotheses of Theorem 1.2, let P and Q be real analytic potentials on M such that $Spec_0(P; L) = Spec_0(Q; L)$. Must it be the case that for each $\alpha \in \mathbb{R}^n$, either $Spec_\alpha(P; L) = Spec_\alpha(Q; L)$ or $Spec_\alpha(P; L) = Spec_{-\alpha}(Q; L)$? If the answer is yes, then Theorem 1.3 would imply for real analytic potentials Q on even-dimensional tori with nondegenerate length spectrum that $Iso(Q; L; \nabla^D) = \{Q, \check{Q}\}$ without any assumptions on the parity of Q .

It seems likely that $Iso(Q; L, \nabla^D) = \{Q, \check{Q}\}$ for generic smooth potentials Q . This remains an open problem.

Finally, we give an extension of Theorem 1.2 in the case of a 2-dimensional torus:

Notation 4.7. — Denote by \mathcal{S} the set of maximal elements in the dual lattice \mathcal{L}' of \mathcal{L} , i.e. the set of elements λ in \mathcal{L}' such that $\{\lambda \cdot l \mid l \in \mathcal{L}\} = \mathbb{Z}$. Let $\mathcal{S}_+ \subset \mathcal{S}$ be a subset with the property that for any $\lambda \in \mathcal{S}$, λ or $-\lambda$ is an element of \mathcal{S}_+ but not both. Then the Fourier decomposition of Q

$$(4.3) \quad Q(x) = c_0 + \sum_{\gamma \in \mathcal{L}', \gamma \neq 0} c_\gamma e^{2i\pi\gamma \cdot x}$$

can be written as (see [4])

$$(4.4) \quad Q(x) = c_0 + \sum_{\lambda \in \mathcal{S}_+} Q_\lambda(\lambda \cdot x)$$

where Q_λ is the one dimensional potential

$$(4.5) \quad Q_\lambda(s) := \sum_{k \in \mathbb{Z} \setminus \{0\}} c_{k\lambda} e^{2\pi i k s}.$$

Note that if Q^\pm are the even and odd parts of the potential Q , then $(Q^\pm)_\lambda = (Q_\lambda)^\pm$ for all $\lambda \in \mathcal{L}'$.

LEMMA 4.8. — Assume that the torus M is two-dimensional. Let $\lambda \in \mathcal{S}_+$ and let $l = S^{-1}(\lambda)$, so that $\Omega(l, x) = \lambda \cdot x$ for all x . Let Q be any smooth potential of mean zero (i.e., the Fourier coefficient c_0 of Q vanishes). Then for $0 \neq k \in \mathbb{Z}$, we have

$$\int_0^{|kl|} d\tau Q\left(x - \tau \frac{l}{|l|}\right) = |kl| Q_\lambda(\lambda \cdot x).$$

Proof of Lemma 4.8. — We have $\lambda \cdot l = \Omega(l, l) = 0$ since Ω is skew-symmetric. Since M is two-dimensional, it follows that for $0 \neq \gamma \in \mathcal{L}'$, we have $\gamma \cdot l = 0$ only when γ is a multiple of λ . Thus, with c_γ , $\gamma \in \mathcal{L}'$, denoting the Fourier coefficients of Q , we have

$$\begin{aligned} \int_0^{|kl|} d\tau Q\left(x - \tau \frac{l}{|l|}\right) &= \sum_{\gamma \in \mathcal{L}'} e^{2\pi i \gamma \cdot x} c_\gamma \int_0^{|kl|} d\tau e^{-2\pi i \tau \gamma \cdot \frac{l}{|l|}} \\ &= \sum_{0 \neq j \in \mathbb{Z}} |kl| e^{2\pi i j \lambda \cdot x} c_{j\lambda} = |kl| Q_\lambda(\lambda \cdot x). \end{aligned}$$

□

THEOREM 4.9. — Assume that M is a 2-dimensional torus with nondegenerate length spectrum and $L \rightarrow M$ is a line bundle with Chern invariant factor $r_1 = 1$. Then the squares of the odd parts of the 1-dimensional potentials Q_λ defined in Notation 4.7 are spectral invariants of $\Delta^D + Q$; i.e., for any $P \in \text{Iso}(Q; L, \nabla^D)$ and any $\lambda \in \mathcal{S}_+$, $(P_\lambda^-)^2 = (Q_\lambda^-)^2$.

Proof. — Fix $\lambda \in \mathcal{S}_+$ and let $l = S^{-1}(\lambda)$. By Theorem C.5 in Appendix C, for every $j \in \mathbb{Z} \setminus \{0\}$, the expression $C_{2,jl}$, given as in Theorem C.9, is a spectral invariant. (Here we are letting the element $jl \in \mathcal{L}$ play the role of l in Theorems C.5 and C.9.) By Lemma C.7 and the assumption that the connection is the distinguished one, the function a_0 appearing in the expression in Theorem C.9 is given by $a_0(|jl|, x + jl, -\frac{l}{|l|}) = \epsilon_{jl} e^{2\pi i e_x(jl)}$, where $\epsilon_{jl} = e^{-\pi i e_{jl}(jl)} = \pm 1$. Since $\Omega(x, y) = e_x(y) - e_y(x)$, we have for any $l \in \mathcal{L} \setminus \{0\}$ that $C_{2,jl}$ is a constant multiple of

$$(4.6) \quad \int_{\mathcal{F}} dx e^{2\pi i \Omega(x, jl)} \left(\int_0^{|jl|} d\tau Q^- \left(x - \tau \frac{l}{|l|} \right) \right)^2.$$

Since $l = S^{-1}(\lambda)$, we have $\Omega(x, jl) = -j\lambda \cdot x$. Thus by Equation (4.6), Lemma 4.8 and the fact that odd potentials always have mean zero, the following expression is a spectral invariant for each j :

$$(4.7) \quad |jl|^2 \int_{\mathcal{F}} dx e^{-2\pi i j\lambda \cdot x} (Q_{\lambda}^-(\lambda \cdot x))^2$$

Writing $(Q_{\lambda}^-(\lambda \cdot x))^2 = \sum_{j \in \mathbb{Z}} b_{j\lambda} e^{2\pi i j\lambda \cdot x}$, the spectral invariance of the expression (4.7) implies that the $b_{j\lambda}$, for $0 \neq j \in \mathbb{Z}$, are spectral invariants. Thus the function $h(x) := (Q_{\lambda}^-(\lambda \cdot x))^2 - b_{0,\lambda}$ is spectrally determined. Note that $h(0) = b_{0,\lambda}$ since Q_{λ}^- is an odd function. Thus $b_{0,\lambda}$ is also a spectral invariant and hence so is the function $(Q_{\lambda}^-(\lambda \cdot x))^2$. This completes the proof of Theorem 4.9. □

COROLLARY 4.10. — *Assume that M is a 2-dimensional torus with non-degenerate length spectrum and $L \rightarrow M$ is a line bundle with Chern invariant factor $r_1 = 1$. If $\{Q_t\}_{t \in [0, \epsilon]}$ is any continuous family of smooth mutually isospectral potentials on M with respect to the distinguished connection, then $Q_t = Q_0$ for all t ; i.e., within the space of smooth potentials, there are no nontrivial continuous isospectral deformations.*

Proof. — By Theorem 1.2, we have $Q_t^+ = Q_0^+$ for all t , and by Theorem 4.9, for each $\lambda \in \mathcal{S}_+$ and $x \in M$, we have

$$(4.8) \quad (Q_t)_{\lambda}^-(\lambda \cdot x) = \pm (Q_0)_{\lambda}^-(\lambda \cdot x)$$

for all t (where a priori the sign may depend on x and t). In particular, the functions $(Q_t)_{\lambda}^-(\lambda \cdot x)$ and $(Q_0)_{\lambda}^-(\lambda \cdot x)$ have the same zeroes for all t . By continuity of the map $t \rightarrow (Q_t)_{\lambda}^-(\lambda \cdot x)$ and Equation (4.8), it follows that $(Q_t)_{\lambda}^-(\lambda \cdot x) = (Q_0)_{\lambda}^-(\lambda \cdot x)$ for all t, x, λ and thus that $Q_t = Q_0$. □

COROLLARY 4.11. — Assume that M is a 2-dimensional torus with non-degenerate length spectrum and $L \rightarrow M$ is a line bundle with Chern invariant factor $r_1 = 1$. Suppose that the odd part Q^- of Q is a 1-dimensional potential, i.e., there exists $\lambda \in \mathcal{S}_+$ such that $Q^-(x) = Q_\lambda^-(\lambda \cdot x)$. Let $P \in \text{Iso}(Q; L, \nabla^D)$. Then

(i) $P^-(x) = P_\lambda^-(\lambda \cdot x)$ and $(P^-)^2 = (Q^-)^2$.

(ii) If Q^- and P^- are real analytic, then $P = Q$ or $P = \check{Q}$.

Proof. — The first statement follows from Theorem 4.9. For the second statement, since P^- and Q^- are both real analytic and $(P^-)^2 = (Q^-)^2$, we have $P^- = \pm Q^-$. By Theorem 1.2, we also have $P^+ = Q^+$ and thus either $P = Q$ or $P = \check{Q} = Q^+ - Q^-$. \square

In contrast, for the rectangular two-dimensional tori considered in Theorem 1.1, the examples of L -isospectral, noncongruent potentials include real analytic odd one-dimensional potentials.

Appendices

Throughout these appendices, we use the notation of Subsection 2B. In particular, L is a fixed Hermitian line bundle over the torus M , ∇ is a connection on L compatible with the Hermitian structure, φ is the 1-form on \mathbb{R}^n associated with ∇ as in Definition 2.7, and Δ_φ is the associated Laplacian, given as in Proposition 2.20. We are following closely the constructions developed in [4], adapting them to the case of non-trivial line bundles.

While we are interested primarily in translation-invariant connections, we make no assumption on the connection in Appendices A-C, except where specified at the end of Appendix C.

Appendix A. Wave kernel on \mathbb{R}^n

Consider the wave equation on $\mathbb{R} \times \mathbb{R}^n$, $n \geq 2$, with smooth potential $Q : \mathbb{R}^n \rightarrow \mathbb{R}$

$$(A.1) \quad (\square_\varphi + Q(x))u(t, x) = 0 \quad (x \in \mathbb{R}^n, t \in \mathbb{R})$$

with initial conditions

$$(A.2) \quad u(0, x) = u_0(x); \quad \partial_t u(0, x) = 0,$$

where $u_0 \in C^\infty(\mathbb{R} \times \mathbb{R}^n, \mathbb{C})$ and

$$(A.3) \quad \square_\varphi = \frac{\partial^2}{\partial t^2} + \Delta_\varphi.$$

Denote by $K(t, x, y)$ the distributional wave kernel and $K(t)$ the wave operator corresponding to (A.1) and (A.2); i.e., for any choice of u_0 , the function

$$u(t, x) = (K(t)u_0)(x) := \int_{\mathbb{R}^n} K(t, x, y)u_0(y)dy$$

solves (A.1) - (A.2). The operator $K(t)$ can be written as the sum of two integral operators $\frac{1}{2}(K_+(t) + K_-(t))$ where K_\pm has a symbol of the form $\frac{1}{(2\pi)^n}e^{iS^\pm}a^\pm$ with $S^\pm = S^\pm(t, x, y, \zeta)$ denoting the phase function of K_\pm and $a^\pm = a^\pm(t, x, \zeta)$ its amplitude. (Here $t \in \mathbb{R}$ and $x, y, \zeta \in \mathbb{R}^n$.) As $S^\pm(t, x, y, \zeta)$ only depends on the principal symbol of $\Delta_\varphi + Q$, which is independent of φ and Q , S^\pm are the corresponding phase functions for the free wave equation on \mathbb{R}^n , i.e.

$$S^\pm(t, x, y, \zeta) = (x - y) \cdot \zeta \pm t|\zeta|.$$

Thus

$$(K_\pm(t)u_0)(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta \int_{\mathbb{R}^n} dy e^{i(x-y)\cdot\zeta \pm it|\zeta|} a^\pm(t, x, \zeta)u_0(y).$$

The amplitudes a^\pm satisfy the following estimates: for any choice of multi-indices $\alpha = (\alpha_j)_{1 \leq j \leq n}$, $\beta = (\beta_j)_{1 \leq j \leq n}$ and any choice of $R > 0$ and $T > 0$, there exists a constant $C_{\alpha, \beta} = C_{\alpha, \beta}(R, T)$ such that for any $\zeta \in \mathbb{R}^n$,

$$\sup_{0 \leq t \leq T, |x| < R} \left| \partial_x^\alpha \partial_\zeta^\beta a^\pm(t, x, \zeta) \right| \leq C_{\alpha, \beta}(1 + |\zeta|)^{-|\beta|}$$

where $|\beta| = \beta_1 + \dots + \beta_n$. Moreover they satisfy the initial conditions $a^\pm(0, x, \zeta) = 1$.

Following [13] (see [4] for the case $\varphi \equiv 0$), we approximate $K_\pm(t)$ by

$$(A.4) \quad (K_\pm^N(t)u_0)(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta \int_{\mathbb{R}^n} dy e^{i(x-y)\cdot\zeta \pm it|\zeta|} \times (a_0^\pm + \dots + a_N^\pm)(t, x, \zeta)\chi(|\zeta|)u_0(y)$$

where $a_j^\pm = a_j^\pm(t, x, \zeta)$ is positive-homogeneous of degree $-j$ in ζ , i.e. $\forall c > 0, t > 0, x \in \mathbb{R}^n, \zeta \in \mathbb{R}^n, a_j^\pm(t, x, c\zeta) = c^{-j}a_j^\pm(t, x, \zeta)$, and a_j^\pm satisfies the initial conditions

$$(A.5) \quad a_0^\pm(0, x, \zeta) = 1 \text{ and } a_j^\pm(0, x, \zeta) = 0 \text{ (} 1 \leq j \leq N \text{)}.$$

Here $\chi(r)$ is a smooth cut-off function of the form

$$(A.6) \quad \chi(r) = \begin{cases} 0 & \text{for } -\infty < r \leq 1/2 \\ 1 & \text{for } 1 \leq r < \infty. \end{cases}$$

The functions $a_j^\pm(t, x, \zeta)$ are determined inductively by solving transport equations. For any given $M \geq 0$ we will choose the integer $N \geq 0$ so large that the integral operators $\tilde{K}_\pm(t)$ defined by

$$(A.7) \quad (\square_\varphi + Q) K_\pm^N(t)u_0 = \tilde{K}_\pm(t)u_0,$$

have C^M kernels. Indeed, $(\square_\varphi + Q) K_\pm^N(t)u_0(x)$ equals

$$\begin{aligned} & \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta \int_{\mathbb{R}^n} dy (\square_\varphi + Q(x)) \\ & \quad \times \left(e^{i(x-y)\cdot\zeta \pm it|\zeta|} (a_0^\pm + \dots + a_N^\pm)(t, x, \zeta) \right) \chi(|\zeta|)u_0(y) \end{aligned}$$

where \square_φ is computed with respect to the variables t and x . By Proposition 2.20, using the notation $\varphi|_x$ established in Remark 2.8,

$$\begin{aligned} & \square_\varphi \cdot e^{i(x-y)\cdot\zeta \pm it|\zeta|} \\ & \quad = \left(4\pi\varphi|_x(\zeta) + 4\pi^2 |\varphi|_x|^2 - 2\pi i \operatorname{div}(\varphi^\#)(x) \right) e^{i(x-y)\cdot\zeta \pm it|\zeta|}. \end{aligned}$$

Further, one has the product formula

$$(A.8) \quad \begin{aligned} \square_\varphi(AB) &= (\square_\varphi A)B + A(\square_\varphi B) - 4\pi^2 |\varphi|^2 AB + 2 \frac{\partial A}{\partial t} \frac{\partial B}{\partial t} \\ & \quad - 2 \operatorname{grad} A \cdot \operatorname{grad} B + 2\pi i AB \operatorname{div}(\varphi^\#) \end{aligned}$$

Hence

$$\begin{aligned} & (\square_\varphi + Q(x)) \left(e^{i(x-y)\cdot\zeta \pm it|\zeta|} (a_0^\pm + \dots + a_N^\pm)(t, x, \zeta) \right) \\ &= e^{i(x-y)\cdot\zeta \pm it|\zeta|} \left(4\pi\varphi|_x(\zeta) \pm 2i|\zeta| \frac{\partial}{\partial t} - 2i\zeta \cdot \operatorname{grad} + (\square_\varphi + Q(x)) \right) \\ & \quad \cdot (a_0^\pm + \dots + a_N^\pm)(t, x, \zeta) \\ &= e^{i(x-y)\cdot\zeta \pm it|\zeta|} \left[\left(4\pi\varphi|_x(\zeta) a_0^\pm(t, x, \zeta) \pm 2i|\zeta| \frac{\partial a_0^\pm}{\partial t}(t, x, \zeta) - 2i\zeta \cdot \operatorname{grad} a_0^\pm(t, x, \zeta) \right) \right. \\ & \quad + \sum_{j=1}^N \left(4\pi\varphi|_x(\zeta) a_j^\pm(t, x, \zeta) \pm 2i|\zeta| \frac{\partial a_j^\pm}{\partial t}(t, x, \zeta) - 2i\zeta \cdot \operatorname{grad} a_j^\pm(t, x, \zeta) \right) \\ & \quad \left. + (\square_\varphi + Q(x)) a_{j-1}^\pm(t, x, \zeta) \right] + (\square_\varphi + Q(x)) a_N^\pm(t, x, \zeta) \end{aligned}$$

where $\text{grad } a_j^\pm(t, x, \zeta)$ refers to the gradient with respect to x . Thus we are led to the transport equations

$$(A.9) \quad \begin{cases} 4\pi\varphi|_x(\zeta)a_0^\pm(t, x, \zeta) \pm 2i|\zeta| \frac{\partial a_0^\pm}{\partial t}(t, x, \zeta) - 2i\zeta \cdot \text{grad } a_0^\pm(t, x, \zeta) = 0 \\ a_0^\pm(0, x, \zeta) = 1 \end{cases}$$

and for $1 \leq j \leq N$,

$$(A.10) \quad \begin{cases} 4\pi\varphi|_x(\zeta) a_j^\pm(t, x, \zeta) \pm 2i|\zeta| \frac{\partial a_j^\pm}{\partial t}(t, x, \zeta) - 2i\zeta \cdot \text{grad } a_j^\pm(t, x, \zeta) \\ = -(\square_\varphi + Q(x)) a_{j-1}^\pm(t, x, \zeta) \\ a_j^\pm(0, x, \zeta) = 0 \end{cases}$$

Note that the solution $a_0^+(t, x, \zeta)$ of (A.9) with $+$ has the property that $a_0^+(-t, x, \zeta)$ satisfies (A.9) with $-$. By uniqueness, it then follows that $a_0^-(t, x, \zeta) = a_0^+(-t, x, \zeta)$. As a consequence, $\square_\varphi a_0^+(-t, x, \zeta) = \square_\varphi a_0^-(t, x, \zeta)$ and hence by the same argument, Equation (A.10) implies that $a_1^-(t, x, \zeta) = a_1^+(-t, x, \zeta)$ and, by induction, that $a_j^-(t, x, \zeta) = a_j^+(-t, x, \zeta)$ for any $j \geq 0$. Hence, for any $j \geq 0$,

$$(A.11) \quad \frac{\partial}{\partial t} \Big|_{t=0} (a_j^+(t, x, \zeta) + a_j^-(t, x, \zeta)) = 0$$

Equation (A.7) is then satisfied with the kernel of the integral operator $\tilde{K}_\pm(t)$ given by

$$(A.12) \quad \tilde{K}_\pm(t, x, y) := \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta e^{i(x-y)\cdot\zeta \pm it|\zeta|} ((\square_\varphi + Q(x)) a_N^\pm(t, x, \zeta)) \chi(|\zeta|).$$

As a_N^\pm is positive-homogeneous of degree $-N$ in ζ , so is $(\square_\varphi + Q) a_N^\pm$. Hence, for each of \tilde{K}_\pm , it follows that the integral in (A.12) is convergent for $N \geq n + 1$ and, given any M , there exists $N \geq n + 1$ so that it is a C^M function of x and y , as required.

We are now able to check the accuracy of the approximation of K by $K^N := \frac{1}{2}(K_+^N + K_-^N)$. Since

$$\begin{aligned} & \frac{d}{dt} \Big|_{t=0} [K^N(t)u_0](x) \\ &= \frac{1}{2} \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta \int_{\mathbb{R}^n} dy e^{i(x-y)\cdot\zeta} \sum_{j=0}^N \left(\frac{\partial}{\partial t} a_j^+ + \frac{\partial}{\partial t} a_j^- \right) (0, x, \zeta) \chi(|\zeta|) u_0(y), \end{aligned}$$

Equation (A.11) implies that

$$(A.13) \quad \frac{\partial}{\partial t} \Big|_{t=0} K^N(t)u_0 = 0.$$

Moreover, by (A.5)

$$K^N(0)u_0(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta \int_{\mathbb{R}^n} dy e^{i(x-y)\cdot\zeta} \chi(|\zeta|) u_0(y).$$

Hence, with \widehat{u}_0 denoting the Fourier transform of u_0 ,

$$\begin{aligned} K^N(0)u_0(x) &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta e^{ix\cdot\zeta} \widehat{u}_0(\zeta) + \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta \int_{\mathbb{R}^n} dy e^{i(x-y)\cdot\zeta} (\chi(|\zeta|) - 1) u_0(y) \\ &= u_0(x) + \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} d\zeta \int_{\mathbb{R}^n} dy e^{i(x-y)\cdot\zeta} (\chi(|\zeta|) - 1) u_0(y), \end{aligned}$$

by the inverse Fourier transform theorem in $L^2(\mathbb{R}^n)$. Thus

$$(A.14) \quad K^N(0)u_0 = u_0 + K_0^N u_0$$

where K_0^N is an integral operator with a smooth kernel.

Denoting by \underline{K}^N the difference $K^N - K$, we are led to the following Cauchy problem

$$(A.15) \quad \begin{cases} (\square_\varphi + Q) \underline{K}^N(t)u_0 = \widetilde{K}(t)u_0 \\ \underline{K}^N(0)u_0 = K_0^N u_0 \\ \frac{\partial}{\partial t} \Big|_{t=0} \underline{K}^N(t)u_0 = 0, \end{cases}$$

where $\widetilde{K}(t) = \frac{1}{2}(\widetilde{K}_+ + \widetilde{K}_-)$ has a C^M kernel for a given M , provided N is sufficiently large, and where K_0^N has a smooth kernel. By regularity theory for solutions of (A.15), $\underline{K}^N(t)$ is an integral operator with a $C^{M'}$ kernel, for any given M' , provided N is large enough. This will be useful in Appendix C (see the proof of Proposition C.2).

We now solve (A.9) and (A.10). As we will only need the a_j^+ 's we compute only these coefficients. Let

$$(A.16) \quad a_j = a_j^+ \Big|_{\mathbb{R}^+ \times \mathbb{R}^n \times S^{n-1}}.$$

Since $a_j^+(t, x, \zeta)$ is positive-homogeneous in ζ , it suffices to solve for the functions a_j . Dividing (A.9) by $2i$ and restricting to $\mathbb{R}^+ \times \mathbb{R}^n \times S^{n-1}$, one

gets

$$(A.17) \quad \begin{cases} \frac{\partial a_0}{\partial t}(t, x, \omega) - \omega \cdot \text{grad } a_0(t, x, \omega) - 2\pi i \varphi|_x(\omega) a_0(t, x, \omega) = 0 \\ a_0(0, x, \omega) = 1 \end{cases}$$

LEMMA A.1. — Equation (A.17) has the following solution

$$(A.18) \quad a_0(t, x, \omega) = \exp \left(2\pi i \int_0^t \varphi|_{(x+\tau\omega)}(\omega) \, d\tau \right).$$

The function a_0 is smooth in all its arguments.

Proof. — First note that if $h \in C^\infty(\mathbb{R}^n)$, then for fixed ω , the function $g(t, x) = \exp \left(2\pi i \int_0^t h(x + \tau\omega) d\tau \right)$ satisfies (with grad denoting the gradient with respect to x):

$$\begin{aligned} \omega \cdot \text{grad } g(t, x) &= g(t, x) 2\pi i \int_0^t \omega \cdot \text{grad}(h)(x + \tau\omega) \, d\tau \\ &= g(t, x) 2\pi i \int_0^t \frac{\partial}{\partial \tau} h(x + \tau\omega) \, d\tau \\ &= g(t, x) 2\pi i h(x + t\omega) - g(t, x) 2\pi i h(x) \\ &= \frac{\partial g}{\partial t}(t, x) - 2\pi i h(x) g(t, x). \end{aligned}$$

Moreover $g(0, x) = 1$. Let $h(t, x) = \varphi|_{(x+\tau\omega)}(\omega)$ (again viewing ω as arbitrary but fixed) to see that the function defined in Equation (A.18) satisfies (A.17). □

Similarly, dividing (A.10) by $2i$, we obtain in view of (A.16)

$$(A.19) \quad \begin{cases} \frac{\partial a_j}{\partial t}(t, x, \omega) - \omega \cdot \text{grad } a_j(t, x, \omega) - 2\pi i \varphi|_x(\omega) a_j(t, x, \omega) \\ \hspace{15em} = \frac{i}{2} (\square_\varphi + Q(x)) a_{j-1}(t, x, \omega) \\ a_j(0, x, \omega) = 0 \end{cases}$$

We solve (A.19) by the method of variation of constants. First for $0 \leq j \leq N$, define $f_j : \mathbb{R}^+ \times \mathbb{R}^n \times S^{n-1} \rightarrow \mathbb{R}$ by

$$(A.20) \quad f_j(t, x, \omega) = \frac{a_j(t, x, \omega)}{a_0(t, x, \omega)}.$$

By (A.17) and (A.20), we may rewrite the system (A.19) for $1 \leq j \leq N$ as

$$(A.21) \quad \begin{cases} \frac{\partial f_j}{\partial t} - \omega \cdot \text{grad } f_j = \frac{i}{2a_0} (\square_\varphi + Q) a_0 f_{j-1} \\ f_j(0, x, \omega) = 0. \end{cases}$$

To solve (A.21) we notice that

$$\frac{d}{dt} f_j(t, x - \omega t, \omega) = \left(\frac{\partial f_j}{\partial t} - \omega \cdot \text{grad } f_j \right) (t, x - \omega t, \omega).$$

Hence

$$f_j(t, x - \omega t, \omega) = \frac{i}{2} \int_0^t \left(\frac{\square_\varphi(a_0 f_{j-1})}{a_0} + Q f_{j-1} \right) (\tau, x - \tau \omega, \omega) d\tau$$

or

$$(A.22) \quad f_j(t, x, \omega) = \frac{i}{2} \int_0^t \left(\frac{\square_\varphi(a_0 f_{j-1})}{a_0} + Q f_{j-1} \right) (\tau, x + (t - \tau)\omega, \omega) d\tau.$$

Inductively, one proves that f_j is smooth in all its arguments.

As $f_0 \equiv 1$, formula (A.22) leads for $j = 1$ to the decomposition

$$(A.23) \quad f_1 = f_{1,0} + f_{1,1}$$

of f_1 , where

$$(A.24) \quad f_{1,0}(t, x, \omega) := \frac{i}{2} \int_0^t \frac{\square_\varphi a_0}{a_0} (\tau, x + (t - \tau)\omega, \omega) d\tau$$

and

$$(A.25) \quad f_{1,1}(t, x, \omega) := \frac{i}{2} \int_0^t Q(x + (t - \tau)\omega) d\tau \stackrel{\text{change of variable}}{=} \frac{i}{2} \int_0^t Q(x + \tau\omega) d\tau.$$

Note that $f_{1,0}$ is independent of Q whereas $f_{1,1}$ is linear in Q .

Equations (A.20) and (A.23) yield a decomposition for a_1 :

$$(A.26) \quad a_1 = a_{1,0} + a_{1,1}$$

where for $j = 1, 0$,

$$(A.27) \quad a_{1,j} := a_0 f_{1,j}.$$

Similarly, one has

$$(A.28) \quad f_2 = f_{2,0} + f_{2,1} + f_{2,2}$$

and

$$(A.29) \quad a_2 = a_{2,0} + a_{2,1} + a_{2,2}$$

where $a_{2,j} := a_0 f_{2,j}$ ($\forall j = 0, 1, 2$) and

$$(A.30) \quad f_{2,0}(t, x, \omega) := \frac{i}{2} \int_0^t \left(\frac{\square_\varphi(a_0 f_{1,0})}{a_0} \right) (\tau, x + (t - \tau)\omega, \omega) d\tau,$$

$$(A.31) \quad f_{2,1}(t, x, \omega) = \frac{i}{2} \int_0^t \left(\frac{\square_\varphi(a_0 f_{1,1})}{a_0} + Q f_{1,0} \right) (\tau, x + (t - \tau)\omega, \omega) d\tau$$

and

$$(A.32) \quad f_{2,2}(t, x, \omega) = \frac{i}{2} \int_0^t Qf_{1,1}(\tau, x + (t - \tau)\omega, \omega) d\tau.$$

Substituting (A.25) into (A.32) and changing variables, we obtain
(A.33)

$$\begin{aligned} f_{2,2}(t, x, \omega) &= -\frac{1}{4} \int_0^t \left[Q(x + (t - \tau)\omega) \int_0^\tau Q(x + (t - s)\omega) ds \right] d\tau \\ &= -\frac{1}{4} \int_0^t \frac{1}{2} \frac{d}{d\tau} \left(\int_0^\tau Q(x + (t - s)\omega) ds \right)^2 d\tau \\ &= -\frac{1}{8} \left(\int_0^t Q(x + (t - \tau)\omega) d\tau \right)^2 \\ &= -\frac{1}{8} \left(\int_0^t Q(x + \tau\omega) d\tau \right)^2 \end{aligned}$$

Appendix B. Wave kernel acting on sections of a line bundle

In this appendix we show how to construct the wave kernel acting on sections of the complex line bundle L over the n -dimensional torus $M = \mathcal{L} \backslash \mathbb{R}^n$, $n \geq 2$, from the wave kernel on \mathbb{R}^n constructed in Appendix A.

Recall that sections of $L \rightarrow M$ may be viewed equivalently as functions $f : \mathbb{R}^n \rightarrow \mathbb{C}$ satisfying Equation (2.6) for all $l \in \mathcal{L}$, where e_l is defined as in Notation 2.4.

We introduce the following distributional kernel

$$(B.1) \quad K_L(t, x, y) := \sum_{l \in \mathcal{L}} e^{-2\pi i e_l(x)} K(t, x + l, y),$$

where $K(t, x, y)$ denotes the wave kernel on \mathbb{R}^n introduced in the previous section. For any section u_0 of L , again viewed as a function $u_0 : \mathbb{R}^n \rightarrow \mathbb{C}$ satisfying Equation (2.6), define

$$(B.2) \quad \begin{aligned} u(t, x) &:= \int_{\mathcal{F}} K_L(t, x, y) u_0(y) dy \\ &= \sum_{l \in \mathcal{L}} e^{-2\pi i e_l(x)} \int_{\mathcal{F}} K(t, x + l, y) u_0(y) dy, \end{aligned}$$

where \mathcal{F} denotes a fixed fundamental domain of the action of \mathcal{L} on \mathbb{R}^n . By a standard domain of dependence argument the sum in (B.2) is finite for any given x and t .

We claim that $u(t, x)$ satisfies (A.1) - (A.2) as well as (2.6), i.e., one has the following:

PROPOSITION B.1. — For any $x \in \mathbb{R}^n, t > 0$,

- (i) $(\square_\varphi + Q)(x) \cdot u(t, x) = 0$
- (ii) $u(0, x) = u_0(x); u_t(0, x) = 0$
- (iii) $u(t, x + l) = e^{2\pi i e_l(x)} u(t, x)$ for all $l \in \mathcal{L}$.

Proof. —

- (i) The group \mathcal{L} acts on $C^\infty(\mathbb{R}^+ \times \mathbb{R}^n, \mathbb{C})$ by $(l.v)(t, x) = e^{2\pi i e_l(x)} v(t, x - l)$. By Remark 2.19, this action commutes with Δ_φ and hence with the Schrödinger operator $(\square_\varphi + Q)$. Assertion (i) follows.
- (ii) Letting t approach 0 in $\sum_{l \in \mathcal{L}} e^{-2\pi i e_l(x)} \int_{\mathcal{F}} K(t, x + l, y) u_0(y) dy$, one gets by (B.2) that for any $x \in \mathbb{R}^n$,

$$u(0, x) = \sum_{l \in \mathcal{L}} e^{-2\pi i e_l(x)} \int_{\mathcal{F}} \delta_{x+l}(y) u_0(y) dy = e^{-2\pi i e_{l_x}(x)} u_0(x + l_x)$$

where δ denotes the Dirac delta function and l_x is the unique element in \mathcal{L} such that $x + l_x \in \mathcal{F}$. Since $u_0(x + l_x) = e^{2\pi i e_{l_x}(x)} u_0(x)$, one thus obtains $u(0, x) = u_0(x)$. Moreover, as $\frac{\partial}{\partial t} K|_{t=0} \equiv 0$, one has

$$\frac{\partial u}{\partial t}(0, x) = \sum_{l \in \mathcal{L}} e^{-2\pi i e_l(x)} \int_{\mathcal{F}} \frac{\partial}{\partial t} K|_{t=0}(t, x + l, y) u_0(y) dy = 0.$$

- (iii) This is immediate since u is invariant under the action of \mathcal{L} defined in (i). □

Appendix C. Expansion of the wave trace at singularities

In this appendix, we analyze the singularities of the wave trace W_L . This is a distribution in the variable t which when applied to $f \in C_0^\infty(\mathbb{R})$ is given by

$$(C.1) \quad \langle W_L, f \rangle = \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt f(t) K_L(t, x, x)$$

where \mathcal{F} again denotes a fixed fundamental domain of the action of \mathcal{L} on \mathbb{R}^n invariant under the \mathbb{Z}_2 -action.

Eskin, Ralston and Trubowitz [4] analyzed the singularities of the wave trace in the case of trivial line bundles over tori (i.e., for the Schrödinger operator acting on smooth functions on the torus). We will use the same

approach in our setting. We continue to assume throughout this section that $n \geq 2$.

In view of Proposition B.1, we have

$$(C.2) \quad \langle W_L, f \rangle = \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt f(t) \sum_{l \in \mathcal{L}} e^{-2\pi i e_l(x)} K(t, x + l, x),$$

where by a standard domain of dependence argument, the sum in (C.2) is finite for any $x \in \mathbb{R}^n$ and $t > 0$. Hence

$$(C.3) \quad \langle W_L, f \rangle = \int_{\mathcal{F}} dx \sum_{l \in \mathcal{L}} e^{-2\pi i e_l(x)} \int_{\mathbb{R}} dt f(t) K(t, x + l, x),$$

where the sum is again finite, as f has compact support.

The singular support of the distribution W_L is given by the set $\{|l| : l \in \mathcal{L}\}$. The contributions to the singularity of W_L at $t = |d|$, $d \in \mathcal{L} - \{0\}$, come from

$$(C.4) \quad \int_{\mathcal{F}} dx \sum_{l \in \mathcal{L}, |l|=|d|} e^{-2\pi i e_l(x)} K(t, x + l, x).$$

For $d \in \mathcal{L} \setminus \{0\}$, let $\beta_{|d|}$ be a smooth, compactly supported function on \mathbb{R} such that $\beta_{|d|}(t) = 1$ for t near $|d|$, $\text{supp}(\beta_{|d|}) \subset \mathbb{R}^+$, and $\text{supp}(\beta_{|d|}) \cap |\mathcal{L}| = \{|d|\}$. Then the singular support of the distribution $\beta_{|d|}W_L$ is given by $\{\pm |d|\}$. Hence, by (C.4) and the assumption that $\mathcal{L} \setminus \mathbb{R}^n$ has nondegenerate length spectrum, the Fourier transform $\widehat{\beta_{|d|}W_L}$ of $\beta_{|d|}W_L$ with respect to time t satisfies

$$(C.5) \quad \widehat{\beta_{|d|}W_L}(\eta) \underset{\eta \rightarrow +\infty}{\sim} \sum_{t=\pm d} \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt e^{-it\eta} \beta_{|d|}(t) e^{-2\pi i e_l(x)} K(t, x + l, x) + O(\eta^{-\infty}).$$

DEFINITION C.1. — For $l \in \mathcal{L} \setminus \{0\}$, $x \in \mathbb{R}^n$, $\omega \in S^{n-1}$ and k any nonnegative integer, set

$$(C.6) \quad b_{k,l,\beta}(x, \omega) = \sum_{j=0}^k \binom{n-1-j}{k-j} \left(i \frac{\partial}{\partial t} \right)^{k-j} \Big|_{t=\omega \cdot l} \beta_{|l|}(t) a_j(t, x + l, -\omega)$$

and let

$$(C.7) \quad b_{k,l}(x, \omega) = \sum_{j=0}^k \binom{n-1-j}{k-j} \left(i \frac{\partial}{\partial t} \right)^{k-j} \Big|_{t=\omega \cdot l} a_j(t, x + l, -\omega).$$

(To review the notion of the binomial coefficients $\binom{-m}{r}$ when m and r are positive, see Equation (D.2) in Appendix D.)

In particular,

$$(C.8) \quad b_{0,l}(x, \omega) = a_0(\omega \cdot l, x + l, -\omega)$$

For any integer $R \geq 1$, set

$$(C.9) \quad J_{l,R}(\eta) := \frac{1}{(2\pi)^{n-1}} \sum_{k=0}^{R+n-2} \eta^{n-1-k} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \\ \times \int_{S_+^{n-1}(l)} d_{vol} \omega e^{-i\eta \omega \cdot l} b_{k,l,\beta}(x, \omega)$$

where $d_{vol} \omega$ is the volume form on S^{n-1} and

$$S_+^{n-1}(l) := \{ \omega \in S^{n-1} \mid \omega \cdot l > 0 \}.$$

PROPOSITION C.2. — Assume that $\mathcal{L} \setminus \mathbb{R}^n$ has nondegenerate length spectrum. Then for any $R \geq 1$,

$$\widehat{\beta_{|d|} W_L}(\eta) \stackrel{\eta \rightarrow +\infty}{=} \frac{1}{2} (J_{d,R}(\eta) + J_{-d,R}(\eta)) + O(\eta^{-R})$$

The proof of Proposition C.2 can be found in Appendix D.

To get an asymptotic expansion of $J_{l,R}$ ($l = \pm d$) we use the method of stationary phase. Note that the map

$$\begin{aligned} S_+^{n-1}(l) &\rightarrow \mathbb{R} \\ \omega &\mapsto -\omega \cdot l \end{aligned}$$

has $\omega = \frac{l}{|l|}$ as the unique critical point and that this point is a minimum. Thus the asymptotic expansion as $\eta \rightarrow +\infty$ of the integral $\int_{S_+^{n-1}(l)} e^{-i\eta \omega \cdot l} b_{k,l,\beta}(x, \omega) d_{vol} \omega$ is related to the behavior of $b_{k,l,\beta}(x, \omega)$ with ω restricted to an arbitrarily small neighborhood of this point. Since $\beta_{|l|}(\omega \cdot l)$ is identically one for ω in some neighborhood U of $\frac{l}{|l|}$ in S^{n-1} , we have $b_{k,l}(x, \omega) = b_{k,l,\beta}(x, \omega)$ when $\omega \in U$. Thus in computing the asymptotic expansion, we may work with the functions $b_{k,l}$ rather than $b_{k,l,\beta}$.

We can give explicit Morse coordinates on the open hemisphere $S_+^{n-1}(l)$. First, let (y_1, \dots, y_n) be the standard rectangular coordinates on \mathbb{R}^n , rotated so that the coordinates of $\frac{l}{|l|}$ are $(0, \dots, 0, 1)$. Define coordinates $z = (z_1, \dots, z_{n-1})$ on $S_+^{n-1}(l)$ by setting

$$(z_1, \dots, z_{n-1}) = \sqrt{\frac{2}{1 + y_n}} (y_1, \dots, y_{n-1}).$$

These coordinates map $S_+^{n-1}(l)$ onto a ball of radius $\sqrt{2}$ about the origin in \mathbb{R}^{n-1} and carry $\frac{l}{|l|}$ to 0. We will denote by $\omega(z)$ the point in $S_+^{n-1}(l)$ with coordinates z .

LEMMA C.3. — Let $z = (z_1, \dots, z_{n-1})$ be the coordinate chart on $S_+^{n-1}(l)$ defined above. Then:

- (i) $\omega(z) \cdot l = |l| (1 - |z|^2 / 2)$ where $|z|^2 = z_1^2 + \dots + z_{n-1}^2$;
- (ii) Letting $v(z) dz_1 \wedge \dots \wedge dz_{n-1}$ be the expression for the volume form of S^{n-1} with respect to these coordinates, then $v(0) = 1$.

Lemma C.3 says that $z = (z_1, \dots, z_{n-1})$ are Morse coordinates for the functional $\omega \mapsto -\omega \cdot \frac{l}{|l|}$.

Proof of Lemma C.3. — For $\omega = (y_1, \dots, y_n) \in S_+^{n-1}(l)$, we have $\omega \cdot l = |l| y_n$. Since $y_1^2 + \dots + y_{n-1}^2 = 1 - y_n^2$, we have $|z|^2 = 2(1 - y_n)$ and thus $y_n = 1 - \frac{|z|^2}{2}$. Statement (i) follows.

With respect to the “standard” coordinates (y_1, \dots, y_{n-1}) on S_+^{n-1} , the volume form is given by $\frac{1}{y_n} dy_1 \wedge \dots \wedge dy_{n-1}$. Since $z_j = \sqrt{\frac{2}{1+y_n}} y_j$, we have $dz_j = dy_j$ when $z = 0$ (and $y_n = 1$). Thus (ii) follows. □

Define the functions $\tilde{b}_{k,l}(x, z)$ by

$$(C.10) \quad \tilde{b}_{k,l}(x, z) = b_{k,l}(x, \omega(z))v(z)$$

where $v(z)$ is given as in Lemma C.3(ii). In view of Lemma C.3 (ii), $\tilde{b}_{k,l}(x, 0) = b_{k,l}(x, l/|l|)$ for any k, x .

By the method of stationary phase (see, e.g., [9] pp. 19-24), one then gets for $l = \pm d$ and any given integer $M \geq 1$,

$$\int_{S_+^{n-1}(l)} d_{vol} \omega e^{-i\eta \omega \cdot l} b_{k,l,\beta}(x, \omega) \underset{\eta \rightarrow +\infty}{=} \frac{e^{-i\eta|l|}}{\eta^{\frac{n-1}{2}}} \sum_{s=0}^{M-1} \frac{(2\pi)^{\frac{n-1}{2}} e^{i\frac{\pi}{4}(n-1)}}{s! |l|^{s+\frac{n-1}{2}} (2i)^s} \frac{1}{\eta^s} \Delta_z^s|_{z=0} \tilde{b}_{k,l}(x, z) + O\left(\frac{1}{\eta^{M+\frac{n-1}{2}}}\right),$$

where $\Delta_z := \frac{\partial^2}{\partial z_1^2} + \dots + \frac{\partial^2}{\partial z_{n-1}^2}$. (Here we are using the fact that $b_{k,l,\beta} = b_{k,l}$ on a neighborhood of $t = |l|$.) This leads to

$$J_{l,R}(\eta) \underset{\eta \rightarrow +\infty}{=} \frac{e^{-i\eta|l|}}{(2\pi)^{n-1}} \sum_{k+s=0}^{R-1+\lceil \frac{n}{2} \rceil} \frac{1}{\eta^{k+s-\frac{n-1}{2}}} \times \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \frac{(2\pi)^{\frac{n-1}{2}} e^{i\frac{\pi}{4}(n-1)}}{s! |l|^{s+\frac{n-1}{2}} (2i)^s} \Delta_z^s|_{z=0} \tilde{b}_{k,l}(x, z) + O\left(\frac{1}{\eta^R}\right)$$

where k and s are understood to be nonnegative integers. For $m \geq 0$, define

$$(C.11) \quad C_m(x, l) := \sum_{j=0}^m \frac{1}{j!} \left(\frac{1}{2i|l|}\right)^j e^{-2\pi i e_l(x)} \Delta_z^j|_{z=0} \tilde{b}_{m-j,l}(x, z).$$

By setting $m = k + s$ in the asymptotic expansion above, we obtain:

PROPOSITION C.4. — For $l = \pm d$,

$$J_{l,R}(\eta) \underset{\eta \rightarrow +\infty}{=} \frac{(2\pi)^{-\frac{n+1}{2}} e^{i\frac{\pi}{4}(n-1)}}{|l|^{\frac{n-1}{2}}} e^{-i\eta|l|} \times \sum_{m=0}^{R-1+\lceil \frac{n}{2} \rceil} \frac{1}{\eta^{m-\frac{n-1}{2}}} \int_{\mathcal{F}} C_m(x, l) dx + O\left(\frac{1}{\eta^R}\right).$$

THEOREM C.5. — Assume that \mathcal{L} has nondegenerate length spectrum. Then for any $m \geq 0$ and any $l \in \mathcal{L} \setminus \{0\}$, the expression $C_{m,l}$ given by

$$(C.12) \quad C_{m,l} := \int_{\mathcal{F}} C_m(x, l) dx + \int_{\mathcal{F}} C_m(-x, -l) dx$$

is a spectral invariant of $\Delta_\varphi + Q$.

The theorem follows from Propositions C.2 and C.4 and the fact that the fundamental domain \mathcal{F} was chosen to be \mathbb{Z}_2 -invariant.

PROPOSITION C.6. — For $l \in \mathcal{L}$, we have in the notation of (C.11) that

$$C_1(x, l) = g(x, l) + \frac{i}{2} e^{-2\pi i e_l(x)} a_0 \left(|l|, x + l, -\frac{l}{|l|} \right) \int_0^{|l|} Q \left(x - \tau \frac{l}{|l|} \right) d\tau$$

where $g(x, l)$ is a function independent of Q .

Proof. — By Definition C.8 and Lemma A.1, $b_{0,l}$ is independent of Q . Thus, by (C.11), there exists a function $g_1(x, l)$ which does not depend on Q such that

$$C_1(x, l) = g_1(x, l) + e^{-2\pi i e_l(x)} b_{1,l} \left(x, \frac{l}{|l|} \right).$$

Hence, by Definition C.1 and Equations (A.20), (A.23), (A.24), and (A.25), there exists a function $g(x, l)$ independent of Q such that

$$C_1(x, l) = g(x, l) + \frac{i}{2} e^{-2\pi i e_l(x)} a_0 \left(|l|, x + l, -\frac{l}{|l|} \right) \int_0^{|l|} Q \left(x + l - \tau \frac{l}{|l|} \right) d\tau$$

The proposition follows since Q is \mathcal{L} -periodic. □

We now assume that n is even, say $n = 2m$, and that the Chern class of $L \rightarrow M$ is nondegenerate. By Remark 2.14 and Proposition 2.16, we may express the connection for the bundle $L \rightarrow M$ as a smooth real-valued 1-form φ on \mathbb{R}^n of the form

$$(C.13) \quad \varphi = \varphi^D + \alpha + d^* \mu$$

where φ^D is the connection form of the distinguished connection, α is a harmonic 1-form, and μ is the pullback to \mathbb{R}^n of a smooth 2-form on M .

For any nonzero vector $l \in \mathcal{L}$, write

$$(C.14) \quad \mathcal{G}_l : \mathbb{R}^{2m} \rightarrow \mathbb{C}, x \mapsto \exp \left[-\frac{2\pi i}{|l|} \int_0^{|l|} d^* \mu \Big|_{(x-\tau \frac{l}{|l|})} (l) d\tau \right].$$

LEMMA C.7. — Given φ as in Equation (C.13), define a_0 as in Lemma A.1. For any $x \in \mathbb{R}^{2m}$ and $l \in \mathcal{L}$, we have

$$a_0(|l|, x + l, -l/|l|) = \pm e^{2\pi i(e_x(l) - \alpha(l))} \mathcal{G}_l(x)$$

where the sign is given by $e^{i\pi e_l(l)}$.

Proof. — By Lemma A.1, we have

$$(C.15) \quad a_0 \left(|l|, x + l, -\frac{l}{|l|} \right) = \exp \left[-\frac{2\pi i}{|l|} \int_0^{|l|} \varphi \Big|_{(x+l-\tau \frac{l}{|l|})} (l) d\tau \right].$$

Denote the right hand side of Equation (C.15) by I_φ . By Equation (C.13), we have

$$(C.16) \quad a_0 \left(|l|, x + l, -\frac{l}{|l|} \right) d\tau = I_{\varphi^D} I_\alpha I_{d^* \mu}.$$

Recall that for $w \in \mathbb{R}^n$, we have $\varphi^D|_w(l) = -e_w(l)$; in particular, φ^D depends linearly on w . Thus $\varphi^D|_{(x+l-\tau \frac{l}{|l|})}(l) = -e_x(l) + \left(\frac{\tau - |l|}{|l|^2} \right) e_l(l)$ and

$$\begin{aligned} I_{\varphi^D} &= e^{2\pi i e_x(l)} \exp \left[\frac{-2\pi i e_l(l)}{|l|^2} \int_0^{|l|} (\tau - |l|) d\tau \right] \\ &= e^{2\pi i e_x(l)} e^{\pi i e_l(l)} \\ &= \pm e^{2\pi i e_x(l)}, \end{aligned}$$

where the last equality uses the fact that $e_l(l) \in \mathbb{Z}$ for all $l \in \mathcal{L}$.

Next, the harmonic 1-form α may be viewed as a linear functional on \mathbb{R}^n , and we have

$$I_\alpha = e^{-2\pi i \alpha(l)}.$$

Finally, since μ is the pull-back of a 2-form on M , it is \mathcal{L} -periodic. Hence

$$I_{d^* \mu} = \mathcal{G}_l(x).$$

The lemma thus follows from Equation (C.16). □

Proof of Proposition 4.1. — In the notation of Proposition C.6, let

$$\begin{aligned} \tilde{C}_1(x, l) &= \frac{2}{i} (C_1(x, l) - g(x, l)) \\ &= e^{-2\pi i e_l(x)} a_0 \left(|l|, x + l, -\frac{l}{|l|} \right) \int_0^{|l|} Q \left(x - \tau \frac{l}{|l|} \right) d\tau. \end{aligned}$$

Since $g(x, l)$ is independent of Q , Theorem C.5 implies that $\int_{\mathcal{F}} (\tilde{C}_1(x, l) + \tilde{C}_1(x, l)) dx$ is a spectral invariant of $\Delta_\varphi + Q$. In the setting of Proposition 4.1, $\varphi = \varphi^D + \alpha$. Thus by Lemma C.7,

$$a_0(|l|, x + l, -l/|l|) = \pm e^{2\pi i(e_x(l) - \alpha(l))},$$

and hence $e^{-2\pi i e_l(x)} a_0\left(|l|, x + l, -\frac{l}{|l|}\right) = \pm e^{2\pi i \Omega(x, l)} e^{-2\pi i \alpha \cdot l}$, where we used the fact that $\Omega(x, l) = e_x(l) - e_l(x)$ (cf (2.1)). Thus W_l^α , as defined in Proposition 4.1, is precisely $\int_{\mathcal{F}} \tilde{C}_1(x, l) dx$. The proposition now follows from Theorem C.5. \square

In the remainder of this appendix, we assume that the connection ∇ is \mathbb{Z}_2 -invariant in order to obtain the expression for $C_{2,l}$ used in the proof of Theorem 4.9. While we are especially interested in the distinguished connection, i.e., the unique \mathbb{Z}_2 -invariant connection that is also translation invariant, the results below are stated for any \mathbb{Z}_2 -invariant connection, not necessarily translation invariant.

PROPOSITION C.8. — *Assume that the connection ∇ is \mathbb{Z}_2 -invariant. Then for any $t > 0$, $x \in \mathbb{R}^n$ and $\omega \in S^{n-1}$,*

- (i) $a_0(t, -x, -\omega) = a_0(t, x, \omega)$.
- (ii) *If in addition Q is even, one has for any $j \geq 1$, $f_j(t, -x, -\omega) = f_j(t, x, \omega)$.*

Proof. — Statement (i) follows from formula (A.18) and Definition 2.12. Statement (ii) follows from an induction argument using (A.22), Remark 2.19, and statement (i). \square

THEOREM C.9. — *Assume that \mathcal{L} has nondegenerate length spectrum and that the connection is \mathbb{Z}_2 -invariant. Let Q^- denote the odd part of Q ; i.e., $Q^- = \frac{1}{2}(Q - \tilde{Q})$. Then in the notation of Theorem C.5,*

$$C_{2,l} = -\frac{1}{4} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} a_0\left(|l|, x + l, -\frac{l}{|l|}\right) \left(\int_0^{|l|} d\tau Q^-\left(x - \tau \frac{l}{|l|}\right)\right)^2 + \text{terms which do not involve } Q^-.$$

Proof. — In view of (C.11), one has

$$(C.17) \quad C_2(x, l) = \sum_{j=0}^2 \frac{1}{j!} \left(\frac{1}{2i|l|}\right)^j B_j(x, l)$$

where

$$(C.18) \quad B_j(x, l) := e^{-2\pi i e_l(x)} \Delta_z^j|_{z=0} \tilde{b}_{2-j,l}(x, z).$$

Given any function $F(x, l)$, write $F^+(x, l) = \frac{1}{2} (F(x, l) + F(-x, -l))$. We have by Equation (C.12) that

$$(C.19) \quad C_{2,l} = 2 \int_{\mathcal{F}} C_2^+(x, l) dx = \sum_{j=0}^2 \frac{2}{j!} \left(\frac{1}{2i|l|} \right)^j \int_{\mathcal{F}} B_j^+(x, l) dx.$$

In the following, a function of any or all of the variables $x \in \mathbb{R}^n, l \in \mathcal{L}$ and $\omega \in S^{n-1}$ will be said to be *even* (respectively, *odd*) if it remains unchanged (respectively, is changed into its negative) when all the variables are replaced by their negatives. Note that in this sense, the function $(x, l) \mapsto e^{-2\pi i e_l(x)}$ is even. By Proposition C.8, the function $(x, \omega) \mapsto a_0(t, x, \omega)$ is also even for any fixed $t \in \mathbb{R}$. Hence by Remark 2.19, the map $(x, \omega) \mapsto f_{1,0}(t, x, \omega)$ defined in (A.24) is even as well.

The functions B_j in (C.18) are given as follows:

- $j = 0$: Using (C.7) with $k = 2$ along with (A.26), (A.29), and the fact that $a_{1,0}$ and $a_{2,0}$ are independent of Q (cf. (A.20), (A.24) and (A.30)), we have

$$(C.20) \quad B_0(x, l) = e^{-2\pi i e_l(x)} \left[a_{2,2} \left(|l|, x + l, -\frac{l}{|l|} \right) + a_{2,1} \left(|l|, x + l, -\frac{l}{|l|} \right) + i(n-2) \frac{\partial}{\partial t} \Big|_{t=|l|} a_{1,1} \left(t, x + l, -\frac{l}{|l|} \right) \right] + \text{terms independent of } Q.$$

- $j = 1$: By (C.7), (A.20), (A.23), (A.24), and (A.27),

$$(C.21) \quad B_1(x, l) = e^{-2\pi i e_l(x)} \Delta_z \Big|_{z=0} a_{1,1}(\omega \cdot l, x + l, -\omega(z)) v(z) + \text{terms independent of } Q$$

with $v(z)$ as in Lemma C.3.

- $j = 2$: By (C.8), (A.18), and (C.18), $B_2(x, l)$ is independent of Q .

We will show below that Q^- contributes to $C_{2,l}$ only through the first term in $B_0^+(x, l)$ (the term involving $a_{2,2}$). Assuming this statement for now, we examine the dependence on Q of the first term in the expression for $B_0(x, l)$,

Write $A_{2,2}(x, l) = a_{2,2} \left(|l|, x + l, -\frac{l}{|l|} \right)$ and $A_0(x, l) = a_0 \left(|l|, x + l, -\frac{l}{|l|} \right)$. By (A.18), (A.20) and (A.33) we have

$$(C.22) \quad A_{2,2}(x, l) = -\frac{1}{8} A_0(x, l) \left(\int_0^{|l|} Q \left(x - \tau \frac{l}{|l|} \right) d\tau \right)^2.$$

Since $A_0(x, l) = A_0(-x, -l)$, we have

$$A_{2,2}^+(x, l) = -\frac{1}{8}A_0(x, l) \left(\int_0^{|l|} Q^- \left(x - \tau \frac{l}{|l|} \right) d\tau \right)^2 + \dots$$

where \dots is independent of Q^- . Since $e_l(x) = e_{-l}(-x)$, the theorem then follows from Equations (C.19) and (C.20).

It remains only to show that Q^- does not contribute to $C_{2,l}$ through $B_1^+(x, l)$ nor through the other terms in $B_0^+(x, l)$.

By (A.18), (A.20) and (A.31),

$$\begin{aligned} & a_{2,1}(t, x + l, -\omega) \\ &= \frac{i}{2} a_0(t, x + l, -\omega) \int_0^t \left(\frac{\square_\varphi(a_0 f_{1,1})}{a_0} + Q f_{1,0} \right) (\tau, x + l - (t - \tau)\omega, -\omega) d\tau. \end{aligned}$$

By (A.18) and (A.24), a_0 and $f_{1,0}$ are independent of Q , while by (A.25), $f_{1,1}$ depends linearly on Q . Moreover, for each fixed t , the functions $(x, \omega) \mapsto a_0(t, x, \omega)$ and $(x, \omega) \mapsto f_{1,0}(t, x, \omega)$ are both even. Thus $a_{2,1}(t, x + l, -\omega) + a_{2,1}(t, -x - l, \omega)$ is independent of Q^- . Thus Q^- does not contribute to $B_0^+(x, l)$ through $a_{2,1}$.

We next show that Q^- will not contribute to $C_{2,l}$ through $a_{1,1}$. At this stage, it is simpler to prove the stronger statement that Q^- does not contribute to the whole sum $J_{l,R} + J_{-l,R}$ through $a_{1,1}$. We come back to the expression for $J_{l,R} + J_{-l,R}$ obtained before using the method of stationary phase. By (C.9), for any $l \neq 0$,

$$\begin{aligned} & J_{l,R}(\eta) + J_{-l,R}(\eta) \\ &= \frac{1}{(2\pi)^{n-1}} \sum_{k=0}^{R+n-2} \eta^{n-1-k} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S_+^{n-1}(l)} d_{vol} \omega e^{-i\eta \omega \cdot l} b_{k,l,\beta}(x, \omega) \\ &+ \frac{1}{(2\pi)^{n-1}} \sum_{k=0}^{R+n-2} \eta^{n-1-k} \int_{\mathcal{F}} dx e^{2\pi i e_l(x)} \int_{S_+^{n-1}(-l)} d_{vol} \omega e^{i\eta \omega \cdot l} b_{k,-l,\beta}(x, \omega). \end{aligned}$$

After changing x into $-x$ and ω into $-\omega$ in the second sum and taking into account that \mathcal{F} was chosen to be symmetric, one gets

$$\begin{aligned} \text{(C.23)} \quad J_{l,R}(\eta) + J_{-l,R}(\eta) &= \frac{1}{(2\pi)^{n-1}} \sum_{k=0}^{R+n-2} \eta^{n-1-k} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \\ &\times \int_{S_+^{n-1}(l)} d_{vol} \omega e^{-i\eta \omega \cdot l} [b_{k,l,\beta}(x, \omega) + b_{k,-l,\beta}(-x, -\omega)]. \end{aligned}$$

By (A.25) and the fact that Q is periodic,

$$a_{1,1}(t, x + l, -\omega) = \frac{i}{2} a_0(t, x + l, -\omega) \int_0^t Q(x - \tau\omega) d\tau.$$

Thus

$$a_{1,1}^-(t, x + l, -\omega) = \frac{i}{2} a_0(t, x + l, -\omega) \int_0^t Q^-(x - \tau\omega) d\tau$$

where $a_{1,1}^-(t, x, \omega) := \frac{1}{2} (a_{1,1}(t, x, \omega) - a_{1,1}(t, -x, -\omega))$ for all $t \in \mathbb{R}$, $x \in \mathbb{R}^n$, $\omega \in S^{n-1}$. In view of (C.6) and (C.23), we have to show that for any $s \geq 0$,

$$(x, l, \omega) \mapsto \left. \frac{\partial^s}{\partial t^s} \right|_{t=\omega \cdot l} a_{1,1}^-(t, x + l, -\omega)$$

is odd.

As the map $(x, l, \omega) \mapsto a_0(t, x + l, -\omega)$ is even for any $t \in \mathbb{R}$, it suffices to prove that for any $s \geq 0$, the function $g^{(s)}$ given by

$$g^{(s)}(x, l, \omega) = \left. \frac{\partial^s}{\partial t^s} \right|_{t=\omega \cdot l} \int_0^t Q^-(x - u\omega) du$$

is odd. Write

$$h_{(x,\omega)}(t) = \int_0^t Q^-(x - u\omega) du.$$

Then $h_{(-x,-\omega)}(t) = -h_{(x,\omega)}(t)$ for all t , so all the t -derivatives also satisfy $h_{(-x,-\omega)}^{(s)}(t) = -h_{(x,\omega)}^{(s)}(t)$ for all t . We thus have

$$g^{(s)}(x, l, \omega) = h_{(x,\omega)}^{(s)}(\omega \cdot l) = -h_{(-x,-\omega)}^{(s)}(\omega \cdot l) = -g^{(s)}(-x, -l, -\omega).$$

I.e., $g^{(s)}$ is odd, thus completing the proof of Theorem C.9. □

Appendix D. Proof of Proposition C.2

In this appendix we prove Proposition C.2 stated in Appendix C.

We fix $d \in \mathcal{L}$ and write β for $\beta_{|d|}$.

Recall from Appendix A that $K = \frac{1}{2}(K_+ + K_-)$. We first prove that the contribution of K_- to $\widehat{\beta W_L}(\eta)$ is $O(\eta^{-\infty})$ as $\eta \rightarrow +\infty$. Indeed, as the symbol of $K_-(t, x + l, x)$ is given by $e^{il \cdot \zeta - it|\zeta|} a^-(t, x + l, \zeta)$ (see the beginning of Appendix C), the sum

$$\sum_{l=\pm d} \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt e^{-it\eta} \beta(t) K_-(t, x + l, x) e^{-2\pi i e_l(x)}$$

equals

$$\begin{aligned} & \frac{1}{(2\pi)^n} \sum_{l=\pm d} \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt e^{-it\eta} \beta(t) e^{-2\pi i e_l(x)} \int_{\mathbb{R}^n} d\zeta e^{il \cdot \zeta - it|\zeta|} a^-(t, x + l, \zeta) \\ &= \frac{1}{(2\pi)^n} \sum_{l=\pm d} \int_{\mathcal{F}} dx \int_{\mathbb{R}^n} d\zeta e^{il \cdot \zeta} e^{-2\pi i e_l(x)} \int_{\mathbb{R}} dt \beta(t) a^-(t, x + l, \zeta) e^{-it(\eta + |\zeta|)}, \end{aligned}$$

which, by integrations by parts with respect to t , is $O(\eta^{-\infty})$ when $\eta \rightarrow +\infty$. Thus it suffices to consider K_+ .

By the construction of K_+^N (see Equation (A.4)) and regularity results for the Cauchy problem for the operator $\square_\varphi + Q$, we obtain for any $l \in \mathcal{L}$ the following property: for any $R \in \mathbb{N}$ there exists $N_R \in \mathbb{N}$ such that for any $N \geq N_R$ one can find $g_{R,N} \in C^R(\mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^n)$ with the property that

$$\begin{aligned} & \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt e^{-it\eta} \beta(t) K_+(t, x + l, x) e^{-2\pi i e_l(x)} \\ &= \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt e^{-it\eta} \beta(t) K_+^N(t, x + l, x) e^{-2\pi i e_l(x)} \\ & \quad + \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt e^{-it\eta} \beta(t) g_{R,N}(t, x + l, x) e^{-2\pi i e_l(x)}. \end{aligned}$$

It follows that for any $R \in \mathbb{N}$ there exists $N_R \in \mathbb{N}$ such that for any $N \geq N_R$,

$$\begin{aligned} \widehat{\beta W}_L(\eta) & \stackrel{\eta \rightarrow +\infty}{=} \sum_{l=\pm d} \frac{1}{2} \int_{\mathcal{F}} dx \int_{\mathbb{R}} dt e^{-it\eta} \beta(t) K_+^N(t, x + l, x) e^{-2\pi i e_l(x)} + O(\eta^{-R}) \\ & \stackrel{\eta \rightarrow +\infty}{=} \sum_{l=\pm d} \frac{1}{2} \frac{1}{(2\pi)^n} \int_{\mathcal{F}} dx \int_{\mathbb{R}^n} d\zeta \int_{\mathbb{R}} dt e^{-it\eta} e^{-2\pi i e_l(x)} e^{il \cdot \zeta + it|\zeta|} \\ & \quad \times \sum_{j=0}^N \beta(t) a_j^+(t, x + l, \zeta) \chi(|\zeta|) + O(\eta^{-R}) \end{aligned}$$

Hence

$$(D.1) \quad \widehat{\beta W}_L(\eta) \stackrel{\eta \rightarrow +\infty}{=} \frac{1}{2} (I_d(\eta) + I_{-d}(\eta)) + O(\eta^{-R})$$

where, using Notation (A.16) and the fact that $a_j^+(t, x, \zeta)$ is positive-homogeneous of degree $-j$ in ζ , $I_l(\eta)$ is given by

$$\begin{aligned} & \frac{1}{(2\pi)^n} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S^{n-1}} d_{vol} \omega \int_{\mathbb{R}_+} d\rho \rho^{n-1} \chi(\rho) e^{i\rho\omega \cdot l} \\ & \quad \times \int_{\mathbb{R}} dt e^{-it(\eta - \rho)} \beta(t) \sum_{j=0}^N \frac{a_j(t, x + l, \omega)}{\rho^j}. \end{aligned}$$

with $N \geq N_R$. Denoting by $\widehat{\beta a_j}$ the Fourier transform of βa_j with respect to the variable t , we then obtain for $l = \pm d$,

$$I_l(\eta) = \frac{1}{(2\pi)^n} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S^{n-1}} d_{vol} \omega \times \sum_{j=0}^N \int_{\mathbb{R}} d\rho \rho^{n-1-j} \chi(\rho) e^{i\rho\omega \cdot l} \widehat{\beta a_j}(\eta - \rho, x + l, \omega)$$

where we used the fact that $\chi(\rho) = 0$ for $\rho \leq \frac{1}{2}$ (see (A.6)). Setting $\tau := \eta - \rho$, we have

$$I_l(\eta) = \frac{1}{(2\pi)^n} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S^{n-1}} d_{vol} \omega e^{i\eta\omega \cdot l} \times \sum_{j=0}^N \int_{\mathbb{R}} d\tau e^{-i\tau\omega \cdot l} (\eta - \tau)^{n-1-j} \chi(\eta - \tau) \widehat{\beta a_j}(\tau, x + l, \omega).$$

Note that the integral with respect to τ in the latter expression is actually an integral over the interval $] -\infty, \eta - 1/2]$, as $\text{supp}(\chi) \subset [1/2, +\infty[$.

In order to expand the expression $(\eta - \tau)^{n-1-j}$, recall that the binomial coefficients $\binom{-m}{k}$ are defined when m and k are positive by

$$(D.2) \quad \binom{-m}{k} = \frac{-m(-m-1)\dots(-m-k+1)}{k!} = (-1)^k \binom{m+k-1}{k}$$

and that $\binom{-m}{0} = 1$. For p a negative integer, the power series expansion for $(a + b)^p = a^p(1 + \frac{b}{a})^p$ is then given by the binomial expansion

$$(D.3) \quad (a + b)^p = \sum_{k=0}^{\infty} \binom{p}{k} a^{p-k} b^k.$$

Interpreting $\binom{p}{k}$ to be zero when p is a nonnegative integer and $k > p$, then the expansion (D.3) is valid for all integers p .

Since $\eta \pm \tau$ is the symbol of the differential operator $\eta \pm i \frac{d}{dt}$, we define the operator $(\eta \pm i \frac{d}{dt})^p$ by

$$(D.4) \quad \left(\eta \pm i \frac{d}{dt}\right)^p = \sum_{k=0}^{\infty} \binom{p}{k} \eta^{p-k} \left(\pm i \frac{d}{dt}\right)^k.$$

We then have the following expression for $I_l(\eta)$:

$$(D.5) \quad \frac{1}{(2\pi)^n} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S^{n-1}} d_{vol} \omega e^{i\eta \omega \cdot l} \\ \times \sum_{j=0}^N \left(\eta - i \frac{d}{dt} \right)^{n-1-j} \Big|_{t=\omega \cdot l} \int_{\mathbb{R}} d\tau e^{-i\tau t} \chi(\eta - \tau) \widehat{\beta a_j}(\tau, x + l, \omega).$$

For $f \in C_0^\infty(\mathbb{R})$, Equation (D.4) implies that

$$(D.6) \quad \left(\eta \pm i \frac{d}{dt} \right)^{n-1-j} (f) \\ = \sum_{r=0}^{R+n-2-j} \binom{n-1-j}{r} \eta^{n-1-j-r} \left(\pm i \frac{d}{dt} \right)^r (f) + O(\eta^{-R})$$

when $\eta \rightarrow +\infty$.

As $\beta \in C_0^\infty(\mathbb{R})$, the Fourier transform of $\beta(\cdot) a_j(\cdot, x + l, \omega)$ is in $\mathcal{S}(\mathbb{R})$ for any $x \in \mathbb{R}^n$, $l \in \mathcal{L}$ and $\omega \in S^{n-1}$. It follows that for any integer $k \geq 1$ there exists a constant $C = C(k, l) > 0$ such that for any $\tau \in \mathbb{R}$, $x \in \mathcal{F}$, and $\omega \in S^{n-1}$,

$$|\tau|^{k-1} \left| \widehat{\beta a_j}(\tau, x + l, \omega) \right| \leq C.$$

Thus for any $r \in \mathbb{N}$ and $l = \pm d$, it follows from the fact that $\chi(s) = 1$ for $s \geq 1$ (see A.6) that

$$\frac{d^r}{dt^r} \Big|_{t=\omega \cdot l} \int_{\mathbb{R}} d\tau e^{-i\tau t} (1 - \chi(\eta - \tau)) \widehat{\beta a_j}(\tau, x + l, \omega) \\ \stackrel{\text{definition of } \chi}{=} \frac{d^r}{dt^r} \Big|_{t=\omega \cdot l} \int_{\eta-1}^{+\infty} d\tau e^{-i\tau t} (1 - \chi(\eta - \tau)) \widehat{\beta a_j}(\tau, x + l, \omega) \\ \stackrel{\eta \rightarrow \pm\infty}{=} O(\eta^{-\infty})$$

uniformly in $x \in \mathcal{F}$ and $\omega \in S^{n-1}$. Hence, in view of (D.5), $I_l(\eta)$ admits for $\eta \rightarrow +\infty$ the asymptotic expansion

$$\frac{1}{(2\pi)^n} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S^{n-1}} d_{vol} \omega e^{i\eta \omega \cdot l} \\ \times \sum_{j=0}^N \left(\eta - i \frac{d}{dt} \right)^{n-1-j} \Big|_{t=\omega \cdot l} \int_{\mathbb{R}} d\tau e^{-i\tau t} \widehat{\beta a_j}(\tau, x + l, \omega) + O(\eta^{-\infty}).$$

The integral $\frac{1}{2\pi} \int_{\mathbb{R}} d\tau e^{i\tau t} \widehat{\beta a_j}(\tau, x+l, \omega)$ is the inverse Fourier transform of $\widehat{\beta a_j}$. Hence for $\eta \rightarrow +\infty$, $I_l(\eta)$ equals

$$\frac{1}{(2\pi)^{n-1}} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S^{n-1}} d_{vol} \omega e^{i\eta \omega \cdot l} \times \sum_{j=0}^N \left(\eta - i \frac{d}{dt} \right)^{n-1-j} \Big|_{t=\omega \cdot l} \beta(-t) a_j(-t, x+l, \omega) + O(\eta^{-\infty})$$

or, changing t into $-t$ and ω into $-\omega$, this equals

$$\frac{1}{(2\pi)^{n-1}} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S_+^{n-1}(l)} d_{vol} \omega e^{-i\eta \omega \cdot l} \times \sum_{j=0}^N \left(\eta + i \frac{d}{dt} \right)^{n-1-j} \Big|_{t=\omega \cdot l} \beta(t) a_j(t, x+l, -\omega) + O(\eta^{-\infty})$$

(we integrate on $S_+^{n-1}(l)$ instead of S^{n-1} because $\text{supp}(\beta) \subset \mathbb{R}_+$).

In Equation (D.6), make the change of index $k = j + r$ to obtain

$$\left(\eta \pm i \frac{d}{dt} \right)^{n-1-j} (f) = \sum_{k=j}^{R+n-2} \binom{n-1-j}{k-j} \eta^{n-1-k} \left(\pm i \frac{d}{dt} \right)^{k-j} (f) + O(\eta^{-R}).$$

Denote by g_j the function $t \mapsto \beta(t) a_j(t, x+l, -\omega)$ (with x, l and ω being fixed). By interchanging the order of summation, we have for any integer R such that $N \geq R + n - 2$,

$$\sum_{j=0}^N \left(\eta + i \frac{d}{dt} \right)^{n-1-j} \Big|_{t=\omega \cdot l} g_j(t) \stackrel{\eta \rightarrow +\infty}{=} \sum_{k=0}^{R+n-2} \sum_{j=0}^k \binom{n-1-j}{k-j} \eta^{n-1-k} \left(i \frac{d}{dt} \right)^{k-j} \Big|_{t=\omega \cdot l} g_j(t) + O(\eta^{-R})$$

where the error term is uniform in $x \in \mathcal{F}$ and $\omega \in S^{n-1}$. Thus

$$(D.7) \quad I_l(\eta) \stackrel{\eta \rightarrow +\infty}{=} \frac{1}{(2\pi)^{n-1}} \int_{\mathcal{F}} dx e^{-2\pi i e_l(x)} \int_{S_+^{n-1}(l)} d_{vol} \omega e^{-i\eta \omega \cdot l} \times \sum_{k=0}^{R+n-2} b_{k,l,\beta}(x, \omega) \eta^{n-1-k} + O(\eta^{-R}),$$

where $b_{k,l,\beta}(x, \omega)$ is given in Definition C.1. Proposition C.2 now follows from Equations (D.1) and (D.7).

BIBLIOGRAPHY

- [1] J. DUISTERMAAT & V. GUILLEMIN, “The spectrum of positive elliptic operators and periodic bicharacteristics”, *Invent. Math.* **29** (1975), p. 39-79.
- [2] J. L. DUPONT, *Curvature and Characteristic Classes*, Lecture Notes in Mathematics, vol. 640, Springer-Verlag, 1968.
- [3] G. ESKIN, “Inverse spectral problem for the Schrödinger equation with periodic vector potential”, *Commun. Math. Phys.* **125** (1989), no. 2, p. 263-300.
- [4] G. ESKIN, J. RALSTON & E. TRUBOWITZ, “On Isospectral Periodic Potentials in \mathbb{R}^n , I, II.”, *Comm. in Pure and Appl. Math.* **37** (1984), p. 647-676, 715-753.
- [5] P. B. GILKEY, “Recursion relations and the asymptotic behavior of the eigenvalues of the Laplacian”, *Comp. Math.* **38** (1978), no. 2, p. 201-240.
- [6] C. GORDON & T. KAPPELER, “On isospectral potentials on tori”, *Duke Math. J.* **63** (1991), no. 1, p. 217-233, Part II, *Comm. in PDE* 20 (1995), p. 709-728.
- [7] C. GORDON & E. N. WILSON, “The spectrum of the Laplacian on Riemannian Heisenberg manifolds”, *Mich. Math. J.* **33** (1986), p. 253-271.
- [8] P. GRIFFITHS & J. HARRIS, *Principles of Algebraic Geometry*, John Wiley & Sons, 1978.
- [9] A. GRIGIS & J. SJÖSTRAND, *Microlocal Analysis for Differential Operators*, Lecture Note Series, vol. 196, London Mathematical Society, Cambridge University Press, 1994.
- [10] V. GUILLEMIN, “Inverse spectral results on two-dimensional tori”, *J. Am. Math. Soc.* **3** (1990), no. 2, p. 375-387.
- [11] T. KAPPELER, “On isospectral periodic potentials on a discrete lattice II.”, *Adv. in Appl. Math.* **9** (1988), p. 428-438.
- [12] P. KUCHMENT, *Floquet theory for partial differential equations*, Birkhäuser, 1993.
- [13] P. LAX, “Asymptotic solutions of oscillatory initial value problems”, *Duke Math. J.* **24** (1957), p. 624-646.
- [14] R. S. PALAIS & T. E. STEWART, “Torus bundles over a torus”, *Proc. Amer. Math. Soc.* **12** (1961), p. 26-29.
- [15] J. ROE, *Elliptic operators, topology, and asymptotic methods*, Research Notes in Mathematics, vol. 395, Pitman, Addison Wesley Longman, 1998.

Manuscrit reçu le 23 juin 2006,
 accepté le 30 novembre 2006.

Carolyn S. GORDON
 Dartmouth College
 Department of Mathematics
 6188 Kemeny Hall
 Hanover, NH 03755-3551 (USA)
 carolyn.s.gordon@dartmouth.edu

Pierre GUERINI
 CPGE Dumont d'Urville
 83056 Toulon cedex (France)
 pguerini@ac-nice.fr

Thomas KAPPELER
 Institut für Mathematik
 Universität Zürich-Irchel
 Winterthurerstrasse 190

8057 Zürich (Switzerland)
tk@math.unizh.ch

David L. WEBB
Dartmouth College
Department of Mathematics
6188 Kemeny Hall
Hanover, NH 03755-3551 (USA)
david.l.webb@dartmouth.edu