



ANNALES

DE

L'INSTITUT FOURIER

Gady KOZMA & Alexander OLEVSKIĪ

Singular distributions, dimension of support, and symmetry of Fourier transform

Tome 63, n° 4 (2013), p. 1205-1226.

http://aif.cedram.org/item?id=AIF_2013__63_4_1205_0

© Association des Annales de l'institut Fourier, 2013, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

SINGULAR DISTRIBUTIONS, DIMENSION OF SUPPORT, AND SYMMETRY OF FOURIER TRANSFORM

by Gady KOZMA & Alexander OLEVSKIĪ

ABSTRACT. — We study the “Fourier symmetry” of measures and distributions on the circle, in relation with the size of their supports. The main results of this paper are:

- (i) A one-side extension of Frostman’s theorem, which connects the rate of decay of Fourier transform of a distribution with the Hausdorff dimension of its support;
- (ii) A construction of compacts of “critical” size, which support distributions (even pseudo-functions) with anti-analytic part belonging to l^2 .

We also give examples of non-symmetry which may occur for measures with “small” support. A number of open questions are stated.

RÉSUMÉ. — On étudie la “symétrie de Fourier” des mesures et des distributions sur le cercle en rapport avec la dimension de leurs supports. Les résultats essentiels du présent travail sont les suivants :

- (i) L’extension unilatérale du théorème de Frostman qui met en rapport la vitesse de décroissance de la transformation de Fourier d’une distribution et la dimension de Hausdorff de son support.
- (ii) La construction des compacts d’une taille “critique” qui peut supporter des distributions (voire des pseudo-fonctions) avec une partie anti-analytique appartenant à l^2 .

On donne également quelques exemples de l’asymétrie qui peut se produire pour des mesures à “petit” support. Plusieurs questions ouvertes sont formulées.

I

Let K be a compact subset of the circle group \mathbb{T} . Frostman’s theorem allows to characterize the Hausdorff dimension of K by examining (non-trivial) measures supported on K . The most common version states that

$$\dim K = \sup \left\{ \alpha : \exists \mu \text{ supported on } K \text{ with } \iint \frac{d\mu(x) d\mu(y)}{|x - y|^\alpha} < \infty \right\}.$$

Keywords: Hausdorff dimension, Frostman’s theorem, Fourier symmetry.

Math. classification: 42A63, 42A50, 42A20, 28A80.

It is not difficult to translate this theorem to the language of Fourier coefficients. One then gets

$$\dim K = \sup \left\{ \alpha : \exists \mu \text{ supported on } K \text{ with } \sum_{n=-\infty}^{\infty} \frac{|\widehat{\mu}(n)|^2}{|n|^{1-\alpha} + 1} < \infty \right\}. \tag{1.1}$$

Beurling showed that one may replace measures with arbitrary (Schwartz) distributions. The following version of Frostman’s theorem thus holds ([2] or [7, p. 40]),

THEOREM A. — *Let $0 < \alpha \leq 1$. Then*

(i) *If there is a (non-trivial) distribution S supported on K such that*

$$\sum_{n=-\infty}^{\infty} \frac{|\widehat{S}(n)|^2}{|n|^{1-\alpha} + 1} < \infty, \tag{1.2}$$

then $\dim K \geq \alpha$.

(ii) *If $\dim K > \alpha$ then there is probability measure S , satisfying (1.2).*

We will show that a one-side estimate of \widehat{S} is already sufficient in (i).

THEOREM 1.1. — *Suppose that there is a non-trivial Schwartz distribution S supported on K with*

$$\sum_{n=-\infty}^{-1} \frac{|\widehat{S}(n)|^2}{|n|^{1-\alpha}} < \infty. \tag{1.3}$$

Then $\dim K \geq \alpha$.

We denote by \mathcal{D}^* the space of Schwartz distributions, and by $\mathcal{D}^*(K)$ the space of Schwartz distributions supported on some compact K .

COROLLARY. — *If there is an $S \in \mathcal{D}^*(K)$ with $\widehat{S}(n) = O(|n|^{-\alpha/2})$, $n < 0$ then $\dim K \geq \alpha$.*

Somewhat reminiscent results have to do with symmetry properties of individual measures. For example, a theorem of Rajchman ([19] or [10, §1.4]) states that for any given complex measure μ , if $\lim_{n \rightarrow \infty} \widehat{\mu}(n) = 0$ then it follows that $\lim_{n \rightarrow -\infty} \widehat{\mu}(n) = 0$. See [14] for an interesting generalization. Another interesting result is that of Hruščev and Peller [5, Corollary 3.26], which state that for any complex measure μ , if

$$\sum_{n=1}^{\infty} \frac{|\widehat{\mu}(n)|^2}{n} < \infty$$

then also

$$\sum_{n=-\infty}^{-1} \frac{|\widehat{\mu}(n)|^2}{|n|} < \infty.$$

Such a result cannot hold with $|n|$ replaced by $|n|^{1-\alpha}$ for some $\alpha > 0$. Indeed,

$$\sum_{n=1}^{\infty} \frac{e^{int}}{n^{1-\alpha}} \in L^1 \quad \forall \alpha > 0$$

(see e.g. [21, chap. V] where the cos and sin terms are handled independently). Hence one can construct L^1 examples with any desired polynomial tail decay on the left and independently on the right.

Question. Does this symmetry result still hold if n is replaced by $n \log n$?

It is occasionally useful to have the following terminology

DEFINITION. — For a Schwartz distribution S we define its analytic part as the distribution S' satisfying

$$\widehat{S}'(n) = \begin{cases} \widehat{S}(n) & n \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$

The anti-analytic part is $S - S'$.

Proof of theorem 1.1. Let us first reduce the general case to the case that $\alpha = 1$. Suppose by contradiction that

$$\dim K < \alpha. \tag{1.4}$$

By convolution with an appropriate measure we want to prepare a new distribution, S' supported by a compact of dimension < 1 such that the anti-analytic part of S' would belong to $L^2(\mathbb{T})$. This will lead to a contradiction. The measure will be taken from the following result of Salem:

LEMMA 1.2. — For every δ , $0 < \delta < 1$ there is a compact set $E \subset \mathbb{T}$ such that

- (i) $\dim_{\text{Mink}}(E) \leq \delta$;
- (ii) For every $\beta < \delta$ there exists a positive measure μ_β supported on E such that $|\widehat{\mu}_\beta(n)| \leq C|n|^{-\beta/2}$.

This result is essentially due to Salem but all references we could find were for the usual Hausdorff dimension while we need the Minkowski dimension. We recall that \dim_{Mink} , the Minkowski dimension (also known as

the upper box dimension) is defined by

$$\dim_{\text{Mink}}(E) = \overline{\lim}_{\epsilon \rightarrow 0} \frac{\log \text{Cov}(E; [0, \epsilon])}{\log 1/\epsilon}$$

where $\text{Cov}(E; A)$ is the cover number of the set E by the set A i.e. the minimal number of translates of A required to cover E . Directly from the definition we see that $\dim_{\text{Mink}}(E) \geq \dim_{\text{Haus}} E$. Of course, by Frostman’s theorem clause (ii) implies that $\dim_{\text{Haus}} \geq \delta$ so in fact the set E of lemma 1.2 satisfies $\dim_{\text{Haus}}(E) = \dim_{\text{Mink}}(E) = \delta$. A common way to refer to Frostman and Salem’s results in the literature is via the notion of the Fourier dimension [6, §17]:

DEFINITION. — For a compact set $K \subset \mathbb{T}$ the supremum of numbers a such that K supports a positive (non-zero) measure μ with $|\widehat{\mu}(u)|^2 \leq C/|u|^a$ is known as the Fourier dimension of K . We denote it by $\dim_{\text{F}}(K)$.

With this definition we have the following succinct statements

Frostman’s theorem: $\dim_{\text{F}}(E) \leq \dim_{\text{Haus}}(E)$.

Salem’s theorem: For every $0 < \delta < 1$ there exists some E with

$$\dim_{\text{F}}(E) = \dim_{\text{Haus}}(E) = \delta.$$

Lemma 1.2: For every $0 < \delta < 1$ there exists some E with

$$\dim_{\text{F}}(E) = \dim_{\text{Haus}}(E) = \dim_{\text{Mink}}(E) = \delta.$$

Proof of lemma 1.2. — By [6, chap. 17, thm. 4] the image of any set of dimension $\delta/2$ by a Brownian bridge is a Salem set. Recall that a Brownian bridge is $B = W + L$ where W is usual Brownian motion while L is a linear term that makes B continuous on the circle (see e.g. [6, §16.3 & §17.5] where the Brownian bridge is called “the Wiener function”). Thus it remains to show that the Brownian image of, for example, a Cantor-like set has the required Minkowski dimension, almost surely. Now, our Cantor set K can be covered by intervals I_1, \dots, I_N (N being a power of 2) of length $N^{-2/\delta}$. By definition, the image of each I_i by Brownian motion is an interval, whose length has expectation $\leq CN^{-1/\delta}$. Let X_i be the number of intervals of length $N^{-1/\delta}$ required to cover WI_i . Then X_i are i.i.d. random variables with expectation some constant C . We get

$$\mathbb{E} \text{Cov} \left(\bigcup_{i=1}^N WI_i; [0, N^{-1/\delta}] \right) \leq \sum_i \mathbb{E} X_i = NC$$

The linear term L is negligible in comparison: if $\text{Cov}(BI_i; [0, N^{-1/\delta}]) > \text{Cov}(WI_i; [0, N^{-1/\delta}]) + 1$ then this means that the derivative of the linear

term must be $> N^{-1/\delta}$ which can only happen for a finite number of N . Hence for all N sufficiently large

$$\mathbb{E} \operatorname{Cov}(BK; [0, N^{-1/\delta}]) \leq \mathbb{E} \operatorname{Cov}\left(\bigcup_{i=1}^N BI_i; [0, N^{-1/\delta}]\right) \leq NC$$

and by Tchebyshev’s inequality, for any $\eta > 0$,

$$\mathbb{P}(\operatorname{Cov}(BK; [0, N^{-1/\delta}]) > N^{1+\eta}) \leq CN^{-\eta}.$$

Summing this over $N = 2^n$ for all n we get that with probability 1, for all N sufficiently large $\operatorname{Cov}(BK; [0, N^{-1/\delta}]) \leq N^{1+\eta}$. By the monotonicity of the covering numbers we can move from a specific sequence of ϵ , $\epsilon = 2^{-n/\delta}$ to a general ϵ and lose only a constant. Hence

$$\operatorname{Cov}(BK; [0, \epsilon]) \leq C\epsilon^{-\delta-\eta\delta}$$

So $\dim_{\text{Mink}}(BK) \leq \delta + \eta\delta$. Since $\eta > 0$ was arbitrary, the lemma is proved. \square

We remark that we use Kahane’s approach to the construction of a Salem set for convenience only. In fact one may check that Salem’s original approach ([7, chap. VIII] or [17, §3]) as well as Kaufman’s [9] also give sets with the correct Minkowski dimension.

We need to consider the Minkowski dimension due to the following lemma

LEMMA 1.3. — *For any compacts K, E ,*

$$\dim_{\text{Haus}}(K + E) \leq \dim_{\text{Haus}} K + \dim_{\text{Mink}}(E).$$

(The Minkowski dimension on the right hand side can not be replaced by the Hausdorff one, see e.g. [4, example 7.8]).

Proof. — By [16, §8.10 (4)],

$$\dim_{\text{Haus}}(K \times E) \leq \dim_{\text{Haus}}(K) + \dim_{\text{Mink}}(E)$$

and the set $K + E$ is a Lipschitz map of $K \times E$. \square

Returning to the proof of Theorem 1.1 take

$$1 - \dim K > \delta > 1 - \alpha \tag{1.5}$$

and find E and μ from lemma 1.2 with $\beta_{\text{lemma 1.2}} = 1 - \alpha$. What we would like to do next is to define a distribution T by $T = S * \mu$. However, we have to consider the unlikely situation that this convolution is 0. Definitely, though, we may find some m such that $T = S * (e^{imt}\mu)$ is not zero and we define T this way. $e^{imt}\mu$ still satisfies the crucial property of μ i.e. $|\widehat{e^{imt}\mu}(n)| \leq C|n|^{(\alpha-1)/2}$. We get

$$T \in \mathcal{D}^*(K') \text{ where } K' := K + E.$$

Due to (1.5) and lemma 1.3 we get:

$$\dim K' < 1.$$

(when we do not state which dimension it is, we mean the Hausdorff dimension). On the other hand,

$$\sum_{n < 0} |\widehat{T}(n)|^2 \leq \sum_{n < 0} |\widehat{S}(n)|^2 \cdot C|n|^{\alpha-1} < \infty.$$

In other words we have reduced the problem to the case that $\alpha = 1$. Hence the following lemma concludes the proof:

LEMMA 1.4. — *Let K be a compact with the Hausdorff measure $\Lambda_{x \log 1/x}(K) = 0$. In particular, this is satisfied if $\dim K < 1$. Let $S \in \mathcal{D}^*(K)$. If $\widehat{S}(n) \in l^2(\mathbb{Z}^-)$ then $S = 0$.*

(the notation $\widehat{S}(n) \in l^2(\mathbb{Z}^-)$ is short for $\sum_{n < 0} |\widehat{S}(n)|^2 < \infty$). We remind the reader that the Hausdorff measure Λ_h for an increasing function h with $h(0) = 0$ is the natural generalization of the usual α -Hausdorff measure ones gets by examining $\sum h(\text{diam } B_i)$ for a covering of a set by balls B_i . See e.g. [6, §10.2, remark 4].

Lemma 1.4 is a consequence of results by Dahlberg [3] and Berman [1]. For the convenience of the reader, let us quote the relevant result from [1] (almost) literally and then show how it implies lemma 1.4.

THEOREM B. — *Let $\omega : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ satisfy that $\omega(x)/x$ is decreasing and $\omega'(0) = \infty$. Let F be an analytic function on the disk $\mathbb{D} = \{|z| < 1\}$ satisfying*

$$\log |F(z)| \leq C \frac{\omega(1 - |z|)}{1 - |z|}. \tag{1.6}$$

Let $K \subset \partial\mathbb{D}$ be a set satisfying.

- (a) $\Lambda_\omega(K) = 0$
- (b) For any $\zeta \in \partial\mathbb{D} \setminus K$ there exists a path ρ in \mathbb{D} ending in ζ such that F is bounded on ρ .

Assume $g \in L^1(\mathbb{T})$ and

$$\liminf_{r \rightarrow 1} \log |F(re^{it})| \leq g(t) \quad \text{a.e.} \tag{1.7}$$

Then $\log |F(z)| \leq G(z)$ for all $z \in \mathbb{D}$, where $G(z)$ is the harmonic extension of g to \mathbb{D} .

See [1], the corollary to theorem 5 on page 479. Actually, the theorem there is more general: all conditions are necessary only on an arc γ (except for a very mild global growth condition) and the result is that $\log |F| \leq G$ in a neighborhood of γ .

Proof of lemma 1.4. — Examine the analytic function

$$F(z) = \sum_{n=0}^{\infty} \widehat{S}(n)z^n.$$

We need two properties of F :

- (i) For every $u \notin K$, the limit $\lim_{z \rightarrow e^{iu}} F(z)$ exists and is finite.
- (ii) The (a.e. defined) limit function $f(u) = \lim_{z \rightarrow e^{iu}} F(z)$ is in $L^2(\mathbb{T})$.

We remark on (i) that we need much less — we only need that the limit exists along some path ρ , e.g. radially. But as we will see, the function F is continuous in a neighborhood of u .

To show property (i) write, for $z \in \mathbb{D}$

$$F(z) = \sum_{n=0}^{\infty} z^n \int S(t)e^{-int} dt = \int S(t)Q_z(t) dt$$

where $Q_z = \sum_0^{\infty} z^n e^{-int} = \frac{1}{1 - ze^{-it}}$ (this is the kernel of the Cauchy transform, which is also the sum of the Poisson kernel and the conjugate Poisson kernel, but this is not important at this point). Note that because S is a distribution, the notation $S(t)$ is slightly misleading — the integral $\int S(t)Q_z(t) dt$ though is well defined. Let ψ be a C^∞ function supported on $[u - \epsilon, u + \epsilon]$ with $\psi \equiv 1$ on $[u - \frac{1}{2}\epsilon, u + \frac{1}{2}\epsilon]$, where $\epsilon = \frac{1}{2} \text{dist}(u, K)$. Then

$$\int S(t)Q_z(t)\psi(t) dt = 0$$

because ψ is 0 on a neighborhood of K and S is supported on K . On the other hand,

$$\lim_{z \rightarrow e^{iu}} \int S(t)Q_z(t)(1 - \psi(t)) dt = \int S(t)(Q_{e^{iu}}(t)(1 - \psi(t))) dt$$

since after multiplication by $1 - \psi(t)$ we get that $Q_z(1 - \psi(t))$ is smooth and its derivatives depend continuously on z all the way up to e^{iu} . This shows that the limit function f exists and is finite outside K i.e. proves property (i).

For property (ii) examine the usual Poisson kernel

$$P_z(t) = \sum_{n=0}^{\infty} z^n e^{-int} + \sum_{n=-\infty}^{-1} \bar{z}^n e^{int} = \frac{1 - |z|^2}{1 - 2|z| \cos(t - \arg z) + |z|^2}.$$

Any distribution is Abel summable to 0 outside its support [7, Proposition 2, appendix I, p. 162] which means that for any $u \notin K$,

$$\lim_{r \rightarrow 1} \int S(t)P_{re^{iu}}(t) = 0.$$

and hence

$$f(u) = \lim_{r \rightarrow 1} \int S(t) (Q_{re^{iu}}(t) - P_{re^{iu}}(t)) dt.$$

However, $Q - P = -\sum_1^\infty \bar{z}^n e^{int}$ so we get

$$f(u) = \lim_{r \rightarrow 1} \sum_{n=1}^\infty -\widehat{S}(-n) (re^{-iu})^n$$

But $\sum_1^\infty |\widehat{S}(-n)|^2 < \infty!$ We get that $f \in L^2$, showing property (ii).

We now apply Berman's theorem. Let $\omega(x) = x \log(2/x)$. We first note that F satisfies (1.6) because

$$\log |F(z)| = \log \left(\sum_{n=0}^\infty \widehat{S}(n) z^n \right) \leq \log \left(\sum_{n=0}^\infty (C + n^C) |z|^n \right) \leq C \log \left(\frac{1}{1 - |z|} \right)$$

(the middle inequality holds because any Schwarz distribution S has that \widehat{S} grows no faster than polynomially). Next, K satisfies (a) by assumption and (b) from property (i) which shows that we may take ρ to be simply the radius. Define now

$$g(t) = \max\{0, \log |f(t)|\}.$$

g is in L^1 because $f \in L^2$ — we cannot control $(\log f)^-$ but $(\log f)^+$ is definitely well-behaved. We get (1.7) again from the continuity of F . Hence Berman's theorem holds and we get $\log |F| \leq G$, where G is as in Berman's theorem namely the harmonic extension of g to \mathbb{D} . To estimate G we note that by definition $G(z) = \int P_z(t) g(t) dt$, and hence by Jensen's inequality,

$$e^{2G(z)} = \exp \left(\int_{\mathbb{T}} P_z(t) 2g(t) dt \right) \leq \int_{\mathbb{T}} P_z(t) e^{2g(t)} dt. \tag{1.8}$$

This observation gives that F is in the Hardy space H^2 because

$$\begin{aligned} \int_{|z|=r} |F(z)|^2 d|z| &\leq \int_{|z|=r} e^{2G(z)} d|z| \stackrel{(1.8)}{\leq} \int_{|z|=r} \int_{\mathbb{T}} P_z(t) e^{2g(t)} dt d|z| = \\ &= \int_{\mathbb{T}} e^{2g(t)} dt \leq 2\pi + \int_{\mathbb{T}} |f(t)|^2 dt < \infty. \end{aligned}$$

This however is impossible: if $F \in H^2$ then by the definition of F we get that $\sum_{n=0}^\infty |\widehat{S}(n)|^2 < \infty$ and since we already know that for the negative Fourier coefficients we get that S is an L^2 function. But it is supported on K which has zero measure. Hence it is identically 0. \square

Question. Frostman's theorem for the case $\alpha = 0$ states that a compact K supports a distribution S with

$$\sum_n \frac{|\widehat{S}(n)|^2}{|n|+1} < \infty$$

if and only if it has positive logarithmic capacity [7, chapter III, theorem V]. Does this theorem have a one-sided analog? We recall again the theorem of Hruščev and Peller which states that for any measure μ , the one-sided estimate $\sum_{n>0} |\widehat{\mu}(n)|^2/n < \infty$ implies the two-sided estimate above. So the question is really only about distributions.

II

There is a delicate difference between the two-sided and one-sided results. The condition (1.2) in fact implies that the compact K has positive α -measure, see [6, §10.3, theorem 2, p. 132]. The condition (1.3) does not (at least for $\alpha = 1$). Indeed, the following, quite surprising, result was proved in [11]

THEOREM 2.1. — *There exist a singular (non-trivial) pseudo-function S such that $\widehat{S} \in l^2(\mathbb{Z}^-)$.*

(recall that a pseudo-function is a distribution with $\lim_{|n| \rightarrow \infty} \widehat{S}(n) = 0$). This is surprising because it follows that the L^2 function $f(t) = \sum_{n<0} \widehat{S}(n)e^{int}$ has an a.e. converging representation by an analytic sum which is different from its Fourier expansion. However, such representations are unique (by Privalov's theorem). Thus theorem 2.1 contradicted a long-established view that *if a method of representation by a trigonometric expansion is unique, it must be the Fourier expansion*. We named the space of functions that have an analytic representation of this sort PLA. See [11] for more details.

Theorem 2.1 can be strengthened to show that the anti-analytic amplitudes of S may vanish faster than any power of $1/|n|$, see [12], where the precise threshold for the decay is found. The class PLA was further analyzed in [13].

Our goal now is to find the threshold for the size of support in Theorem 2.1 in terms of Hausdorff Λ -measure. By lemma 1.4 we see that if $\Lambda_{x \log 1/x}(K) = 0$, no such example may exist, and the condition that the positive Fourier coefficients tend to 0 is not needed for this (the assumption that S is a Schwartz distribution does imply that the positive coefficients

do not grow faster than polynomial). We now show that this condition is precise.

THEOREM 2.2. — *There exist a non-trivial pseudo-function S with $\widehat{S}(n) \in l^2(\mathbb{Z}^-)$ supported on a compact of finite $t \log 1/t$ -Hausdorff measure.*

The idea of the proof is as follows. Take the natural measure μ on some symmetric compact of the exact size. Extend it to a harmonic function on the disk, and let $F = e^{\mu+i\widetilde{\mu}}$ be an analytic function inside the disk. Let f be its almost-everywhere boundary limit. $S = F - f$ is the required object, except it might not be a pseudo-function. To solve this problem we introduce a random perturbation in the compact (and hence in μ , F , f and S). If one does not require from S to be a pseudo-function then the probabilistic part of the proof is not needed. We shall therefore note in the proof what is needed to get the simpler result, which we shall refer to as “the corollary”:

COROLLARY. — *There exist a non-trivial Schwartz distribution S with $\widehat{S}(n) \in l^2(\mathbb{Z}^-)$ supported on a compact of finite $t \log 1/t$ -Hausdorff measure.*

Question. What about the analog for $\alpha < 1$? Namely, is it true that for any $\alpha < 1$ there exists a pseudo-function S supported on a set of zero Λ_{x^α} -Hausdorff measure with

$$\sum_{n < 0} \frac{|\widehat{S}(n)|^2}{|n|^{1-\alpha}} < \infty?$$

Proof of theorem 2.2 and of the corollary. As mentioned above, our construction is similar to that of [11] in that we take a singular measure μ and define $F = e^{\mu+i\widetilde{\mu}}$. For the theorem we will take μ to be random, an idea used also in [12], to get improved smoothness of the Fourier transform, but for the corollary that will not be needed. Here are the details. Define

$$\sigma_n = \frac{2\pi}{n2^n}, \quad \tau_n = \frac{1}{6}(\sigma_{n-1} - 2\sigma_n) \quad n \geq 0. \quad (2.1)$$

Hence,

$$\frac{\tau_n}{\sigma_n} = \frac{1}{3(n-1)}. \quad (2.2)$$

Let $s(n, k)$ be a collection of numbers between 0 and 1, for each $n \in \mathbb{N}$ and each $0 \leq k < 2^n$. Most of the proof will hold for any choice of $s(n, k)$, but in the last part we shall make them random, and prove that the constructed function will have the required properties for almost any choice of $s(n, k)$.

For the corollary one may take them to be 0. Define now inductively intervals $I(n, k) = [a(n, k), a(n, k) + \sigma_n]$ (we call these $I(n, k)$ “intervals of rank n ”) using the following: $I(0, 0) = [0, 2\pi]$ and for $n \geq 0, 0 \leq k < 2^n$

$$\begin{aligned} a(n + 1, 2k) &= a(n, k) + \tau_{n+1}(1 + s(n + 1, 2k)) \\ a(n + 1, 2k + 1) &= a(n, k) + \frac{1}{2}\sigma_n + \tau_{n+1}(1 + s(n + 1, 2k + 1)) \end{aligned} \tag{2.3}$$

In other words, at the n^{th} step, inside each interval of rank n (which has length σ_n), situate two disjoint intervals of rank $n + 1$ of lengths σ_{n+1} in random places (but not too near the boundary of $I(n, k)$ or its middle). Define

$$\begin{aligned} K &:= \bigcap_{n=1}^{\infty} K_n, & K_n &:= \bigcup_{k=0}^{2^n-1} I(n, k). \\ K^\circ &:= e^{iK} & K_n^\circ &:= e^{iK_n}. \end{aligned}$$

Note that $|K_n| = \frac{2\pi}{n}$ and hence K has zero measure and finite $\Lambda_{t \log 1/t}$ -measure. Define a measure μ by the weak limit of the measures with density $\frac{1}{|K_n|} \mathbf{1}_{K_n}$,

$$\mu = \lim_{n \rightarrow \infty} \frac{1}{|K_n|} \mathbf{1}_{K_n}.$$

It is easy to see that the limit exists and is supported on K . However, we will need later on a more quantitative version of this convergence. Define therefore

$$g_n = \frac{1}{|K_n|} \mathbf{1}_{K_n}$$

and let G_n be the harmonic extension of g_n to the disk $\overline{\mathbb{D}}$.

LEMMA 2.3. — For any $z \in \overline{\mathbb{D}} \setminus K_n^\circ$

$$|G_{n+1}(z) - G_n(z)| \leq \frac{C}{2^n d(z, K_n^\circ)}. \tag{2.4}$$

Further, this holds also for the conjugate harmonic functions \widetilde{G}_n ,

$$|\widetilde{G}_{n+1}(z) - \widetilde{G}_n(z)| \leq \frac{C}{2^n d(z, K_n^\circ)}. \tag{2.5}$$

Proof. — For any n and k , $\int_{I(n,k)} g_n = 2^{-n}$. Subtracting we get

$$\int_{I(n,k)} (g_{n+1}(x) - g_n(x)) dx = 0. \tag{2.6}$$

and because $g_{n+1} - g_n$ is non-zero only on the intervals $I(n, k)$,

$$\left| \int_t^u g_{n+1}(x) - g_n(x) dx \right| \leq 2^{-n} \quad \forall t, u \in [0, 1], \forall n \tag{2.7}$$

Write $G(z) = \int_{\mathbb{T}} g(t)P_z(t)$, where P_z is the Poisson kernel. We divide into two cases: if $1 - |z| > \frac{1}{2}d(z, K_n^\circ)$ then

$$\int_{\mathbb{T}} |P'_z| \leq \frac{C}{(1 - |z|)} \leq \frac{C}{d(z, K_n^\circ)}.$$

(the first inequality is a well known property of the Poisson kernel). On the other hand, if $1 - |z| \leq \frac{1}{2}d(z, K_n^\circ)$ then $g_{n+1} - g_n$ is zero in an interval $J := [t - cd(z, K_n^\circ), t + cd(z, K_n^\circ)]$ for some c sufficiently small, where t is given by $e^{it} = z/|z|$, and

$$\int_{\mathbb{T} \setminus J} |P'_z| \leq \frac{C}{d(z, K_n^\circ)}.$$

In either case, a simple integration by parts gives (2.4) on \mathbb{D} . Finally, on $\partial\mathbb{D} \setminus K$ we have $G_{n+1}(e^{it}) - G_n(e^{it}) = g_{n+1}(t) - g_n(t) = 0$ for every $t \notin K_n$.

The proof for \tilde{G}_n is identical except the Poisson kernel P_z has to be replaced with the conjugate kernel Q_z . □

Let G be the harmonic extension of μ into the unit disc i.e. the Poisson transform of μ (equivalently you may define G as the limit of the G_n). Let \tilde{G} be the harmonic conjugate of G . Let $\delta \in (0,1)$ be some sufficiently small number (we will fix its value later). Let $F = \exp(\delta(G + i\tilde{G}))$. We note the following properties of F

- (i) F is unbounded in the unit disc.
- (ii) F has a boundary limit at almost every point of the boundary of the disk, and this limit is uniform on every closed interval disjoint from K° . Denote the boundary limit by f ,

$$f(t) := \lim_{z \rightarrow e^{it}} F(z).$$

- (iii) The function f is bounded.

All properties are simple. The first follows from the fact that for any $t \in K$, $\lim_{z \rightarrow e^{it}} G(z) = \infty$ and $|F(z)| = e^{\delta G(z)}$. The second follows because G and \tilde{G} have boundary limits outside of K — recall that G is the harmonic extension of a measure μ supported on the compact K . The third follows because for any $t \notin K$ $\lim_{z \rightarrow e^{it}} G(z) = 0$ and hence $|f(t)| = 1$.

We now define our distribution S by “ $F - f$ ” or formally by

$$\widehat{S}(m) = \widehat{F}(m) - \widehat{f}(m)$$

where \widehat{F} are the Taylor coefficients of the analytic function F at 0 namely

$$F(z) = \sum_{m=0}^{\infty} \widehat{F}(m)z^m \quad \widehat{F}(-m) = 0 \forall m \in \mathbb{N}$$

and \widehat{f} are the Fourier coefficients of f . Because $\widehat{F}(-m) = 0$ and $f \in L^2$ we immediately get that $\widehat{S} \in l^2(\mathbb{Z}^-)$. The uniform convergence of F outside K shows that S is supported on K . Thus, for the corollary it is enough to show that S is a Schwartz distribution i.e. that $\widehat{S}(m) \leq Cm^C$ for some constant C , while for the theorem it is necessary to show that S is a pseudo-function i.e. that

$$\lim_{m \rightarrow \infty} \widehat{S}(m) = 0.$$

Since this holds for f , we need only verify that $\widehat{F}(m) \rightarrow 0$, with probability 1.

The Taylor coefficients of F . We will now show that with probability 1, $\widehat{F}(m) \rightarrow 0$ as $m \rightarrow \infty$. The first step is to define $F_n = e^{\delta(G_n + i\widetilde{G}_n)}$ and find some n such that $\widehat{F}_n(m)$ approximates $\widehat{F}(m)$. Summing (2.4) and (2.5) over n we get

$$|(G_n + i\widetilde{G}_n)(z) - (G + i\widetilde{G})(z)| \leq \frac{C}{(1 - |z|)2^n}.$$

Fix, therefore, $n = n(m) := \lceil C \log m \rceil$ for some C sufficiently large, and get, for every z with $|z| = 1 - \frac{1}{m}$ that $|(G_n + i\widetilde{G}_n)(z) - (G + i\widetilde{G})(z)| \leq 1/m$. Now,

$$\sup_{z \in \mathbb{D}} |G_n(z)| = \sup_{t \in [0, 2\pi]} |g_n(t)| = \frac{1}{|K_n|} = n$$

so $|F_n(z)| \leq e^{\delta n}$. Hence for $|z| = 1 - \frac{1}{m}$,

$$\begin{aligned} |F_n(z) - F(z)| &\leq |F_n(z)| |1 - \exp(\delta((G_n + i\widetilde{G}_n)(z) - (G + i\widetilde{G})(z)))| \\ &\leq C \frac{e^{\delta n}}{m} \end{aligned} \tag{2.8}$$

and if δ is taken sufficiently small, this is $\leq Cm^{-1/2}$. Finally we use

$$\widehat{F}(m) = \frac{1}{m!} F^{(m)}(0) = \int_{|z|=1-1/m} z^{-m-1} F(z) dz$$

so

$$|\widehat{F}_n(m) - \widehat{F}(m)| = \left| \int_{|z|=1-1/m} z^{-m-1} (F_n(z) - F(z)) dz \right| \leq Cm^{-1/2} \tag{2.9}$$

and we see that it is enough to calculate $\widehat{F}_n(m)$. At this point the proof of the corollary is complete. Indeed, we may write

$$\widehat{F}_n(m) \leq \|F_n\|_2 \leq e^{\delta n} \leq m^{\delta C}$$

and hence $\widehat{F}(m) \leq Cm^C$, S is a Schwartz distribution and the corollary is proved. From now on we focus on proving the theorem.

Since F_n is a bounded function on the disk, its Taylor coefficients at 0 are identical to the Fourier coefficients of its boundary value. Denote the boundary value by $f_n(t) = F_n(e^{it})$. Thus we have reduced our problem to that of estimating

$$\widehat{f_n}(m) = \int f_n(t)e^{-imt} dt.$$

Now, $f_n = e^{\delta(g_n + i\widetilde{g_n})}$ so on K_n it is large ($|f_n(t)| = e^{\delta n}$ for any $t \in K_n$) while outside of K_n it has absolute value 1. Let us first show that the part outside of K_n is irrelevant. As in (2.8) above we use lemma 2.3 and sum over n . We get

$$|f_n(t) - f(t)| \leq \min\left(2, \frac{C}{2^n d(t, K)}\right) \quad \forall t \notin K_n$$

and by integrating

$$\int_{[0, 2\pi] \setminus K_n} |f_n(t) - f(t)| dt \leq \int_0^{2\pi} \min\left(2, \frac{C}{2^n d(t, K)}\right) dt \xrightarrow{n \rightarrow \infty} 0$$

since K is a compact of measure zero. The convergence above is uniform in the choice of the translations $s(n, k)$ that we used to construct K . Since $\widehat{f}(n) \rightarrow 0$, f being a bounded function, we get that it is enough to show

$$\int_{K_n} f_n(t)e^{-imt} dt \rightarrow 0 \tag{2.10}$$

where the limit is as $m \rightarrow \infty$ and $n = \lceil C \log m \rceil$. Here is where we will use that K_n was a random set, and we will show that this convergence holds for almost every choice of $s(n, k)$.

Probability. Take $s(n, k)$ to be independent and uniformly distributed on $[0, 1]$. We shall estimate the integral (2.10) by moment methods. Unfortunately, it seems we need the fourth moment. We start with a lemma that contains the calculation we need without referring to analytic functions

LEMMA 2.4. — *Let I_i be 4 intervals and let $\tau, \alpha, \beta > 0$ be some numbers. Let h_1, h_2, h_3 be functions satisfying*

$$\int_{I_i} |h_j| \leq \alpha, \quad i = j \text{ or } i = 4 \text{ and } j = 3 \tag{2.11}$$

$$|h_j(x)| = 1, |h'_j(x)| \leq \beta, |h''_j(x)| \leq \beta^2 \tag{2.12}$$

$$\forall x \in I_i + [-\tau, \tau] \quad \text{otherwise}$$

where “+” stands for regular set addition. See table 1. Let t_1 and t_2 be two random variables, uniformly distributed on $[0, \tau]$, and let $t_3 = t_4 = 0$.

Define

$$f(x) = f_{t_1, t_2}(x) := h_1(x - t_1)h_2(x - t_2)h_3(x). \tag{2.13}$$

Then

$$E := \left| \mathbb{E} \left(\prod_{i=1}^4 \int_{I_i+t_i} f(x_i) e^{-imx_i} dx_i \right) \right| \leq C \frac{\alpha^4}{m^2} \left(\max \beta, \frac{1}{\tau} \right)^2. \tag{2.14}$$

Proof. — See [12, lemma 9, page 1050]. The proof is essentially nothing more than two integrations by parts, each one giving a $\frac{1}{m}$ factor (from integrating e^{-imx}) and a β factor from differentiating the h -s. The $\frac{1}{\tau}$ factors come from the boundary conditions. \square

	I_1	I_2	I_3	I_4
h_1	$\int_{I_1} h_1 \leq \alpha$	$h_1^{(k)} \leq \beta^k$	$h_1^{(k)} \leq \beta^k$	$h_1^{(k)} \leq \beta^k$
h_2	$h_2^{(k)} \leq \beta^k$	$\int_{I_2} h_2 \leq \alpha$	$h_2^{(k)} \leq \beta^k$	$h_2^{(k)} \leq \beta^k$
h_3	$h_3^{(k)} \leq \beta^k$	$h_3^{(k)} \leq \beta^k$	$\int_{I_3} h_3 \leq \alpha$	$\int_{I_4} h_3 \leq \alpha$

Table 1. Relations between the functions h and the intervals I . k is 0, 1 or 2.

Continuing the proof of the theorem, for every $0 \leq k < 2^n$ denote

$$\mathcal{I}_k = \int_{I(n,k)} f_n(x) e^{-imx} dx$$

Recall that on $I(n, k)$ $|f_n| = e^{\delta n}$, so

$$|\mathcal{I}_k| \leq \int_{I(n,k)} |f_n(x)| = \frac{2\pi}{n2^n} e^{\delta n} =: \alpha. \tag{2.15}$$

In other words, $\alpha = \alpha(n)$ is a bound for $|\mathcal{I}_k|$ independent of k .

LEMMA 2.5. — Let $0 \leq k_1, k_2, k_3, k_4 < 2^n$ and let $1 \leq r < n$, and assume that the $I(n, k_i)$ belong to at least three different intervals of rank r . Then

$$|\mathbb{E}(\mathcal{I}_{k_1} \mathcal{I}_{k_2} \mathcal{I}_{k_3} \mathcal{I}_{k_4})| \leq \alpha^4 \frac{Cn^2}{m^2 \tau_r^3}.$$

(recall that τ_r were defined in (2.1))

Proof. — Define q_1, \dots, q_4 using $I(n, k_i) \subset I(r, q_i)$. We may assume without loss of generality that the two q_i -s which may be equal are q_3 and q_4 . Let \mathcal{X} be the σ -field spanning all s except $s(r, q_1)$ and $s(r, q_2)$. We shall show

$$|\mathbb{E}(\mathcal{I}_{k_1}\mathcal{I}_{k_2}\mathcal{I}_{k_3}\mathcal{I}_{k_4}|\mathcal{X})| \leq \alpha^4 \frac{Cn^2}{m^2\tau_r^3}$$

and then integrating over \mathcal{X} will give the result. We note that conditioning on \mathcal{X} is in effect fixing everything except the positions of $I(r, q_1)$ and $I(r, q_2)$ inside $I(r-1, \lfloor q_i/2 \rfloor)$. Denote $J_j := I(r, q_j)$ ($j = 1, 2$) and $J_3 = \mathbb{T} \setminus (J_1 \cup J_2)$. Assume for a moment that $s(r, q_1) = s(r, q_2) = 0$ and define, using this assumption,

$$\begin{aligned} \eta_j &:= g_n|_{J_j} & j = 1, 2, 3, & \quad h_j &:= e^{\eta_j + i\tilde{\eta}_j}, \\ I_i &:= I(n, k_i) & i = 1, 2, 3, 4. \end{aligned} \tag{2.16}$$

Under the assumption $s(r, q_1) = s(r, q_2) = 0$ we clearly have $f_n = h_1 h_2 h_3$ and when we remove this assumption, the only change is a translation of h_1 and h_2 . In other words, if we define $t_i = s(r, q_i)\tau_r$ then $f_n(x) = h_1(x - t_1)h_2(x - t_2)h_3(x)$. Examining (2.13) we see that $|\mathbb{E}(\mathcal{I}_{k_1}\mathcal{I}_{k_2}\mathcal{I}_{k_3}\mathcal{I}_{k_4}|\mathcal{X})| = E$ where E is defined by (2.14); where the I_i of (2.14) are the same as those of (2.16); and where the τ of (2.14) is τ_r and where the α of (2.11) is our α . To make (2.14) concrete we need to specify a value for the β of (2.12) and prove that it holds. We define

$$\beta = C_1 \frac{n}{\tau_r^{3/2}}$$

for some sufficiently large constant C_1 to be fixed later. Notice that β is obviously larger than $1/\tau_r$. With all these, lemma 2.5 would follow from lemma 2.4 once we show (2.12).

Examining the definitions of η_j and $I(n, k)$ it is easy to see that $\eta_j(x) = 0$ for $x \in I_i + [-\tau, \tau]$ when $i \neq j$ except when $j = 3$ and $i = 4$ (recall that in (2.3) we left a little space in the sides of the intervals — this is the reason). This immediately shows $|h_j(x)| = 1$. Further, $h'_j = h_j(\eta'_j + i\tilde{\eta}'_j)$ gives $|h'_j| = |\tilde{\eta}'_j|$ and $h''_j = h_j((\eta'_j + i\tilde{\eta}'_j)^2 + \eta''_j + i\tilde{\eta}''_j)$ gives $|h''_j| \leq |\tilde{\eta}'_j|^2 + |\tilde{\eta}''_j|$. Now, the derivatives of $\tilde{\eta}_j$ have the representations

$$\tilde{\eta}'_j(x) = \int_{\mathbb{T}} \eta_j(x-t)H'(t) dt \quad \tilde{\eta}''_j(x) = \int_{\mathbb{T}} \eta_j(x-t)H''(t) dt$$

where H is the Hilbert kernel. This again works because η_j is zero in a neighborhood of x , otherwise there would be extra terms (the derivatives of H are distributions with a singular part at 0, but here we may consider them as just functions). We may therefore estimate

$$\tilde{\eta}'_j(x) \leq \|\eta\|_2 \|H'\|_{\mathbb{T} \setminus [-\tau_r, \tau_r]} \quad \tilde{\eta}''_j(x) \leq \|\eta\|_2 \|H''\|_{\mathbb{T} \setminus [-\tau_r, \tau_r]} \tag{2.17}$$

Now, it is well known that for any $D \geq 0$, the D^{th} derivative of H satisfies

$$|H^{(D)}(t)| \leq \frac{C(D)}{|e^{it} - 1|^{D+1}}. \tag{2.18}$$

So

$$\|H'\|_{[-\tau_r, \tau_r]^c} \approx \tau_r^{-3/2} \quad \|H''\|_{[-\tau_r, \tau_r]^c} \approx \tau_r^{-5/2}.$$

and since $\|\eta\|_2 \leq Cn$ we get the estimate we need:

$$|h'_j| \leq C\tau_r^{-3/2}n \quad |h''_j| \leq C\tau_r^{-3}n^2.$$

We may now fix the value of C_1 from the definition of β . With this the conditions of lemma 2.4 are fulfilled and we are done. \square

End of the proof of theorem 2.2. — We return to the bound on

$$X := \int_{K_n} f_n(t)e^{-imt} dt.$$

We shall do so by estimating $\mathbb{E}|X|^4$. Let

$$E(k_1, k_2, k_3, k_4) := \left| \mathbb{E} \prod \mathcal{I}_{k_i} \right|$$

let $r(k_1, \dots, k_4)$ be the minimal r such that the $I(n, k_i)$ -s are contained in at least 3 distinct intervals of rank r . A simple calculation shows

$$\#\{(k_1, \dots, k_4) : r(k_1, \dots, k_4) = r\} \approx 2^{4n-2r}.$$

If τ_r is too small then the estimate of lemma 2.5 is useless and it would be better to estimate $|E(k_1, \dots, k_4)| \leq \alpha^4$. Let R be some number. Then for large r we have the estimate

$$E_1 := \sum_{r(k_1, \dots, k_4) \geq R} E(k_1, \dots, k_4) \leq C\alpha^4 2^{4n-2R} \stackrel{(2.15)}{\leq} Ce^{4\delta n} 2^{-2R} \leq Cm^{C\delta} 2^{-2R}. \tag{2.19}$$

For small r we use the lemma to get a better estimate. Examine one such k_1, \dots, k_4 and let $r = r(k_1, \dots, k_4)$. Lemma 2.5 gives

$$E(k_1, \dots, k_4) \leq \alpha^4 \frac{Cn^2}{m^2\tau_r^3} = \frac{\alpha^4}{m^{2-o(1)}\tau_r^3}.$$

Therefore

$$\begin{aligned} E_2 := \sum_{r(k_1, \dots, k_4) < R} E(k_1, \dots, k_4) &\leq \alpha^4 2^{4n} m^{-2+o(1)} \sum_{r=1}^R 2^{-2r} \tau_r^{-3} \\ &\stackrel{(2.1, 2.15)}{=} m^{-2+C\delta+o(1)} \sum_{r=1}^R 2^{r+o(r)} = m^{-2+C\delta+o(1)} 2^{R+o(R)}. \end{aligned} \tag{2.20}$$

Taking $R = \lfloor \frac{2}{3} \log_2 m \rfloor$ and summing (2.19) and (2.20) we get

$$\mathbb{E}|X|^4 \leq m^{-4/3+C\delta+o(1)}. \tag{2.21}$$

With δ sufficiently small this is summable. Hence

$$\mathbb{E} \sum_m |X_m|^4 = \sum \mathbb{E} |X_m|^4 < \infty$$

In particular, with probability 1, $X_m \rightarrow 0$. As remarked above, this shows that $\widehat{f}_n(m) \rightarrow 0$ and hence $\widehat{F}(m) \rightarrow 0$ which concludes the theorem. \square

Let us remark that we do not know if the probabilistic part of the proof is really necessary. In fact it is quite possible that the non-probabilistic construction (i.e. setting all $s(n, k)$ to 0) also satisfies that $\widehat{F}(m) \rightarrow 0$. What we do know is that the theorem does not follow formally from the corollary, i.e. there exists a singular distribution S with $\widehat{S}(n) \in l^2(\mathbb{Z}^-)$ which is not a pseudo-function. The construction is very similar to that of the theorem except one takes δ — the δ from the definition $F = \exp(\delta(G + i\widetilde{G}))$ — large rather than small. It then follows that $\|F_n\|_2$ is a large power of m leading to $\widehat{F}(m)$ being unbounded. We omit all other details.

III

Let now K be some fixed compact. Suppose K supports a non-trivial measure with some one-sided smoothness property. Does it imply that K in fact supports a measure with a two-sided property? What about distributions? We summarize some relations of this sort in the tables in figure 1.

Each arrow denotes an implication which holds for any compact. For example, the arrow labelled “B” in the top right corner is a result of Beurling that any compact K which supports a distribution S with

$$\sum |\widehat{S}(n)|^2 |n|^{\alpha-1} < \infty$$

also supports a measure μ with the same property [2]. In the top-right table we assume $\alpha < 1$ and in the bottom-left $2 < q < \infty$. The abbreviations are KO for [11], Ri for the Riesz analyticity theorem, PS for Piatetski-Shapiro [18] and Ra for Rajchman [19]. The arrow marked LO comes from [15] and noting that the set constructed in [15] is a Helson set, so it cannot support even a measure μ with $\widehat{\mu}(n) \rightarrow 0$. The unmarked arrow on the top-left is trivial, while the unmarked arrow on the bottom-right and the two unmarked arrows on the bottom left follow from the other arrows in their diagrams. All missing arrows are unknown to us.




Figure 1. Properties for specific compacts.

Our last result is to show that a certain “non-symmetry” is possible for singular measures supported by compacts of dimension $\alpha < 1$. It does not fit in any of the tables above, but it is very close in spirit.

THEOREM 3.1. — *Let K be a compact on \mathbb{T} , $\dim_{\mathbb{F}}(K) = d > 0$. Then for every $p > 2/d$ there exists a (complex) measure $\nu \in \mathcal{M}(K)$ such that*

$$\widehat{\nu} \in l^p(\mathbb{Z}_-), \quad \widehat{\nu} \notin l^p(\mathbb{Z}_+).$$

(recall the definition of the Fourier dimension on page 1208).

LEMMA 3.2. — *Let μ be a non-atomic positive measure, g be a function in $C(\mathbb{T})$. Then for every $\delta > 0$ there is a positive integer l such that*

$$\int |g(lt)| d\mu < \int |g(t)| d\mu + \delta \tag{3.1}$$

Proof. — Clearly it is enough to consider g as a positive trigonometric polynomial. Assume first that $\widehat{\mu}(n) = o(1)$ as $|n| \rightarrow \infty$ (the only case needed for the proof of Theorem 3.1). Then we have

$$\int g(lt) d\mu = \sum \widehat{g}(n) \overline{\widehat{\mu}(ln)}$$

and for a large l the right-hand side is $< \widehat{g}(0)\widehat{\mu}(0) + \delta$ which gives (3.1).

For a general non-atomic μ we can use Wiener’s theorem [8, §I.7.11] to get

$$\frac{1}{2N+1} \sum_{|n| \leq N} |\widehat{\mu}(n)|^2 = o(1).$$

From this it follows that for a random l_N between $N/2K$ and N/K ,

$$\lim_{N \rightarrow \infty} \sum_{0 < |n| \leq K} |\widehat{\mu}(l_N n)| = 0$$

and the argument can be completed in the same way. □

Proof of Theorem 3.1. — Let $\dim_{\mathbb{F}} K = d$, $p > 2/d$. According to the definition of the Fourier dimension we can find a positive measure $\mu \in \mathcal{M}(K)$ such that $\widehat{\mu} \in l^p(\mathbb{Z})$. We can assume that $\|\widehat{\mu}\|_{l^p(\mathbb{Z})} = 1$. Now we define inductively a sequence of (complex) measures ν_k . Let us describe the k^{th} step of the induction. Choose a number $s = s_k > 0$ such that $\|\widehat{\mu}\|_{l^p(-\infty, -s)} < 2^{-k}$. Set

$$g_k(t) := 4^{-k} \sum_{j=1}^{4^k} e^{iq(j)t}$$

By choosing the frequencies $s < q(1) < \dots < q(4^k)$ sparse enough one can satisfy the inequality:

$$\|\widehat{\mu g_k}\|_{l^p(\mathbb{Z}_+)} > 1/2 \|\widehat{\mu}\|_{l^p} \|\widehat{g_k}\|_{l^1} = 1/2$$

and it will hold true when replace $g_k(t)$ by $g_k(lt)$ for $l > 1$. So for $\nu_k := g_k(l)\mu$ we will have $\|\nu_k\|_{l^p(I_k)} > 1/2$ for a certain finite interval $I_k \subset \mathbb{Z}_+$, and $\|\nu_k\|_{l^p(\mathbb{Z}_-)} < 2^{-k}$. In addition, choosing l from lemma 3.2 for sufficiently small δ we get from (3.1)

$$\|\nu_k\|_{\mathcal{M}(\mathbb{T})} \stackrel{(3.1)}{<} 2 \|g_k\|_1 \|\mu\|_{\mathcal{M}(\mathbb{T})} \leq 2 \|g_k\|_2 = 2^{-k+1}$$

Certainly in the above induction we can get the intervals I_k to be disjoint. It follows that the measure $\nu = \sum \nu_k$ satisfies the requirements of the theorem. □

Remark. — The restriction $p > 2/d$ in theorem 3.1 is essentially sharp. Indeed, take a Salem set K of given dimension, that is $\dim K = \dim_{\mathbb{F}} K = d$. Suppose there is a measure $\nu \in \mathcal{M}(K)$ such that $\widehat{\nu} \in l^p(\mathbb{Z}_-)$ for some $p < 2/d$. Then the condition (1.3) is satisfied for some $\alpha > d$. So Theorem 1.1 implies $\dim K > d$. Which is a contradiction.

Questions.

- (i) In the spirit of Theorem 3.1 one may ask the following. Let K be a compact with $\dim K = d$. Does it follow that for any $0 < \alpha < d$ there is a measure μ supported on K with

$$\sum_{n < 0} \frac{|\widehat{\mu}(n)|^2}{|n|^{1-\alpha}} < \infty \quad \sum_{n > 0} \frac{|\widehat{\mu}(n)|^2}{|n|^{1-\alpha}} = \infty? \tag{3.2}$$

The following strict version is also open: Let K be some compact which supports a measure μ with $\sum_{n \neq 0} |\widehat{\mu}(n)|^2 |n|^{\alpha-1} < \infty$. Is it always true that it also supports a measure μ with (3.2)?

- (ii) Does there exist a compact of uniqueness K (see e.g. the book [10] for sets of uniqueness) which supports a pseudo-measure S with $\widehat{S}(n) = o(1)$ as $n \rightarrow +\infty$? One can show that Cantor sets with any ratio of dissection do not (in fact any $H^{(n)}$ -set does not support such a pseudo-measure, the argument of Piatetski-Shapiro applies, see [10, chap. III, thm. 4]).

- (iii) One can introduce $\dim_1(K)$ as

$$\sup\{a : K \text{ supports a measure } \mu \text{ such that } \widehat{\mu} \in l^{2/a}(\mathbb{Z})\}.$$

Then $\dim_{\mathbb{F}} K \leq \dim_1 K \leq \dim K$. Is it any easier to construct a set with $\dim K = \dim_1 K$ than a Salem set? One may call such sets “quasi-Salem sets”.

Acknowledgements. Work partially supported by the authors’ respective Israel Science Foundation grants.

BIBLIOGRAPHY

- [1] R. D. BERMAN, “Boundary limits and an asymptotic Phragmén-Lindelöf theorem for analytic functions of slow growth”, *Indiana University Mathematics Journal* **41/2** (1992), p. 465-481.
- [2] A. BEURLING, “Sur les spectres des fonctions”, [French, on the spectrum of functions], *Analyse Harmonique, Colloq. Internat. CNRS* 15, Paris, 1949, p. 9-29.
- [3] B. E. J. DAHLBERG, “On the radial boundary values of subharmonic functions”, *Math. Scand.* **40** (1977), p. 301-317.
- [4] K. FALCONER, *Fractal geometry*, second ed., Mathematical foundations and applications, John Wiley & Sons, Inc., Hoboken, New Jersey, 2003.
- [5] S. V. HRUŠČEV & V. V. PELLER, “Hankel operators of Schatten-von Neumann class and their application to stationary processes and best approximations”, Appendix to the English edition of: N. K. Nikol’skii, *Treatise on the shift operator*, Translated from the Russian by Jaak Peetre. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 273, Springer-Verlag, Berlin, 1986, p. 399-454.
- [6] J.-P. KAHANE, *Some random series of functions*, second ed., Cambridge Studies in Advanced Mathematics 5, Cambridge University Press, Cambridge, 1985.
- [7] J.-P. KAHANE & R. SALEM, *Ensembles parfaits et séries trigonométriques*, [French, Perfect sets and trigonometric series], Second ed., With notes by Kahane, Thomas W. Körner, Russell Lyons and Stephen William Drury. Hermann, Paris, 1994.
- [8] Y. KATZNELSON, *An introduction to harmonic analysis*, Second corrected ed., Dover Publications, Inc., New York, 1976.
- [9] R. KAUFMAN, “On the theorem of Jarník and Besicovitch”, *Acta Arith.* **39:3** (1981), p. 265-267.

- [10] A. S. KECHRIS & A. LOUVEAU, *Descriptive set theory and the structure of sets of uniqueness*, London Mathematical Society Lecture Note Series, vol. 128, Cambridge University Press, Cambridge, 1987.
- [11] G. KOZMA & A. OLEVSKIĬ, “A null series with small anti-analytic part”, *Comptes Rendus de l’Académie des Sciences Paris, Série I Mathématique* **336:6** (2003), p. 475-478.
- [12] ———, “Analytic representation of functions and a new quasi-analyticity threshold”, *Annals of Math.* **164:3** (2006), p. 1033-1064.
- [13] ———, “Is PLA large?”, *Bull. Lond. Math. Soc.* **39:2** (2007), p. 173-180.
- [14] K. DE LEEUW & Y. KATZNELSON, “The two sides of a Fourier-Stieltjes transform and almost idempotent measures”, *Israel J. Math.* **8** (1970), p. 213-229.
- [15] N. LEV & A. OLEVSKIĬ, “Wiener’s ‘closure of translates’ problem and Piatetski-Shapiro’s uniqueness phenomenon”, To appear in *Ann. Math.* <http://arxiv.org/abs/0908.0447>.
- [16] P. MATTLA, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*, Cambridge Studies in Advanced Mathematics, vol. 44, Cambridge University Press, Cambridge, 1995.
- [17] G. MOCKENHAUPT, “Salem sets and restriction properties of Fourier transforms”, *Geom. Funct. Anal.* **10:6** (2000), p. 1579-1587.
- [18] I. I. РYАТЕСКИĬ-ШАПИРО, “Дополнение к работе “К проблеме единственности разложения функции в тригонометрический ряд””, *Moskov. Gos. Univ. Uč. Zap. Mat.* **165** (1954), p. 79-97, [Russian, Supplement to the work “On the problem of uniqueness of expansion of a function in a trigonometric series”] English translation in *Selected Works of Ilya Piatetski-Shapiro*, AMS Collected Works, vol. 15, 2000.
- [19] A. RAJCHMAN, “Une classe de séries trigonométriques qui convergent presque partout vers zéro”, [French, A class of trigonometric series converging almost everywhere to zero] *Math. Ann.* **101:1** (1929), p. 686-700.
- [20] M. TAMASHIRO, “Dimensions in a separable metric space”, *Kyushu J. Math.* **49:1** (1995), p. 143-162.
- [21] A. ZYGMUND, *Trigonometric series. Vol. I, II*, third ed., With a foreword by Robert A. Fefferman. Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2002.

Manuscrit reçu le 26 avril 2010,
 accepté le 21 septembre 2010.

Gady KOZMA
 Department of Mathematics,
 The Weizmann Institute of Science,
 Rehovot POB 76100,
 Israel.
gady.kozma@weizmann.ac.il

Alexander OLEVSKIĬ
 School of Mathematics,
 Tel Aviv University,
 Tel Aviv 69978,
 Israel.
olevskii@post.tau.ac.il