



# ANNALES

DE

# L'INSTITUT FOURIER

Jonathan PILA

**A correction to “Counting rational points on a certain exponential-algebraic surface”**

Tome 67, n° 3 (2017), p. 1277-1278.

[http://aif.cedram.org/item?id=AIF\\_2017\\_\\_67\\_3\\_1277\\_0](http://aif.cedram.org/item?id=AIF_2017__67_3_1277_0)



© Association des Annales de l'institut Fourier, 2017,

*Certains droits réservés.*



Cet article est mis à disposition selon les termes de la licence  
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.  
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »  
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales  
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

*Article mis en ligne dans le cadre du*  
*Centre de diffusion des revues académiques de mathématiques*  
<http://www.cedram.org/>

## A CORRECTION TO “COUNTING RATIONAL POINTS ON A CERTAIN EXPONENTIAL-ALGEBRAIC SURFACE”

*Annales de l'institut Fourier*, vol. 60 (2010), n°2, 489–514

by Jonathan PILA (\*)

---

ABSTRACT. — We correct a mistake in the paper. All the theorems remain correct.

RÉSUMÉ. — Nous corrigeons une erreur dans la papier. Tous les théorèmes restent vrais.

This note is to correct a part of the paper [1]. All the theorems remain correct (though some of the implied constants will be adversely affected). I am very grateful to John Armitage for pointing out my error.

The mistake occurs in the early lines of the proof of Theorem 6.1, p. 507. The exhibited parameterization of the surface

$$\mathcal{X} = \{(x, y, z) \in (0, 1)^3 : \log x \log y = -\log z\}$$

is not *mild* (as defined in the paper), as claimed there, because the function

$$z = \exp\left(\left(1 - \frac{1}{s^g}\right)\left(1 - \frac{1}{t^g}\right)\right), \quad (s, t) \in (0, 1)^2$$

is not mild. The seemingly innocuous change from  $-1/s^g$  to  $1 - 1/s^g$  (and the same for  $t$ ) to cover all of  $\mathcal{X}$  spoils the mildness, and indeed the surface needs to be non-trivially reparameterized near the  $x = 1$  boundary for small  $y$  (and near the  $y = 1$  boundary for small  $x$ ), as well as near the origin.

---

*Keywords:* O-minimal structure, rational points.

*Math. classification:* 11G99, 03C64.

(\*) Preparation of this correction was partially supported by a grant from EPSRC, number EP/N008359/1.

We should therefore use instead the following parameterization of  $\mathcal{X}$  by four maps  $\theta_i : (0, 1)^2 \rightarrow (0, 1)^3$ ,  $i = 1, \dots, 4$  (here  $g \in \mathbb{Z}$ ,  $g > 0$ ):

$$\theta_1(s, t) = \left( \exp\left(-\frac{1}{s^g}\right), \exp\left(-\frac{1}{t^g}\right), \exp\left(-\frac{1}{(st)^g}\right) \right),$$

$$\theta_2(s, t) = \left( \exp(-s^g), \exp\left(-\frac{1}{(st)^g}\right), \exp\left(-\frac{1}{t^g}\right) \right),$$

$$\theta_3(s, t) = \left( \exp\left(-\frac{1}{(st)^g}\right), \exp(-t^g), \exp\left(-\frac{1}{s^g}\right) \right),$$

$$\theta_4(s, t) = (\exp(-s^g), \exp(-t^g), \exp(-(st)^g)).$$

These maps cover, respectively, the  $(x, y)$ -regions:  $(0, \frac{1}{e}) \times (0, \frac{1}{e})$ ,  $(\frac{1}{e}, 1) \times (0, \frac{1}{e})$ ,  $(0, \frac{1}{e}) \times (\frac{1}{e}, 1)$ , and  $(\frac{1}{e}, 1) \times (\frac{1}{e}, 1)$ . The first three maps are  $(A, 1+1/g)$ -mild for some suitable  $A = A(g) > 0$ , by Proposition 2.8, and the last is  $(A(g), 0)$ -mild, being analytic on the closure of  $(0, 1)^2$ . The omitted lines are in the “algebraic part” of  $\mathcal{X}$  and play no role.

## BIBLIOGRAPHY

- [1] J. PILA, “Counting rational points on a certain exponential-algebraic surface”, *Ann. Inst. Fourier* **60** (2010), no. 2, p. 489-514.

Manuscrit reçu le 24 janvier 2017,  
accepté le 1<sup>er</sup> mars 2017.

Jonathan PILA  
Mathematical Institute,  
University of Oxford,  
Oxford, OX2 6RT (United Kingdom)  
pila@maths.ox.ac.uk