



ANNALES

DE

L'INSTITUT FOURIER

Gilles CARRON

Riesz transforms on connected sums

Tome 57, n° 7 (2007), p. 2329-2343.

http://aif.cedram.org/item?id=AIF_2007__57_7_2329_0

© Association des Annales de l'institut Fourier, 2007, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

RIESZ TRANSFORMS ON CONNECTED SUMS

by Gilles CARRON

ABSTRACT. — Assume that M_0 is a complete Riemannian manifold with Ricci curvature bounded from below and that M_0 satisfies a Sobolev inequality of dimension $\nu > 3$. Let M be a complete Riemannian manifold isometric at infinity to M_0 and let $p \in (\nu/(\nu - 1), \nu)$. The boundedness of the Riesz transform of $L^p(M_0)$ then implies the boundedness of the Riesz transform of $L^p(M)$.

RÉSUMÉ. — Soit M_0 une variété riemannienne complète à courbure de Ricci bornée inférieurement et qui vérifie l'inégalité Sobolev de dimension $\nu > 3$. Si M est une variété riemannienne complète isométrique à M_0 en dehors d'un compact et si $p \in (\nu/(\nu - 1), \nu)$ alors lorsque la transformée de Riesz est bornée sur $L^p(M_0)$ elle est également bornée sur $L^p(M)$.

1. Introduction

Let (M, g) be a complete Riemannian manifold with infinite volume, we denote by $\Delta = \Delta^g$ its Laplace operator, it has an unique self-adjoint extension on $L^2(M, d \text{vol}_g)$ which is also denoted by Δ . The Green formula and the spectral theorem show that for any $\varphi \in C_0^\infty(M)$:

$$\|d\varphi\|_{L^2}^2 = \langle \Delta\varphi, \varphi \rangle = \|\Delta^{1/2}\varphi\|_{L^2}^2;$$

hence the Riesz transform $T := d\Delta^{-1/2}$ extends to a bounded operator

$$T : L^2(M) \rightarrow L^2(M; T^*M).$$

On the Euclidean space, it is well known that the Riesz transform has also a bounded extension $L^p(M) \rightarrow L^p(M; TM)$ for any $p \in]1, \infty[$. However, this is not a general feature of the Riesz transform on complete Riemannian manifolds, as a matter of fact, on the connected sum of two copies of the Euclidean space \mathbb{R}^n , the Riesz transform is not bounded on L^p for any

Keywords: Riesz transform, Sobolev inequalities.

Math. classification: 58J37, 58J35, 42B20.

$p \in [n, \infty[\cap]2, \infty[$ ([7, 5]). It is of interest to figure out the range of p for which T extends to a bounded map $L^p(M) \rightarrow L^p(M; T^*M)$. The main result of [5] answered to this question for manifolds with Euclidean ends :

THEOREM 1.1. — *Let M be a complete Riemannian manifold of dimension $n \geq 3$ which is the union of a compact part and a finite number of Euclidean ends. Then the Riesz transform is bounded from $L^p(M)$ to $L^p(M; T^*M)$ for $1 < p < n$, and is unbounded on L^p for all other values of p if the number of ends is at least two.*

The proof of this result used an asymptotic expansion of the Schwarz kernel of the resolvent $(\Delta + k^2)^{-1}$ near $k \rightarrow 0$. In [5] using L^p cohomology, we also find a criterion which insures that the Riesz transform is unbounded on L^p :

THEOREM 1.2. — *Assume that (M, g) is a complete Riemannian manifold with Ricci curvature bounded from below such that for some $\nu > 2$ and $C > 0$, (M, g) satisfies the Sobolev inequality*

$$\forall \varphi \in C_0^\infty(M), \|\varphi\|_{L^{\frac{2\nu}{\nu-2}}} \leq C \|d\varphi\|_{L^2}$$

and

$$(1.1) \quad \forall x \in M, \forall r > 1, \text{ vol } B(x, r) \leq Cr^\nu.$$

If M has at least two ends, then the Riesz transform is not bounded on L^p for any $p \geq \nu$.

Let (N, g_0) be a simply connected nilpotent Lie group of dimension $n > 2$ (endowed with a left invariant metric). According to [1] we know that the Riesz transform on (N, g_0) is bounded on L^p for every $p \in]1, \infty[$. Let ν be the homogeneous dimension of N ; for instance we can set

$$\nu = \lim_{R \rightarrow \infty} \frac{\log \text{vol } B(o, R)}{\log R},$$

$o \in N$ being a fixed point. Let (M, g) be a manifold isometric at infinity to $k \geq 1$ copies of (N, g_0) . That is to say there are compact sets $K \subset M$ and $K_0 \subset N$ such that $(M \setminus K, g)$ is isometric to k copies of $(N \setminus K_0, g_0)$. According to [7] we know that on (M, g) the Riesz transform is bounded on L^p for $p \in]1, 2]$. And the theorem 1.2 says that the Riesz transform is not bounded on L^p when $p \geq \nu$. In [5], we make the following conjecture : *show that the Riesz transform on (M, g) is bounded on L^p for $p \in]1, \nu[$.* The main result of this paper gives a positive answer to this conjecture ; in fact we obtain a more general result concerning the boundedness of Riesz transform for connected sums, under some mild geometrical conditions :

THEOREM 1.3. — *Let (M_0, g_0) be a complete Riemannian manifold, we assume that the Ricci curvature of (M_0, g_0) is bounded from below and that for some $\nu > 3$ and $C > 0$, (M_0, g_0) satisfies the Sobolev inequality*

$$\forall \varphi \in C_0^\infty(M_0), \quad \|\varphi\|_{L^{\frac{2\nu}{\nu-2}}} \leq C \|d\varphi\|_{L^2}.$$

Let $p \in]\nu/(\nu - 1), \nu[$, if on (M_0, g_0) the Riesz transform is bounded on L^p then the Riesz transform is also bounded on L^p for any manifold M isometric at infinity to several copies of (M_0, g_0) .

Moreover under a uniform upper growth control of the volume of geodesic balls (such as (1.1)), the result of [7] implies that under the assumption of the theorem 1.3, the Riesz transform is bounded on M for any $p \in]1, 2]$; hence the restriction of $p > \nu/(\nu - 1)$ is not really a serious one. Our method is here less elaborate than the one of [5], its gives a more general result but it is less sharp ; there are two restrictions : the first one is the dimension restriction $\nu > 3$ which is unsatisfactory, and the second concerns the limitation $p < \nu$ which is perhaps also unsatisfactory when M has only one end. However there are recent results of T. Coulhon and N. Dungey in this direction [6].

There is now a long list of complete Riemannian manifolds (M_0, g_0) satisfying our hypothesis and on which the Riesz transform is known to be bounded on L^p for every $p \in]1, \infty[$. For instance Cartan-Hadamard manifolds with a spectral gap [17], non-compact symmetric spaces [2] and Lie groups of polynomial growth [1], manifolds with nonnegative Ricci curvature and maximal volume growth [3] (see the discussion at the end of the proof of theorem 1.3 about the case of manifolds with nonnegative Ricci curvature and non maximal volume growth). Also H.-Q. Li [16] proved that the Riesz transform on n -dimensional cones with compact basis is bounded on L^p for $p < p_0$, where

$$p_0 = \begin{cases} n\left(\frac{n}{2} - \sqrt{\left(\frac{n-2}{2}\right)^2 + \lambda_1}\right)^{-1}, & \lambda_1 < n - 1 \\ +\infty, & \lambda_1 \geq n - 1, \end{cases}$$

where λ_1 is the smallest nonzero eigenvalue of the Laplacian on the basis. Note that $p_0 > n$. Our proof also applies to a manifold isometric at infinity to several copies of cones, hence our theorem 1.3 also gives a partial answer to the open problem 8.1 of [5] :

COROLLARY 1.4. — *If (M, g) is a smooth Riemannian n -manifold of dimension $n \geq 4$ with conic ends, then the Riesz transform is bounded on L^p for any $p \in]1, n[$.*

Our manifold (M_0, g_0) is not assumed to be connected, for instance the theorem 1.3 implies that on the connected sum of a hyperbolic space and a euclidean space of dimension $n > 3$, the Riesz transform is bounded on L^p , for $p \in]n/(n-1), n[$.

Acknowledgements. I thank T. Coulhon for his interest and for his nice comments which have improved the previous version of this paper.

2. Analytic preliminaries

2.1. A Sobolev inequality

PROPOSITION 2.1. — *Let (M, g) be a complete Riemannian manifold with Ricci curvature bounded from below then for any $p \in [1, \infty[$, there is a constant C such that for all $\varphi \in C_0^\infty(M)$*

$$\|df\|_{L^p} \leq C [\|\Delta f\|_{L^p} + \|f\|_{L^p}].$$

Remark 2.2.

- i) In [8], T. Coulhon and X. Duong have shown that for every complete Riemannian manifolds and any $p \in]1, 2]$, there is a constant C such that

$$\forall f \in C_0^\infty(M), \|df\|_{L^p}^2 \leq C \|\Delta f\|_{L^p} \|f\|_{L^p}.$$

When $p \in]1, 2]$, this is clearly a stronger result.

- ii) When the injectivity radius is assumed moreover to be positive, this result is due to B. Davies (see corollary 10 in [10]) ; in this setting, another proof along the idea of [14] can be given.

Proof. — According to (theorem 4.1 in [3]) we know that if (M, g) is a complete manifold with Ricci curvature bounded from below then for any $p \in]1, \infty[$ there is a constant C such that

$$\forall f \in C_0^\infty(M), \|df\|_{L^p} \leq C \left[\left\| \Delta^{1/2} f \right\|_{L^p} + \|f\|_{L^p} \right].$$

Then an interpolation argument (see for instance proposition 5.5 in [15]) implies that

$$\left\| \Delta^{1/2} f \right\|_{L^p}^2 \leq \|\Delta f\|_{L^p} \|f\|_{L^p},$$

the proposition is now straightforward. □

2.2. Some estimates on the Poisson operator

LEMMA 2.3. — *Let (M, g) be a complete Riemannian manifold which for some $\nu > 2$ and $C > 0$ satisfies the Sobolev inequality :*

$$\forall \varphi \in C_0^\infty(M), \|\varphi\|_{L^{\frac{2\nu}{\nu-2}}} \leq C \|d\varphi\|_{L^2}$$

then the Schwarz kernel $P_\sigma(x, y)$ of the Poisson operator $e^{-\sigma\sqrt{\Delta}}$ satisfies

$$P_\sigma(x, y) \leq \frac{C\sigma}{(\sigma^2 + d(x, y)^2)^{\frac{\nu+1}{2}}}.$$

Moreover if $1 \leq r \leq p \leq +\infty$ then

$$\left\| e^{-\sigma\sqrt{\Delta}} \right\|_{L^r \rightarrow L^p} \leq \frac{C}{\sigma^{\nu(\frac{1}{r} - \frac{1}{p})}}.$$

We know that the heat operator $e^{-t\Delta}$ and the Poisson operator are related through the subordination identity :

$$e^{-\sigma\sqrt{\Delta}} = \frac{\sigma}{2\sqrt{\pi}} \int_0^\infty e^{-\frac{\sigma^2}{4t}} e^{-t\Delta} \frac{dt}{t^{3/2}}.$$

Hence these properties follow directly from the corresponding ones for the heat operator $e^{-t\Delta}$ and its Schwarz kernel $H_t(x, y)$:

$$(2.1) \quad H_t(x, y) \leq \frac{c}{t^{\nu/2}} e^{-\frac{d(x,y)^2}{5t}}$$

and if $1 \leq r \leq p \leq +\infty$ then

$$\left\| e^{-t\Delta} \right\|_{L^r \rightarrow L^p} \leq \frac{C}{t^{\frac{\nu}{2}(\frac{1}{r} - \frac{1}{p})}},$$

which are consequences of the Sobolev inequality [18, 9].

We will also need an estimate for the derivative of the Poisson kernel :

LEMMA 2.4. — *Under the assumptions of lemma (2.3), let $\Omega \subset M$ be an open subset and K be a compact set in the interior of $M \setminus \Omega$ then*

$$\begin{aligned} \left\| e^{-\sigma\sqrt{\Delta}} \right\|_{L^p(\Omega) \rightarrow L^\infty(K)} &\leq \frac{C}{(1 + \sigma)^{\nu/p}}, \\ \left\| \nabla e^{-\sigma\sqrt{\Delta}} \right\|_{L^p(\Omega) \rightarrow L^\infty(K)} &\leq \frac{C}{(1 + \sigma)^{\nu/p}}. \end{aligned}$$

Proof. — The first estimate is only a consequence of the lemma 2.3 because by assumption there is a constant $\varepsilon > 0$ such that

$$(2.2) \quad (x, y) \in K \times \Omega \Rightarrow d(x, y) \geq \varepsilon.$$

To prove the second inequality, we will again only show the corresponding estimate for the heat operator. First, according to the local Harnack

inequality (see V.4.2 in [9]), there is a constant C such that for any $x \in K$, $t \in]0, 1]$ and $y \in M$:

$$(2.3) \quad |\nabla_x H_t(x, y)| \leq \frac{C}{\sqrt{t}} H_{2t}(x, y).$$

But hence by (2.2) and (2.1), we get : for all $(x, y) \in K \times \Omega$ then

$$H_{2t}(x, y) \leq \frac{c}{t^{\nu/2}} e^{-\frac{c^2}{10t}}.$$

It follows easily that there is a certain constant C such that

$$\forall t \in]0, 1] : \|\nabla e^{-t\Delta}\|_{L^p(\Omega) \rightarrow L^\infty(K)} \leq C.$$

Now assume that $t > 1$:

$$\|\nabla e^{-t\Delta}\|_{L^p(\Omega) \rightarrow L^\infty(K)} \leq \left\| \nabla e^{-\frac{1}{2}\Delta} \right\|_{L^\infty(M) \rightarrow L^\infty(K)} \left\| e^{-(t-\frac{1}{2})\Delta} \right\|_{L^p(\Omega) \rightarrow L^\infty(M)}.$$

But we have

$$\left\| e^{-(t-\frac{1}{2})\Delta} \right\|_{L^p(\Omega) \rightarrow L^\infty(M)} \leq \frac{C}{(t-1/2)^{\nu/2p}}.$$

But with 2.3, we obtain :

$$\begin{aligned} \left\| \nabla e^{-\frac{1}{2}\Delta} \right\|_{L^\infty(M) \rightarrow L^\infty(K)} &\leq \sup_{x \in K} \int_M |\nabla_x H_{1/2}(x, y)| dy \\ &\leq C \sup_{x \in K} \int_M H_1(x, y) dy \leq C. \end{aligned}$$

Hence for all $t > 0$, we obtain

$$\|\nabla e^{-t\Delta}\|_{L^p(\Omega) \rightarrow L^\infty(K)} \leq \frac{C}{(1+t)^{\frac{\nu}{2p}}}$$

and the second estimate follows from the subordination identity. □

3. Proof of the main theorem

Let (M_0, g_0) be a complete Riemannian manifold, we assume that the Ricci curvature of (M_0, g_0) is bounded from below and that for some $\nu > 3$ and $C > 0$, that (M, g) satisfies the Sobolev inequality

$$\forall \varphi \in C_0^\infty(M_0), \|\varphi\|_{L^{\frac{2\nu}{\nu-2}}} \leq C \|d\varphi\|_{L^2}.$$

We assume that on (M_0, g_0) the Riesz transform is bounded on L^p for some $p \in]\nu/(\nu-1), \nu[$. And we consider M a complete Riemannian manifold such that outside compact sets $K \subset M$ and $K_0 \subset M_0$, $M \setminus K$ is isometric to $M_0 \setminus K_0$, the case where $M \setminus K$ is isometric to several copies of $M_0 \setminus K_0$ can

be done similarly by considering the disjoint union of k copies of M_0 . We are going to prove that on M the Riesz transform is also bounded on L^p . The first step is to build a good parametrix for the Poisson operator on M . The first problem is that the operator $\sqrt{\Delta}$ is not a differential operator, we circumvent these difficulties by working on $\mathbb{R}_+ \times M$. As a matter of fact, the Poisson operator solves the Dirichlet problem :

$$(3.1) \quad \begin{cases} \left(-\frac{\partial^2}{\partial \sigma^2} + \Delta\right) u(\sigma, x) = 0 & \text{on }]0, \infty[\times M \\ u(0, x) = u(x) \\ \lim_{\sigma \rightarrow \infty} u(\sigma, \cdot) = 0. \end{cases}$$

The construction of the parametrix will be standard, the non locality nature of the operator $\sqrt{\Delta}$ implies that we can not use the Duhamel formula, instead we used the Green operator. The idea is to find $E_\sigma(u)$ an approximate solution for (3.1) and then to use the fact that, if G is the Green operator of the operator $-\frac{\partial^2}{\partial \sigma^2} + \Delta$ for the Dirichlet boundary condition, then

$$e^{-\sigma\sqrt{\Delta}}u = E_\sigma(u) - G\left(-\frac{\partial^2}{\partial \sigma^2} + \Delta\right)E_\sigma(u).$$

3.1. The parametrix construction

Let \tilde{K} be another compact set in M containing K in its interior. We identify

$$\Omega = M \setminus K = M_0 \setminus K_0.$$

Let ρ_0, ρ_1 a smooth partition of unity such that

$$\text{supp } \rho_0 \subset \Omega \quad \text{and} \quad \text{supp } \rho_1 \subset \tilde{K},$$

let also φ_0, φ_1 be smooth functions, such that

$$\text{supp } \varphi_0 \subset \Omega \quad \text{and} \quad \text{supp } \varphi_1 \subset \tilde{K}$$

Moreover we require that $\varphi_i = 1$ on a neighborhood of the support of ρ_i so that we have :

$$\varphi_i \rho_i = \rho_i.$$

Let Δ_1 be the realization of the Laplace operator on \tilde{K} for the Dirichlet boundary condition and let Δ_0 be the Laplace operator on M_0 . Let $e^{-\sigma\sqrt{\Delta_i}}$ their associated Poisson operator then we define for $u \in L^p(M)$:

$$E_\sigma(u) = \sum_{i=0}^1 \varphi_i(e^{-\sigma\sqrt{\Delta_i}} \rho_i u),$$

where we think of $\rho_0 u$ as a function on $\Omega \subset M_0$ and of $\varphi_0(e^{-\sigma\sqrt{\Delta_0}}\rho_0 u)$ as a function on $\Omega \subset M$.

We can easily compute :

$$\left(-\frac{\partial^2}{\partial\sigma^2} + \Delta\right) E_\sigma(u) = \sum_{i=0}^1 [\Delta, \varphi_i](e^{-\sigma\sqrt{\Delta_i}}\rho_i u) = f(\sigma, x) = \sum_{i=0}^1 f_i(\sigma, x),$$

where

$$(3.2) \quad \begin{aligned} f_i(\sigma, x) &= [\Delta, \varphi_i](e^{-\sigma\sqrt{\Delta_i}}\rho_i u)(x) \\ &= \Delta\varphi_i(x)(e^{-\sigma\sqrt{\Delta_i}}\rho_i u)(x) - 2\left\langle d\varphi_i(x), d(e^{-\sigma\sqrt{\Delta_i}}\rho_i u)(x) \right\rangle. \end{aligned}$$

From lemma 2.4 and the fact that the support of $d\varphi_0$ and ρ_0 are disjoint, we easily get that for all $\sigma \geq 0$:

$$(3.3) \quad \|f_0(\sigma)\|_{L^1} + \|f_0(\sigma)\|_{L^p} \leq \frac{C}{(1 + \sigma)^{\nu/p}} \|\rho_0 u\|_{L^p}.$$

Let us explain why this estimate also holds for f_1 . Note that the operator

$$\mathcal{S}(\sigma) = [\Delta, \varphi_1]e^{-\sigma\sqrt{\Delta_1}}\rho_1$$

is an operator with smooth Schwarz kernel and compact support, moreover because the corresponding estimate of the lemma (2.4) also holds for $\sigma \in [0, 1]$ on a compact manifold, the Schwarz kernel of $\mathcal{S}(\sigma)$ is uniformly bounded when $\sigma \rightarrow 0$. Hence there is a constant C such that

$$\forall \sigma \in [0, 1], \|\mathcal{S}(\sigma)u\|_{L^\infty} \leq C\|\rho_1 u\|_{L^p}.$$

Now the operator Δ_1 has a spectral gap on L^p (its L^p spectrum is also its L^2 spectrum), hence there is a constant C such that for all $\sigma \geq 0$ then

$$\|e^{-\sigma\sqrt{\Delta_1}}\|_{L^p \rightarrow L^p} \leq Ce^{-\sigma/C}.$$

Hence for $\sigma \geq 1$:

$$\begin{aligned} \|\mathcal{S}(\sigma)u\|_{L^\infty} &\leq \|[\Delta, \varphi_1]e^{\frac{1}{2}\sqrt{\Delta_1}}\|_{L^p \rightarrow L^\infty} \|e^{-(\sigma-1/2)\sqrt{\Delta_1}}\rho_1 u\|_{L^p} \\ &\leq Ce^{-\sigma/C} \|\rho_1 u\|_{L^p}. \end{aligned}$$

The result follows by noticing that the f_i 's have compact support in $\tilde{K} \setminus K$. Eventually we obtain the estimate :

LEMMA 3.1. — *When $u \in L^p(M)$ and we define an operator S_σ by $S_\sigma u = f = f_0 + f_1$ where f_0, f_1 are defined by (3.2) then*

$$\forall \sigma \geq 0, \|S_\sigma(u)\|_{L^1} + \|S_\sigma(u)\|_{L^p} \leq \frac{C}{(1 + \sigma)^{\nu/p}} \|u\|_{L^p}.$$

3.2. The Riesz transform on M

We introduce now G , the Green operator of the operator $(-\frac{\partial^2}{\partial \sigma^2} + \Delta)$ on $\mathbb{R}_+ \times M$ for the Dirichlet boundary condition. Its Schwarz kernel is given by

$$G(\sigma, s, x, y) = \int_0^\infty \left[\frac{e^{-\frac{(\sigma-s)^2}{4t}} - e^{-\frac{(\sigma+s)^2}{4t}}}{\sqrt{4\pi t}} \right] H_t(x, y) dt$$

where H_t is the heat kernel on M and

$$\frac{e^{-\frac{(\sigma-s)^2}{4t}} - e^{-\frac{(\sigma+s)^2}{4t}}}{\sqrt{4\pi t}}$$

the heat kernel on the half-line \mathbb{R}_+ for the Dirichlet boundary condition. We have

$$e^{-\sigma\sqrt{\Delta}}u = E_\sigma(u) - G(S_\sigma(u)).$$

Hence

$$\begin{aligned} \Delta^{-1/2}u &= \int_0^\infty e^{-\sigma\sqrt{\Delta}}u d\sigma = \sum_{i=0}^1 \varphi_i \Delta_i^{-1/2} \rho_i u \\ &\quad - \int_{\mathbb{R}_+^2 \times M} G(\sigma, s, x, y) f(s, y) d\sigma ds dy. \end{aligned}$$

Let

$$g(x) = \int_{\mathbb{R}_+^2 \times M} G(\sigma, s, x, y) f(s, y) d\sigma ds dy$$

then we have

$$(3.4) \quad \Delta^{-1/2}u = \sum_{i=0}^1 \varphi_i \Delta_i^{-1/2} \rho_i u - g.$$

But

$$\begin{aligned} \int_0^\infty G(\sigma, s, x, y) d\sigma &= \frac{1}{\sqrt{4\pi}} \int_0^\infty \left[\int_{-s}^s e^{-\frac{v^2}{4t}} dv \right] H_t(x, y) \frac{dt}{\sqrt{t}} \\ &= \frac{2}{\sqrt{\pi}} \int_0^\infty e^{-r^2} \left[\int_0^{\frac{s^2}{4r^2}} H_t(x, y) dt \right] dr. \end{aligned}$$

It follows from the above computation that

$$g(x) = \frac{2}{\sqrt{\pi}} \int_{\mathbb{R}_+^2} e^{-r^2} \left[\int_0^{\frac{s^2}{4r^2}} (e^{-t\Delta} f(s))(x) dt \right] dr ds.$$

The following lemma is now the last crucial estimate :

LEMMA 3.2. — *There is a constant C such that*

$$\|\Delta g\|_{L^p} + \|g\|_{L^p} \leq C\|u\|_{L^p}.$$

Proof. — Recall that according to [4], (M, g) itself satisfies the same Sobolev inequality :

$$\forall \varphi \in C_0^\infty(M), \|\varphi\|_{L^{\frac{2\nu}{\nu-2}}} \leq C\|d\varphi\|_{L^2}.$$

Hence the heat operator satisfies the following mapping properties : for $1 \leq q \leq p \leq +\infty$ we have

$$\|e^{-t\Delta}\|_{L^q \rightarrow L^p} \leq \frac{C}{t^{\frac{\nu}{2}(\frac{1}{q}-\frac{1}{p})}}.$$

As a consequence, for all $t \in [0, 1]$, then

$$\|(e^{-t\Delta} f(s))\|_{L^p} \leq \|f(s)\|_{L^p} \leq \frac{C}{(1+s)^{\nu/p}} \|u\|_{L^p}$$

and if $t > 1$, then

$$\|(e^{-t\Delta} f(s))\|_{L^p} \leq \|e^{-t\Delta}\|_{L^1 \rightarrow L^p} \|f(s)\|_{L^1} \leq \frac{1}{t^{\frac{\nu}{2}(1-\frac{1}{p})}} \frac{C}{(1+s)^{\nu/p}} \|u\|_{L^p}.$$

Hence

$$\begin{aligned} \|g\|_{L^p} &\leq \frac{2}{\sqrt{\pi}} \int_{\mathbb{R}_+^2} e^{-r^2} \left[\int_0^{\frac{s^2}{4r^2}} \|(e^{-t\Delta} f(s))\|_{L^p} dt \right] ds dr \\ &\leq \frac{2}{\sqrt{\pi}} \left(\int_{\mathbb{R}_+^2} e^{-r^2} \left[\int_0^{\frac{s^2}{4r^2}} \frac{C}{\max\left(1, t^{\frac{\nu}{2}(1-\frac{1}{p})}\right)} \frac{1}{(1+s)^{\nu/p}} dt \right] ds dr \right) \|u\|_{L^p}. \end{aligned}$$

But because $p < \nu$, we have

$$\begin{aligned} &\int_{\{2r\sqrt{t} \leq s\}} e^{-r^2} \frac{1}{\max\left(1, t^{\frac{\nu}{2}(1-\frac{1}{p})}\right)} \frac{1}{(1+s)^{\nu/p}} ds dt dr \\ &= \frac{\nu}{\nu-p} \int_{\mathbb{R}_+^2} e^{-r^2} \frac{1}{\max\left(1, t^{\frac{\nu}{2}(1-\frac{1}{p})}\right)} \frac{1}{(1+2r\sqrt{t})^{\nu/p-1}} dt dr \end{aligned}$$

and this integral is finite exactly when $p > \nu/(\nu - 1)$ and $\nu > 3$.

It remains to estimate $\|\Delta g\|_{L^p}$, which is easier because

$$\begin{aligned} \Delta g &= \frac{2}{\sqrt{\pi}} \int_{\mathbb{R}_+^2} e^{-r^2} \left[\int_0^{\frac{s^2}{4r^2}} \Delta(e^{-t\Delta} f(s)) dt \right] dr ds \\ &= -\frac{2}{\sqrt{\pi}} \int_{\mathbb{R}_+^2} e^{-r^2} \left[\int_0^{\frac{s^2}{4r^2}} \frac{d}{dt} (e^{-t\Delta} f(s)) dt \right] dr ds \\ &= \frac{2}{\sqrt{\pi}} \int_{\mathbb{R}_+^2} e^{-r^2} \left[f(s) - (e^{-\frac{s^2}{4r^2}\Delta} f(s)) \right] dr ds. \end{aligned}$$

Hence

$$\begin{aligned} \|\Delta g\|_{L^p} &\leq \frac{4}{\sqrt{\pi}} \int_{\mathbb{R}_+^2} e^{-r^2} \|f(s)\|_{L^p} dr ds \\ &\leq \frac{4}{\sqrt{\pi}} \left(\int_{\mathbb{R}_+^2} e^{-r^2} \frac{C}{(1+s)^{\nu/p}} dr ds \right) \|u\|_{L^p}. \end{aligned}$$

□

Now we can finish the proof of the main theorem : let T_i be the Riesz transform associated with the operator Δ_i . With the formula (3.4), we obtain

$$d\Delta^{-1/2}u = \sum_{i=0}^1 \varphi_i T_i \rho_i u + \sum_{i=0}^1 d\varphi_i (\Delta_i^{-1/2} \rho_i u) - dg.$$

By hypothesis, T_0 is bounded on L^p . Moreover since $\varphi_1 T_1 \rho_1$ is a pseudo differential operator of order 0 with compact support it is also bounded on L^p . The operator $d\varphi_1 (\Delta_i^{-1/2} \rho_1 u)$ has a smooth kernel with compact support, hence it is bounded on L^p . Moreover, the Sobolev inequality

$$\forall \varphi \in \mathbb{C}_0^\infty(M_0), \|\varphi\|_{L^{\frac{2\nu}{\nu-2}}} \leq C \|d\varphi\|_{L^2}.$$

also implies the following mapping properties of the $\Delta_0^{-1/2}$ ([18]) :

$$\left\| \Delta_0^{-1/2} \right\|_{L^p \rightarrow L^{\frac{p\nu}{\nu-p}}} \leq C.$$

Hence

$$\begin{aligned} \left\| d\varphi_0 (\Delta_0^{-1/2} \rho_0 u) \right\|_{L^p} &\leq C \|\Delta_0^{-1/2} \rho_0 u\|_{L^p(\tilde{K} \setminus K)} \leq C' \|\Delta_0^{-1/2} \rho_0 u\|_{L^{\frac{p\nu}{\nu-p}}(\tilde{K})} \\ &\leq C \|\rho_0 u\|_{L^p}. \end{aligned}$$

Moreover the lemmas (3.2) and (2.1) imply that

$$\|dg\|_{L^p} \leq C \|u\|_{L^p}.$$

All these estimates yield the fact that the Riesz transform is bounded on L^p .

3.3. A comment on manifolds with non negative Ricci curvature

The proof of theorem 1.3 is fairly general, we can easily make a list of the properties which makes it runs ; let (M_i, g_i) $i = 1, \dots, b$ be complete Riemannian manifolds and let (M, g) be isometric at infinity to the disjoint union $M_1 \cup \dots \cup M_b$. That is to say there are compact sets $K \subset M$, $K_i \subset M_i$ such that $M \setminus K$ is isometric to $(M_1 \setminus K_1) \cup \dots \cup (M_b \setminus K_b)$. Let $\tilde{K} \subset \hat{K}$ such that \tilde{K} (resp. \hat{K}) contains K in its interior (resp. \tilde{K}). And let $\hat{K}_i, \tilde{K}_i \subset M_i$ such that :

$$M \setminus \tilde{K} \simeq (M_1 \setminus \tilde{K}_1) \cup \dots \cup (M_b \setminus \tilde{K}_b), \quad M \setminus \hat{K} \simeq (M_1 \setminus \hat{K}_1) \cup \dots \cup (M_b \setminus \hat{K}_b),$$

let Δ_i be the Laplace operator on M_i . We assume that on each M_i , the Ricci curvature is bounded from below such that on each M_i and M , we get the estimate induced by the Sobolev inequality (2.1). Assume that for some functions $f, g : \mathbb{R}_+ \rightarrow \mathbb{R}_+^*$ we have the estimate :

$$\left\| e^{-\sigma\sqrt{\Delta_i}} \right\|_{L^p(M_i \setminus \hat{K}_i) \rightarrow L^\infty(\tilde{K}_i)} + \left\| \nabla e^{-\sigma\sqrt{\Delta_i}} \right\|_{L^p(M_i \setminus \hat{K}_i) \rightarrow L^\infty(\tilde{K}_i)} \leq \frac{1}{f(\sigma)},$$

and that on the manifold M :

$$\left\| e^{-t\Delta} \right\|_{L^1(\hat{K}) \rightarrow L^p(M)} \leq \frac{1}{g(t)}.$$

with

$$(3.5) \quad \int_0^\infty \frac{ds}{f(s)} < \infty$$

$$(3.6) \quad \int_{\mathbb{R}_+^2} e^{-u^2} \min\left(1, \frac{1}{g(t)}\right) \left[\int_{2u\sqrt{t}}^\infty \frac{ds}{f(s)} \right] dudt < \infty.$$

Then if for all i , the Riesz transform $T_i := d\Delta_i^{-1/2}$ is bounded on L^p , then on M , the Riesz transform is also bounded on L^p .

A natural and well study class of manifolds satisfying such estimates are manifolds satisfying the so called relative Faber-Krahn inequality : for some $\alpha > 0$ and $c > 0$, we have :

$$\forall B(x, R), \forall \Omega \subset B(x, R), \lambda_1(\Omega) \geq \frac{c}{R^2} \left(\frac{\text{vol } \Omega}{\text{vol } B(x, R)} \right)^{-\alpha}$$

where

$$\lambda_1(\Omega) = \inf_{f \in C_0^\infty(\Omega)} \frac{\int_\Omega |df|^2}{\int_\Omega f^2}$$

is the first eigenvalue of the Laplace operator on Ω for the Dirichlet boundary condition. According to A. Grigor'yan [11] this inequality is equivalent

to the conjunction of the doubling property : uniformly in x and $R > 0$ we have

$$\frac{\text{vol } B(x, 2R)}{\text{vol } B(x, R)} \leq C$$

and of the upper bound on the heat operator

$$H_t(x, y) \leq \frac{C}{\text{vol } B(x, \sqrt{t})} e^{-\frac{d(x,y)^2}{5t}}.$$

Manifolds with non negative Ricci curvature are examples of manifolds satisfying this relative Faber-Krahn inequalities.

Assume that each M_i satisfies this relative Faber-Krahn inequality and if we assume that for $i = 1, \dots, b$, there is a point $o_i \in K_i$ and all $R \geq 1$

$$\text{vol } B(o_i, R) := V_i(R) \geq CR^\nu$$

then we get easily from the subordination identity :

$$\left\| e^{-\sigma\sqrt{\Delta_i}} \right\|_{L^p(M_i \setminus \widehat{K}_i) \rightarrow L^\infty(\widehat{K}_i)} + \left\| \nabla e^{-\sigma\sqrt{\Delta_i}} \right\|_{L^p(M_i \setminus \widehat{K}_i) \rightarrow L^\infty(\widehat{K}_i)} \leq \frac{1}{(1 + \sigma)^{\nu/p}}.$$

Now the problem comes from the fact that we don't know how to obtain a relative Faber-Krahn inequality on M from the one we assume on the M_i 's. However, recently in (page 877 of [13]), A. Grigor'yan and L. Saloff-Coste have announced the following very useful result (see also [12]) : when the M_i 's satisfy the relative Faber-Krahn inequality then

$$\forall B(x, R) \subset M, \forall \Omega \subset B(x, R), \quad \lambda_1(\Omega) \geq \frac{c}{R^2} \left(\frac{\text{vol } \Omega}{\mu(x, R)} \right)^{-\alpha},$$

where

$$\mu(x, R) = \begin{cases} \text{vol } B(x, R) & \text{if } B(x, R) \subset M \setminus K \\ \inf_i V_i(R) & \text{else.} \end{cases}$$

Hence from our volume growth estimate, we will obtain (see [11]) when $t \geq 1$:

$$\left\| e^{-t\Delta} \right\|_{L^1(\widehat{K}) \rightarrow L^p(M)} \leq \frac{C}{t^{\frac{\nu}{2} \frac{p-1}{p}}}.$$

With this result of A. Grigor'yan and L. Saloff-Coste and with the result of D. Bakry [3], we will obtain :

PROPOSITION 3.3. — *Let $(M_1, g_1), \dots, (M, g_b)$ be complete Riemannian manifolds with non negative Ricci curvature. Assume that on all M_i 's we have the volume growth lower bound :*

$$\text{vol } B(o_i, R) \geq CR^\nu.$$

Then assume that $\nu > 3$ and $p \in]\nu/(\nu - 1), \nu[$ then on any manifold isotropic at infinity to the disjoint union of the M_i 's, the Riesz transform is bounded on L^p .

BIBLIOGRAPHY

- [1] G. ALEXOPOULOS, "An application of homogenization theory to harmonic analysis: Harnack inequalities and Riesz transforms on Lie groups of polynomial growth", *Canad. J. Math.* **44** (1992), p. 691-727.
- [2] J.-P. ANKER, "Sharp estimates for some functions of the Laplacian on noncompact symmetric spaces", *Duke Math. J.* **65** (1992), no. 2, p. 257-297.
- [3] D. BAKRY, "Étude des transformations de Riesz dans les variétés riemanniennes à courbure de Ricci minorée", *Lecture Notes in Math.* **1247** (1987), p. 137-172, Séminaire de Probabilités, XXI.
- [4] G. CARRON, "Une suite exacte en L^2 -cohomologie", *Duke Math. J.* **95** (1998), p. 343-372.
- [5] G. CARRON, T. COULHON & A. HASSELL, "Riesz transform for manifolds with Euclidean ends", to appear in *Duke Math. Journal*.
- [6] T. COULHON & N. DUNGEY, "Riesz transform and perturbation", preprint, 2006.
- [7] T. COULHON & X. T. DUONG, "Riesz transforms for $1 \leq p \leq 2$ ", *Trans. Amer. Math. Soc.* **351** (1999), p. 1151-1169.
- [8] ———, "Riesz transform and related inequalities on non-compact Riemannian manifolds", *Comm. in Pure and Appl. Math.* **56** (2003), no. 12, p. 1728-1751.
- [9] T. COULHON, L. SALOFF-COSTE & N. T. VAROPOULOS, "Analysis and geometry on groups", in *Cambridge Tracts in Mathematics*, vol. 100, Cambridge University Press, Cambridge, 1992.
- [10] E. B. DAVIES, "Pointwise bounds on the space and time derivatives of heat kernels", *Operator Theory* **21** (1989), no. 2, p. 367-378.
- [11] A. GRIGOR'YAN, "Heat kernel upper bounds on a complete non-compact manifold", *Rev. Mat. Iberoamericana* **10** (1994), no. 2, p. 395-452.
- [12] A. GRIGOR'YAN & L. SALOFF-COSTE, "Heat kernel on connected sums of Riemannian manifolds", *Math. Res. Lett.* **6** (1999), no. 3-4, p. 307-321.
- [13] ———, "Stability results for Harnack inequalities", *Ann. Inst. Fourier* **55** (2005), no. 3, p. 825-890.
- [14] E. HEBEY, "Optimal Sobolev inequalities on complete Riemannian manifolds with Ricci curvature bounded below and positive injectivity radius", *Amer. J. Math.* **118** (1996), no. 2, p. 291-300.
- [15] H. KOMATSU, "Fractional powers of operators", *Pacific J. Math.* **19** (1966), p. 285-346.
- [16] H.-Q. LI, "La transformation de Riesz sur les variétés coniques", *J. Funct. Anal.* **168** (1999), no. 1, p. 145-238.
- [17] N. LOHOUE, "Comparaison des champs de vecteurs et des puissances du laplacien sur une variété riemannienne à courbure non positive", *J. Funct. Anal.* **61** (1985), no. 2, p. 164-201.

- [18] N. T. VAROPOULOS, “Hardy-Littlewood theory for semigroups”, *J. Funct. Anal.* **63** (1985), no. 2, p. 240-260.

Manuscrit reçu le 23 juin 2006,
accepté le 30 novembre 2006.

Gilles CARRON
Université de Nantes
Laboratoire de Mathématiques Jean Leray
(UMR 6629)
2, rue de la Houssinière
B.P. 92208
44322 Nantes Cedex 3 (France)
Gilles.Carron@math.univ-nantes.fr