



ANNALES

DE

L'INSTITUT FOURIER

Peter SJÖGREN & Maria VALLARINO

Boundedness from H^1 to L^1 of Riesz transforms on a Lie group of exponential growth

Tome 58, n° 4 (2008), p. 1117-1151.

http://aif.cedram.org/item?id=AIF_2008__58_4_1117_0

© Association des Annales de l'institut Fourier, 2008, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

BOUNDEDNESS FROM H^1 TO L^1 OF RIESZ TRANSFORMS ON A LIE GROUP OF EXPONENTIAL GROWTH

by Peter SJÖGREN & Maria VALLARINO (*)

ABSTRACT. — Let G be the Lie group $\mathbb{R}^2 \rtimes \mathbb{R}^+$ endowed with the Riemannian symmetric space structure. Let X_0, X_1, X_2 be a distinguished basis of left-invariant vector fields of the Lie algebra of G and define the Laplacian $\Delta = -(X_0^2 + X_1^2 + X_2^2)$. In this paper we consider the first order Riesz transforms $R_i = X_i \Delta^{-1/2}$ and $S_i = \Delta^{-1/2} X_i$, for $i = 0, 1, 2$. We prove that the operators R_i , but not the S_i , are bounded from the Hardy space H^1 to L^1 . We also show that the second-order Riesz transforms $T_{ij} = X_i \Delta^{-1} X_j$ are bounded from H^1 to L^1 , while the transforms $S_{ij} = \Delta^{-1} X_i X_j$ and $R_{ij} = X_i X_j \Delta^{-1}$, for $i, j = 0, 1, 2$, are not.

RÉSUMÉ. — On considère le groupe de Lie $G = \mathbb{R}^2 \rtimes \mathbb{R}^+$ muni de la structure Riemannienne d'espace symétrique. On choisit une base X_0, X_1, X_2 de champs vectoriels invariants à gauche de l'algèbre de Lie de G et on définit le Laplacien $\Delta = -(X_0^2 + X_1^2 + X_2^2)$. Dans cet article nous considérons les transformées de Riesz du premier ordre $R_i = X_i \Delta^{-1/2}$ et $S_i = \Delta^{-1/2} X_i$, avec $i = 0, 1, 2$. Nous prouvons que les opérateurs R_i , mais non pas les S_i , sont bornés de l'espace de Hardy H^1 à L^1 . Nous démontrons aussi que les transformées de Riesz du deuxième ordre $T_{ij} = X_i \Delta^{-1} X_j$ sont bornées de H^1 à L^1 , tandis que les transformées $S_{ij} = \Delta^{-1} X_i X_j$ et $R_{ij} = X_i X_j \Delta^{-1}$, $i, j = 0, 1, 2$, ne sont pas bornées.

1. Introduction

Let G be the Lie group $\mathbb{R}^2 \rtimes \mathbb{R}^+$ where the product rule is the following:

$$(x_1, x_2, a) \cdot (x'_1, x'_2, a') = (x_1 + a x'_1, x_2 + a x'_2, a a')$$

Keywords: Singular integrals, Riesz transforms, Hardy space, Lie groups, exponential growth.

Math. classification: 43A80, 42B20, 42B30, 22E30.

(*) Work partially supported by the European Commission via the Network HARP, "Harmonic analysis and related problems".

for $(x_1, x_2, a), (x'_1, x'_2, a') \in G$. The group G is not unimodular; the right and left Haar measures are given by

$$d\rho(x_1, x_2, a) = a^{-1} dx_1 dx_2 da \quad \text{and} \quad d\lambda(x_1, x_2, a) = a^{-3} dx_1 dx_2 da,$$

respectively. The modular function is thus $\delta(x_1, x_2, a) = a^{-2}$. Throughout this paper, unless explicitly stated, we consider the right measure ρ on G and we denote by $L^p, \|\cdot\|_p$ and $\langle \cdot, \cdot \rangle$ the L^p -space, the L^p -norm and the L^2 -scalar product with respect to the measure ρ .

The group G has a Riemannian symmetric space structure, and the corresponding metric, which we denote by d , is that of the three-dimensional hyperbolic half-space. The metric d is invariant under left translation and it is given by

$$(1.1) \quad \cosh r(x_1, x_2, a) = \frac{a + a^{-1} + a^{-1}(x_1^2 + x_2^2)}{2} \quad \forall (x_1, x_2, a) \in G,$$

where $r(x_1, x_2, a) = d((x_1, x_2, a), e)$ denotes the distance of the point (x_1, x_2, a) from the identity $e = (0, 0, 1)$ of G . It is easy to verify that if $r(x_1, x_2, a) < 1$, then $r(x_1, x_2, a) \sim |(x_1, x_2, \log a)|$, where $|\cdot|$ denotes the euclidean norm in \mathbb{R}^3 . The measure of a hyperbolic ball B_r , centered at the identity and of radius r , behaves like

$$\lambda(B_r) = \rho(B_r) \sim \begin{cases} r^3 & \text{if } r < 1 \\ e^{2r} & \text{if } r \geq 1. \end{cases}$$

Thus G is a group of exponential growth. In this context, the classical Calderón–Zygmund theory and the classical definition of the atomic Hardy space H^1 (see [8, 23]) do not apply. Recently W. Hebisch and T. Steger [17] constructed a new Calderón–Zygmund theory which holds in some spaces of exponential growth, in particular in the space (G, d, ρ) defined above. The main idea is to replace the family of balls which is used in the classical Calderón–Zygmund theory by a suitable family of parallelepipeds which we call *Calderón–Zygmund sets*. The definition appears in [17] and implicitly in [16], and reads as follows.

DEFINITION 1.1. — *A Calderón–Zygmund set is a parallelepiped $R = [b_1 - L/2, b_1 + L/2] \times [b_2 - L/2, b_2 + L/2] \times [ae^{-r}, ae^r]$, where $L > 0, r > 0$ and $a \in \mathbb{R}^+$ are related by*

$$\begin{aligned} e^2 a r \leq L < e^8 a r & \quad \text{if } r < 1, \\ a e^{2r} \leq L < a e^{8r} & \quad \text{if } r \geq 1. \end{aligned}$$

We let \mathcal{R} denote the family of all Calderón–Zygmund sets, and observe that \mathcal{R} is invariant under left translation. Given $R \in \mathcal{R}$, we define its dilated set as $R^* = \{x \in G : d(x, R) < r\}$. There exists a constant C_0 such that $\rho(R^*) \leq C_0 \rho(R)$ and $R \subset B((b_1, b_2, a), C_0 r)$.

In [17] it is proved that every integrable function on G admits a Calderón–Zygmund decomposition involving the family \mathcal{R} , and that a new Calderón–Zygmund theory can be developed in this context. This makes it natural to introduce an atomic Hardy space H^1 on the group G , as follows (see [24] for details).

We define an *atom* as a function a in L^1 such that

- (i) a is supported in a Calderón–Zygmund set R ;
- (ii) $\|a\|_\infty \leq \rho(R)^{-1}$;
- (iii) $\int a \, d\rho = 0$.

The atomic Hardy space is now defined in a standard way.

DEFINITION 1.2. — *The Hardy space H^1 is the space of all functions f in L^1 which can be written as $f = \sum_j \lambda_j a_j$, where a_j are atoms and λ_j are complex numbers such that $\sum_j |\lambda_j| < \infty$. We denote by $\|f\|_{H^1}$ the infimum of $\sum_j |\lambda_j|$ over such decompositions.*

The new Calderón–Zygmund theory is used to study the boundedness of some singular integral operators related to a distinguished Laplacian on G , which is defined as follows.

Let X_0, X_1, X_2 denote the left-invariant vector fields

$$X_0 = a \partial_a \quad X_1 = a \partial_{x_1} \quad X_2 = a \partial_{x_2},$$

which span the Lie algebra \mathfrak{g} of G . The Laplacian $\Delta = -(X_0^2 + X_1^2 + X_2^2)$ is a left-invariant operator which is essentially selfadjoint on $L^2(\rho)$. Since Δ is positive definite and one-to-one [13], its powers Δ^α , $\alpha \in \mathbb{R}$, have dense domains and are selfadjoint. This makes it possible to form the Riesz transforms of the first order associated with Δ , defined by

$$(1.2) \quad R_i = X_i \Delta^{-1/2} \quad \text{and} \quad S_i = \Delta^{-1/2} X_i, \quad i = 0, 1, 2,$$

and the Riesz transforms of the second order, defined by

$$(1.3) \quad R_{ij} = X_i X_j \Delta^{-1} \quad \text{and} \quad S_{ij} = \Delta^{-1} X_i X_j \quad \text{and} \quad T_{ij} = X_i \Delta^{-1} X_j,$$

for $i, j = 0, 1, 2$. The boundedness properties of the Riesz transforms associated with the distinguished Laplacian Δ defined above have been considered by many authors. Actually, some results in the literature concern

the Riesz transforms associated with a distinguished right-invariant Laplacian Δ^r , which is related to Δ as follows. Let $X_i^r, i = 0, 1, 2$, be the right-invariant vector fields on G which agree with X_i at the identity, i.e.,

$$X_0^r = x_1 \partial_{x_1} + x_2 \partial_{x_2} + a \partial_a \quad X_1^r = \partial_{x_1} \quad X_2^r = \partial_{x_2} .$$

It is well known that $X_i^r f = (X_i \check{f})^\vee$ for any $f \in C^\infty(G)$, where $\check{f}(x) = f(x^{-1})$ for $x \in G$. The Laplacian $\Delta^r = -(X_0^r)^2 - (X_1^r)^2 - (X_2^r)^2$ is a right-invariant operator which is essentially selfadjoint on $L^2(\lambda)$. We denote by $R_i^r, S_i^r, R_{ij}^r, S_{ij}^r, T_{ij}^r$ the Riesz transforms defined as above by using the right-invariant vector fields and the right-invariant Laplacian instead of the left-invariant ones. It is easy to see that for any $f \in C_c^\infty(G)$, $\Delta^r f = (\Delta \check{f})^\vee$,

$$R_i^r f = (R_i \check{f})^\vee, \quad S_i^r f = (S_i \check{f})^\vee$$

and

$$R_{ij}^r f = (R_{ij} \check{f})^\vee, \quad S_{ij}^r f = (S_{ij} \check{f})^\vee, \quad T_{ij}^r f = (T_{ij} \check{f})^\vee .$$

Since $f \rightarrow \check{f}$ is an isometry between $L^p(\lambda)$ and $L^p(\rho)$ for $p \in [1, \infty]$, results concerning the boundedness of the right-invariant Riesz transforms with respect to the left Haar measure λ may be reformulated in terms of the left-invariant Riesz transforms with respect to the right Haar measure ρ . We now summarize some results formulated in terms of the left-invariant Riesz transforms defined by (1.2) and (1.3).

In [15, 22] G. Gaudry and P. Sjögren studied Riesz transforms of the type $X\Delta^{-1/2}$ and $\Delta^{-1/2}X$, where Δ is a distinguished Laplacian and X is a distinguished vector field, in the context of the group $\mathbb{R} \ltimes \mathbb{R}^+$, also known as affine group of the real line. They proved that these operators are of weak type 1 and bounded on L^p , for $1 < p < \infty$. In the sequel we sometimes refer to their papers: even if their setting is different, their arguments may be applied also to our context with some slight changes, and so their results carry over.

Hebisch and Steger then proved that all the operators R_i are of weak type 1 and bounded on L^p when $1 < p \leq 2$ [17, Theorem 6.4]. This result was obtained as an application of the Calderón-Zygmund theory on the group G .

The operators S_i are bounded on L^2 , for $i = 0, 1, 2$. For $i \neq 0$ the operators S_i are of weak type 1 and bounded on L^p when $1 < p \leq 2$, while the operator S_0 is not of weak type 1 but bounded on L^p for $1 < p \leq 2$ (Hebisch, private communication).

Since R_i and S_i are bounded on L^p for $p < 2$, it follows by duality that R_i and S_i are also bounded on L^p when $2 < p < \infty$.

The second-order Riesz transforms have been studied first in [13] in the context of the affine group of the real line, then in [14] in the general setting of NA groups of rank 1, including the group G . The operators T_{ij} are of weak type 1 and bounded on L^p when $1 < p < \infty$. The operators R_{ij} and S_{ij} are not of weak type p , for any $1 \leq p < \infty$.

In this paper we study the $H^1 - L^1$ boundedness of the Riesz transforms on the group G . Our main results are the following:

- (1) the operators R_i , $i = 0, 1, 2$, are bounded from H^1 to L^1 (Section 3);
- (2) the operators S_i , $i = 0, 1, 2$, are not bounded from H^1 to L^1 (Sections 4, 5);
- (3) the operators T_{ij} are bounded from H^1 to L^1 (Section 7);
- (4) the operators S_{ij} and R_{ij} are not bounded from H^1 to L^1 (Sections 8, 9).

We remark that since the interpolation spaces between H^1 and L^2 for the real interpolation method are the L^p spaces for $1 < p < 2$ (see [24]), the boundedness of R_i and T_{ij} from H^1 to L^1 implies their boundedness on L^p , for $1 < p < 2$.

The Riesz operators, and in particular their boundedness on L^p and on the Hardy space H^1 , have been studied on various Lie groups and Riemannian manifolds. Many results in the literature concern “doubling spaces”, i.e., measured metric spaces where the volume of balls satisfies the doubling condition. In this context, the Hardy space H^1 is defined as in [8].

In the classical setting of \mathbb{R}^n , the Riesz transforms are bounded on L^p for $1 < p < \infty$, of weak type 1 and bounded on H^1 [23, III.3].

For nilpotent Lie groups and first-order Riesz operators, the L^p -boundedness, for $1 < p < \infty$, the weak type 1 and the H^1 -boundedness were proved by N. Lohoué and N. Varopoulos [18]. Subsequently, this was extended to all connected Lie groups of polynomial growth by L. Saloff-Coste [21] and G. Alexopoulos [1].

In the setting of symmetric spaces of noncompact type, J.-P. Anker [2] considered Riesz transforms associated with the Laplace–Beltrami operator. He proved the weak type 1 estimate for the first-order operators and the L^p -estimates for operators of arbitrary order.

On a Riemannian manifold the Riesz transform $R = \nabla \Delta^{-1/2}$, where ∇ is the gradient and Δ is the Laplace–Beltrami operator, has been considered. If the manifold has nonnegative Ricci curvature, then the Riesz transform R is bounded on L^p , $1 < p < \infty$, of weak type 1 and bounded from H^1

to L^1 [6, 7]. Subsequently, T. Coulhon and X.T. Duong proved that on a Riemannian manifold with the doubling property whose heat kernel verifies an upper estimate on the diagonal, R is of weak type 1 and bounded on L^p , for $1 < p \leq 2$ [9]. The connection between the L^p -boundedness of the Riesz transform, Poincaré inequalities and heat kernel estimates is also studied in [4, 5, 10, 11]. In Riemannian manifolds satisfying the doubling condition and the Poincaré inequality, E. Russ [20] proved that R is bounded from H^1 to L^1 ; then M. Marias and Russ [19] proved the boundedness on H^1 of the linearized Riesz transforms.

The previous results do not apply to our space (G, d, ρ) , since it is of exponential growth and the doubling condition fails.

Our paper is organized as follows: Section 2 contains an analysis of the kernels of the Riesz transforms. The $H^1 - L^1$ -boundedness of the operators R_i is proved in Section 3, as a consequence of a general boundedness theorem for integral operators. In Section 4, we prove the unboundedness from H^1 to L^1 of the operators S_1 and S_2 , and in Section 5 that of S_0 . We analyze the local part of the second-order Riesz transforms in Section 6, proving that they are bounded from H^1 to L^1 . In Section 7 we show that the operators T_{ij} are bounded from H^1 to L^1 . Finally, we show that the global part of the operators S_{ij} and R_{ij} are not bounded from H^1 to L^1 in Sections 8 and 9.

In the following, C denotes a positive, finite constant which may vary from line to line and may depend on parameters according to the context. Given two quantities f and g , by $f \sim g$ we mean that there exists a constant C such that $1/C \leq f/g \leq C$.

2. The convolution kernels of the Riesz transforms

In this section, we analyze the convolution kernels of the Riesz transforms of the first and the second order. First recall that the definition of the convolution of two functions f, g on G is

$$f * g(x) = \int_G f(xy^{-1})g(y) \, d\rho(y) \quad \forall x \in G.$$

Let V denote the space $\{\Delta u : u \in C_c^\infty(G)\}$. In [14] it is verified that V is a dense subspace of L^2 and that $V \subset D(\Delta^{-1}) \subset D(\Delta^{-1/2})$. For $\alpha > 0$, we denote by U_α the convolution kernel of $\Delta^{-\alpha/2}$, in the sense that

$\Delta^{-\alpha/2}f = f * U_\alpha$ for all $f \in V$. Since

$$\Delta^{-\alpha/2} = \frac{1}{\Gamma(\alpha/2)} \int_0^\infty t^{\alpha/2-1} e^{-t\Delta} dt,$$

we have that

$$U_\alpha = \frac{1}{\Gamma(\alpha/2)} \int_0^\infty t^{\alpha/2-1} p_t dt,$$

where p_t denotes the heat kernel of Δ . It is well known [12, Theorem 5.3, Proposition 5.4], [3, Formula (5.7)] that

$$p_t(x) = \frac{1}{8\pi^{3/2}} \delta^{1/2}(x) \frac{r(x)}{\sinh r(x)} t^{-3/2} e^{-\frac{r^2(x)}{4t}} \quad \forall x \in G,$$

where $r(x)$ denotes as before the distance of x from the identity. Hence, for $\alpha < 3$

$$\begin{aligned} U_\alpha(x) &= \frac{1}{\Gamma(\alpha/2)} \frac{1}{8\pi^{3/2}} \delta^{1/2}(x) \frac{r(x)}{\sinh r(x)} \int_0^\infty t^{\alpha/2-1} t^{-3/2} e^{-\frac{r^2(x)}{4t}} dt \\ &= \frac{1}{\Gamma(\alpha/2)} \frac{2^{1-\alpha}}{\pi^{3/2}} \delta^{1/2}(x) \frac{r(x)}{\sinh r(x)} \int_0^\infty r(x)^{\alpha-3} v^{2-\alpha} e^{-v^2} dv \\ &= C_\alpha \delta^{1/2}(x) \frac{r^{\alpha-2}(x)}{\sinh r(x)} \quad \forall x \in G. \end{aligned}$$

We consider the cases $\alpha = 1$ and $\alpha = 2$ and get that $C_1 = \frac{1}{2\pi^2}$ and $C_2 = \frac{1}{4\pi}$. We denote by $U = U_1$ the convolution kernel of $\Delta^{-1/2}$ given by

$$(2.1) \quad U(x) = \frac{1}{2\pi^2} \delta^{1/2}(x) \frac{1}{r(x) \sinh r(x)} \quad \forall x \in G,$$

and by $W = U_2$ the convolution kernel of Δ^{-1} given by

$$(2.2) \quad W(x) = \frac{1}{4\pi} \delta^{1/2}(x) \frac{1}{\sinh r(x)} \quad \forall x \in G.$$

Since $R_i = X_i \Delta^{-1/2}$, we get for all $f \in V$ and $x \in G$

$$\begin{aligned} (2.3) \quad R_i f(x) &= X_i(f * U)(x) = \int X_{i,x} f(xy^{-1}) U(y) d\rho(y) \\ &= \lim_{\varepsilon \rightarrow 0} \int_{r(y) > \varepsilon} X_{i,x} f(xy^{-1}) U(y) d\rho(y) \\ &= - \lim_{\varepsilon \rightarrow 0} \int_{r(y) > \varepsilon} X_{i,y} f(xy^{-1}) U(y) d\rho(y) \\ &= \lim_{\varepsilon \rightarrow 0} \int_{r(y) > \varepsilon} f(xy^{-1}) X_{i,y} U(y) d\rho(y), \end{aligned}$$

where the last step follows by integration by parts, as in [22, Section 3]. Thus the convolution kernel of R_i is the distribution $\text{pv } k_i$, where $k_i = X_i U$. Moreover, for $f \in V$ and $x \notin \text{supp } f$

$$\begin{aligned}
 R_i f(x) &= \int_G f(xy) k_i(y^{-1}) \, d\lambda(y) \\
 &= \int_G f(y) k_i(y^{-1}x) \delta(y) \, d\rho(y) \\
 (2.4) \qquad &= \int_G f(y) R_i(x, y) \, d\rho(y),
 \end{aligned}$$

where $R_i(\cdot, \cdot)$ denotes the integral kernel of R_i , related to k_i by

$$(2.5) \qquad R_i(x, y) = \delta(y) k_i(y^{-1}x) \quad \forall x, y \in G, \quad x \neq y.$$

We now consider the operators S_i . By arguing as in [15, page 246-247], it is easy to see that if $f \in C_c^\infty(G)$, then $X_i f \in D(\Delta^{-1/2})$, so that S_i is well defined on $C_c^\infty(G)$. Moreover, for all $f \in C_c^\infty(G)$ and $g \in V$

$$\langle S_i f, g \rangle = \langle \Delta^{-1/2} X_i f, g \rangle = \langle X_i f, \Delta^{-1/2} g \rangle = -\langle f, X_i \Delta^{-1/2} g \rangle = -\langle f, R_i g \rangle.$$

Thus by (2.5) we deduce that the integral kernel of S_i is given by

$$(2.6) \quad S_i(x, y) = -\overline{R_i(y, x)} = -\delta(x) k_i(x^{-1}y) \quad \forall x, y \in G, \quad x \neq y.$$

We now compute k_i explicitly. To do so, we shall need the following simple lemma.

LEMMA 2.1. — *At any point $(x_1, x_2, a) \neq (0, 0, 1)$ in G , the derivatives of r along the vector fields X_i are given by*

$$X_i r(x_1, x_2, a) = \begin{cases} \frac{a - a^{-1} - a^{-1}(x_1^2 + x_2^2)}{2 \sinh r(x_1, x_2, a)} = \frac{a - \cosh r}{\sinh r} & \text{if } i = 0 \\ \frac{x_i}{\sinh r(x_1, x_2, a)} & \text{if } i = 1, 2. \end{cases}$$

Proof. — It suffices to differentiate the expression

$$\cosh r(x_1, x_2, a) = \frac{a + a^{-1} + a^{-1}(x_1^2 + x_2^2)}{2},$$

with respect to X_i . For $X_0 = a \partial_a$ we obtain

$$\sinh r(x_1, x_2, a) X_0 r(x_1, x_2, a) = a \frac{1 - a^{-2} - a^{-2}(x_1^2 + x_2^2)}{2},$$

which gives the result for $i = 0$. The cases of $X_i = a \partial_{x_i}$, $i = 1, 2$, are similar. □

By (2.1) and Lemma 2.1 for $i = 1, 2$ and $(x_1, x_2, a) \neq (0, 0, 1)$ we get

$$\begin{aligned}
 k_i(x_1, x_2, a) &= X_i U(x_1, x_2, a) \\
 &= -\frac{1}{2\pi^2} a^{-1} \frac{\sinh r + r \cosh r}{r^2 \sinh^2 r} X_i r(x_1, x_2, a) \\
 (2.7) \quad &= -\frac{1}{2\pi^2} a^{-1} x_i \frac{\sinh r + r \cosh r}{r^2 \sinh^3 r}.
 \end{aligned}$$

For $i = 0$ and $(x_1, x_2, a) \neq (0, 0, 1)$ we get

$$\begin{aligned}
 k_0(x_1, x_2, a) &= X_0 U(x_1, x_2, a) \\
 &= \frac{1}{2\pi^2} \left[-a a^{-2} \frac{1}{r \sinh r} - a^{-1} \frac{\sinh r + r \cosh r}{r^2 \sinh^2 r} X_0 r(x_1, x_2, a) \right] \\
 &= \frac{1}{2\pi^2} \left[-a^{-1} \frac{1}{r \sinh r} - a^{-1} \frac{a - a^{-1} - a^{-1}(x_1^2 + x_2^2)}{2} \frac{\sinh r + r \cosh r}{r^2 \sinh^3 r} \right] \\
 (2.8) \quad &= -U(x_1, x_2, a) + \frac{1}{2\pi^2} \frac{-1 + a^{-2} + a^{-2}(x_1^2 + x_2^2)}{2} \frac{\sinh r + r \cosh r}{r^2 \sinh^3 r}.
 \end{aligned}$$

We now consider the second-order Riesz transforms. We shall regard Δ^{-1} as the operator of convolution by the kernel W . The operators $R_{ij} = X_i X_j \Delta^{-1}$, $S_{ij} = \Delta^{-1} X_i X_j$, $T_{ij} = X_i \Delta^{-1} X_j$ are then properly defined on $C_c^\infty(G)$, with values in $C^\infty(G)$. By arguing as in [14, Lemma 6] we may show that there exist distributions k_{ij} , ℓ_{ij} , g_{ij} such that for any $f \in C_c^\infty(G)$

$$R_{ij} f = f * k_{ij} \quad S_{ij} f = f * \ell_{ij} \quad T_{ij} f = f * g_{ij}.$$

To compute these convolution kernels, we recall some simple properties of right- and left-invariant vector fields, which are the analogs of those proved in [14, Section 4.2] with respect to the measure λ .

Given a vector Z in \mathfrak{g} , we here denote by Z^r and Z^ℓ the right-invariant and left-invariant vector fields on G which agree with Z at the identity, defined by

$$Z^r f(x) = \frac{d}{dt} \Big|_{t=0} f(\exp(tZ)x) \quad \text{and} \quad Z^\ell f(x) = \frac{d}{dt} \Big|_{t=0} f(x \exp(tZ))$$

for any $f \in C^\infty(G)$ and $x \in G$. It is easy to check that for any $f \in C^\infty(G)$

$$(2.9) \quad Z^r \check{f} = (Z^\ell f)^\vee.$$

Let k be a distribution on G and $f, g \in C_c^\infty(G)$. Then

$$(2.10) \quad \langle f * k, g \rangle = \langle k, \check{f} * g \rangle.$$

The left-invariant derivative $Z^\ell k$ of k is the distribution such that for any $g \in C_c^\infty(G)$

$$(2.11) \quad \langle Z^\ell k, g \rangle = -\langle k, Z^\ell g \rangle.$$

If $f, g \in C_c^\infty(G)$, then

$$\langle Z^r f, g \rangle = \langle f, -Z^r g \rangle + Z\delta(e) \langle f, g \rangle.$$

So it is natural to define the right-invariant derivative of a distribution k as the distribution $Z^r k$ for which

$$(2.12) \quad \langle Z^r k, g \rangle = \langle k, -Z^r g \rangle + Z\delta(e) \langle k, g \rangle \quad \forall g \in C_c^\infty(G).$$

It is easy to verify that

$$(2.13) \quad Z^\ell(f * k) = f * Z^\ell k \quad \text{and} \quad Z^r(f * k) = Z^r f * k.$$

By (2.10) and (2.12) we deduce that

$$(2.14) \quad Z^\ell f * k = f * (-Z^r k + Z\delta(e)k).$$

Applying (2.13) we get that for any $f \in C_c^\infty(G)$

$$R_{ij}f = X_i X_j \Delta^{-1} f = X_i X_j (f * W) = X_i (f * X_j W) = f * X_i X_j W.$$

Thus the convolution kernel of R_{ij} is

$$(2.15) \quad k_{ij} = X_i X_j W,$$

the derivative taken in the distribution sense. We denote by $R_{ij}(\cdot, \cdot)$ the integral kernel of R_{ij} defined by $R_{ij}(x, y) = \delta(y) k_{ij}(y^{-1}x)$, for $x \neq y$.

Moreover, by (2.11) for all $f, g \in C_c^\infty(G)$

$$\begin{aligned} \langle S_{ij}f, g \rangle &= \langle \Delta^{-1} X_i X_j f, g \rangle = \langle X_i X_j f, \Delta^{-1} g \rangle \\ &= -\langle X_j f, X_i \Delta^{-1} g \rangle = \langle f, R_{ji}g \rangle. \end{aligned}$$

This implies that the integral kernel of S_{ij} is given by

$$(2.16) \quad S_{ij}(x, y) = R_{ji}(y, x) = \delta(x) k_{ji}(x^{-1}y) \quad \forall x \neq y.$$

It easily follows that the convolution kernel of S_{ij} is $\ell_{ij} = \delta \check{k}_{ji}$.

Applying (2.14), we get that for any $f \in C_c^\infty(G)$

$$\begin{aligned} T_{ij}f &= X_i \Delta^{-1} X_j f = X_i (X_j f * W) \\ &= X_j f * X_i W = f * (-X_j^r X_i W + X_j \delta(e) X_i W). \end{aligned}$$

Thus the convolution kernel of T_{ij} is $g_{ij} = -X_j^r X_i W + X_j \delta(e) X_i W$.

To avoid long computations, we do not compute explicitly the kernels of the second-order Riesz transforms, but we shall find their behavior away from the identity, i.e., in the complement of the unit ball B_1 .

In the sequel, we shall denote by $R(r)$ any series of the type $\sum_{k=1}^{\infty} c_k e^{-2kr}$, where the c_k are real numbers and the series is convergent for $r > 1$; we denote by $S(r)$ any function of the type $1 + R(r)$. These functions may vary from occurrence to occurrence. It is easy to see that a function $S(r) = 1 + R(r)$ may be differentiated termwise and its derivative is $S'(r) = R(r)$. Moreover, multiplying two functions of the type S , we obtain a function of the same kind.

Lemma 2.1 implies that for points $(x_1, x_2, a) \in \overline{B}_1^c$

$$(2.17) \quad X_i r(x_1, x_2, a) = \begin{cases} 2x_i e^{-r} S(r) & \text{if } i = 1, 2 \\ 2a e^{-r} S(r) - S(r) & \text{if } i = 0, \end{cases}$$

and by (2.2)

$$(2.18) \quad W(x_1, x_2, a) = \frac{1}{2\pi} a^{-1} e^{-r} S(r).$$

Let \mathbb{Z}_+^3 be the set of $m = (m_0, m_1, m_2)$ in \mathbb{Z}^3 such that $m_1, m_2 \geq 0$ and $m_0 \geq -1$. We denote by $|m|$ the sum $m_0 + m_1 + m_2$ and by x^m the product $x_1^{m_1} x_2^{m_2} a^{m_0}$. The principal term of W is of the type $x^m e^{-pr}$, where $|m| - p = -2$. We shall study the integrability of similar expressions in an elementary lemma, and first split \overline{B}_1^c into two parts, as follows:

$$G_+ = \{(x_1, x_2, a) \in \overline{B}_1^c : a > 1\},$$

and

$$G_- = \{(x_1, x_2, a) \in \overline{B}_1^c : a < 1\}.$$

LEMMA 2.2. — *Let m be in \mathbb{Z}_+^3 and $p \in \mathbb{N}$.*

(i) *The function $x^m e^{-pr}$ is integrable in G_+ if and only if*

$$(2.19) \quad m_1 + m_2 - 2p < -2 \quad \text{and} \quad |m| - p < -2.$$

(ii) *The function $x^m e^{-pr}$ is integrable in G_- if and only if*

$$(2.20) \quad m_1 + m_2 - 2p < -2 \quad \text{and} \quad m_0 + p > 0.$$

Proof. — If $(x_1, x_2, a) \in G_+$, then $e^r \sim a(1 + a^{-2}|(x_1, x_2)|^2)$, so that

$$\begin{aligned} & \int_{G_+} a^{m_0} |x_1|^{m_1} |x_2|^{m_2} e^{-pr} dx_1 dx_2 \frac{da}{a} \\ & \sim \int_1^\infty a^{m_0} \int_{\mathbb{R}^2} |x_1|^{m_1} |x_2|^{m_2} a^{-p} (1 + a^{-2}|(x_1, x_2)|^2)^{-p} dx_1 dx_2 \frac{da}{a}. \end{aligned}$$

Under the change of variables $a^{-1}(x_1, x_2) = (y_1, y_2)$, this transforms into the product

$$\int_1^\infty a^{|m|-p+1} da \int_{\mathbb{R}^2} |y_1|^{m_1} |y_2|^{m_2} (1 + |(y_1, y_2)|^2)^{-p} dy_1 dy_2.$$

Here the integral in a converges if and only if $|m| - p < -2$. By means of polar coordinates, the second integral is seen to converge if and only if $m_1 + m_2 - 2p < -2$. This proves (i).

If $(x_1, x_2, a) \in G_-$, then $e^r \sim a^{-1} (1 + |(x_1, x_2)|^2)$, so that

$$\int_{G_-} a^{m_0} |x_1|^{m_1} |x_2|^{m_2} e^{-pr} dx_1 dx_2 \frac{da}{a} \sim \int_0^1 a^{m_0+p-1} da \int_{\mathbb{R}^2} |x_1|^{m_1} |x_2|^{m_2} (1 + |(x_1, x_2)|^2)^{-p} dx_1 dx_2,$$

and (ii) follows. □

To study the higher order derivatives of W , we start with the derivatives along X_0, X_1, X_2 of an expression $x^m e^{-pr} S(r)$, as above. We shall always have

$$(2.21) \quad m_1 + m_2 - 2p \leq -2 \quad |m| - p \leq -2 \quad \text{and} \quad m_0 + p \geq 0,$$

which does not imply the integrability of $x^m e^{-pr}$. For many remainder terms, we shall denote by $Q(x)$ any finite sum of terms $x^n e^{-qr} R(r)$, where $|n| - q \leq -2$, $n_0 + q \geq 0$ and $n_1 + n_2 - 2q \leq -2$, so that $Q(x)$ is integrable in B_1^c . By (2.17) we get that in \overline{B}_1^c

$$\begin{aligned} X_1(x^m e^{-pr} S(r)) &= m_1 a^{m_0+1} x_1^{m_1-1} x_2^{m_2} e^{-pr} S(r) - \\ &\quad - p a^{m_0} x_1^{m_1} x_2^{m_2} e^{-pr} 2x_1 e^{-r} S(r) + \\ &\quad + a^{m_0} x_1^{m_1} x_2^{m_2} e^{-pr} R(r) 2x_1 e^{-r} S(r) \\ &= m_1 a^{m_0+1} x_1^{m_1-1} x_2^{m_2} e^{-pr} - 2p a^{m_0} x_1^{m_1+1} x_2^{m_2} e^{-(p+1)r} + \\ (2.22) \quad &\quad + Q(x). \end{aligned}$$

By symmetry an analogous formula holds for $i = 2$. From (2.17) we get

$$\begin{aligned} X_0(x^m e^{-pr} S(r)) &= m_0 a^{m_0} x_1^{m_1} x_2^{m_2} e^{-pr} S(r) - \\ &\quad - p a^{m_0} x_1^{m_1} x_2^{m_2} e^{-pr} [2ae^{-r} S(r) - S(r)] + \\ &\quad + a^{m_0} x_1^{m_1} x_2^{m_2} e^{-pr} R(r) [2ae^{-r} S(r) - S(r)] \\ &= (m_0 + p) a^{m_0} x_1^{m_1} x_2^{m_2} e^{-pr} - 2p a^{m_0+1} x_1^{m_1} x_2^{m_2} e^{-(p+1)r} + \\ (2.23) \quad &\quad + Q(x). \end{aligned}$$

Differentiating the expression (2.18) for W and applying (2.22) and (2.23), we get that in \overline{B}_1^c

$$X_j W(x_1, x_2, a) = -\frac{1}{2\pi} a^{-1} x_j e^{-2r} + Q(x) \quad \text{if } j = 1, 2,$$

and

$$X_0 W(x_1, x_2, a) = -\frac{1}{2\pi} e^{-2r} + Q(x).$$

We now differentiate W a second time, applying again (2.22) and (2.23) and also the observation that $X_j Q(x) = Q(x)$, for $i, j = 0, 1, 2$. The result is that there exist constants $\alpha_{ij}, \beta_{ij} \in \mathbb{R}$ and $m, n \in \mathbb{Z}_+^3$ such that in \overline{B}_1^c

$$(2.24) \quad k_{ij}(x) = \alpha_{ij} x^m e^{-2r} + \beta_{ij} x^n e^{-3r} + Q(x),$$

where $\beta_{ij} \neq 0$, $|m| = 0$, $|n| = 1$, $m_1 + m_2 - 4 < -2$, $n_1 + n_2 - 6 < -2$, $m_0 + 2 > 0$ and $n_0 + 3 > 0$. This means that k_{ij} has a principal part in \overline{B}_1^c given by at most two nonintegrable terms, while the remaining part of the kernel is integrable. Finally, we estimate the derivative of k_{ij} along the vector field X_2 . We get that, for $i, j = 0, 1, 2$, there exist constants $\gamma_{ij}, \eta_{ij}, \sigma_{ij}, \theta_{ij} \in \mathbb{R}$ and $h, \ell, m, n \in \mathbb{Z}_+^3$ such that in \overline{B}_1^c

$$(2.25) \quad X_2 k_{ij}(x) = \gamma_{ij} x^h e^{-2r} + \eta_{ij} x^\ell e^{-3r} + \sigma_{ij} x^m e^{-3r} + \theta_{ij} x^n e^{-4r} + Q(x),$$

where $\theta_{ij} \neq 0$, $|h| = 0$, $|\ell| = |m| = 1$, $|n| = 2$.

3. $H^1 - L^1$ -boundedness of R_i

In this section we prove that the Riesz transforms R_i are bounded from H^1 to L^1 , for $i = 0, 1, 2$.

This result is a consequence of the following boundedness theorem for integral operators. Note that the hypotheses of the following proposition are the same as those of [17, Theorem 2.1].

PROPOSITION 3.1. — *Let T be a linear operator bounded on L^2 such that $T = \sum_{j \in \mathbb{Z}} T_j$, where*

- (i) *the series converges in the strong operator topology of L^2 ;*
- (ii) *every T_j is an integral operator with integral kernel T_j ;*
- (iii) *there exist positive constants α, A, ε and $c > 1$ such that*

$$(3.1) \quad \int_G |T_j(x, y)| (1 + c^j d(x, y))^\varepsilon d\rho(x) \leq A \quad \forall y \in G;$$

$$(3.2) \quad \int_G |T_j(x, y) - T_j(x, z)| d\rho(x) \leq A (c^j d(y, z))^\alpha \quad \forall y, z \in G.$$

Then T is bounded from H^1 to L^1 .

Proof. — We first show that there exists a constant C such that for any atom a

$$(3.3) \quad \|Ta\|_1 \leq C.$$

Let R be the Calderón–Zygmund set supporting a , and denote by c_R the center of R and by R^* its dilated set (defined in Section 1). We estimate the integral of Ta on R^* by the Cauchy–Schwarz inequality:

$$(3.4) \quad \int_{R^*} |Ta| \, d\rho \leq \|Ta\|_2 \rho(R^*)^{1/2} \\ \leq C \|T\|_{L^2 \rightarrow L^2} \|a\|_2 \rho(R)^{1/2} \leq C \|T\|_{L^2 \rightarrow L^2}.$$

It is easy to show that from the estimates (3.1) and (3.2) it follows that

$$(3.5) \quad \sup_{R \in \mathcal{R}} \sup_{y, z \in R} \int_{(R^*)^c} |T(x, y) - T(x, z)| \, d\rho(x) < \infty,$$

where T is the integral kernel of T . Thus the integral of Ta on the complementary set of R^* is estimated as follows:

$$\begin{aligned} \int_{(R^*)^c} |Ta| \, d\rho &\leq \int_{(R^*)^c} \left| \int_R T(x, y) a(y) \, d\rho(y) \right| \, d\rho(x) \\ &= \int_{(R^*)^c} \left| \int_R [T(x, y) - T(x, c_R)] a(y) \, d\rho(y) \right| \, d\rho(x) \\ &\leq \int_{(R^*)^c} \int_R |T(x, y) - T(x, c_R)| |a(y)| \, d\rho(y) \, d\rho(x) \\ &= \int_R |a(y)| \left(\int_{(R^*)^c} |T(x, y) - T(x, c_R)| \, d\rho(x) \right) \, d\rho(y) \\ &\leq \|a\|_1 \sup_{y \in R} \int_{(R^*)^c} |T(x, y) - T(x, c_R)| \, d\rho(x) \\ &\leq C. \end{aligned}$$

This concludes the proof of (3.3). We shall deduce from (3.3) that T is bounded from H^1 to L^1 . Indeed, by [17, Theorem 2.1] T is bounded from L^1 to $L^{1, \infty}$. Now take a function f in H^1 and suppose that $f = \sum_{j=1}^{\infty} \lambda_j a_j$ is an atomic decomposition with $\sum_j |\lambda_j| \sim \|f\|_{H^1}$. Define $f_N = \sum_{j=1}^N \lambda_j a_j$. Since f_N converges to f in L^1 , $Tf_N = \sum_{j=1}^N \lambda_j Ta_j$ converges to Tf in $L^{1, \infty}$. On the other hand, by (3.3)

$$\|Tf_N - \sum_{j=1}^{\infty} \lambda_j Ta_j\|_1 \leq \sum_{j=N+1}^{\infty} |\lambda_j| \|Ta_j\|_1 \leq C \sum_{j=N+1}^{\infty} |\lambda_j|,$$

so that Tf_N converges to $\sum_{j=1}^\infty \lambda_j T a_j$ in L^1 . This implies that $Tf = \sum_{j=1}^\infty \lambda_j T a_j \in L^1$ and $\|Tf\|_1 \leq C \|f\|_{H^1}$. □

We now easily obtain the following theorem.

THEOREM 3.2. — *The Riesz transforms R_i , for $i = 0, 1, 2$, are bounded from H^1 to L^1 .*

Proof. — In the proof of [17, Theorem 6.4], it is shown that the operator R_i satisfies the assumptions of Proposition 3.1. Thus R_i is bounded from H^1 to L^1 . □

4. Unboundedness of S_1 and S_2

In this section we prove that the Riesz transforms S_1 and S_2 are not bounded from H^1 to L^1 . To do so, we shall define an atom a on G such that the images of a under these operators are not integrable in a region far from the support of the atom (see Theorem 4.2).

Differentiating the expression (2.7) for k_1 along the vector field X_2 and applying Lemma 2.1, we obtain that

$$\begin{aligned}
 X_2 k_1(x_1, x_2, a) &= -\frac{1}{2\pi^2} a^{-1} x_1 X_2 r(x_1, x_2, a) \left[\frac{r^2 \sinh^3 r (2 \cosh r + r \sinh r)}{r^4 \sinh^6 r} - \right. \\
 &\quad \left. - \frac{(\sinh r + r \cosh r)(2r \sinh^3 r + 3r^2 \sinh^2 r \cosh r)}{r^4 \sinh^6 r} \right] \\
 (4.1) \qquad &= \frac{1}{2\pi^2} a^{-1} \frac{x_1 x_2}{\sinh r} \frac{2r^2 \cosh^2 r + r^2 + 2 \sinh^2 r + 3r \sinh r \cosh r}{r^3 \sinh^4 r}.
 \end{aligned}$$

LEMMA 4.1. — *There exist regions $\Gamma'' \subset \Gamma' \subset \Gamma$, a positive continuous function Φ on Γ and a positive constant C such that*

- (i) $X_2 k_1 \geq C \Phi$ in Γ ;
- (ii) for any (x_1, x_2, a) in Γ' and τ in $[0, 1/4]$, the point $(x_1, x_2, a) \cdot (0, \tau, 1)$ is in Γ and

$$\Phi((x_1, x_2, a) \cdot (0, \tau, 1)) = \Phi(x_1, x_2, a);$$

- (iii) $\int_{\Gamma''} \Phi \, d\rho = \infty$.

Let E be the parallelepiped $(-1/2, 1/2) \times (-1/4, 0) \times (1, 2)$. Then

$$(4.2) \qquad \Gamma'' \cdot E^{-1} \cdot E \subseteq \Gamma'.$$

Proof. — Given $B, A > 1$ and $0 < \varepsilon < 1$ we define Γ as the region

$$\Gamma = \{(x_1, x_2, a) \in G : 1 - \varepsilon < x_2/x_1 < 1 + \varepsilon, x_1 > Ba, a > A\}.$$

For any (x_1, x_2, a) in Γ we have that

$$\frac{a^{-1}x_1^2}{2} < \cosh r(x_1, x_2, a) < C a^{-1} x_1^2$$

and $r(x_1, x_2, a) > 1$, for A and B sufficiently large. Moreover, since $e^r < 2 \cosh r < C a^{-1} x_1^2$,

$$r(x_1, x_2, a) \leq C \log(a^{-1} x_1^2).$$

By the formula (4.1) it is clear that $X_2 k_1$ is positive on Γ . Considering the first term in the numerator of the last fraction in (4.1), we see that for (x_1, x_2, a) in Γ

$$X_2 k_1(x_1, x_2, a) \geq C a^{-1} x_1^2 \frac{1}{r \cosh^3 r} \geq C \frac{a^{-1} x_1^2}{\log(a^{-1} x_1^2) (a^{-1} x_1^2)^3}.$$

We define

$$\Phi(x_1, x_2, a) = \frac{1}{\log(a^{-1} x_1^2) (a^{-1} x_1^2)^2}.$$

The condition (i) is verified. We now define

$$\begin{aligned} \Gamma' &= \{(x_1, x_2, a) \in \Gamma : 1 - \varepsilon' < x_2/x_1 < 1 + \varepsilon', x_1 > B'a\}, \\ \Gamma'' &= \{(x_1, x_2, a) \in \Gamma : 1 - \varepsilon'' < x_2/x_1 < 1 + \varepsilon'', x_1 > B''a, a > 2A\}, \end{aligned}$$

where $B'' > B' > B$, $0 < \varepsilon'' < \varepsilon' < \varepsilon < 1$ have to be chosen.

Let (x_1, x_2, a) be a point in Γ' and τ in $[0, 1/4]$. Then $(x_1, x_2, a) \cdot (0, \tau, 1) = (x_1, x_2 + a\tau, a)$. It is easy to see that we may choose B', ε' such that $(x_1, x_2 + a\tau, a) \in \Gamma$. Moreover,

$$\Phi((x_1, x_2, a) \cdot (0, \tau, 1)) = \frac{1}{\log(a^{-1} x_1^2) (a^{-1} x_1^2)^2} = \Phi(x_1, x_2, a),$$

as required in (ii). To prove (iii), we integrate Φ over Γ'' and obtain

$$\begin{aligned} \int_{\Gamma''} \Phi \, d\rho &= \int_{2A}^{\infty} \int_{B''a}^{\infty} \int_{(1-\varepsilon'')x_1}^{(1+\varepsilon'')x_1} \frac{1}{\log(a^{-1} x_1^2) (a^{-1} x_1^2)^2} \, dx_2 \, dx_1 \, \frac{da}{a} \\ &= C \int_{2A}^{\infty} \int_{B''a}^{\infty} \frac{x_1}{(a^{-1} x_1^2)^2 \log(a^{-1} x_1^2)} \, dx_1 \, \frac{da}{a} \\ &= C \int_{2A}^{\infty} \int_{(B'')^2 a}^{\infty} \frac{du}{u^2 \log u} \, da \\ &\geq C \int_{2A}^{\infty} \frac{1}{a \log a} \, da \\ &= \infty. \end{aligned}$$

Given $(x_1, x_2, a) \in \Gamma''$ and $(y_1, y_2, b), (z_1, z_2, c) \in E$, we have

$$\begin{aligned} & (x_1, x_2, a) \cdot (y_1, y_2, b)^{-1} \cdot (z_1, z_2, c) \\ &= (x_1 + ab^{-1}(z_1 - y_1), x_2 + ab^{-1}(z_2 - y_2), ab^{-1}c), \end{aligned}$$

where $ab^{-1}c > 2A/2 = A$, and

$$x_1 + ab^{-1}(z_1 - y_1) > B''a - ab^{-1} > B''a/2 > B'ab^{-1}c,$$

for B'' sufficiently large. Moreover,

$$\begin{aligned} x_2 + ab^{-1}(z_2 - y_2) &> x_1(1 - \varepsilon'') - ab^{-1}/4 \\ &= x_1(1 - \varepsilon') + (\varepsilon' - \varepsilon'')x_1 - ab^{-1}/4 \\ &> x_1(1 - \varepsilon') + (\varepsilon' - \varepsilon'')B''a - ab^{-1}/4 \\ &> x_1(1 - \varepsilon') + [(\varepsilon' - \varepsilon'')B'' - 1/4]ab^{-1} \\ &> [x_1 + ab^{-1}(z_1 - y_1)](1 - \varepsilon'), \end{aligned}$$

if $\varepsilon'' < \varepsilon'$ and B'' is sufficiently large. In the same way, we can achieve

$$x_2 + ab^{-1}(z_2 - y_2) < [x_1 + ab^{-1}(z_1 - y_1)](1 + \varepsilon').$$

Thus the point $(x_1, x_2, a) \cdot (y_1, y_2, b)^{-1} \cdot (z_1, z_2, c)$ is in Γ' , proving (4.2). \square

THEOREM 4.2. — *The operators S_1 and S_2 are not bounded from H^1 to L^1 .*

Proof. — By symmetry, it is enough to treat the case of S_1 . We shall construct an atom a such that S_1a does not belong to L^1 . Let R be the parallelepiped $[-e^2 \log 2/2, e^2 \log 2/2] \times [-e^2 \log 2/2, e^2 \log 2/2] \times [1/2, 2]$; it is easy to check that R is a Calderón–Zygmund set centered at the identity. Now let E be the parallelepiped defined in Lemma 4.1, and consider the right translate E^σ of E by the point $\exp(\sigma X_2) = (0, \sigma, 1)$ for some $\sigma > 0$, i.e.,

$$\begin{aligned} E^\sigma &= E \cdot (0, \sigma, 1) = \{(y_1, y_2 + b\sigma, b) : (y_1, y_2, b) \in E\} \\ &\subset (-1/2, 1/2) \times (-1/4 + \sigma, 2\sigma) \times (1, 2). \end{aligned}$$

With $\sigma = 1/4$, E and E^σ are disjoint and contained in R .

Let us consider the function $a = \rho(R)^{-1}(\mathbf{1}_E - \mathbf{1}_{E^\sigma})$. It is obvious that a is supported in the Calderón–Zygmund set R and $\|a\|_\infty \leq \rho(R)^{-1}$. Moreover $\int a \, d\rho = 0$ and so a is an atom. We now compute S_1a outside the support

of a . For all $x \notin \overline{E \cup E^\sigma}$

$$\begin{aligned} S_1 a(x) &= \int S_1(x, y) a(y) \, d\rho(y) \\ &= \rho(R)^{-1} \int_E S_1(x, y) \, d\rho(y) - \rho(R)^{-1} \int_{E^\sigma} S_1(x, y) \, d\rho(y). \end{aligned}$$

Changing variable $y = v \cdot (0, \sigma, 1)$ in the last integral, this transforms into

$$\begin{aligned} &\rho(R)^{-1} \int_E S_1(x, y) \, d\rho(y) - \rho(R)^{-1} \int_E S_1(x, v \cdot (0, \sigma, 1)) \, d\rho(v) \\ &= \rho(R)^{-1} \int_E [S_1(x, y) - S_1(x, y \cdot (0, \sigma, 1))] \, d\rho(y). \end{aligned}$$

By (2.6) we know that

$$\begin{aligned} S_1(x, y) - S_1(x, y \cdot (0, \sigma, 1)) &= \delta(x) (-k_1(x^{-1}y) + k_1(x^{-1}y \exp(\sigma X_2))) \\ &= \delta(x) \sigma \left. \frac{d}{dt} \right|_{t=\tau(x,y)} k_1(x^{-1}y \exp(t X_2)) \\ &= \delta(x) \sigma X_2 k_1(x^{-1}y \exp(\tau(x, y) X_2)), \end{aligned}$$

for some $\tau(x, y)$ in $(0, \sigma)$. It follows that for all $x \notin \overline{E \cup E^\sigma}$

$$(4.3) \quad S_1 a(x) = \rho(R)^{-1} \sigma \delta(x) \int_E X_2 k_1(x^{-1}y \exp(\tau(x, y) X_2)) \, d\rho(y).$$

To prove that $S_1 a$ is not in L^1 , we integrate $|S_1 a|$ in the region $E(\Gamma'')^{-1}$, where Γ'' is the set which appears in Lemma 4.1. It is easy to check that $E(\Gamma'')^{-1}$ is disjoint with $\overline{E \cup E^\sigma}$, so that we can apply (4.3) and obtain

$$\begin{aligned} &\int_{E(\Gamma'')^{-1}} |S_1 a(x)| \, d\rho(x) \\ &= \rho(R)^{-1} \sigma \int_{E(\Gamma'')^{-1}} \delta(x) \left| \int_E X_2 k_1(x^{-1}y \exp(\tau(x, y) X_2)) \, d\rho(y) \right| \, d\rho(x) \\ &= \rho(R)^{-1} \sigma \int_{\Gamma'' E^{-1}} \left| \int_E X_2 k_1(xy \exp(\tau(x^{-1}, y) X_2)) \, d\rho(y) \right| \, d\rho(x). \end{aligned}$$

If $x \in \Gamma'' E^{-1}$ and $y \in E$, then $xy \in \Gamma'$, in view of (4.2). Since $0 < \tau(x^{-1}, y) < \sigma = 1/4$, by Lemma 4.1 the point $xy \exp(\tau(x^{-1}, y) X_2)$ is in Γ and

$$X_2 k_1(xy \exp(\tau(x^{-1}, y) X_2)) \geq C \Phi(xy \exp(\tau(x^{-1}, y) X_2)) = C \Phi(xy).$$

Hence, applying Fubini's theorem and using $w = xy$ instead of x , we get

$$\begin{aligned} \int_{E(\Gamma'')^{-1}} |S_1 a(x)| \, d\rho(x) &\geq C \rho(R)^{-1} \sigma \int_{\Gamma''} \int_{E^{-1}} \Phi(xy) \, d\rho(y) \, d\rho(x) \\ &= C \rho(R)^{-1} \sigma \int_E d\rho(y) \int_{\Gamma''} \int_{E^{-1}y} \Phi(w) \, d\rho(w) \\ &\geq C \rho(R)^{-1} \sigma \int_E d\rho(y) \int_{\Gamma''} \Phi(w) \, d\rho(w). \end{aligned}$$

Lemma 4.1 (iii) implies that this integral diverges. □

5. Unboundedness of S_0

To prove that the operator S_0 is not bounded from H^1 to L^1 , we use the same idea as in the previous section. The only difference is that we consider now the derivative $X_0 k_0$ in a slightly different region.

We first compute the derivative of the expression (2.8) for k_0 along the vector field X_0 :

$$\begin{aligned} X_0 k_0(x_1, x_2, a) &= \\ &= \frac{1}{2\pi^2} \frac{a^{-1}}{r \sinh r} + \frac{1}{2\pi^2} \frac{1 - a^{-2} - a^{-2}(x_1^2 + x_2^2)}{2} \frac{\sinh r + r \cosh r}{r^2 \sinh^3 r} - \\ &- \frac{1}{2\pi^2} [a^{-2} + a^{-2}(x_1^2 + x_2^2)] \frac{\sinh r + r \cosh r}{r^2 \sinh^3 r} + \\ &+ \frac{1}{2\pi^2} \frac{-1 + a^{-2} + a^{-2}(x_1^2 + x_2^2)}{2} \frac{a - a^{-1} - a^{-1}(x_1^2 + x_2^2)}{2 \sinh r} \times \\ &\times \left[\frac{(2 \cosh r + r \sinh r)r^2 \sinh^3 r}{r^4 \sinh^6 r} - \right. \\ &\left. - \frac{(\sinh r + r \cosh r)(2r \sinh^3 r + 3r^2 \sinh^2 r \cosh r)}{r^4 \sinh^6 r} \right] \\ &= \frac{1}{2\pi^2} \frac{a^{-1}}{r \sinh r} + \frac{1}{2\pi^2} \frac{1 - 3a^{-2} - 3a^{-2}(x_1^2 + x_2^2)}{2} \frac{\sinh r + r \cosh r}{r^2 \sinh^3 r} + \\ &+ \frac{1}{2\pi^2} a^{-1} \frac{[a - a^{-1} - a^{-1}(x_1^2 + x_2^2)]^2}{4} \times \\ (5.1) \quad &\times \frac{2r^2 \cosh^2 r + r^2 + 2 \sinh^2 r + 3r \sinh r \cosh r}{r^3 \sinh^5 r}. \end{aligned}$$

LEMMA 5.1. — *There exist two regions $\Omega' \subset \Omega$, a positive continuous function Ψ on Ω and a positive constant C such that*

- (i) $X_0 k_0 \geq C \Psi$ in Ω ;
- (ii) for any (x_1, x_2, a) in Ω and τ in $[0, 1]$, the point $(x_1, x_2, a) \cdot (0, 0, e^\tau)$ is in Ω and

$$\Psi((x_1, x_2, a) \cdot (0, 0, e^\tau)) \geq C \Psi(x_1, x_2, a);$$

- (iii) $\int_{\Omega'} \Psi \, d\rho = \infty$.

Let F be the parallelepiped $(-1/16, 1/16) \times (-1/16, 1/16) \times (1, \sqrt{2})$. Then

$$(5.2) \quad \Omega' \cdot F^{-1} \cdot F \subseteq \Omega.$$

Proof. — Let $A > 1$ be a constant to be chosen later and define

$$(5.3) \quad \begin{aligned} \Omega &= \{(x_1, x_2, a) \in G : x_1^2 + x_2^2 < a^2/4, a > A\}, \\ \Omega' &= \{(x_1, x_2, a) \in G : x_1^2 + x_2^2 < a^2/64, a > \sqrt{2}A\}. \end{aligned}$$

For all (x_1, x_2, a) in Ω

$$\frac{a}{2} < \cosh r(x_1, x_2, a) < C a.$$

For A sufficiently large, $r(x_1, x_2, a) > 1$ here, and, since $e^\tau \leq 2 \cosh r \leq C a$, we have $r \leq C \log a$.

It is easy to show that in the region Ω all the summands which appear in the last expression in (5.1) are positive, so that for all (x_1, x_2, a) in Ω

$$X_0 k_0(x_1, x_2, a) \geq C \frac{a^{-1}}{r \sinh r} \geq \frac{C}{a^2 \log a}.$$

We define

$$\Psi(x_1, x_2, a) = \frac{1}{a^2 \log a}.$$

The condition (i) is satisfied.

Let $(x_1, x_2, a) \in \Omega$ and $\tau \in [0, 1]$. It is easy to check that the point $(x_1, x_2, a) \cdot (0, 0, e^\tau) = (x_1, x_2, a e^\tau)$ is in Ω . Moreover,

$$\Psi((x_1, x_2, a) \cdot (0, 0, e^\tau)) = \frac{1}{a^2 e^{2\tau} \log(a e^\tau)} \geq C \frac{1}{a^2 \log a} = C \Psi(x_1, x_2, a),$$

as claimed in (ii). To prove (iii), we integrate Ψ over Ω' and obtain

$$\begin{aligned} \int_{\Omega'} \Psi \, d\rho &= \int_{\sqrt{2}A}^{\infty} \frac{1}{a^2 \log a} \int \int_{x_1^2 + x_2^2 \leq a^2/64} dx_1 dx_2 \frac{da}{a} \\ &= C \int_{\sqrt{2}A}^{\infty} \frac{1}{a \log a} da \\ &= \infty. \end{aligned}$$

Given $(x_1, x_2, a) \in \Omega'$ and $(y_1, y_2, b), (z_1, z_2, c) \in F$, we have that

$$\begin{aligned} & (x_1, x_2, a) \cdot (y_1, y_2, b)^{-1} \cdot (z_1, z_2, c) \\ &= (x_1 + ab^{-1}(z_1 - y_1), x_2 + ab^{-1}(z_2 - y_2), ab^{-1}c), \end{aligned}$$

where $ab^{-1}c > \sqrt{2}A/\sqrt{2} = A$, and

$$\begin{aligned} & [x_1 + ab^{-1}(z_1 - y_1)]^2 + [x_2 + ab^{-1}(z_2 - y_2)]^2 \\ & < (|x_1| + a/8)^2 + (|x_2| + a/8)^2 \\ & < 2(1/8 + 1/8)^2 a^2 \\ & < (ab^{-1}c)^2/4. \end{aligned}$$

Thus $(x_1, x_2, a) \cdot (y_1, y_2, b)^{-1} \cdot (z_1, z_2, c) \in \Omega$, and (5.2) is proved. \square

THEOREM 5.2. — *The operator S_0 is not bounded from H^1 to L^1 .*

Proof. — Following closely the proof of Theorem 4.2, we shall construct an atom a such that S_0a does not belong to L^1 . With R as in that proof, we let F be the parallelepiped defined in Lemma 5.1 and consider

$$\begin{aligned} F^\sigma &= F \cdot (0, 0, e^\sigma) = \{(y_1, y_2, ae^\sigma) : (y_1, y_2, b) \in F\} \\ &= (-1/16, 1/16) \times (-1/16, 1/16) \times (e^\sigma, e^\sigma\sqrt{2}). \end{aligned}$$

With $\sigma = (\log 2)/2$, F and F^σ are disjoint and contained in R .

Let us consider the atom $a = \rho(R)^{-1}(\mathbf{1}_F - \mathbf{1}_{F^\sigma})$. We compute S_0a outside the support of a . For all $x \notin \overline{F \cup F^\sigma}$

$$\begin{aligned} S_0a(x) &= \rho(R)^{-1} \int_F S_0(x, y) d\rho(y) - \rho(R)^{-1} \int_{F^\sigma} S_0(x, y) d\rho(y) \\ &= \rho(R)^{-1} \int_F [S_0(x, y) - S_0(x, y \cdot (0, 0, e^\sigma))] d\rho(y). \end{aligned}$$

By (2.6) we know that

$$\begin{aligned} S_0(x, y) - S_0(x, y \cdot (0, 0, e^\sigma)) &= \delta(x) (-k_0(x^{-1}y) + k_0(x^{-1}y \exp(\sigma X_0))) \\ &= \delta(x) \sigma \frac{d}{dt} \Big|_{t=\tau(x, y)} k_0(x^{-1}y \exp(t X_0)) \\ &= \delta(x) \sigma X_0 k_0(x^{-1}y \exp(\tau(x, y) X_0)), \end{aligned}$$

for some $\tau(x, y)$ in $(0, \sigma)$. It follows that for all $x \notin \overline{F \cup F^\sigma}$

$$(5.4) \quad S_0a(x) = \rho(R)^{-1} \sigma \delta(x) \int_F X_0 k_0(x^{-1}y \exp(\tau(x, y) X_0)) d\rho(y).$$

To prove that S_0a is not in L^1 , we integrate S_0a in the region $F(\Omega')^{-1}$. It is easy to verify that $F(\Omega')^{-1}$ is disjoint with $\overline{F \cup F^\sigma}$, so that we can apply (5.4) and obtain

$$\begin{aligned} & \int_{F(\Omega')^{-1}} |S_0a(x)| \, d\rho(x) = \\ &= \rho(R)^{-1} \sigma \int_{F(\Omega')^{-1}} \delta(x) \left| \int_F X_0 k_0(x^{-1}y \exp(\tau(x, y) X_0)) \, d\rho(y) \right| \, d\rho(x) \\ &= \rho(R)^{-1} \sigma \int_{\Omega' F^{-1}} \left| \int_F X_0 k_0(xy \exp(\tau(x^{-1}, y) X_0)) \, d\rho(y) \right| \, d\rho(x). \end{aligned}$$

If $x \in \Omega' F^{-1}$ and $y \in F$, then $xy \in \Omega$, in view of (5.2). Since $0 < \tau(x^{-1}, y) < \sigma < 1$, by Lemma 5.1(ii) the point $xy \exp(\tau(x^{-1}, y) X_0)$ is in Ω and

$$X_0 k_0(xy \exp(\tau(x^{-1}, y) X_0)) \geq C \Psi(xy \exp(\tau(x^{-1}, y) X_0)) \geq C \Psi(xy).$$

As in the proof of Theorem 4.2, we get

$$\begin{aligned} \int_{F(\Omega')^{-1}} |S_0a(x)| \, d\rho(x) &\geq C \rho(R)^{-1} \sigma \int_{\Omega' F^{-1}} \int_F \Psi(xy) \, d\rho(y) \, d\rho(x) \\ &= C \rho(R)^{-1} \sigma \int_F d\rho(y) \int_{\Omega' F^{-1}y} \Psi(w) \, d\rho(w) \\ &\geq C \rho(R)^{-1} \sigma \int_F d\rho(y) \int_{\Omega'} \Psi(w) \, d\rho(w). \end{aligned}$$

Lemma 5.1 (iii) implies that the last integral diverges. □

6. The local parts of T_{ij}, S_{ij} and R_{ij}

In this section, we study the local parts of the kernels of the second-order Riesz transforms. We shall prove that they behave like standard Calderón–Zygmund kernels in \mathbb{R}^3 and deduce that they correspond to operators which are bounded from H^1 to L^1 .

Let Ψ be a function in $C_c^\infty(S)$ such that $0 \leq \Psi \leq 1$, Ψ is supported in the ball B_2 of radius 2 and $\Psi = 1$ on the ball B_1 . Define

$$\begin{aligned} g_{ij}^0 &= g_{ij} \Psi & \text{and} & & g_{ij}^\infty &= g_{ij} (1 - \Psi), \\ k_{ij}^0 &= k_{ij} \Psi & \text{and} & & k_{ij}^\infty &= k_{ij} (1 - \Psi), \\ \ell_{ij}^0 &= \ell_{ij} \Psi & \text{and} & & \ell_{ij}^\infty &= \ell_{ij} (1 - \Psi), \end{aligned}$$

and let $T_{ij}^0, T_{ij}^\infty, R_{ij}^0, R_{ij}^\infty, S_{ij}^0$ and S_{ij}^∞ be the corresponding convolution operators. We shall prove that the operators T_{ij}^0, R_{ij}^0 and S_{ij}^0 are bounded from H^1 to L^1 . To do so, we use the following lemma.

LEMMA 6.1. — *Let T be a convolution operator which is bounded on L^2 . Suppose that its kernel k is a distribution supported in the ball B_2 and given by a function in $B_2 \setminus \{e\}$. Define*

$$(6.1) \quad \beta((x_1, x_2, s), (y_1, y_2, t)) = \delta(y_1, y_2, e^t) k((y_1, y_2, e^t)^{-1} \cdot (x_1, x_2, e^s))$$

for any $(x_1, x_2, s) \neq (y_1, y_2, t) \in \mathbb{R}^3$. If β satisfies the standard estimate

$$(6.2) \quad |\beta(\mathbf{x}, \mathbf{y})| + |\mathbf{x} - \mathbf{y}| [|\nabla_{\mathbf{x}} \beta(\mathbf{x}, \mathbf{y})| + |\nabla_{\mathbf{y}} \beta(\mathbf{x}, \mathbf{y})|] \leq C |\mathbf{x} - \mathbf{y}|^{-3},$$

for $|\mathbf{y}| < 2A_0$, $\mathbf{x} \neq \mathbf{y}$, where A_0 is a suitable constant, then T is bounded from H^1 to L^1 .

Proof. — We first verify that the operator T is of weak type 1. Via a standard Calderón-Zygmund decomposition argument, the L^2 -boundedness of T and the estimate (6.2) imply that for any $f \in L^1(B_1)$

$$\rho(\{x \in G : |Tf(x)| > t\}) \leq \frac{C}{t} \|f\|_1 \quad \forall t > 0.$$

There exists a sequence of balls B_j , centered at points x_j and of radius 1, such that $G = \bigcup_j B_j$ and each point of G belongs to at most n of the balls B_j (see [13, Lemma 8]). From the left-invariance of the operator T , the right-invariance of the measure and a simple application of a partition of unity $(\psi_j)_j$ such that $\text{supp } \psi_j \subseteq \overline{B_j}$, we may deduce that for any $f \in L^1$ and $t > 0$

$$\begin{aligned} \rho(\{x \in G : |Tf(x)| > t\}) &\leq \rho(\{x \in G : \sum_j |T(\psi_j f)(x)| > t\}) \\ &\leq \frac{C}{t} \sum_j \|\psi_j f\|_1 \leq \frac{C}{t} \|f\|_1. \end{aligned}$$

The inequalities above follow by a standard argument (see [13, Lemma 7] for the details). Thus, T is of weak type 1. As in the proof of Theorem 3.1, the lemma will follow if we show that there exists a constant C such that $\|Tb\|_1 \leq C$ for any atom b .

Any atom b can be transformed by an appropriate left-translation into an atom a supported in a Calderón-Zygmund set centered at the identity, and $\|Tb\|_1 = \|Ta\|_1$ by the left-invariance of T . Thus, it suffices to consider an atom a supported in a Calderón-Zygmund set $R = [-L/2, L/2] \times [-L/2, L/2] \times [e^{-r}, e^r]$ centered at the identity. Recall that the dilated set R^* is defined by $\{x \in G : d(x, R) < r\}$. Since T is bounded on L^2 ,

$$(6.3) \quad \|Ta\|_{L^1(R^*)} \leq \rho(R^*)^{1/2} \|T\|_{L^2 \rightarrow L^2} \|a\|_2 \leq C.$$

Note that $\text{supp}(Ta) \subseteq R \cdot B_2 \subseteq \{x \in S : d(x, R) < 2\}$.

If $r > 2$, then $\text{supp}(Ta) \subseteq R \cdot B_2 \subseteq R^*$, so that $\|Ta\|_1 = \|Ta\|_{L^1(R^*)} \leq C$.

Suppose now that $r < 2$. Since $r(x_1, x_2, a) \sim |(x_1, x_2, \log a)|$ near the identity, there exists an absolute constant A_0 such that

$$R \subset \{(x_1, x_2, a) : |(x_1, x_2, \log a)| < A_0 r\} = B.$$

Notice that B corresponds to a euclidean ball in \mathbb{R}^3 . Since $r < 2$, $\rho(B) \sim \rho(R) \sim r^3$ and, by arguing as in (6.3), we obtain that $\|Ta\|_{L^1(2B)} \leq C$.

It remains to estimate the L^1 -norm of Ta outside $2B$. Since the distributional convolution of T is k , we get

$$\begin{aligned} \int_{(2B)^c} |Ta(x)| \, d\rho(x) &= \int_{(2B)^c} \left| \int_B a(y) \delta(y) k(y^{-1}x) \, d\rho(y) \right| \, d\rho(x) \\ &= \int_{(2B)^c} \left| \int_B a(y) \delta(y) [k(y^{-1}x) - k(x)] \, d\rho(y) \right| \, d\rho(x). \end{aligned}$$

Rewriting this in terms of β and euclidean coordinates, we conclude

$$\begin{aligned} \|Ta\|_{L^1(2B)^c} &\leq \|a\|_\infty \times \int_{|(x_1, x_2, s)| > 2A_0 r} \int_{|(y_1, y_2, t)| < A_0 r} \\ &\quad |\beta((x_1, x_2, s), (y_1, y_2, t)) - \beta((x_1, x_2, s), \mathbf{0})| \, dy_1 \, dy_2 \, dt \, dx_1 \, dx_2 \, ds. \end{aligned}$$

If $|\mathbf{x}| > 2A_0 r$ and $|\mathbf{y}| < A_0 r$, by (6.2) we get

$$|\beta(\mathbf{x}, \mathbf{y}) - \beta(\mathbf{x}, \mathbf{0})| \leq \sup_{|\mathbf{y}'| < A_0 r} |\nabla_{\mathbf{y}} \beta(\mathbf{x}, \mathbf{y}')| |\mathbf{y}| \leq C r |\mathbf{x}|^{-4},$$

so that

$$\begin{aligned} \|Ta\|_{L^1(2B)^c} &\leq \|a\|_\infty \int_{|\mathbf{x}| > 2A_0 r} \int_{|\mathbf{y}| < A_0 r} |\beta(\mathbf{x}, \mathbf{y}) - \beta(\mathbf{x}, \mathbf{0})| \, d\mathbf{y} \, d\mathbf{x} \\ &\leq C r^{-3} \int_{|\mathbf{x}| > 2A_0 r} \int_{|\mathbf{y}| < A_0 r} |\mathbf{x}|^{-4} r \, d\mathbf{y} \, d\mathbf{x} \\ &\leq C. \end{aligned}$$

□

PROPOSITION 6.2. — *The operators T_{ij}^0 , R_{ij}^0 and S_{ij}^0 are bounded from H^1 to L^1 .*

Proof. — It is enough to apply Lemma 6.1 to the operators T_{ij}^0 , R_{ij}^0 and S_{ij}^0 . By [14, Theorem 12] they are bounded on L^2 and their kernels g_{ij}^0 , k_{ij}^0 , ℓ_{ij}^0 are supported in the ball B_2 . Let k denote one of the kernel g_{ij}^0 , k_{ij}^0 , ℓ_{ij}^0 . We must show that the function β , given by

$$\beta((x_1, x_2, s), (y_1, y_2, t)) = e^{-2t} k(e^{-t}(x_1 - y_1), e^{-t}(x_2 - y_2), e^{s-t}),$$

satisfies (6.2). By means of some elementary Taylor expansions in the variable $\mathbf{x} = (x_1, x_2, \log a) \in \mathbb{R}^3$, one finds that near e

$$W(x_1, x_2, a) = \frac{1}{4\pi} \frac{1}{|\mathbf{x}|} (1 + h_1 + h_2 + \dots),$$

where each h_j is a function of \mathbf{x} which is homogeneous of degree j and smooth away from 0, and the series $\sum_j h_j$ converges near 0. Termwise differentiation is possible, and we let ∂^α denotes a differentiation operator with respect to $(x_1, x_2, \log a)$, of order α . Then

$$\partial^\alpha W = \tilde{h}_{-1-|\alpha|} + \tilde{h}_{-|\alpha|} + \dots,$$

with similar smooth homogeneous functions \tilde{h}_j . This implies

$$|\partial^\alpha W(\mathbf{x})| \leq C |\mathbf{x}|^{-1-|\alpha|}$$

for small \mathbf{x} , and thus

$$(6.4) \quad |k(x_1, x_2, a)| + |(x_1, x_2, \log a)| |\nabla k(x_1, x_2, a)| \leq C |(x_1, x_2, \log a)|^{-3}$$

in $B_2 \setminus \{e\}$, where ∇ denotes the gradient with respect to $(x_1, x_2, \log a)$.

If (y_1, y_2, t) is near the origin, we have

$$\begin{aligned} |\beta((x_1, x_2, s), (y_1, y_2, t))| &\leq C |k(e^{-t}(x_1 - y_1), e^{-t}(x_2 - y_2), e^{s-t})| \\ &\leq C |(e^{-t}(x_1 - y_1), e^{-t}(x_2 - y_2), s - t)|^{-3} \\ &\leq C |(x_1 - y_1, x_2 - y_2, s - t)|^{-3}, \end{aligned}$$

and

$$\begin{aligned} &|\nabla_{(x_1, x_2, s)} \beta((x_1, x_2, s), (y_1, y_2, t))| + |\nabla_{(y_1, y_2, t)} \beta((x_1, x_2, s), (y_1, y_2, t))| \\ &\leq C |k(e^{-t}(x_1 - y_1), e^{-t}(x_2 - y_2), e^{s-t})| + \\ &+ C |\nabla k(e^{-t}(x_1 - y_1), e^{-t}(x_2 - y_2), e^{s-t})| \\ &\leq C |(x_1 - y_1, x_2 - y_2, s - t)|^{-4}, \end{aligned}$$

and the theorem follows. □

7. Boundedness of T_{ij}

We shall prove that the operators $T_{ij} = X_i \Delta^{-1} X_j$ are bounded from H^1 to L^1 . Since we already verified the boundedness of their local parts, it remains to consider the global parts.

In [14, Lemma 9] it is proved that the global parts of the kernels $g_{ij}^{r, \infty}$ of the right-invariant Riesz transforms T_{ij}^r are integrable with respect to the measure λ . Since (2.9) implies that $(T_{ij}^r \check{f})^\vee = T_{ij} f$ for any $f \in C_c^\infty(G)$, we

obtain that $g_{ij}^\infty = \check{g}_{ij}^{r,\infty}$. Thus g_{ij}^∞ is integrable with respect to the measure ρ and the corresponding convolution operator T_{ij}^∞ is bounded from H^1 to L^1 .

8. Unboundedness of S_{ij}

In this section we prove that the operators S_{ij} are not bounded from H^1 to L^1 . Again it suffices to consider their global parts. To do so, we use the same idea as in Section 4, defining an atom whose image under the operator S_{ij} is not integrable far from the support of the atom.

We will need to estimate some integrals of derivatives of the kernels k_{ij} . Notice that it is enough to treat the values of (i, j) listed in the following lemma, since the remaining cases will follow by symmetry.

LEMMA 8.1. — *For each pair $(i, j) \in \{(1, 1), (1, 2), (1, 0), (0, 1), (0, 0)\}$, there exist regions $\Gamma'' \subset \Gamma' \subset \Gamma$ in G , a positive continuous function Φ on Γ and positive constants C, τ such that*

- (i) $|X_2 k_{ij}| \geq C \Phi$ in Γ ;
- (ii) for any (x_1, x_2, a) in Γ' and σ in $[0, \tau]$, the point $(x_1, x_2, a) \cdot (0, \sigma, 1)$ is in Γ and

$$\Phi((x_1, x_2, a) \cdot (0, \sigma, 1)) = \Phi(x_1, x_2, a);$$

- (iii) $\int_{\Gamma''} \Phi d\rho = \infty$.

Moreover, there exist constants $0 < \delta < 1$ and $1 < \beta < 2$ such that the parallelepiped $E = (0, \delta) \times (-\delta, 0) \times (1, \beta)$ satisfies the condition

$$(8.1) \quad \Gamma'' \cdot E^{-1} \cdot E \subseteq \Gamma'.$$

Proof. — Let us fix a pair (i, j) . To simplify the notation we write k for the kernel k_{ij} and drop the indices i, j . Because of (2.25), there exist constants $\gamma, \eta, \sigma, \theta \in \mathbb{R}$ and $h, \ell, m, n \in \mathbb{Z}_+^3$ such that for x in \overline{B}_1^c

$$\begin{aligned} X_2 k(x) &= \gamma x^h e^{-2r} + \eta x^\ell e^{-3r} + \sigma x^m e^{-3r} + \theta x^n e^{-4r} + Q(x) \\ &= F(x) + Q(x), \end{aligned}$$

where $F(x)$ is defined by the last equality. Here $\theta \neq 0, |h| = 0, |\ell| = |m| = 1, |n| = 2$. The remainder term $Q(x)$ is as described in Section 2.

For large a , (1.1) implies $e^r \sim a + a^{-1}(x_1^2 + x_2^2)^2$ and more precisely,

$$e^r = a + a^{-1}(x_1^2 + x_2^2) + a^{-1} - e^{-r}$$

$$= (a + a^{-1}(x_1^2 + x_2^2)) \left(1 + \frac{a^{-1} - e^{-r}}{a + a^{-1}(x_1^2 + x_2^2)}\right).$$

Inverting the last factor here and expanding, we see that for $x = (x_1, x_2, a)$ with a large

$$e^{-pr} = a^p (a^2 + x_1^2 + x_2^2)^{-p} (1 + O(e^{-r})) = a^p |x|^{-2p} (1 + O(e^{-r})),$$

where $|\cdot|$ denotes the euclidean norm in \mathbb{R}^3 . Thus, for such x

$$F(x) = \frac{\theta x^n a^4 + |x|^2 [\gamma x^h a^2 |x|^2 + \eta x^\ell a^3 + \sigma x^m a^3]}{|x|^8} + E(x)$$

$$= \frac{P(x)}{|x|^8} + E(x),$$

where P is a polynomial in the variables x_1, x_2, a , homogeneous of degree 6. Further, $E(x)$ is a sum like $F(x)$, but with $e^{-(p+1)r}$ instead of e^{-pr} in each term. We write $P(x) = \theta x^n a^4 + |x|^2 \tilde{P}(x)$, where \tilde{P} is homogeneous of degree 4. Notice that P is not identically 0, since the monomial $x^n a^4$ cannot equal a product $\theta^{-1} |x|^2 \tilde{P}(x)$. We can thus find $q_1, q_2 > 0$ with $P(q_1, q_2, 1) \neq 0$. By continuity and homogeneity, $P(x) \neq 0$ also for x in a narrow cone near the ray in the direction $(q_1, q_2, 1)$, in particular for x in the truncated cone

$$\Gamma = \{(x_1, x_2, a) \in G : a > A, |x_1/a - q_1| < \varepsilon, |x_2/a - q_2| < \varepsilon\},$$

for some small $\varepsilon > 0$. With A sufficiently large, this implies that in the region Γ the quantities $|E|, |Q|$ are much smaller than $|F|$ and so for any $(x_1, x_2, a) \in \Gamma$

$$|X_2 k(x_1, x_2, a)| \geq C |F(x_1, x_2, a)| \geq C \frac{|P(x_1, x_2, a)|}{|(x_1, x_2, a)|^8} \geq C a^{-2}.$$

Defining $\Phi(x_1, x_2, a) = a^{-2}$ in Γ , we have proved (i).

We define

$$\Gamma' = \{(x_1, x_2, a) \in \Gamma : |x_1/a - q_1| < \varepsilon/2, |x_2/a - q_2| < \varepsilon/2\},$$

and

$$\Gamma'' = \{(x_1, x_2, a) \in \Gamma : a > 2A, |x_1/a - q_1| < \varepsilon/4, |x_2/a - q_2| < \varepsilon/4\}.$$

Now choose $\tau < \varepsilon/2$ and $\sigma \in [0, \tau]$, and let (x_1, x_2, a) be in Γ' . Then $(x_1, x_2, a) \cdot (0, \sigma, 1) = (x_1, x_2 + a\sigma, a)$. We have that $|x_1/a - q_1| < \varepsilon/2 < \varepsilon$ and

$$|(x_2 + a\sigma)/a - q_2| < |x_2/a - q_2| + \sigma < \varepsilon/2 + \tau < \varepsilon.$$

Thus $(x_1, x_2 + a\sigma, a) \in \Gamma$ and $\Phi((x_1, x_2, a) \cdot (0, \sigma, 1)) = a^{-2} = \Phi(x_1, x_2, a)$.

To prove (iii), it suffices to note that

$$\int_{\Gamma''} a^{-2} dx_2 dx_1 \frac{da}{a} \geq C \int_{2A}^{\infty} a^{-2} a^2 \frac{da}{a} = \infty.$$

Aiming at (8.1) we take points $(x_1, x_2, a) \in \Gamma''$ and $(y_1, y_2, b), (z_1, z_2, c) \in E$ and consider $(x_1, x_2, a) \cdot (y_1, y_2, b)^{-1} \cdot (z_1, z_2, c) = (x_1 + ab^{-1}(z_1 - y_1), x_2 + ab^{-1}(z_2 - y_2), ab^{-1}c)$. Obviously, $ab^{-1}c > 2A/\beta > 2A/2 = A$. Moreover,

$$\begin{aligned} \left| \frac{x_1 + ab^{-1}(z_1 - y_1)}{ab^{-1}c} - q_1 \right| &\leq \left| \frac{x_1}{ab^{-1}c} - q_1 \right| + \frac{|z_1 - y_1|}{c} \\ &\leq \frac{x_1}{a} |b/c - 1| + \left| \frac{x_1}{a} - q_1 \right| + \delta \\ &\leq 2q_1 |\beta - 1| + \varepsilon/4 + \delta \\ &< \varepsilon/2, \end{aligned}$$

for δ sufficiently small and β sufficiently close to 1. In a similar way, we can achieve

$$\left| \frac{x_2 + ab^{-1}(z_2 - y_2)}{ab^{-1}c} - q_2 \right| < \varepsilon/2,$$

so that $(x_1, x_2, a) \cdot (y_1, y_2, b)^{-1} \cdot (z_1, z_2, c) \in \Gamma'$, proving (8.1). □

THEOREM 8.2. — *The operators S_{ij} , for $i, j = 0, 1, 2$, are not bounded from H^1 to L^1 .*

Proof. — As remarked above, we need only consider the operators $S_{11}, S_{21}, S_{10}, S_{01}, S_{00}$.

We argue as in the proof of Theorem 4.2. This time by (2.16) for all $\sigma > 0$ and $x, y \in G$, with $x \neq y$ and $x \neq y \cdot (0, \sigma, 1)$

(8.2)

$$S_{ij}(x, y) - S_{ij}(x, y \cdot (0, \sigma, 1)) = \delta(x) \sigma X_2 k_{ji}(x^{-1}y \exp(\tau(x, y) X_2)),$$

where $0 < \tau(x, y) < \sigma$ and S_{ij} denotes the integral kernel of the operator S_{ij} .

As in the proof of Theorem 4.2 one constructs an atom a such that $\int S_{ij}(\cdot, y)a(y) d\rho(y)$ does not belong to L^1 : it suffices to apply (8.2) and Lemma 8.1. We omit the details. □

9. Unboundedness of R_{ij}

In this section we prove that the operators R_{ij} are not bounded from H^1 to L^1 , and it suffices to consider their global parts. The proof of the

unboundedness of R_{ij}^∞ is different from the previous cases. We shall now construct a sequence of functions in H^1 such that their images under the operator R_{ij}^∞ lie in L^1 but have large L^1 -norms. To do so, we first analyze the kernels k_{ij}^∞ .

LEMMA 9.1. — *For any $i, j = 0, 1, 2$, there exists a splitting $k_{ij}^\infty = k_{ij}^1 + k_{ij}^2 + k_{ij}^3$ such that*

- (i) $k_{ij}^1 = k_{ij}^\infty \chi_{\{(x_1, x_2, a) \in G: a \leq 1\}}$ is integrable;
- (ii) k_{ij}^2 is supported in the region $\{(x_1, x_2, a) \in B_1^c : a \geq 1\}$ and is integrable;
- (iii) k_{ij}^3 is supported in the region $\{(x_1, x_2, a) \in B_1^c : a \geq 1\}$ and for any $f \in L^1$

$$f * k_{ij}^3(x_1, x_2, a) = [\psi_a *_{\mathbb{R}^2} h](x_1, x_2) \quad \forall (x_1, x_2, a) \in G,$$

where $h(x_1, x_2) = \int_0^\infty f(x_1, x_2, a) da/a$, ψ is a continuous function on \mathbb{R}^2 such that $|\psi(x_1, x_2)| \leq C(1 + |(x_1, x_2)|)^{-3}$ for some C , and $\psi_a(x_1, x_2) = a^{-2}\psi(a^{-1}x_1, a^{-1}x_2)$ for $a > 0$.

Proof. — We fix a pair (i, j) and drop the indices i, j on the kernels. By (2.24) there exist constants α, β and $m, n \in \mathbb{Z}^3$ such that in B_1^c

$$k^\infty(x) = \alpha x^m e^{-2r} + \beta x^n e^{-3r} + Q(x),$$

where Q is integrable, $\beta > 0$, $|m| = 0$, $|n| = 1$ and

$$(9.1) \quad \begin{aligned} m_0 + 2 > 0 & \quad n_0 + 3 > 0 \\ m_1 + m_2 - 4 < -2 & \quad \text{and} \quad n_1 + n_2 - 6 < -2. \end{aligned}$$

We define $k^1 = k^\infty \chi_{\{(x_1, x_2, a) \in G: a \leq 1\}}$. By (9.1) and Lemma 2.2(ii), k^1 is integrable.

We now consider the region $\{(x_1, x_2, a) \in B_1^c : a \geq 1\}$. There we may approximate e^{-r} by $1/(2 \cosh r)$ and $\cosh r$ by $a(1 + |a^{-1}(x_1, x_2)|^2)$. Estimating the errors, we can write the principal terms in the expression for k^∞ above as

$$\begin{aligned} & \alpha x^m e^{-2r} + \beta x^n e^{-3r} \\ &= \alpha a^{|m|} (a^{-1}x_1)^{m_1} (a^{-1}x_2)^{m_2} \frac{1}{4a^2(1 + |a^{-1}(x_1, x_2)|^2)^2} \\ &+ \beta a^{|n|} (a^{-1}x_1)^{n_1} (a^{-1}x_2)^{n_2} \frac{1}{8a^3(1 + |a^{-1}(x_1, x_2)|^2)^3} \\ &+ q(x) \\ &= a^{-2}\psi(a^{-1}(x_1, x_2)) + q(x), \end{aligned}$$

where

$$\psi(x_1, x_2) = \alpha \frac{x_1^{m_1} x_2^{m_2}}{4(1 + |(x_1, x_2)|^2)^2} + \beta \frac{x_1^{n_1} x_2^{n_2}}{8(1 + |(x_1, x_2)|^2)^3}$$

and

$$q(x) = O(x^m e^{-3r} + x^n e^{-4r}).$$

By Lemma 2.2(i) and (9.1), q is integrable in the region where $a \geq 1$, and $|\psi(x_1, x_2)| \leq C(1 + |(x_1, x_2)|)^{-3}$.

Define $k^3(x_1, x_2, a) = a^{-2}\psi(a^{-1}(x_1, x_2))\chi_{\{(x_1, x_2, a) \in B_1^c: a \geq 1\}}$ and $k^2 = k^\infty - k^3 - k^1$. Then

$$k^2(x_1, x_2, a) = Q(x) + q(x),$$

and so k^2 is integrable, which proves (ii).

Given a function f in L^1 , we obtain

$$\begin{aligned} f * k^3(x_1, x_2, a) &= \\ &= \int_0^\infty \int \int_{\mathbb{R}^2} f(x_1 - ab^{-1}y_1, x_2 - ab^{-1}y_2, ab^{-1}) b^{-2} \psi(b^{-1}y_1, b^{-1}y_2) \\ &\quad dy_1 dy_2 db/b \\ &= \int_0^\infty \int \int_{\mathbb{R}^2} f(x_1 - az_1, x_2 - az_2, ab^{-1}) \psi(z_1, z_2) dz_1 dz_2 db/b \\ &= \int \int_{\mathbb{R}^2} a^{-2} \psi(a^{-1}v_1, a^{-1}v_2) \int_0^\infty f(x_1 - v_1, x_2 - v_2, c) dc/c dv_1 dv_2 \\ &= [\psi_a *_{\mathbb{R}^2} h](x_1, x_2), \end{aligned}$$

which proves (iii). □

We remark that in [14, Section 7], the analog of Lemma 9.1 was proved for the operators $R_{ij}^r = X_i^r X_j^r (\Delta^r)^{-1}$. We could also deduce Lemma 9.1 from that result.

We shall need the following technical lemma, which shows how to construct functions in $H^1(G)$ from functions in $H^1(\mathbb{R}^2)$.

LEMMA 9.2. — *For any function h in $H^1(\mathbb{R}^2)$, there exists a function f in $H^1(G)$ such that $\|f\|_{H^1(G)} \leq \|h\|_{H^1(\mathbb{R}^2)}$ and*

$$h(x_1, x_2) = \int_0^\infty f(x_1, x_2, a) \frac{da}{a}.$$

Proof. — Let h be in $H^1(\mathbb{R}^2)$. Take a decomposition of h as $\sum_j \lambda_j b_j$, where $\lambda_j \in \mathbb{C}$, $\sum_j |\lambda_j| < \infty$ and b_j are atoms in \mathbb{R}^2 . The atom b_j is supported in a square Q_j of side L_j , and $\int b_j = 0$ and $\|b_j\|_\infty \leq L_j^{-2}$. We

choose $r_j > 0$ such that either $r_j < 1$ and $e^2 r_j \leq L_j < e^8 r_j$ or $r_j \geq 1$ and $e^{2r_j} \leq L_j < e^{8r_j}$. Define

$$a_j(x_1, x_2, a) = \frac{1}{2} r_j^{-1} \chi_{[e^{-r_j}, e^{r_j}]}(a) b_j(x_1, x_2).$$

The functions a_j are atoms in G supported in the Calderón–Zygmund sets $R_j = Q_j \times [e^{-r_j}, e^{r_j}]$. Now define $f = \sum_j \lambda_j a_j$. It is easy to check that f is $H^1(G)$ and has the required properties. □

We now concentrate on the part of the kernel which is not integrable, i.e., k_{ij}^3 .

LEMMA 9.3. — *The operator $f \mapsto f * k_{ij}^3$ is not bounded from H^1 to L^1 .*

Proof. — The proof will follow those of [13, Lemmata 13, 14]. We will define a sequence of functions h_N in the Hardy space $H^1(\mathbb{R}^2)$ such that $\|\psi_a * h_N\|_1 / \|h_N\|_{H^1(\mathbb{R}^2)}$ is large. From Lemma 9.2, we then obtain a sequence of functions f_N in $H^1(G)$ such that $\|f_N * k_{ij}^3\|_1 / \|f_N\|_{H^1}$ is not uniformly bounded.

Let ϕ be a C^∞ -function in \mathbb{R}^2 supported in $[-1, 1] \times [-1, 1]$ such that $\int \phi = 0$, and $\psi *_{\mathbb{R}^2} \phi(0, 0) \neq 0$. Let $L > 1$ and let N be the greatest natural number with $N < \log L$. Let p, q be large natural numbers to be chosen later. Define

$$(9.2) \quad h_N = \sum_{n=0}^N \sum_{k \in \mathbb{Z}^2, |k_i| < (2^{qn} L - 1)/p} \pm \phi_{nk},$$

where the signs will be chosen later and

$$\phi_{nk}(x_1, x_2) = \phi(2^{qn} x_1 - p k_1, 2^{qn} x_2 - p k_2).$$

Let n and k be as in the double sum. Since $\text{supp } \phi \subset [-1, 1] \times [-1, 1]$, we conclude

$$\text{supp } \phi_{nk} \subset [2^{-nq}(p k_1 - 1), 2^{-nq}(p k_1 + 1)] \times [2^{-nq}(p k_2 - 1), 2^{-nq}(p k_2 + 1)].$$

It follows that h_N is supported in $[-L, L] \times [-L, L]$.

Claim 1. One can choose p, q and $t > 0$ independently of N so that for all sign choices in (9.2)

$$\rho(\{(x_1, x_2, a) : |\psi_a * h_N(x_1, x_2)| > t\}) \geq C N L^2.$$

Claim 2. The signs in (9.2) can be chosen so that $\|h_N\|_2 \leq C \sqrt{N} L$.

Proof of Claim 1. Since $\psi *_{\mathbb{R}^2} \phi(0, 0) \neq 0$, there exists a positive δ such that $|\psi_a *_{\mathbb{R}^2} \phi(x_1, x_2)| > \delta$ for (x_1, x_2, a) in a neighbourhood U of $(0, 0, 1)$

in $\mathbb{R}^2 \times \mathbb{R}^+ = G$. We can take U contained in $[-1, 1] \times [-1, 1] \times [1/2, 2]$. It follows that

$$(9.3) \quad |\psi_a *_{\mathbb{R}^2} \phi_{nk}(x_1, x_2)| > \delta \quad \text{if } (x_1, x_2, a) \in U_{nk},$$

where $U_{nk} = \{(x_1, x_2, a) : (2^{qn}x_1 - pk_1, 2^{qn}x_2 - pk_2, 2^{qn}a) \in U\}$. The sets U_{nk} are mutually disjoint and $\rho(U_{nk}) = 2^{-2qn}\rho(U)$.

Now fix $0 \leq m \leq N$, $\ell \in \mathbb{Z}^2$ such that $|\ell_i| < (2^{qm}L - 1)/p$ and take $(x_1, x_2, a) \in U_{m\ell}$. By (9.3), in the sum

$$\sum_{n=0}^N \sum_{k \in \mathbb{Z}^2, |k_i| < (2^{qn}L-1)/p} \pm \psi_a * \phi_{nk}(x_1, x_2),$$

the term with $n = m$, $k = \ell$ is greater than δ in absolute value. The other terms are much smaller; more precisely, we can choose p, q such that

$$(9.4) \quad \sum_{(n,k) \neq (m,l)} \sum |\psi_a * \phi_{nk}(x_1, x_2)| \leq \delta/2.$$

The proof of (9.4) is the same as [13, Proof of Claim 1, page 277], and we omit it.

This means that

$$\{(x_1, x_2, a) : |\psi_a * h_N(x_1, x_2, a)| > \delta/2\} \supseteq \bigcup_{n=0}^N \bigcup_{k \in \mathbb{Z}^2, |k_i| \leq (2^{qn}L-1)/p} U_{nk}.$$

Thus, choosing $t = \delta/2$,

$$\begin{aligned} \rho(\{(x_1, x_2, a) : |\psi_a * h_N(x_1, x_2, a)| > t\}) &\geq \sum_{n=0}^N \sum_{k \in \mathbb{Z}^2, |k_i| < (2^{qn}L-1)/p} \rho(U_{nk}) \\ &\geq C \rho(U) \sum_{n=0}^N 2^{-2qn} (2^{qn}L - 1)^2 \\ &\geq C N L^2. \end{aligned}$$

Proof of Claim 2. This proof follows the idea of [13, Proof of Claim 2, page 279]. On the set of all sign choices in (9.2), consider the probability measure which makes the signs into independent Bernoulli variables. Denote by \mathbf{E} the corresponding expectation. Then

$$\begin{aligned} \mathbf{E}|h_N|^2(x) &= \sum_{n,k} |\phi_{nk}(x)|^2 \\ &= \sum_{n=0}^N \sum_{k \in \mathbb{Z}^2, |k_i| < (2^{qn}L-1)/p} |\phi(2^{qn}x_1 - pk_1, 2^{qn}x_2 - pk_2)|^2 \end{aligned}$$

and

$$\begin{aligned} & \int \mathbf{E} |h_N(x_1, x_2)|^2 dx_1 dx_2 \\ & \leq C \sum_{n=0}^N \sum_{k \in \mathbb{Z}^2, |k_i| < (2^{qn} L - 1)/p} 2^{-2nq} \\ & \leq C N L^2. \end{aligned}$$

Thus $\mathbf{E} \|h_N\|_2 \leq C \sqrt{N} L$, and Claim 2 follows.

If we choose p, q and the signs in (9.2) as in Claim 1 and 2, the function h_N will be a multiple of a $(1, 2)$ -atom in \mathbb{R}^2 (see [8]). Indeed, it is supported in $[-L, L] \times [-L, L]$, with integral zero, and $\|h_N\|_2 \leq C \sqrt{N} L$. In particular, h_N is in $H^1(\mathbb{R}^2)$ and $\|h_N\|_{H^1(\mathbb{R}^2)} \leq C L^2 \sqrt{N}$.

By Lemma 9.2, there exists f_N in $H^1(G)$ such that $\|f_N\|_{H^1(G)} \leq C L^2 \sqrt{N}$ and

$$h_N(x_1, x_2) = \int_0^\infty f_N(x_1, x_2, a) \frac{da}{a}.$$

Thus by Lemma 9.1 and Claim 1,

$$\begin{aligned} \|f_N * k_{ij}^3\|_1 &= \int_0^\infty \int_{\mathbb{R}^2} |\psi_a * h_N(x_1, x_2, a)| d\rho(x_1, x_2, a) \\ &\geq t \rho(\{(x_1, x_2, a) : |\psi_a * h_N(x_1, x_2, a)| > t\}) \\ &\geq C N L^2. \end{aligned}$$

This shows that $\|f_N * k_{ij}^3\|_1 / \|f_N\|_{H^1(G)}$ is not uniformly bounded, proving the lemma. \square

THEOREM 9.4. — *The operators R_{ij}^∞ , for $i = 0, 1, 2$, are not bounded from H^1 to L^1 .*

Proof. — This is a direct consequence of Lemmata 9.1 and 9.3. \square

BIBLIOGRAPHY

- [1] G. ALEXOPOULOS, “An application of homogenization theory to harmonic analysis: Harnack inequalities and Riesz transforms on Lie groups of polynomial growth”, *Can. J. Math.* **44** (1992), p. 691-727.
- [2] J. ANKER, “Sharp estimates for some functions of the Laplacian on noncompact symmetric spaces”, *Duke Math. J.* **65** (1992), p. 257-297.
- [3] J. ANKER, E. DAMEK & C. YACOUB, “Spherical Analysis on harmonic AN groups”, *Ann. Scuola Nom. Sup. Pisa Cl. Sci.* **23** (1996), p. 643-679.
- [4] P. AUSCHER & T. COULHON, “Riesz transform on manifolds and Poincaré inequalities”, *Ann. Sci. École Norm. Sup.* **4** (2005), p. 531-555.
- [5] P. AUSCHER, T. COULHON, X. DUONG & S. HOFMANN, “Riesz transform on manifolds and heat kernel regularity”, *Ann. Sci. École Norm. Sup.* **37** (2004), p. 911-957.
- [6] D. BAKRY, “Étude des transformations de Riesz dans les variétés riemanniennes à courbure de Ricci minorée”, *Séminaire de Probabilité XXI, Lecture Notes in Math.* **1247** (1987), p. 137-172.
- [7] J. CHEN & C. LUO, “Duality of H^1 and BMO on positively curved manifolds and their characterizations”, *Lecture Notes in Math.* **1494** (1991), p. 23-38.
- [8] R. COIFMAN & G. WEISS, “Extensions of Hardy spaces and their use in Analysis”, *Bull. Am. Math. Soc.* **83** (1977), p. 569-645.
- [9] T. COULHON & X. DUONG, “Riesz transforms for $1 \leq p \leq 2$ ”, *Trans. Amer. Math. Soc.* **351** (1999), p. 1151-1169.
- [10] ———, “Riesz transforms for $p > 2$ ”, *C. R. Acad. Sci. Paris Sér. I Math.* **332** (2001), p. 975-980.
- [11] ———, “Estimates of lower bounds for the heat kernel on conical manifolds and Riesz transform”, *Arch. Math. (Basel)* **83** (2004), p. 229-242.
- [12] M. COWLING, G. GAUDRY, S. GIULINI & G. MAUCERI, “Weak type $(1, 1)$ estimates for heat kernel maximal functions on Lie groups”, *Trans. Amer. Math. Soc.* **323** (1991), p. 637-649.
- [13] G. GAUDRY, T. QIAN & P. SJÖGREN, “Singular integrals associated to the Laplacian on the affine group $ax + b$ ”, *Ark. Mat.* **30** (1992), p. 259-281.
- [14] G. GAUDRY & P. SJÖGREN, “Singular integrals on Iwasawa NA groups of rank 1”, *J. Reine Angew. Math.* **479** (1996), p. 39-66.
- [15] ———, “Haar-like expansions and boundedness of a Riesz operator on a solvable Lie group”, *Math. Z.* **232** (1999), p. 241-256.
- [16] S. GIULINI & P. SJÖGREN, “A note on maximal functions on a solvable Lie group”, *Arch. Math. (Basel)* **55** (1990), p. 156-160.
- [17] W. HEBISCH & T. STEGER, “Multipliers and singular integrals on exponential growth groups”, *Math. Z.* **245** (2003), p. 37-61.
- [18] N. LOHOUE & N. VAROPOULOS, “Remarques sur les transformées de Riesz sur les groupes de Lie nilpotents”, *C. R. Acad. Sci. Paris Sér. I Math.* **301** (1985), p. 559-560.
- [19] M. MARIAS & E. RUSS, “ H^1 -boundedness of Riesz transforms and imaginary powers of the Laplacian on Riemannian manifolds”, *Ark. Mat.* **41** (2003), p. 115-132.
- [20] E. RUSS, “ H^1 - L^1 boundedness of Riesz transforms on Riemannian manifolds and on graphs”, *Potential Anal.* **14** (2001), p. 301-330.
- [21] L. SALOFF-COSTE, “Analyse sur les groupes de Lie à croissance polynomiale”, *Ark. Mat.* **28** (1990), p. 315-331.
- [22] P. SJÖGREN, “An estimate for a first-order Riesz operator on the affine group”, *Trans. Amer. Math. Soc.* **351** (1999), p. 3301-3314.

- [23] E. STEIN, *Harmonic Analysis*, Princeton University Press, 1993.
- [24] M. VALLARINO, "Spaces H^1 and BMO on exponential growth groups", submitted, 2007.

Manuscrit reçu le 19 avril 2007,
accepté le 21 juin 2007.

Peter SJÖGREN
Göteborg University and
Chalmers University of Technology
Department of Mathematical Sciences
412 96 Göteborg (Sweden)
peters@math.chalmers.se

Maria VALLARINO
Università di Milano-Bicocca
Dipartimento di Matematica e Applicazioni
Via R. Cozzi 53
20125 Milano (Italy)
maria.vallarino@unimib.it