



ANNALES

DE

L'INSTITUT FOURIER

Luis E. SOLÁ CONDE & Matei TOMA

Maximal rationally connected fibrations and movable curves

Tome 59, n° 6 (2009), p. 2359-2369.

http://aif.cedram.org/item?id=AIF_2009__59_6_2359_0

© Association des Annales de l'institut Fourier, 2009, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

MAXIMAL RATIONALLY CONNECTED FIBRATIONS AND MOVABLE CURVES

by Luis E. SOLÁ CONDE & Matei TOMA (*)

ABSTRACT. — A well known result of Miyaoka asserts that a complex projective manifold is uniruled if its cotangent bundle restricted to a general complete intersection curve is not nef. Using the Harder-Narasimhan filtration of the tangent bundle, it can moreover be shown that the choice of such a curve gives rise to a rationally connected foliation of the manifold. In this note we show that, conversely, a movable curve can be found so that the maximal rationally connected fibration of the manifold may be recovered as a term of the associated Harder-Narasimhan filtration of the tangent bundle.

RÉSUMÉ. — Un résultat bien connu de Miyaoka affirme qu'une variété projective est uniréglée si son fibré tangent restreint à une courbe intersection complète générale n'est pas nef. De plus, en utilisant la filtration de Harder-Narasimhan, on peut montrer que le choix d'une telle courbe induit des feuilletages rationnellement connexes de la variété. Dans cette note nous montrons qu'une courbe mobile peut être trouvée telle que la fibration rationnellement connexe maximale soit associée à un terme de la filtration de Harder-Narasimhan correspondante du fibré tangent.

1. Introduction

Since the 1980's it has become evident that rational curves play a central role in birational algebraic geometry, bringing uniruled varieties -that is, varieties covered by rational curves- into the area of interest of many researchers.

Several characterizations of uniruledness may be found in the literature, all of them relating this property with positivity properties of the tangent

Keywords: Uniruled variety, maximal rationally connected fibration, movable curve, Harder-Narasimhan filtration.

Math. classification: 14J99, 14D99, 53C12.

(*) The authors wish to thank the Max-Planck-Institut für Mathematik in Bonn and the Institut Elie Cartan Nancy for hospitality and financial support during the preparation of this paper. The collaboration between the authors was partially supported by the project MTM2006-04785 of the Spanish Government.

bundle. In 1987 Miyaoka and Mori proved that a complex projective manifold X is uniruled if and only if K_X is negative on a curve C passing by x for general $x \in X$ (cf. [13]). This result has been recently improved by Boucksom, Demailly, Păun and Peternell, who have proved in [1] that in fact X is uniruled if and only if K_X is not pseudoeffective.

Also in 1987, Miyaoka provided a more geometric criterion of uniruledness: a complex projective manifold X is uniruled if the restriction of T_X to a general complete intersection curve has a vector subbundle of positive degree ([12], see also [8]). In fact, under the above hypotheses, the lower terms of the Harder-Narasimhan filtration of T_X with respect to the chosen complete intersection curve are rationally connected foliations.

On the other side, one may associate to a uniruled variety X its maximally rationally connected fibration (MRCF for brevity), constructed by Campana and by Kollár, Miyaoka and Mori independently, see [2] and [10].

In this note we address the question whether the MRCF of X may be recovered as one of the foliations provided by Miyaoka's construction for a suitably chosen complete intersection curve. Unfortunately it is not even clear whether one may always find a complete intersection curve on which K_X has negative degree (see Example 3.8), so we have decided to work with a broader class of curves.

Given a complex projective manifold X we will denote by N_{amp} the interior of the closed cone $\overline{\text{ME}}(X)$ generated by movable curves, see [1, Def. 1.3]. One has a notion of slope and stability of torsion free sheaves on X with respect to a class in $\overline{\text{ME}}(X)$, cf. [3]. Our main result is the following

THEOREM 1.1. — *Let X be a uniruled complex projective manifold, and let $G \subset T_X$ denote the foliation associated with its MRCF. There exists a class $\alpha \in N_{\text{amp}}$ represented by a reduced movable curve C such that $G|_C$ is ample and G is a member of the Harder-Narasimhan filtration of T_X with respect to α .*

As a corollary we get a positive answer to our initial question when $\dim X = 2$.

Acknowledgement: We wish to thank Frédéric Campana for useful discussions, Carolina Araujo for pointing to us an error in a previous version of the manuscript and the referee for his comments. After having submitted this paper, we were informed that in the two-dimensional case and in the rationally connected case Sebastian Neumann and Thomas Peternell obtained similar results by different methods (arXiv:0811.3398v1, arXiv:0807.0982v1).

2. Preliminaries

Along this paper X will denote a complex projective manifold of dimension at least two.

In this section we first review known facts about stability with respect to movable classes. We refer the interested reader to [3] for further details. Finally we discuss the relation of that concept with the maximal rationally connected fibration of X .

DEFINITION 2.1. — A curve $C \subset X$ is called *movable* if there exists an irreducible algebraic family of curves containing C as a reduced member and dominating X . As usual, we denote by $N_1(X)$ the vector space of numerical classes of real 1-cycles on X and by $N^1(X)$ its dual. The closure in $N_1(X)$ of the convex cone generated by classes of movable curves will be denoted by $\overline{\text{ME}}(X)$, and its interior by N_{amp} . The elements of $\overline{\text{ME}}(X)$ are called *movable classes*.

By a theorem of Boucksom, Demailly, Păun and Peternell (cf. [1]), $\overline{\text{ME}}(X)$ equals the dual of the Neron-Severi part of the *pseudoeffective* cone, i.e. the closure in $N^1(X)$ of the convex cone generated by classes of effective divisors. The full (transcendental) pseudoeffective cone may be described as the cone of classes of positive closed $(1, 1)$ -currents on X .

DEFINITION 2.2. — A movable class α is called *weakly free* if given any codimension 2 subset $B \subset X$ there exists a representative C of α such that $C \cap B = \emptyset$.

Remark 2.3. — A smooth curve $C \subset X$ is usually called *free* (cf. [9, II.3]) if $H^1(C, T_X|_C) = 0$ and $T_X|_C$ is globally generated. Moreover, a rational curve C is called *very free* if $T_X|_C$ is ample. By [9, II.3.7], the classes of free and very free curves are weakly free.

We may associate a notion of stability to every movable class α .

DEFINITION 2.4. — Given a movable class $\alpha \in \overline{\text{ME}}(X)$ and a torsion free sheaf E on X , we define the slope of E with respect to α in the usual way:

$$\mu_\alpha(E) := \frac{c_1(E) \cdot \alpha}{\text{rank}(E)}.$$

When there is no possible confusion about the class we are using, we will denote the slope simply by μ .

We say that a torsion free sheaf E on X is *stable* (resp. *semistable*) with respect to α if the slope of every coherent subsheaf of E is smaller than (resp. smaller than or equal to) $\mu_\alpha(E)$.

Note that if α is the class of a complete intersection curve of ample divisors H_1, \dots, H_{n-1} , these concepts coincide with the usual notions of stability and semistability with respect to the polarizations H_1, \dots, H_{n-1} .

The following remark is well known for the usual concept of stability, and the same proof can be applied verbatim in our setting:

Remark 2.5. — For a fixed movable class α and a torsion free sheaf E on a variety X , the set of slopes of torsion free subsheaves of E has a maximum, and the saturated subsheaf of maximal rank for which the maximal slope is achieved is called the *maximally destabilizing subsheaf* of E . By recursion one obtains the *Harder-Narasimhan filtration* of E with respect to α , which is the only filtration

$$\mathrm{HN}^\alpha(E) : 0 = E_0 \subsetneq E_1 \subsetneq \dots \subsetneq E_r = E$$

such that the quotients E_i/E_{i-1} are torsion free semistable sheaves, and the sequence of slopes $(\mu_i(E) := \mu(E_i/E_{i-1}))_i$ is strictly decreasing. The numbers $\mu_1(E)$ and $\mu_r(E)$ are also called *maximal slope* and *minimal slope* of E , respectively.

If $\alpha = [C]$ we may also consider the Harder-Narasimhan filtration of $E|_C$. Although it is not true in general that this filtration coincides with the restriction of $\mathrm{HN}^\alpha(E)$, it does when C is a general complete intersection curve. This follows from the Mehta-Ramanathan restriction theorem (cf. [11], see also [4]) and as a consequence we obtain in this case that $E_i|_C$ is ample if $\mu_i(E) > 0$.

In this paper we are interested in the Harder-Narasimhan filtration of T_X with respect to movable classes α . Let us introduce the notation that we will use below.

Notation 2.6. — Let X be a smooth complex projective variety and α be a movable class in X . We will denote by

$$\mathrm{HN}(T_X) : 0 = F_0 \subsetneq F_1 \subsetneq \dots \subsetneq F_r = T_X$$

the Harder-Narasimhan filtration of T_X with respect to α and by μ_i the slopes of F_i/F_{i-1} , for all i . We will also denote by s the integer $s = \max(\{0\} \cup \{i \mid \mu_i > 0\})$.

The first property we are concerned with is the behaviour of $\mathrm{HN}(T_X)$ with respect to birational transformations:

LEMMA 2.7. — *Let $\alpha = [C]$ be a movable class on X , and let $\pi : \tilde{X} \rightarrow X$ be projective birational morphism. Assume that C does not intersect the image B of the exceptional locus E of π . Then the Harder-Narasimhan*

filtrations of T_X with respect to α and of $T_{\tilde{X}}$ with respect to the pull-back class $\tilde{\alpha}$ of α coincide over $X \setminus B$. Thus the Harder-Narasimhan filtration of T_X with respect to weakly free classes is birationally invariant.

Proof. — Let $0 = F_0 \subset F_1 \subset \dots \subset F_r = T_X$ denote the Harder-Narasimhan filtration of T_X with respect to α . For every i let us denote by \tilde{F}_i the saturation in $T_{\tilde{X}}$ of the extension by zero of $F_i|_U$, where U denotes the open set $X \setminus B \cong \tilde{X} \setminus E$.

By construction $\mu_{\tilde{\alpha}}(\tilde{F}_i/\tilde{F}_{i-1}) = \mu_{\alpha}(F_i/F_{i-1})$, hence it suffices to show that $\tilde{F}_i/\tilde{F}_{i-1}$ is semistable with respect to $\tilde{\alpha}$ for all i . In fact, given any destabilizing subsheaf $D \subset \tilde{F}_i/\tilde{F}_{i-1}$, the saturation in T_X of the extension by zero of $D|_U$ would destabilize F_i/F_{i-1} , a contradiction. \square

Using the fact that the tensor product of semistable sheaves, modulo torsion, is again semistable ([3, Appendix]), one checks that the minimal slope of $F_i \otimes F_i$ is twice the minimal slope of F_i , hence larger than the maximal slope of T_X/F_i provided $i \leq s$. This implies that $\text{Hom}(F_i \otimes F_i, T_X/F_i) = 0$, in particular the O’Neill-tensor $N : F_i \otimes F_i \rightarrow T_X/F_i$ induced by the Lie bracket has to vanish and thus F_i is integrable for each $i \leq s$.

If $s > 0$ and α is the class of a general complete intersection curve C , then $F_s|_C$ is ample and [8] allows us to say that the leaves of F_s are algebraic and rationally connected. Unfortunately, since the ampleness of $F_s|_C$ is not known for a general movable curve C , the previous argument cannot be used in general. Nevertheless the following property is still fulfilled:

LEMMA 2.8. — *With the notation in 2.6, let G be a subsheaf of T_X and assume further that $\alpha = [C]$, where C is a curve contained in the regular locus of the distributions G and F_s (that is, the locus where T_X/G and T_X/F_s are locally free). If moreover $G|_C$ is ample, then G is contained in F_s .*

Proof. — Assume the contrary. By the ampleness of $G|_C$ we get $\mu(G + F_s/F_s) = \mu(G/G \cap F_s) > 0$, therefore $\mu_{s+1} > 0$, contradicting the choice of s . \square

The following result will be used in the case of the MRCF.

LEMMA 2.9. — *Let X and Z be smooth projective varieties of positive dimension, $\pi : X \dashrightarrow Z$ a rational map, G the foliation on X associated to π and α a weakly free class on X . If $G \subsetneq F_s^\alpha$, then Z is uniruled.*

Proof. — Since by Lemma 2.7 the Harder-Narasimhan filtration of T_X with respect to weakly free classes is birationally invariant, we may assume that π is a morphism. The same property will allow us to reduce ourselves

to the situation when π is also equidimensional. Indeed, over some open subset V of Z the morphism π coincides with the universal family over some locally closed subvariety U of the corresponding Chow scheme of X . Let Z_1 be the closure of U and X_1 be the universal family over Z_1 . Let further $Z_2 \rightarrow Z_1$ denote a desingularization of Z_1 and X_2 the normalization of $X_1 \times_{Z_1} Z_2$. We get the following diagram

$$\begin{array}{ccccccc}
 & & & \text{bir} & & & \\
 X_2 & \longrightarrow & X_1 & \dashrightarrow & U^{\subset} & \longrightarrow & X \\
 \downarrow \pi_2 & & \downarrow & & \downarrow & & \downarrow \pi \\
 Z_2 & \longrightarrow & Z_1 & \dashrightarrow & V^{\subset} & \longrightarrow & Z
 \end{array}$$

where the composition of the rational maps in the upper row is a birational morphism. The pullback of α by this morphism is a weakly free class on X_2 .

We now replace $\pi : X \rightarrow Z$ by $\pi_2 : X_2 \rightarrow Z_2$ which is equidimensional and change notation accordingly. The fact that X could be singular will not bother us since α is weakly free.

As before the leaves of the foliation F_s contain the leaves of G .

We claim that the foliation F_s descends to a foliation E on Z . Indeed, consider a small analytic neighborhood W around a general point x of a fixed general fiber of π . The foliation defined by F_s is regular around x and is a product in W . Each leaf L of F_s in W projects onto a smooth closed submanifold S of $\pi(U)$ with $\dim S = \text{rank}(F_s/G)$. Therefore the closure of the leaf of F_s in $\pi^{-1}(\pi(W))$ containing L is $\pi^{-1}(S)$. Since $\pi^{-1}(\pi(W))$ is a neighbourhood of $\pi^{-1}(\pi(x))$ we see that S is independent of the choice of $x \in \pi^{-1}(\pi(x))$. This defines the desired foliation on Z .

Choose now a movable curve C representing α and avoiding $\text{Sing}X$, the singular sets of the foliations F_s and G as well as the pull-back of the singular set of the foliation E . Then in a neighbourhood of C it is easy to check that F_s/G and π^*E are locally free and that there exists an injective morphism $F_s/G \rightarrow \pi^*E$. This implies $\text{deg}_\alpha(\pi^*E) \geq \text{deg}_\alpha(F_s/G) > 0$. Thus E has positive degree with respect to the movable curve $C' = \pi(C)$ and by [3, Thm. 1.9] it follows that Z is uniruled. \square

3. Movable curves and rationally connected foliations

In this section we prove Theorem 1.1 and make some comments on the two dimensional case.

3.1. The rationally connected case

For rationally connected varieties the statement of the main theorem takes the following form.

PROPOSITION 3.1. — *Let X be a rationally connected manifold. There exists a class $\alpha \in N_{\text{amp}}$ represented by a movable curve C such that $T_X|_C$ is ample.*

In order to prove this result we need some preparations.

DEFINITION 3.2. — Let T be a positive closed current of bidegree $(1, 1)$ on X and $f : C \rightarrow X$ a non-constant morphism from a smooth connected curve C to X . Choose local $i\partial\bar{\partial}$ -potentials u_i for T on open sets $U_i \subset X$ and suppose that $f(C) \cap U_i$ is not contained in the polar set of u_i for some i . Then define locally f^*T as $i\partial\bar{\partial}(u_i \circ f)$.

LEMMA 3.3. — *Under the above conditions f^*T is a well defined positive closed current of bidegree $(1, 1)$ on C and $[f^*T] = f^*[T]$ where $[\]$ denotes taking cohomology classes.*

Proof. — There exists a smooth closed $(1, 1)$ -form η in the cohomology class of T . Therefore $T - \eta = i\partial\bar{\partial}v$. We may assume U_i to be biholomorphic to open balls, hence $\eta|_{U_i} = i\partial\bar{\partial}v_i$ for some smooth functions v_i . Thus f^*T is given locally as $i\partial\bar{\partial}((v_i + v|_{U_i}) \circ f)$ showing that $[f^*T] = [f^*\eta] = f^*[T]$. \square

DEFINITION 3.4. — We shall say that a (connected) family of curves $(C_s)_{s \in S}$ on X is good if there exists some closed subset $A \subset X$ of codimension two such that for all $x \in X \setminus A$ there exist C_{s_1}, \dots, C_{s_n} curves in the family passing through x and analytically open connected local branches $C_{s_i}^0$ of C_{s_i} at x with the following properties

- (1) $C_{s_i}^0$ is smooth at x for all i ,
- (2) $\sum_{i=1}^n T_x C_{s_i}^0 = T_x X$,
- (3) for all i there are $n - 1$ -dimensional subfamilies in S of curves whose connected local branches give foliations locally around x which contain $C_{s_i}^0$ as a leaf.

LEMMA 3.5. — *If $(C_s)_{s \in S}$ is a good family of curves on X and T is a positive closed current of bidegree $(1, 1)$ such that the intersection product $[T][C_s]$ vanishes, then $T = 0$.*

Proof. — Take $f : C'_s \rightarrow C_s$ the morphism of normalization. Since the family is good f^*T is well defined for general $s \in S$ and we see that the hypothesis $[T][C_s] = 0$ implies $f^*T = 0$. Now adopting the notations of the

above definition, let $x \in X \setminus A$ and $\pi_i : U_i \rightarrow B_i$ be a smooth map from an open neighborhood U_i of x onto a $n - 1$ -dimensional base B_i defining the local foliation which contains $C_{s_i}^0$ as a leaf. Let further ω_i be the pull-back to U_i of a volume form on B_i . Then our assumption on T provides $T \wedge \omega_i = 0$ on U_i and thus $T \wedge \sum_{i=1}^n \omega_i = 0$ around x . But $\sum_{i=1}^n \omega_i$ is a strictly positive $(n - 1, n - 1)$ -form and thus the $(n - 1)$ -st power of a $(1, 1)$ -positive form Ω on $\cap_{i=1}^n U_i$, cf. [3, Appendix]. Since the trace measure of T with respect to Ω vanishes, the support of T will be contained in A , hence $T = 0$. \square

COROLLARY 3.6. — *If $(C_s)_{s \in S}$ is a good family on X then $[C_s] \in N_{\text{amp}}$.*

Proof. — By [1] the class α of a curve belongs to N_{amp} if for every non-zero positive closed current T on X one has $[T]\alpha > 0$. For any non-zero positive closed current T on X there exists an element C_s such that f^*T is well defined, where $f : C'_s \rightarrow C_s$ denotes normalization. Then the above Lemma yields $[T][C_s] > 0$. \square

Proof of Proposition 3.1. — From the previous discussion, it is enough to show that any rationally connected manifold X admits good families of very free rational curves.

Take x_1 any point of X . By [9, Thm. IV.3.9] there is an immersion f of \mathbb{P}^1 such that $f(0) = x_1$ and f^*T_X is ample. Let us denote by \mathcal{M}_1 the family of curves determined by the component of the Hom-scheme $\text{Hom}(\mathbb{P}^1, X)$ containing $[f]$. It satisfies condition (1) of Definition 3.4 at x_1 by construction. Arguing as in [7, Prop. 2.3] one sees that the tangent directions to general deformations of f sending 0 to x_1 fill in almost all of $\mathbb{P}(\Omega_X|_{x_1})$, hence we get condition (2). Finally, note that the differential of the evaluation map is surjective at $(f, 0)$, by the ampleness of f^*T_X , hence the last condition at x_1 is satisfied as well.

Let us denote by $U_1 \subset X$ an open set where the family \mathcal{M}_1 satisfies (1), (2) and (3). For any $x_2 \in X \setminus U_1$ we can consider \mathcal{M}_2 and an open set U_2 constructed as above. Consider now the family \mathcal{M} of curves of the form $C_1 \cup C_2$, for $[C_i] \in \mathcal{M}_i$, $i = 1, 2$. It verifies the properties (1), (2) and (3) for every point in $U_1 \cup U_2$. By noetherian recursion the assertion follows. \square

3.2. The general case

Proof of Theorem 1.1. — Suppose that X is uniruled but not rationally connected and denote by G the foliation induced by the MRCF of X . We

first show that a weakly free curve C exists in X such that $G|_C$ is ample and $[C] \in N_{\text{amp}}$.

We take a general smooth complete intersection curve D on X avoiding the singularities of G . Following [9, II 7], we will consider combs with handle D and m teeth R_i which are smooth free rational curves contained in distinct general fibers of the rationally connected fibration, each R_i meeting D transversely in one point P_i . By Proposition 3.1 we may also assume that $G|_{R_i}$ is ample for all $1 \leq i \leq m$. By [9, II 7.9-7.10], for arbitrarily large m one can find such a comb C_0 which deforms to give a covering family $(C_t)_{t \in T}$ of curves on X such that the general member C_t is smooth and free. Moreover, $[C_t]$ is weakly free and the family $(C_t)_{t \in T}$ is good since the family of complete intersection curves containing D was already good. In particular $[C_t] \in N_{\text{amp}}$ by Corollary 3.6.

We now prove that $G|_{C_t}$ is ample. Let us denote by

$$\begin{array}{ccc} S & \xrightarrow{q} & X \\ p \downarrow & & \\ & & T \end{array}$$

the desingularization of the smoothing described above, where T is a smooth curve. In particular the central fiber $S_0 \cong C_0 \cup \bigcup_j E_j$, where $q(E_j)$ is zero-dimensional, hence $q^*(G|_{E_j})$ is trivial for every j .

We will consider the relative Harder-Narasimhan filtration

$$0 = G_0 \subsetneq G_1 \subsetneq \dots \subsetneq G_k = q^*G$$

of q^*G , see [6, 2.3]. Over some open subset $U \subset T$, it induces the usual Harder-Narasimhan filtration of $q^*(G|_{C_t})$, $t \in U$. Let G' denote the quotient q^*G/G_{k-1} modulo torsion. It will be enough to show that $\text{deg}_{C_t}(G')$ is positive.

In fact, the morphism $q^*G \rightarrow G'$ induces a surjective morphism

$$\bigwedge^r q^*G \rightarrow \det G' \otimes \mathcal{I}_Z,$$

where r denotes the rank of G' and \mathcal{I}_Z is the ideal sheaf of a zero-dimensional subscheme $Z \subset S$. Now, for general $t \in U$

$$\begin{aligned} \text{deg}_{C_t}(G') &= \text{deg}_{C_t}(\det G') = \\ &= \text{deg}_D(\det G') + \sum_{i=1}^m \text{deg}_{R_i}(\det G') + \sum_j \text{deg}_{E_j}(\det G'). \end{aligned}$$

For each component B of the curve S_0 we may consider the image P_B of the natural map $(\det G' \otimes \mathcal{I}_Z)|_B \rightarrow (\det G')|_B$, which is a quotient of

$(q^* \wedge^r G)|_B$ and thus

$$\deg_B(\det G') \geq \deg(P_B) \geq \mu_{\min} \left(q^* \wedge^r G \right) |_B.$$

The last term is nonnegative for $B = E_j$ and it is at least $1/r$ when $B = R_i$, hence

$$\deg_{C_t}(G') \geq \mu_{\min} \left(\wedge^r G \right) |_D + \frac{m}{r}$$

is strictly positive for $m \gg 0$.

Let as usual $\alpha = [C_t]$ and consider the Harder-Narasimhan filtration of T_X with respect to α . By Lemma 2.8 it follows that $G \subset F_s$. If $G \neq F_s$, then by Lemma 2.9 the target of the MRCF of X would be uniruled. But this is impossible by [5], thus $G = F_s$ and the Theorem is proved. \square

3.3. Comments on surfaces

In the case of surfaces we get the following

COROLLARY 3.7. — *Let X be a uniruled complex projective smooth surface and let G denote the foliation associated with the MRCF of X . There exists an ample divisor C on X such that $G|_C$ is ample and G is a member of the Harder-Narasimhan filtration of T_X with respect to $[C]$.*

Proof. — Since N_{amp} coincides in this case with the cone of ample divisors of X , the statement follows from Theorem 1.1. \square

For a rationally connected surface X , the above corollary asserts that there exists an ample curve C such that $T_X|_C$ is ample. But one cannot expect this property to hold for every ample curve C on X in general. This can be seen in the following example, shown to us by F. Campana.

Example 3.8. — Let X be the blow-up of \mathbb{P}^2 at sixteen general points P_1, \dots, P_{16} and let C be the strict transform of a general quintic passing through P_1, \dots, P_{16} . It is easy to check that C is a very ample divisor on X but $-K_X \cdot C = -1$.

BIBLIOGRAPHY

- [1] S. BOUCKSOM, J.-P. DEMAILLY, M. PĂUN & T. PETERNELL, “The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension”, preprint math.AG/0405285.

- [2] F. CAMPANA, “Connexité rationnelle des variétés de Fano”, *Ann. Sci. École Norm. Sup. (4)* **25** (1992), no. 5, p. 539-545.
- [3] F. CAMPANA & T. PETERNELL, “Geometric stability of the cotangent bundle and the universal cover of a projective manifold”, preprint arXiv:math/0405093.
- [4] H. FLENNER, “Restrictions of semistable bundles on projective varieties”, *Comment. Math. Helv.* **59** (1984), no. 4, p. 635-650.
- [5] T. GRABER, J. HARRIS & J. STARR, “Families of rationally connected varieties”, *J. Amer. Math. Soc.* **16** (2003), no. 1, p. 57-67.
- [6] D. HUYBRECHTS & M. LEHN, *The geometry of moduli spaces of sheaves*, Aspects of Mathematics, E31, Friedr. Vieweg & Sohn, Braunschweig, 1997.
- [7] J.-M. HWANG, “Geometry of minimal rational curves on Fano manifolds”, in *School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000)*, ICTP Lect. Notes, vol. 6, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001, p. 335-393.
- [8] S. KEBEKUS, L. SOLÁ CONDE & M. TOMA, “Rationally connected foliations after Bogomolov and McQuillan”, *J. Algebraic Geom.* **16** (2007), no. 1, p. 65-81.
- [9] J. KOLLÁR, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, vol. 32, Springer-Verlag, Berlin, 1996.
- [10] J. KOLLÁR, Y. MIYAOKA & S. MORI, “Rationally connected varieties”, *J. Algebraic Geom.* **1** (1992), no. 3, p. 429-448.
- [11] V. B. MEHTA & A. RAMANATHAN, “Semistable sheaves on projective varieties and their restriction to curves”, *Math. Ann.* **258** (1981/82), no. 3, p. 213-224.
- [12] Y. MIYAOKA, “Deformations of a morphism along a foliation and applications”, in *Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985)*, Proc. Sympos. Pure Math., vol. 46, Amer. Math. Soc., Providence, RI, 1987, p. 245-268.
- [13] Y. MIYAOKA & S. MORI, “A numerical criterion for uniruledness”, *Ann. of Math. (2)* **124** (1986), no. 1, p. 65-69.

Manuscrit reçu le 8 novembre 2008,
 accepté le 27 mars 2009.

Luis E. SOLÁ CONDE
 Universidad Rey Juan Carlos
 Departamento de Matemáticas
 C. Tulipán s.n. 28933
 Móstoles, Madrid (Spain)
 luis.sola@urjc.es

Matei TOMA
 Institut de Mathématiques Elie Cartan
 Nancy-Université
 B.P. 239
 54506 Vandoeuvre-lès-Nancy Cedex (France)
 and
 Institute of Mathematics of the Romanian Academy
 toma@iecn.u-nancy.fr