



ANNALES

DE

L'INSTITUT FOURIER

Nicolas MONOD & Henrik Densing PETERSEN

An obstruction to ℓ^p -dimension

Tome 64, n° 4 (2014), p. 1363-1371.

http://aif.cedram.org/item?id=AIF_2014__64_4_1363_0

© Association des Annales de l'institut Fourier, 2014, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

AN OBSTRUCTION TO ℓ^p -DIMENSION

by Nicolas MONOD & Henrik Densing PETERSEN (*)

ABSTRACT. — Let G be any group containing an infinite elementary amenable subgroup and let $2 < p < \infty$. We construct an exhaustion of $\ell^p G$ by closed invariant subspaces which all intersect trivially a fixed non-trivial closed invariant subspace. This is an obstacle to ℓ^p -dimension and gives an answer to a question of Gaboriau.

RÉSUMÉ. — Soit G un groupe contenant un sous-groupe infini élémentairement moyennable et soit $2 < p < \infty$. Nous construisons des sous- G -modules fermés de $\ell^p G$ d'union croissante dense mais qui rencontrent trivialement un sous-module fermé non trivial. Ce phénomène est un obstacle à la quête d'une dimension ℓ^p et répond à une question de Gaboriau.

Introduction and Statement of the Result

The Murray–von Neumann dimension, also called ℓ^2 -dimension, has proved tremendously valuable since its introduction in 1936 [11]. This is particularly true for its use through ℓ^2 -Betti numbers of groups since the seminal article [1] by Cheeger and Gromov. This has prompted recent work in pursuit of a more general ℓ^p -dimension for $1 < p < \infty$; see e.g. [9, 8, 5, 6]. The purpose of this short note is to establish a fundamental obstruction to this endeavour.

One of the central motivations to define a notion of ℓ^p -dimension is to take advantage of the situations where ℓ^p -cohomology carries information not accessible through to ℓ^2 -methods, see e.g. [12]. However, this involves typically p large, in particular $p > 2$; unfortunately this range appears to be much more difficult than $p \leq 2$.

Keywords: ℓ^p -dimension, abstract harmonic analysis.

Math. classification: 43A15.

(*) NM was supported in part by the Swiss National Science Foundation and the ERC; HDP acknowledges the support of the Danish National Research Foundation (DNRF) through the Centre for Symmetry and Deformation.

Given a (discrete) group G , a concrete test for the existence of an ℓ^p -dimension with at least some positivity and continuity properties is the following question of Gaboriau, see Question 1.1 in [6]. Let $(E_i)_{i \in \mathbb{N}}$ be an *invariant exhaustion* of $\ell^p G$, i.e., an increasing sequence of (left-) G -invariant closed subspaces $E_i \subseteq \ell^p G$ with dense union. If a G -invariant closed subspace $F \subseteq \ell^p G$ meets each E_i trivially, does it follow that F is trivial?

This question also seems to be the main missing point to establish the measure-equivalence invariance of the vanishing of ℓ^p -cohomology in parallel to [4]. Such invariance would notably settle the well-known open problem of the vanishing of reduced ℓ^p -cohomology for amenable groups.

The case of *amenable* groups was expected to be much more tractable, and indeed Gournay [6] has given a positive answer to Gaboriau's question when G is amenable and $p \leq 2$. The present note shows that the situation is very much opposite for $p > 2$, even when $G = \mathbb{Z}$.

THEOREM 0.1. — *Let G be an infinite elementary amenable group and let $2 < p < \infty$.*

There exists a closed invariant subspace $0 \neq F \subseteq \ell^p G$ and an invariant exhaustion $(E_i)_{i \in \mathbb{N}}$ of $\ell^p G$ such that $E_i \cap F = 0$ for all $i \in \mathbb{N}$.

Our current proof of this theorem involves notably a cameo appearance of the Feit–Thompson Theorem [3] and a result of Saeki [14] in classical harmonic analysis.

In general, we will say that an invariant exhaustion $(E_i)_i$ of $\ell^p G$ is *thin*, if there is a closed invariant subspace $F \neq 0$ satisfying $E_i \cap F = 0$ for all i . It is not hard to see (and will be needed in the proof anyway) that the existence of such exhaustions can be induced up from subgroups. We thus record the following formally stronger statement:

COROLLARY 0.2. — *Thin invariant exhaustions of $\ell^p G$ for all $p > 2$ exist more generally for every group G containing an infinite elementary amenable subgroup. In particular, this holds for every non-trivial group which is not torsion.*

Thus, for instance, any infinite linear group G over any field of any characteristic admits thin invariant exhaustions of $\ell^p G$ for all $p > 2$ because the assumptions of Corollary 0.2 are satisfied due to the Tits alternative (Theorems 1 and 2 in [16]).

Perhaps one should find another proof of Theorem 0.1. In any case, this would be necessary to address the following.

CONJECTURE 0.3. — Any infinite group G admits a thin invariant exhaustion of $\ell^p G$ for all $p > 2$.

Proof of the Result

First we observe that the property of *not* having a thin invariant exhaustion is hereditary:

LEMMA 0.4. — Let G be a group and $1 \leq p < \infty$. Suppose that G contains a subgroup H which has a thin invariant exhaustion $(E_i)_i$ of $\ell^p H$. Then G has a thin invariant exhaustion of $\ell^p G$.

Thus Corollary 0.2 follows indeed from Theorem 0.1.

Proof of Lemma 0.4. — Let G and p be given and fix a subgroup H of G and a thin invariant exhaustion $(E_i)_i$ of $\ell^p H$. We have a G -equivariant isomorphism of ℓ^p -spaces

$$\ell^p G \cong \ell^p(G/H, \ell^p H),$$

where the G -action on the right-hand side is given by $(g.\xi)(x) = c(g, g^{-1}.x) \cdot \xi(g^{-1}.x)$ for $g \in G, x \in G/H, \xi \in \ell^p(G/H, \ell^p H)$, and $c : G \times G/H \rightarrow H$ a cocycle representative for the inclusion $H < G$.

Now take $F \neq 0$ a closed invariant subspace of $\ell^p H$ such that $E_i \cap F = 0$ for all i , and define $\tilde{E}_i := \ell^p(G/H, E_i)$ and similarly $\tilde{F} := \ell^p(G/H, F)$. Then $(\tilde{E}_i)_i$ is a thin invariant exhaustion of $\ell^p G$. \square

From here on, the proof splits into two cases:

- (A) G contains an element of infinite order.
- (B) G is a torsion group.

Proof in case (A)

By Lemma 0.4, we can assume $G \cong \mathbb{Z}$. For a measure μ on a Borel space we denote the support of μ by $\text{supp } \mu$. We denote the Lebesgue measure on \mathbb{T} by λ and the Fourier-Stieltjes coefficients of μ by $\hat{\mu}$.

PROPOSITION 0.5. — Let $2 < p < \infty$ and suppose that μ is a measure on \mathbb{T} such that $\lambda(\text{supp } \mu) = 0$ and $\hat{\mu} \in \ell^p \mathbb{Z}$. Denote by F the closed, invariant subspace of $\ell^p \mathbb{Z}$ generated by $\hat{\mu}$. Then there is an invariant exhaustion $(E_i)_{i \in \mathbb{N}}$ of $\ell^p \mathbb{Z}$ such that $F \cap E_i = 0$ for all $i \in \mathbb{N}$.

Proof. — Let p, μ be given as in the statement and denote $K := \text{supp } \mu$. We can choose a decreasing family $(U_i)_{i \in \mathbb{N}}$ of open neighbourhoods of K , such that

- (i) $K = \bigcap_i U_i$.
- (ii) $\overline{U_{i+1}} \subseteq U_i$ for all $i \in \mathbb{N}$.

By the existence of a smooth partition of unity, there exist smooth functions σ_i on \mathbb{T} such that for every $i \in \mathbb{N}, t \in \mathbb{T}$ we have

$$\sigma_i(t) = \begin{cases} 0 & \text{if } t \in \overline{U_{i+1}}, \\ 1 & \text{if } t \in U_i^c. \end{cases}$$

On $U_i \setminus \overline{U_{i+1}}$, we do not care about the value $\sigma_i(t)$ as long as σ_i is smooth.

By the smoothness of σ_i , the Fourier transforms $S_i := \widehat{\sigma}_i$ are in $\ell^1\mathbb{Z}$ for all $i \in \mathbb{N}$. Further, we observe that in $\ell^1\mathbb{Z}$ we have for every $i \in \mathbb{N}$ that

$$(0.1) \quad S_{i+1} * S_i = S_i,$$

since the identity $\sigma_{i+1}(t) \cdot \sigma_i(t) = \sigma_i(t)$ holds pointwise on \mathbb{T} . Indeed, fixing some i we observe that both sides of the latter equality are zero if $t \in U_{i+1}$, and that if $t \in U_{i+1}^c$ then $\sigma_{i+1}(t) = 1$ so that the equality is tautological.

From this point on we consider the S_i as translation-invariant operators on $\ell^p\mathbb{Z}$, acting by convolution.

Since μ and σ_i have disjoint support, $\widehat{\mu}$ is in $\ker S_i$ (compare [13, Lemma 2]). It follows that $F \subseteq \ker S_i$ for all i . We define $E_i := \ker(\text{Id}_{\ell^p\mathbb{Z}} - S_i)$. The E_i are then closed, invariant subspaces, and it is obvious that $F \cap E_i = 0$ for all $i \in \mathbb{N}$.

The fact that $E_i \subseteq E_{i+1}$ for all $i \in \mathbb{N}$ follows directly from the identity (0.1). Indeed, take $x \in E_i$. Then we get $x = S_i \cdot x = (S_{i+1} S_i) \cdot x = S_{i+1} \cdot (S_i \cdot x) = S_{i+1} \cdot x$ whence $x \in E_{i+1}$.

Finally, we need to see that $\bigcup_i E_i$ is dense in $\ell^p\mathbb{Z}$. For this denote by $\iota_p : \ell^2\mathbb{Z} \rightarrow \ell^p\mathbb{Z}$ the canonical embedding. For each $i \in \mathbb{N}$ there is an orthogonal projection $p_i \in LZ$, the von Neumann algebra generated by the left-regular representation of \mathbb{Z} on $\ell^2\mathbb{Z}$, such that under the spatial isomorphism $L\mathbb{Z} \cong L^\infty\mathbb{T}$, p_i corresponds to the indicator function, $\mathbb{1}_{U_i^c}$, of the complement of U_i . Then, denoting by $S_i^{(2)}$ the operator of convolution by S_i on $\ell^2\mathbb{Z}$ we have $p_i(\ell^2\mathbb{Z}) \subseteq \ker(\text{Id}_{\ell^2\mathbb{Z}} - S_i^{(2)})$. It follows that $\iota_p(p_i(\ell^2\mathbb{Z})) \subseteq \ker(\text{Id}_{\ell^p\mathbb{Z}} - S_i) = E_i$.

Then it just remains to note that, since $\lambda(U_i^c) \nearrow_i 1$, the union $\bigcup_i p_i(\ell^2\mathbb{Z})$ is dense in $\ell^2\mathbb{Z}$. Thus $\bigcup_i \iota_p(p_i(\ell^2\mathbb{Z}))$ is dense in $\ell^p\mathbb{Z}$. \square

To complete the proof of Theorem 0.1 in case (A), we now need to know that a *non-zero* measure μ as in the statement of the previous proposition does indeed exist ($\mu \neq 0$ implies $F \neq 0$). This follows from the main

theorem of [14] as we shall now recall. Let $\varphi : c_0(\mathbb{Z}) \rightarrow c_0(\mathbb{Z})$ be a continuous map (with respect to sup-norm); for instance, φ can be the composition by any continuous map $\varphi_0 : \mathbb{R} \rightarrow \mathbb{R}$ with $\varphi_0(0) = 0$. Then Saeki proves in [14] that there is a probability measure μ with Lebesgue-null support such that the Fourier–Stieltjes coefficients $\widehat{\mu}(n)$ satisfy

$$(0.2) \quad \sum_{n \in \mathbb{Z}} |\widehat{\mu}(n)^2 \varphi(\widehat{\mu})(n)| < \infty.$$

Thus, implicitly, $\widehat{\mu}$ vanishes at infinity; see Remark (ii) below. If we choose for instance $\varphi_0(x) = |x|^{p-2}$ for $p > 2$, then (0.2) shows that $\widehat{\mu}$ is in ℓ^p , as desired. We can also obtain $\widehat{\mu}$ in all ℓ^p simultaneously if we arrange $\lim_{x \rightarrow 0} |x|^\epsilon / \varphi_0(x) = 0$ for all $\epsilon > 0$; an explicit example is $\varphi_0(x) = \exp(-\sqrt{|\log |x||})$. This finishes the proof of case (A).

Proof in case (B)

Since G is an elementary amenable torsion group, it is locally finite by Theorem 2.3 in [2]. We can now appeal to a result of Hall–Kulatilaka [7], relying in turn on the Feit–Thompson Theorem [3], to conclude that G contains an infinite abelian subgroup. Therefore, using again Lemma 0.4, we can assume that G is a countably infinite locally finite abelian group. The proof of Theorem 0.1 in this case follows the overall strategy of the case of \mathbb{Z} , if one redefines a “smooth function” to mean a locally constant function on a totally disconnected compact space. We shall therefore need to address the question of the existence of a suitable measure μ on the Pontryagin dual \widehat{G} , Proposition 0.7 below. We first record a version of Proposition 0.5 in this setup. Let $\lambda_{\widehat{G}}$ denote the normalized Haar measure on the profinite group \widehat{G} .

PROPOSITION 0.6. — *Let $2 < p < \infty$ and suppose that μ is a measure on \widehat{G} such that $\lambda_{\widehat{G}}(\text{supp } \mu) = 0$ and $\widehat{\mu} \in \ell^p G$. Denote by F the closed, invariant subspace of $\ell^p G$ generated by $\widehat{\mu}$. Then there is an invariant exhaustion $(E_i)_{i \in \mathbb{N}}$ of $\ell^p G$ such that $F \cap E_i = 0$ for all $i \in \mathbb{N}$.*

Proof. — We first observe that Lemma 2 of [13] holds in general for G countable abelian and the proof goes through verbatim in this generality. This forms the basis for imitating the proof of proposition 0.5; the only point that needs justification is the “smoothness” of σ_i to ensure $\widehat{\sigma}_i \in \ell^1 G$. In fact, we claim that we can even arrange $\widehat{\sigma}_i \in \mathbb{C}G$.

Indeed, since \widehat{G} is profinite, we can take U_i to be a compact-open subset given as the pre-image $\pi^{-1}(V_i)$ of a subset $V_i \subseteq \widehat{H}_i$ in some finite quotient $\pi : \widehat{G} \rightarrow \widehat{H}_i$, where H_i is a finite subgroup of G . The indicator function $\mathbb{1}_{U_i^c}$ of the complement is $\mathbb{1}_{V_i^c} \circ \pi$ and hence $\widehat{\mathbb{1}}_{U_i^c}$ is supported on H_i . Therefore, defining $\sigma_i = \mathbb{1}_{U_i^c}$, the transform $\widehat{\sigma}_i$ is finitely supported. Now (0.1) holds and the rest of the proof is unchanged. \square

We now need again to show that a non-zero measure μ as in Proposition 0.6 exists to finish the proof of the theorem. Saeki states in [14, Remark (II)] that his method still works as long as \widehat{G} does not contain an open subgroup of bounded order (in Saeki’s notation, the desired conclusion is obtained by choosing f to be the constant function one on our compact group \widehat{G} , which he denotes by G). He indicates however that a different technique is required when \widehat{G} does contain an open subgroup of bounded order, although that technique is not provided. We shall therefore propose a proof in the proposition below.

PROPOSITION 0.7. — *If \widehat{G} contains an open subgroup of bounded order, then it admits a probability measure μ such that $\lambda_{\widehat{G}}(\text{supp } \mu) = 0$ and $\widehat{\mu} \in \ell^p G$ for all $p > 2$.*

Proof. — The assumption allows us to apply Corollary 25.10 in [10] and deduce that \widehat{G} is isomorphic to a (countably infinite) product of finite cyclic groups $\mathbb{Z}/n_i\mathbb{Z}$ where only finitely many distinct integers n_i occur. Therefore, regrouping factors and using the Chinese remainder theorem, we can write $\widehat{G} \cong A_0 \times B^{\mathbb{N}}$ for some finite abelian groups A_0 and $B \neq 0$. Let $k \geq 2$ be the cardinality of B . Regrouping factors some more, we write

$$\widehat{G} \cong A_0 \times \prod_{n=1}^{\infty} (B^n)^{k^n} = \prod_{j=0}^{\infty} A_j,$$

where for $j \geq 1$ each A_j is a group of size c_j , a power of k , in such a way that $c_j = k^n$ occurs exactly k^n times for each $n \in \mathbb{N}$. We now define μ explicitly as the infinite product $\mu = \prod_{j=0}^{\infty} \mu_j$, where μ_0 is the uniform distribution on A_0 and μ_j is the uniform distribution on the set $A_j \setminus \{0\}$ for all $j \geq 1$.

Since $\lambda_{\widehat{G}}$ is the product of the normalized measures on each factor, the support of μ has Haar measure $\prod_{j=1}^{\infty} (1 - 1/c_j)$. This product is zero because $\sum_{j=1}^{\infty} 1/c_j = \infty$ since this is a sum of terms k^{-n} , each of which occurs k^n times.

We now fix $p > 2$ and proceed to prove that $\widehat{\mu}$ is in $\ell^p G$. We have $G \cong \bigoplus_{j=0}^{\infty} \widehat{A}_j$ and the dual Haar measure on \widehat{A}_j is the (non-renormalized)

counting measure. Therefore, it suffices to prove that the product of all ℓ^p -norms of $\widehat{\mu}_j$ over $j \geq 1$ is finite. Writing $\mathbb{1}_X$ for the indicator function of a subset X of A_j or of \widehat{A}_j , we have $\widehat{\mathbb{1}}_{\{0\}} = c_j^{-1} \mathbb{1}_{\widehat{A}_j}$ for $0 \in A_j$ and $\widehat{\mathbb{1}}_{A_j} = \mathbb{1}_{\{0\}}$ for $0 \in \widehat{A}_j$. The density of μ_j can be written as $c_j(c_j - 1)^{-1}(\mathbb{1}_{A_j} - \mathbb{1}_{\{0\}})$, which implies $\widehat{\mu}_j = \mathbb{1}_{\{0\}} - (c_j - 1)^{-1} \mathbb{1}_{\widehat{A}_j \setminus \{0\}}$. We conclude that $\|\widehat{\mu}_j\|_p^p = 1 + (c_j - 1)^{1-p}$. In other words, it remains to verify that the product $\prod_{j=1}^{\infty} (1 + (c_j - 1)^{1-p})$ is finite, or equivalently that $\sum_{j=1}^{\infty} (c_j - 1)^{1-p}$ is finite. The latter series is $\sum_{n=1}^{\infty} k^n (k^n - 1)^{1-p}$ which is indeed finite if and only if $p > 2$. \square

This completes the proof of Theorem 0.1.

Remarks

- (i) Saeki's measure μ on \mathbb{T} satisfies that $\mu * \mu$ has a continuous density. He points out that this cannot be achieved for certain groups in case (B) above, see Remark (III) of [14].
- (ii) Stating the above property (0.2) of μ requires to know a priori that $\widehat{\mu}$ vanishes at infinity. This can be verified in Saeki's proof: it follows e.g. from condition (1) on p. 230 in [14].
- (iii) If G is any amenable group, then a necessary condition for the existence of a thin invariant exhaustion $(E_i)_i$ as in Theorem 0.1, is that the invariant subspace F have ℓ^p -dimension zero in the sense of Gournay [6]. In fact, we conjecture that the converse is also true:

CONJECTURE 0.8. — *Let G be a (countable, discrete) amenable group, $1 \leq p < \infty$, and $F \subseteq \ell^p G$ a closed invariant subspace. Then $\dim_{\ell^p} F = 0$ if and only if there is an invariant exhaustion $(E_i)_i$ of $\ell^p G$ such that $E_i \cap F = 0$ for all i .*

More generally one can also consider this conjecture for sofic groups [9, 8].

This conjecture is motivated by a result of Sauer [15, Theorem 2.4] for Lück's extended dimension function.

- (iv) For $G = \mathbb{Z}^n$, $n \geq 2$ and $p > \frac{2n}{n-1}$ the situation in Theorem 0.1 is particularly nice. In this case one can take the subspace F to be the kernel of an operator $T \in \mathbb{C}\mathbb{Z}^n$ which acts without kernel on $\ell^2 \mathbb{Z}^n$, and it was noticed already by Gournay [6, Remark 9.4] that in this case the kernel in $\ell^p \mathbb{Z}^n$ must have ℓ^p -dimension zero, in the sense of [6].

For a proof that there exist a T with these properties see [13]. The proof that we can take the kernel of T as our F is analogous to the proof of Proposition 0.5.

- (v) Using an Euler characteristic argument, Gaboriau had previously observed that for certain non-amenable groups G one cannot hope to have a notion of ℓ^p -dimension for which $(\ell^p G)^{\oplus n}$ has dimension n , and which also satisfies additivity for short exact sequences (see the introduction of [5]). The fact that we produce thin exhaustions for *amenable* groups should further increase the doubts that there be any reasonable ℓ^p -dimension for large p .

Acknowledgements. Question [6, 1.1] was suggested to us by Kate Juschenko. We also thank Antoine Gournay and Antoine Derighetti for some nice discussions, and Damien Gaboriau for reading a preliminary version of this note.

BIBLIOGRAPHY

- [1] J. CHEEGER & M. GROMOV, “ L_2 -cohomology and group cohomology”, *Topology* **25** (1986), no. 2, p. 189-215.
- [2] C. CHOU, “Elementary amenable groups”, *Illinois J. Math.* **24** (1980), no. 3, p. 396-407.
- [3] W. FEIT & J. G. THOMPSON, “Solvability of groups of odd order”, *Pacific J. Math.* **13** (1963), p. 775-1029.
- [4] D. GABORIAU, “Invariants l^2 de relations d’équivalence et de groupes”, *Publ. Math. Inst. Hautes Études Sci.* (2002), no. 95, p. 93-150.
- [5] A. GOURNAY, “A dynamical approach to von Neumann dimension”, *Discrete Contin. Dyn. Syst.* **26** (2010), no. 3, p. 967-987.
- [6] ———, “Further properties of ℓ^p dimension”, *J. Funct. Anal.* **266** (2014), no. 2, p. 487-513.
- [7] P. HALL & C. R. KULATILAKA, “A property of locally finite groups”, *J. London Math. Soc.* **39** (1964), p. 235-239.
- [8] B. HAYES, “An ℓ^p -version of von Neumann dimension for Banach space representations of sofic groups II”, Preprint, arXiv:1302.2286v2.
- [9] ———, “An ℓ^p -version of von Neumann dimension for Banach space representations of sofic groups”, *J. Funct. Anal.* **266** (2014), no. 2, p. 989-1040.
- [10] E. HEWITT & K. A. ROSS, *Abstract harmonic analysis. Vol. I: Structure of topological groups. Integration theory, group representations*, Die Grundlehren der mathematischen Wissenschaften, Bd. 115, Academic Press, Inc., Publishers, New York; Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963, viii+519 pages.
- [11] F. J. MURRAY & J. VON NEUMANN, “On rings of operators”, *Ann. of Math. (2)* **37** (1936), no. 1, p. 116-229.
- [12] P. PANSU, “ L^p -cohomology of symmetric spaces”, in *Geometry, analysis and topology of discrete groups*, Adv. Lect. Math. (ALM), vol. 6, Int. Press, Somerville, MA, 2008, p. 305-326.
- [13] M. J. PULS, “Zero divisors and $L^p(G)$ ”, *Proc. Amer. Math. Soc.* **126** (1998), no. 3, p. 721-728.

- [14] S. SAEKI, “On convolution squares of singular measures”, *Illinois J. Math.* **24** (1980), no. 2, p. 225-232.
- [15] R. SAUER, “ L^2 -Betti numbers of discrete measured groupoids”, *Internat. J. Algebra Comput.* **15** (2005), no. 5-6, p. 1169-1188.
- [16] J. TITS, “Free subgroups in linear groups”, *J. Algebra* **20** (1972), p. 250-270.

Manuscrit reçu le 30 janvier 2013,
révisé le 4 février 2014,
accepté le 13 juin 2014.

Nicolas MONOD
École Polytechnique Fédérale de Lausanne
Station 8, CH-1015 Lausanne (Switzerland)
nicolas.monod@epfl.ch

Henrik Densing PETERSEN
University of Copenhagen
Department of Mathematical Sciences
Universitetsparken 5
2100 København Ø (Denmark)
hdp@math.ku.dk

Current address:
École Polytechnique Fédérale de Lausanne
Station 8, CH-1015 Lausanne (Switzerland)
henrik.petersen@epfl.ch