



ANNALES

DE

L'INSTITUT FOURIER

Mark BLUME

Toric orbifolds associated to Cartan matrices

Tome 65, n° 2 (2015), p. 863-901.

http://aif.cedram.org/item?id=AIF_2015__65_2_863_0



© Association des Annales de l'institut Fourier, 2015,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

TORIC ORBIFOLDS ASSOCIATED TO CARTAN MATRICES

by Mark BLUME (*)

ABSTRACT. — We investigate moduli stacks of pointed chains of \mathbb{P}^1 related to the Losev-Manin moduli spaces and show that these moduli stacks coincide with certain toric stacks which can be described in terms of the Cartan matrices of root systems of type A . We also consider variants of these stacks related to root systems of type B and C .

RÉSUMÉ. — Nous étudions les champs de modules des chaînes de \mathbb{P}^1 marquées, reliés aux espaces de modules de Losev-Manin, et montrons que ces champs de modules coïncident avec certains champs toriques qui peuvent être décrits en termes de matrices de Cartan de systèmes de racines de type A . Nous considérons également les variantes de ces champs liés aux systèmes de racines de type B et C .

Introduction

The Losev-Manin moduli spaces \overline{L}_n , introduced in [16], parametrise isomorphism classes of stable n -pointed chains of \mathbb{P}^1 . The space \overline{L}_n forms a compactification of the torus $(\mathbb{G}_m)^n/\mathbb{G}_m$ that parametrises n points s_1, \dots, s_n in $\mathbb{P}^1 \setminus \{0, \infty\} = \mathbb{G}_m$ up to automorphisms of \mathbb{P}^1 fixing the two points $0, \infty$. It is a smooth projective toric variety isomorphic to the toric variety $X(A_{n-1})$ associated with the root system A_{n-1} , see [2].

In the present paper we are concerned with a variant of the Losev-Manin moduli spaces which arises as a compactification of the moduli space of n indistinguishable points in $\mathbb{P}^1 \setminus \{0, \infty\}$, or equivalently, finite subschemes of degree n in $\mathbb{P}^1 \setminus \{0, \infty\}$. Isomorphism classes of such subschemes correspond to polynomials of the form $y^n + a_{n-1}y^{n-1} + \dots + a_1y + 1$ up to multiplication

Keywords: Losev-Manin moduli spaces, toric stacks, root systems, Cartan matrices, permutohedron.

Math. classification: 14M25, 14D23, 14H10.

(*) Supported by DFG-Schwerpunkt 1388 Darstellungstheorie.

of the variable y by an n -th root of unity. The torus $(\mathbb{G}_m)^{n-1}$, parametrising polynomials with non-zero coefficients a_1, \dots, a_{n-1} , is compactified by the moduli stack of chains of \mathbb{P}^1 with finite subschemes of degree n . On the boundary both the coefficients of the polynomials may become zero and the curve may become a reducible chain of \mathbb{P}^1 . The category of these pointed curves, which we call degree- n -pointed chains of \mathbb{P}^1 , forms an orbifold $\overline{\mathcal{L}}_n$.

The orbifold $\overline{\mathcal{L}}_n$ is related to the Losev-Manin moduli space \overline{L}_n by an S_n -equivariant morphism $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$, \overline{L}_n with the operation of the symmetric group S_n that permutes the n sections and $\overline{\mathcal{L}}_n$ with trivial operation, which is given by mapping an n -pointed chain of \mathbb{P}^1 to the corresponding degree- n -pointed chain by forgetting the labels of the sections. The moduli stack $\overline{\mathcal{L}}_n$ is defined such that the morphism $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$ is closely related to morphisms of the form $C_0^n \rightarrow C_0^n/S_n = C_0^{(n)} = \text{Div}_{C_0/Y}^n$ from the n -fold product over Y to the scheme of relative effective Cartier divisors of degree n for $C_0 \rightarrow Y$ a relative smooth curve over Y , here a chain of \mathbb{P}^1 over Y without the poles of the components of the fibres. Therefore the morphism $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$ inherits properties like being faithfully flat and finite of degree $n!$ and being ramified exactly in the points corresponding to curves with coinciding marked points, see proposition 1.5. The stack $\overline{\mathcal{L}}_n$ differs from the quotient stack $[\overline{L}_n/S_n]$, it has the same points but different automorphism groups. The coarse moduli space of $\overline{\mathcal{L}}_n$ coincides with the quotient \overline{L}_n/S_n .

A main result of this paper, theorem 3.1, is an explicit description of the structure of the stacks $\overline{\mathcal{L}}_n$: we show that $\overline{\mathcal{L}}_n$ is a toric orbifold and we determine the associated combinatorial data.

Toric Deligne-Mumford stacks over fields of characteristic 0 were introduced in [4] and constructed from combinatorial data called (simplicial) stacky fans, consisting of a simplicial fan and some extra data, as quotient stacks $[U/T]$ of an open subscheme U of some affine space by a diagonalisable group scheme G , generalising the quotient construction of a smooth toric variety described in [6]. Over more general base schemes in the same way these data give rise to toric stacks which are not necessarily Deligne-Mumford stacks but tame stacks in the sense of [1]. As our moduli problem results in stacks which are orbifolds, in this paper we are mainly concerned with toric orbifolds, i.e. toric tame stacks with trivial generic stabiliser. We will work with toric orbifolds over the integers, considering the fact that our moduli problem is naturally defined over the integers.

It turns out that the moduli stacks $\overline{\mathcal{L}}_n$ can be described in terms of the Cartan matrices of root systems of type A , more precisely, $\overline{\mathcal{L}}_n$ is isomorphic

to the toric orbifold $\mathcal{Y}(A_{n-1})$ which corresponds to the stacky fan $\Upsilon(A_{n-1})$ defined in section 2 using the Cartan matrix of the root system A_{n-1} . For the proof of the isomorphism $\overline{\mathcal{L}}_n \cong \mathcal{Y}(A_{n-1})$ we make use of a generalisation of the description of the functor of toric varieties [5] for toric stacks, which allows to characterise $\mathcal{Y}(A_{n-1})$ as a stack $\mathcal{C}_{\Upsilon(A_{n-1})}$ of $\Upsilon(A_{n-1})$ -collections, i.e. collections of pairs of a line bundle with a section and additional data.

We also characterise the morphism $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$, determined by forgetting the labels of the n sections, in terms of the combinatorial data by specifying the $\Upsilon(A_{n-1})$ -collection on $X(A_{n-1}) \cong \overline{L}_n$ corresponding to this morphism, see theorem 4.14. In doing this, in section 4 we compare the description of the functor of the toric varieties $X(A_{n-1})$ associated with root systems of type A after Cox [5] in terms of $\Sigma(A_{n-1})$ -collections to two other descriptions: the description of [2] in terms of A_{n-1} -data and a new description involving S_n -invariant line bundles on $X(A_{n-1})$. Both of these are related to Minkowski sum decompositions of the permutohedron: the first is a decomposition into line segments and the second corresponds to an embedding $X(A_{n-1}) \rightarrow \prod_{j=1}^{n-1} \mathbb{P}^{\binom{n}{j}-1}$ and expresses the permutohedron as sum of S_n -symmetric polytopes.

Generalisations of the Losev-Manin moduli spaces were investigated in [3]. We considered $(2n+1)$ -pointed and $2n$ -pointed chains of \mathbb{P}^1 with involution and showed that the moduli spaces $\overline{L}_n^{0,\pm}$ and \overline{L}_n^{\pm} of these objects coincide with the toric varieties $X(B_n)$ and $X(C_n)$ associated with the root systems B_n and C_n , see [3, Thm. 4.1 and 6.15].

In the present setting it makes sense to investigate similar generalisations of the moduli stacks $\overline{\mathcal{L}}_n$ and to relate these to the toric orbifolds $\mathcal{Y}(R)$ for root systems R belonging to other classical families as well as to the moduli spaces $\overline{L}_n^{0,\pm} \cong X(B_n)$ and $\overline{L}_n^{\pm} \cong X(C_n)$. In section 5 we consider moduli stacks of stable degree- $(2n+1)$ -pointed and degree- $2n$ -pointed chains of \mathbb{P}^1 with involution, $\overline{\mathcal{L}}_n^{0,\pm}$ and $\overline{\mathcal{L}}_n^{\pm}$. We show that $\overline{\mathcal{L}}_n^{\pm}$ has a main component $\overline{\mathcal{L}}_{n,+}^{\pm}$ isomorphic to $\mathcal{Y}(C_n)$ and that $\overline{\mathcal{L}}_n^{0,\pm}$ is isomorphic to $\mathcal{Y}(B_n)^{\text{can}}$, the canonical stack associated to $\mathcal{Y}(B_n)$ (see [7]). We have morphisms $\overline{L}_n^{0,\pm} \rightarrow \overline{\mathcal{L}}_n^{0,\pm}$ and $\overline{L}_n^{\pm} \rightarrow \overline{\mathcal{L}}_{n,+}^{\pm}$, defined by forgetting the labels of the sections, which are equivariant under the Weyl group.

Acknowledgements. Thanks to Victor Batyrev.

1. Moduli stacks of degree- n -pointed chains

We define moduli stacks of stable degree- n -pointed chains of \mathbb{P}^1 . Compared to the Losev-Manin moduli spaces considered in [16], [2], we replace the n marked points s_1, \dots, s_n of an n -pointed chain of \mathbb{P}^1 by a finite closed subscheme S of degree n .

DEFINITION 1.1. — *A stable degree- n -pointed chain of \mathbb{P}^1 over an algebraically closed field K is a tuple (C, s_-, s_+, S) , where C is a chain of \mathbb{P}^1 over K with two distinct closed points s_-, s_+ on the outer components such that on each component the number of intersection points together with s_-, s_+ adds up to 2 (cf. [2, Def. 3.1]), and $S \subset C$ a finite closed subscheme of degree n that does neither meet the intersection points of components nor s_-, s_+ , but that does meet every component of C . We define the category $\overline{\mathcal{L}}_n$ of stable degree- n -pointed chains of \mathbb{P}^1 over the category of schemes. The objects over a scheme Y are stable degree- n -pointed chains of \mathbb{P}^1 over Y , i.e. tuples $\mathcal{C} = (C \rightarrow Y, s_-, s_+, S)$, where $C \rightarrow Y$ is a locally finitely presented, flat, proper morphism of schemes, $s_-, s_+ : Y \rightarrow C$ are sections and $S \subset C$ is a subscheme finite flat over Y , such that the geometric fibres are stable degree- n -pointed chains of \mathbb{P}^1 . We have the natural notion of isomorphism of degree- n -pointed chains of \mathbb{P}^1 over the same scheme Y and of pullback of an object over a scheme Y with respect to a morphism $f : Y' \rightarrow Y$. A morphism in $\overline{\mathcal{L}}_n$ over a morphism $f : Y' \rightarrow Y$ is a cartesian diagram of stable degree- n -pointed chains of \mathbb{P}^1 over f .*

Remark 1.2. — (1) For a chain of \mathbb{P}^1 (C, s_-, s_+) over a field K any component is isomorphic to \mathbb{P}_K^1 since it contains a point with residue field K .

(2) As the morphisms $C \rightarrow Y$ are locally finitely presented, by [9, IV, (8.9.1)] we can use some results which originally require some noetherian hypothesis.

Remark 1.3. — The automorphism group of a chain of \mathbb{P}^1 (C, s_-, s_+) of length l over a field K is a torus $(\mathbb{G}_m)_K^l$. A stable degree- n -pointed chain of \mathbb{P}^1 (C, s_-, s_+, S) of length l over K has a finite automorphism group scheme which is a subgroup scheme of $(\mathbb{G}_m)_K^l$. There are objects (C, s_-, s_+, S) having nontrivial automorphisms: consider for example \mathbb{P}_K^1 with homogeneous coordinates z_0, z_1 , two poles $s_- = (1 : 0)$, $s_+ = (0 : 1)$ and a subscheme S of degree k given by the equation $z_0^k - z_1^k = 0$; in this example we have an automorphism group scheme isomorphic to μ_k .

PROPOSITION 1.4. — *The category $\overline{\mathcal{L}}_n$ is a category fibred in groupoids over the category of schemes. It forms a stack over the fpqc site of schemes with representable, finite diagonal. Over fields of characteristic 0 the diagonal is unramified.*

Proof. — The category $\overline{\mathcal{L}}_n$ together with the natural functor to the category of schemes is a fibred category, the cartesian arrows being cartesian diagrams of degree- n -pointed chains, and moreover the fibres $\overline{\mathcal{L}}_n(Y)$ over schemes Y form a groupoid.

The fibred category $\overline{\mathcal{L}}_n$ is a prestack in the fpqc topology, i.e. descent data for morphisms are effective, see for example [18, Prop. 4.31]. To show that $\overline{\mathcal{L}}_n$ is a stack, it remains to show that descent data for objects are effective. Let $(\pi: C \rightarrow Y, s_-, s_+, S)$ be a stable degree- n -pointed chain of \mathbb{P}^1 over a scheme Y . The subscheme $S \subset C$ is an effective Cartier divisor in C because this is true on the fibres, see [13, Lemma 9.3.4], and so its ideal sheaf $\mathcal{I} \subset \mathcal{O}_C$ is a line bundle. The line bundle $\mathcal{O}_C(S) = \mathcal{I}^{-1}$ is relatively ample with respect to π since it is ample on the fibres, see [9, III, (4.7.1)], [9, IV, (9.6.5)]. In fact, $\mathcal{O}_C(S)$ defines a closed embedding in the projective bundle $\mathbb{P}_Y(\pi_*\mathcal{O}_C(S))$, see proposition 3.4. Given a morphism $F: (C' \rightarrow Y', s'_-, s'_+, S') \rightarrow (C \rightarrow Y, s_-, s_+, S)$ of two degree- n -pointed chains over a morphism $f: Y' \rightarrow Y$ forming a cartesian diagram, we have a natural isomorphism $F^*\mathcal{O}_C(S) \cong \mathcal{O}_{C'}(S')$, and further, given morphisms F and G over $f: Y' \rightarrow Y$ and $g: Y'' \rightarrow Y'$, after identifying $(FG)^*\mathcal{O}_C(S)$ with $G^*F^*\mathcal{O}_C(S)$ the isomorphisms $(FG)^*\mathcal{O}_C(S) \rightarrow \mathcal{O}_{C''}(S'')$ and $G^*F^*\mathcal{O}_C(S) \rightarrow G^*\mathcal{O}_{C'}(S') \rightarrow \mathcal{O}_{C''}(S'')$ coincide. Then, by descent theory of flat proper morphisms of schemes with a relatively ample invertible sheaf, see [18, Thm. 4.38], descent data for objects of $\overline{\mathcal{L}}_n$ are effective.

We show that the diagonal $\overline{\mathcal{L}}_n \rightarrow \overline{\mathcal{L}}_n \times \overline{\mathcal{L}}_n$ is representable and finite. For a scheme Y and a morphism $Y \rightarrow \overline{\mathcal{L}}_n \times \overline{\mathcal{L}}_n$ given by two objects $\mathcal{C}, \mathcal{C}' \in \overline{\mathcal{L}}_n(Y)$, the category $Y \times_{\overline{\mathcal{L}}_n \times \overline{\mathcal{L}}_n} \overline{\mathcal{L}}_n$ fibred over the category of Y -schemes is isomorphic to the functor on Y -schemes $\text{Isom}(\mathcal{C}, \mathcal{C}')(f: Z \rightarrow Y) = \text{Mor}_{\overline{\mathcal{L}}_n(Z)}(f^*\mathcal{C}, f^*\mathcal{C}')$. Using the embedding via $\mathcal{O}_C(S)$ into $\mathbb{P}_Y(\pi_*\mathcal{O}_C(S))$ described in proposition 3.4 we see that $\text{Isom}(\mathcal{C}, \mathcal{C}')$ is a finite closed subgroup scheme of the open dense torus of $\mathbb{P}_Y(\pi_*\mathcal{O}_C(S))$. In characteristic 0 it is unramified over Y , because then the fibres are reduced. \square

The stack $\overline{\mathcal{L}}_n$ is related to the Losev-Manin moduli space \overline{L}_n by a morphism $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$ that arises by considering the n sections of an n -pointed chain over Y as a relative effective Cartier divisor of degree n over Y .

PROPOSITION 1.5. — *The morphism $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$ is faithfully flat and finite of degree $n!$. It is ramified exactly in the points of \overline{L}_n corresponding to n -pointed chains with some coinciding marked points.*

Proof. — Note that the morphism is representable, since \overline{L}_n has representable diagonal. We show that for any morphism $Y \rightarrow \overline{\mathcal{L}}_n$, Y a scheme, the morphism of schemes $Y \times_{\overline{\mathcal{L}}_n} \overline{L}_n \rightarrow Y$ has the properties in question. The morphism $Y \rightarrow \overline{\mathcal{L}}_n$ corresponds to an object $\mathcal{C} = (C \rightarrow Y, s_-, s_+, S)$ over Y and the functor $Y \times_{\overline{\mathcal{L}}_n} \overline{L}_n$ maps a scheme T to the set $\{(f: T \rightarrow Y, (C' \rightarrow T, s_-, s_+, s_1, \dots, s_n), \alpha) \mid \alpha: f^* \mathcal{C} \rightarrow (C' \rightarrow T, s_-, s_+, s_1 + \dots + s_n)\}$, where $s_1 + \dots + s_n$ denotes the divisor of degree n associated to the n sections and α is a morphism in $\overline{L}_n(T)$. We denote by C_0 the open subscheme of C obtained by excluding the poles and intersection points of components on the fibres. Then C_0 is a quasi-projective curve over Y , which is smooth over Y since it is flat with smooth fibres (see [9, IV, (17.5.1)]). We may, for any T , identify the chains C' over T occurring in the above sets with $C \times_Y T$ via the specified isomorphisms. The additional data given by the subscheme $S \subset C_0$ are equivalent to a section $s: Y \rightarrow \text{Div}_{C_0/Y}^n = C_0^{(n)}$ of the scheme of relative effective divisors of degree n , which coincides with the n -fold symmetric product of C_0 over Y , see [10, Exposé XVII, 6.3.9, p. 186]. Likewise, the data given by the sections s_1, \dots, s_n are equivalent to a section $s': T \rightarrow (C_0^n)_T$ of the n -fold product such that its composition with $(C_0^n)_T \rightarrow (C_0^{(n)})_T$ is the base extension s_T of s , or equivalently, to a morphism $s': T \rightarrow C_0^n$ whose composition with $C_0^n \rightarrow C_0^{(n)}$ coincides with $s \circ f$. Thus the functor $Y \times_{\overline{\mathcal{L}}_n} \overline{L}_n$ is isomorphic to the functor of the scheme $Y \times_{C_0^{(n)}} C_0^n$, and this concludes the proof because the morphism $C_0^n \rightarrow C_0^{(n)}$ has the required properties. □

Remark 1.6. — With proposition 1.5 and some general theory we can derive some properties of the stack $\overline{\mathcal{L}}_n$: by [15, Thm. 10.1], making use of the proposition, $\overline{\mathcal{L}}_n$ is an algebraic stack (Artin stack); in characteristic 0, by [15, Thm. 8.1] and the fact that it has unramified diagonal, it is a Deligne-Mumford stack. However, the result will follow independently later in section 3 together with a more detailed description of the structure of $\overline{\mathcal{L}}_n$.

On the Losev-Manin moduli space \overline{L}_n we have an operation of the symmetric group S_n permuting the n sections. Any S_n -equivariant morphism $\overline{L}_n \rightarrow Z$, Z a scheme with trivial S_n -action, factors through $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$. This implies that the quotient morphism $\overline{L}_n \rightarrow \overline{L}_n/S_n$ factors as

$$\overline{L}_n \rightarrow \overline{\mathcal{L}}_n \rightarrow \overline{L}_n/S_n,$$

and moreover, as $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$ is an epimorphism, $\overline{\mathcal{L}}_n \rightarrow \overline{L}_n/S_n$ forms the coarse moduli space of $\overline{\mathcal{L}}_n$.

Remark 1.7. — There is the quotient stack $[\overline{L}_n/S_n]$, which has the same geometric points as $\overline{\mathcal{L}}_n$. However, the automorphism groups of objects of $[\overline{L}_n/S_n]$ differ from those of $\overline{\mathcal{L}}_n$ which are always abelian.

In the case of the Losev-Manin moduli spaces, the boundary divisors arise as images of closed embeddings $\overline{L}_m \times \overline{L}_n \rightarrow \overline{L}_{m+n}$. For the stacks $\overline{\mathcal{L}}_n$ we also have embeddings $\overline{\mathcal{L}}_m \times \overline{\mathcal{L}}_n \rightarrow \overline{\mathcal{L}}_{m+n}$, defined as in the Losev-Manin case by concatenation of chains, and the diagrams

$$\begin{array}{ccc} \overline{L}_m \times \overline{L}_n & \longrightarrow & \overline{L}_{m+n} \\ \downarrow & & \downarrow \\ \overline{\mathcal{L}}_m \times \overline{\mathcal{L}}_n & \longrightarrow & \overline{\mathcal{L}}_{m+n} \end{array} \tag{1.1}$$

are commutative.

The morphism $\overline{L}_n \rightarrow \overline{\mathcal{L}}_n$ maps the open dense torus of $\overline{L}_n = X(A_{n-1})$, the moduli space of irreducible n -pointed chains, onto the moduli stack of irreducible degree- n -pointed chains. This open substack of $\overline{\mathcal{L}}_n$ parametrises subschemes S of degree n in $\mathbb{P}^1 \setminus \{0, \infty\}$ modulo automorphisms of \mathbb{P}^1 fixing 0 and ∞ . An object over an algebraically closed field K can be described by a monic polynomial $\prod_{i=1}^n (y - s_i)$ of degree n with $s_1, \dots, s_n \in K^*$ determined up to scaling by a common factor $\lambda \in K^*$ and permutations. We can write this polynomial as

$$y^n - (s_1 + \dots + s_n)y^{n-1} + \dots + (-)^n s_1 \cdots s_n$$

where the coefficients are the symmetric polynomials in s_1, \dots, s_n . Assuming $s_1 \cdots s_n = (-1)^n$, we have a polynomial of the form

$$y^n + a_1 y^{n-1} + a_2 y^{n-2} + \dots + a_{n-1} y + 1$$

with coefficients $a_1, \dots, a_{n-1} \in K$. The isomorphism class of the object determines these coefficients up to the equivalence $(a_1, \dots, a_{n-1}) \sim (\xi^{n-1} a_1, \dots, \xi a_{n-1})$, ξ an n -th root of unity. The moduli stack of such objects is the quotient stack $[\mathbb{A}^{n-1}/\mu_n]$, where the group scheme μ_n of n -th roots of unity acts with weights $(n-1, \dots, 1)$.

It contains an $(n-1)$ -dimensional algebraic torus T parametrising classes of polynomials with non-zero coefficients. A K -valued point of T is given by an $(n-1)$ -tuple

$$b_1 = \frac{a_2}{a_1^2}, b_2 = \frac{a_1 a_3}{a_2^2}, \dots, b_k = \frac{a_{k-1} a_{k+1}}{a_k^2}, \dots, b_{n-2} = \frac{a_{n-3} a_{n-1}}{a_{n-2}^2}, b_{n-1} = \frac{a_{n-2}}{a_{n-1}^2}$$

of elements $b_i \in K^*$. These expressions in the a_i form a set of generators of μ_n -invariants in the coordinate ring of the torus $(\mathbb{G}_m)^{n-1} \subset$

\mathbb{A}^{n-1} . Equivalently, we can express a K -valued point of T as a collection $(a_1, \dots, a_{n-1}, b_1, \dots, b_{n-1}) \in (K^*)^{2n-2}$ up to the equivalence


$$(a_1, \dots, a_{n-1}, b_1, \dots, b_{n-1}) \sim (\kappa_1 a_1, \dots, \kappa_{n-1} a_{n-1}, \lambda_1 b_1, \dots, \lambda_{n-1} b_{n-1})$$

for $\kappa_i \in K^*$ and $\lambda_i = \kappa_i^2 / (\kappa_{i-1} \kappa_{i+1})$, putting $\kappa_0 = \kappa_n = 1$. Then $(1, \dots, 1, b_1, \dots, b_{n-1}) \sim (a_1, \dots, a_{n-1}, 1, \dots, 1)$ if a_i, b_i satisfy the above equations.


Allowing certain subsets of the coordinates a_i, b_i to become zero, we obtain a toric tame stack which compactifies the moduli stack $[\mathbb{A}^{n-1}/\mu_n]$ of irreducible chains. Its definition and properties are contained in section 2 and we will show in section 3 that it coincides with the moduli stack $\overline{\mathcal{L}}_n$. In particular, degenerating some of the b_i to zero can be interpreted as degenerating \mathbb{P}^1 to a reducible chain of \mathbb{P}^1 . These additional divisors arise as in diagram (1.1).

Example 1.8. — We illustrate some results of this paper in the case $n = 2$ (see also examples 2.2, 3.8, 4.16). There is a natural embedding of degree-2-pointed chains (C, s_-, s_+, S) into \mathbb{P}^2 determined by the line bundle $\mathcal{O}_C(S)$ (for arbitrary n see section 3): in $\mathbb{P}^2 = \mathbb{P}(H^0(C, \mathcal{O}_C(S)))$ we can choose homogeneous coordinates y_0, y_1, y_2 such that C is given by an equation $y_0 y_2 = b_1 y_1^2$, the subscheme $S \subset C$ by an additional equation $y_0 + a_1 y_1 + y_2 = 0$, and the two sections s_-, s_+ are $(1 : 0 : 0), (0 : 0 : 1)$. Over an algebraically closed field K , data $(a_1, b_1) \in K^2 \setminus \{(0, 0)\}$ up to the equivalence $(a_1, b_1) \sim (\kappa_1 a_1, \lambda_1 b_1)$ for $\kappa_1^2 = \lambda_1 \in K^*$ correspond to isomorphism classes of degree-2-pointed chains over K .

The moduli stack $\overline{\mathcal{L}}_2$ is isomorphic to the quotient stack $[(\mathbb{A}^2 \setminus \{(0, 0)\})/\mathbb{G}_m]$ for the operation with weights $(1, 2)$, i.e. the weighted projective line $\mathbb{P}(1, 2)$ (which coincides with the toric orbifold $\mathcal{Y}(A_{n-1})$ for $n = 2$ defined in section 2). The open substack parametrising irreducible curves, the locus where $b_1 \neq 0$, is the quotient stack $[\mathbb{A}^1/\mu_2]$ with coordinate a_1 on \mathbb{A}^1 . The open substack parametrising objects without isomorphisms, the locus where $a_1 \neq 0$, is isomorphic to \mathbb{A}^1 with coordinate b_1 .



The morphism $\bar{\mathcal{L}}_2 \rightarrow \bar{\mathcal{L}}_2$ is faithfully flat and finite of degree 2. We introduce homogeneous coordinates z_-, z_+ of $\bar{\mathcal{L}}_2 = \mathbb{P}^1$ that measure the position of one of the marked points of a 2-pointed chain with respect to the other marked point at $(1:1)$ of its component isomorphic to \mathbb{P}^1 , such that the two points $(0:1), (1:0)$ correspond to reducible chains (cf. [2]). Then the point $(1:-1)$ corresponds to a 2-pointed curve \mathbb{P}^1 with marked points $(1:1), (1:-1)$ giving rise to a degree-2-pointed curve with automorphism group μ_2 . The point $(1:1)$ corresponds to the point of $\bar{\mathcal{L}}_2$ with nonreduced S , the morphism is ramified here and étale elsewhere.



We will see in section 4 that the morphism $\bar{\mathcal{L}}_2 \rightarrow \bar{\mathcal{L}}_2$ is given as

$$(z_- + z_+ : z_- z_+) : \mathbb{P}^1 \rightarrow \mathbb{P}(1, 2).$$

2. The toric orbifolds $\mathcal{Y}(A_n)$

In this section we will consider a family of toric orbifolds associated to the Cartan matrices of root systems of type A , but also comment on some generalities on toric stacks.

We use the definitions and notations of [4]. A stacky fan $\Sigma = (N, \Sigma, \beta)$ defining a toric orbifold has the property that the abelian group N is free; it consists of the data of a simplicial fan Σ in the lattice N and elements $n_\varrho \in \varrho \cap N$ for the one-dimensional cones $\varrho \in \Sigma(1)$. Here we assume them to span the ambient space $N_{\mathbb{Q}}$. The homomorphism $\beta: \mathbb{Z}^{\Sigma(1)} \rightarrow N$ maps the elements of the standard basis to the elements n_ϱ . Dually we have the exact sequence

$$0 \longrightarrow M = \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z}) \xrightarrow{\beta^*} \mathbb{Z}^{\Sigma(1)} \xrightarrow{\beta^\vee} \text{DG}(\beta) \longrightarrow 0$$

giving rise, as sequence of character groups, to the exact sequence of diagonalisable group schemes

$$1 \longrightarrow G \longrightarrow T_{\Sigma(1)} \longrightarrow T_M \longrightarrow 1.$$

The toric orbifold \mathcal{X}_Σ is defined as the quotient stack $[U/G]$ with $U \subseteq \mathbb{A}^{\Sigma(1)}$ the open subset defined by the information which of the one-dimensional cones form higher dimensional cones of Σ . The constructions make sense over the integers, however, working with G -torsors, in general one may have to choose an appropriate Grothendieck topology on the base category possibly finer than the étale topology (see also remark 2.8). The resulting algebraic stacks \mathcal{X}_Σ are tame stacks in the sense of [1].

DEFINITION 2.1. — We define the toric orbifold $\mathcal{Y}(A_n)$ associated to the Cartan matrix of the root system A_n in terms of the stacky fan $\Upsilon(A_n) = (N, \Upsilon(A_n), \beta)$: let $N = \mathbb{Z}^n$ and let the linear map $\beta: \mathbb{Z}^{2n} \rightarrow N$ be given by the $n \times 2n$ matrix

$$\begin{pmatrix} -2 & 1 & 0 & \cdots & 1 & 0 & 0 & \cdots \\ 1 & -2 & 1 & \cdots & 0 & 1 & 0 & \cdots \\ 0 & 1 & -2 & \cdots & 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

i.e. the matrix consisting of two blocks $(-C(A_n) \ I_n)$, where $C(A_n)$ is the Cartan matrix of the root system A_n and I_n the $n \times n$ identity matrix. The fan $\Upsilon(A_n)$ has the $2n$ one-dimensional cones $\varrho_1, \dots, \varrho_n, \tau_1, \dots, \tau_n$ generated by the columns of the above matrix. A subset of one-dimensional cones generates a higher dimensional cone of $\Upsilon(A_n)$ if it does not contain one of the sets $\{\varrho_1, \tau_1\}, \dots, \{\varrho_n, \tau_n\}$. This defines a fan containing 2^n n -dimensional cones σ_I generated by sets $\{\varrho_i : i \notin I\} \cup \{\tau_i : i \in I\}$ for subsets $I \subseteq \{1, \dots, n\}$.

For the stacky fan $\Upsilon(A_n)$ the map $\beta: \mathbb{Z}^{2n} \rightarrow N$ gives rise to the exact sequence of lattices

$$0 \longrightarrow M \cong \mathbb{Z}^n \xrightarrow{\begin{pmatrix} -C \\ I_n \end{pmatrix}} \mathbb{Z}^{2n} \xrightarrow{\begin{pmatrix} I_n & C \end{pmatrix}} \text{DG}(\beta) \cong \mathbb{Z}^n \longrightarrow 0$$


where $C = C(A_n)^\top = C(A_n)$ is (the transpose of) the Cartan matrix, and the exact sequence of tori

$$1 \longrightarrow G \cong (\mathbb{G}_m)^n \longrightarrow (\mathbb{G}_m)^{2n} \longrightarrow T_N \cong (\mathbb{G}_m)^n \longrightarrow 1$$


where $G \cong (\mathbb{G}_m)^n \longrightarrow (\mathbb{G}_m)^{2n}, (\kappa_1, \dots, \kappa_n) \mapsto (\kappa_1, \dots, \kappa_n, \lambda_1, \dots, \lambda_n)$ with $\lambda_i = \kappa_i^2 / (\kappa_{i-1} \kappa_{i+1})$ setting $\kappa_0 = \kappa_{n+1} = 1$ (cf. last section). Note that the toric orbifold $\mathcal{Y}(A_n)$ arises as quotient $[U/G]$ by a torus G .

Example 2.2. — The toric orbifold $\mathcal{Y}(A_1)$ is isomorphic to the weighted projective line $\mathbb{P}(1, 2)$: we have the matrix $\begin{pmatrix} -2 & 1 \end{pmatrix}$ and the stacky fan looks

as follows:



Example 2.3. — The toric orbifold $\mathcal{Y}(A_2)$ arises from the matrix $\begin{pmatrix} -2 & 1 & 1 & 0 \\ 1 & -2 & 0 & 1 \end{pmatrix}$. We have the stacky fan



The description of the functor of a smooth toric variety given by Cox [5] in terms of collections of line bundles with sections determined by the combinatorial data has been extended to toric Deligne-Mumford stacks by Iwanari [11] and Perroni [17]. For the stacky fan $\Upsilon(A_n)$ we have:

DEFINITION 2.4. — A $\Upsilon(A_n)$ -collection on a scheme Y is a collection

$$\mathcal{L} = ((\mathcal{L}_{\rho_i}, a_i)_{i=1, \dots, n}, (\mathcal{L}_{\tau_i}, b_i)_{i=1, \dots, n}, (c_i)_{i=1, \dots, n})$$

where $(\mathcal{L}_{\rho_i}, a_i)$ and $(\mathcal{L}_{\tau_i}, b_i)$ are line bundles with a section and

$$c_1: \mathcal{L}_{\tau_1} \otimes \mathcal{L}_{\rho_1}^{\otimes -2} \otimes \mathcal{L}_{\rho_2} \rightarrow \mathcal{O}_Y, c_2: \mathcal{L}_{\tau_2} \otimes \mathcal{L}_{\rho_1} \otimes \mathcal{L}_{\rho_2}^{\otimes -2} \otimes \mathcal{L}_{\rho_3} \rightarrow \mathcal{O}_Y, \dots, c_{n-1}: \mathcal{L}_{\tau_{n-1}} \otimes \mathcal{L}_{\rho_{n-2}} \otimes \mathcal{L}_{\rho_{n-1}}^{\otimes -2} \otimes \mathcal{L}_{\rho_n} \rightarrow \mathcal{O}_Y, c_n: \mathcal{L}_{\tau_n} \otimes \mathcal{L}_{\rho_{n-1}} \otimes \mathcal{L}_{\rho_n}^{\otimes -2} \rightarrow \mathcal{O}_Y$$

are isomorphisms. These data are subject to the nondegeneracy condition that for every point $y \in Y$ and $i = 1, \dots, n$ not both $a_i(y) = 0$ and $b_i(y) = 0$.

A morphism $\mathcal{L}' \rightarrow \mathcal{L}$ between two $\Upsilon(A_n)$ -collections $\mathcal{L} = ((\mathcal{L}_{\rho_i}, a_i)_i, (\mathcal{L}_{\tau_i}, b_i)_i, (c_i)_i)$ on Y and $\mathcal{L}' = ((\mathcal{L}'_{\rho_i}, a'_i)_i, (\mathcal{L}'_{\tau_i}, b'_i)_i, (c'_i)_i)$ on Y' over a morphism of schemes $f: Y' \rightarrow Y$ is a collection $((r_i)_{i=1, \dots, n}, (t_i)_{i=1, \dots, n})$ consisting of isomorphisms of line bundles $r_i: f^* \mathcal{L}_{\rho_i} \rightarrow \mathcal{L}'_{\rho_i}, t_i: f^* \mathcal{L}_{\tau_i} \rightarrow$

\mathcal{L}'_{τ_i} such that $r_i(f^*a_i) = a'_i$, $t_i(f^*b_i) = b'_i$ and the diagrams

$$\begin{array}{ccc}
 f^*\mathcal{L}'_{\tau_i} \otimes f^*\mathcal{L}'_{\varrho_{i-1}} \otimes f^*\mathcal{L}'_{\varrho_i} \otimes f^*\mathcal{L}'_{\varrho_{i+1}} & \xrightarrow{f^*c_i} & f^*\mathcal{O}_{Y'} \\
 \downarrow & & \downarrow \\
 \mathcal{L}'_{\tau_i} \otimes \mathcal{L}'_{\varrho_{i-1}} \otimes \mathcal{L}'_{\varrho_i} \otimes \mathcal{L}'_{\varrho_{i+1}} & \xrightarrow{c_i} & \mathcal{O}_{Y'}
 \end{array} \tag{2.1}$$

($i = 1, \dots, n$; for $i = 1, n$ omit the factors indexed by ϱ_0, ϱ_{n+1}) commute.

We denote the fibred category of $\Upsilon(A_n)$ -collections over the category of schemes by $\mathcal{C}_{\Upsilon(A_n)}$. It comes with the cleavage given by pull-back of line bundles: for $f: Y' \rightarrow Y$ we have an arrow $f^*\mathcal{L} \rightarrow \mathcal{L}$ in $\mathcal{C}_{\Upsilon(A_n)}$. The definition describes a morphism $\mathcal{L}' \rightarrow \mathcal{L}$ in $\mathcal{C}_{\Upsilon(A_n)}$ as composition of a morphism $\mathcal{L}' \rightarrow f^*\mathcal{L}$ over $id_{Y'}$ with $f^*\mathcal{L} \rightarrow \mathcal{L}$ over $f: Y' \rightarrow Y$.

Remark 2.5. — A morphism of $\Upsilon(A_n)$ -collections $\mathcal{L}' \rightarrow \mathcal{L}$ over id_Y Zariski-locally for some open $Y' \subseteq Y$, after fixing isomorphisms of the line bundles with the structure sheaf such that the isomorphisms c_i become $id_{\mathcal{O}_{Y'}}$, corresponds to a collection $\kappa_1, \dots, \kappa_n, \lambda_1, \dots, \lambda_n \in \mathcal{O}_{Y'}^*(Y')$ such that the isomorphisms $\mathcal{O}_{Y'} \cong \mathcal{L}'_{\varrho_i}|_{Y'} \rightarrow \mathcal{L}'_{\varrho_i}|_{Y'} \cong \mathcal{O}_{Y'}$ and $\mathcal{O}_{Y'} \cong \mathcal{L}'_{\tau_i}|_{Y'} \rightarrow \mathcal{L}'_{\tau_i}|_{Y'} \cong \mathcal{O}_{Y'}$ are given by multiplication by κ_i and λ_i . The condition expressed in diagram (2.1) translates into the equations $\lambda_i = \kappa_i^2 / (\kappa_{i-1}\kappa_{i+1})$, putting $\kappa_0 = \kappa_{n+1} = 1$.

The category of Σ -collections \mathcal{C}_{Σ} for a stacky fan Σ is a category fibred in groupoids (CFG) over the base category of schemes. By descent theory for quasi-coherent sheaves the CFG \mathcal{C}_{Σ} forms a stack in the fpqc topology, see [18, Thm. 4.23]. By Iwanari [11, Thm. 1.4] (for toric orbifolds) and Perroni [17, Thm. 2.6] (for toric Deligne-Mumford stacks) over fields of characteristic 0, working with the étale topology, there is an isomorphism of stacks $\mathcal{X}_{\Sigma} \cong \mathcal{C}_{\Sigma}$. Also over more general base schemes we have an isomorphism $\mathcal{X}_{\Sigma} \cong \mathcal{C}_{\Sigma}$; we make some comments on this issue.

CONSTRUCTION 2.6. — Explicitly, one can construct an isomorphism $\mathcal{X}_{\Sigma} \cong \mathcal{C}_{\Sigma}$ as follows, here for simplicity we stick to the orbifold case and assume that the one-dimensional cones generate the ambient space $N_{\mathbb{Q}}$.

Note that we have a natural G -equivariant Σ -collection $((\mathcal{O}_U \otimes V_{\varrho}, x_{\varrho})_{\varrho}, (id)_m)$ on U , where V_{ϱ} is the one-dimensional representation such that the coordinate x_{ϱ} of $U \subset \mathbb{A}^{\Sigma(1)}$ is an invariant section of $\mathcal{O}_U \otimes V_{\varrho}$ (in the case of smooth toric varieties as considered in [6] this G -equivariant collection descends to the universal collection on the toric variety).

Starting with an object of \mathcal{X}_{Σ} over Y , that is a G -torsor $p: E \rightarrow Y$ together with a G -equivariant morphism $t: E \rightarrow U$, the pull-back $t^*((\mathcal{O}_U \otimes V_{\varrho}, x_{\varrho})_{\varrho}, (id)_m)$ is a G -equivariant Σ -collection on E and gives rise to the

Σ -collection $p_*^G t^*((\mathcal{O}_U \otimes V_\varrho, x_\varrho)_\varrho, (id)_m)$ on Y (the functor p_*^G takes the G -invariant part of the push-forward).

On the other hand, for a given Σ -collection $((\mathcal{L}_\varrho, u_\varrho)_\varrho, (c_m)_m)$ on a scheme Y we construct a G -torsor with a G -equivariant morphism to U . Let \underline{E} be the contravariant functor on the category of Y -schemes

$$\underline{E}: (q: Y' \rightarrow Y) \mapsto \left\{ \begin{array}{l} \Sigma\text{-collections } ((\mathcal{O}_{Y'} \otimes V_\varrho, u'_\varrho), (id)_m) \text{ on } Y' \\ \text{with an isomorphism of } \Sigma\text{-collections} \\ q^*((\mathcal{L}_\varrho, u_\varrho)_\varrho, (c_m)_m) \cong ((\mathcal{O}_{Y'} \otimes V_\varrho, u'_\varrho)_\varrho, (id)_m) \end{array} \right\}$$

where V_ϱ , the one-dimensional representation as above, is used to define an operation of G on this functor. Then one can show that the functor \underline{E} with this G -action is represented by a G -torsor $p: E \rightarrow Y$ together with a universal isomorphism $p^*((\mathcal{L}_\varrho, u_\varrho)_\varrho, (c_m)_m) \cong ((\mathcal{O}_E \otimes V_\varrho, u_\varrho^E)_\varrho, (id)_m)$ of G -equivariant Σ -collections, provided that the original Σ -collection is locally trivial in the sense that there is a covering $f: Y' \rightarrow Y$ such that $f^*((\mathcal{L}_\varrho, u_\varrho)_\varrho, (c_m)_m)$ is isomorphic to a collection of the form $((\mathcal{O}_{Y'}, u'_\varrho)_\varrho, (id)_m)$; collections of this form correspond to trivial G -torsors. We will assume that the topology on the base category is such that any Σ -collection has this property, see also the following remarks. The sections $(u_\varrho^E)_\varrho$ of the universal Σ -collection on E then define a G -equivariant morphism $E \rightarrow U \subset \mathbb{A}^{\Sigma(1)}$.

Making use of the fact that for a G -torsor $p: E \rightarrow Y$ we have the equivalence $\text{QCoh}(Y) \leftrightarrow \text{QCoh}^G(E)$ given by the functors p^* and p_*^G , one can show that these constructions define functors $\mathcal{X}_\Sigma \leftrightarrow \mathcal{C}_\Sigma$ whose compositions are isomorphic to the identity functors.

Remark 2.7. — Zariski-locally we can interpret the construction of the G -torsor as the coboundary homomorphism d in the exact sequence (see [8, Ch. III, §3])

$$0 \rightarrow H^0(Y, G) \rightarrow H^0(Y, T_{\Sigma(1)}) \rightarrow H^0(Y, T_M) \xrightarrow{d} H^1(Y, G)$$

where elements of $H^1(Y, G)$ are isomorphism classes of G -torsors over Y : given a Σ -collection $((\mathcal{O}_Y, u_\varrho)_\varrho, (c_m)_m)$ on Y , the automorphisms $(c_m)_m$ of the structure sheaf can be interpreted as a morphism $Y \rightarrow T_M$ or section of $T_M \times Y \rightarrow Y$, and fitting in the cartesian diagram

$$\begin{array}{ccc} E & \longrightarrow & T_{\Sigma(1)} \\ \downarrow & & \downarrow \\ Y & \longrightarrow & T_M \end{array}$$

we obtain a G -torsor $E \rightarrow Y$ which is trivial if and only if $(c_m)_m \in H^0(Y, T_M)$ comes from an element of $H^0(Y, T_{\Sigma(1)})$.

Remark 2.8. — Working with G -torsors, we usually assume that the Grothendieck topology on the base category is fine enough in the sense that we have the same G -torsors as we have with respect to the canonical topology. We have seen that this assumption was necessary to derive the isomorphism $\mathcal{X}_\Sigma \cong \mathcal{C}_\Sigma$: whereas the notion of G -torsor depends on the topology, this is not the case for the notion of Σ -collections. For Σ -collections we have the corresponding assumption that Σ -collections are locally trivial with respect to the topology (in the sense of construction 2.6). In characteristic 0 this is always true for the étale topology. In general we may have to take a finer topology, for example the fppf topology.

In the case of the stacky fan $\Upsilon(A_n)$ the lattice M is a direct summand of $\mathbb{Z}^{\Sigma(1)}$ and the group scheme G a torus, so the following result also holds in weaker topologies like étale or Zariski.

COROLLARY 2.9. — *There is an isomorphism of stacks $\mathcal{Y}(A_n) \cong \mathcal{C}_{\Upsilon(A_n)}$.*

In particular, a K -valued point of $\mathcal{Y}(A_n)$ corresponds to $(a_1, \dots, a_n, b_1, \dots, b_n) \in K^{2n}$ such that for any i not both $a_i = 0$ and $b_i = 0$, up to the equivalence relation given by multiplication by a collection $(\kappa_1, \dots, \kappa_n, \lambda_1, \dots, \lambda_n) \in (K^*)^{2n}$ as in remark 2.5.

3. $\mathcal{Y}(A_{n-1})$ as moduli stack of degree- n -pointed chains

In this section we will prove the following theorem.

THEOREM 3.1. — *There is an isomorphism of stacks $\overline{\mathcal{L}}_n \cong \mathcal{Y}(A_{n-1})$.*

We will relate families of pointed chains to $\Upsilon(A_{n-1})$ -collections and prove an equivalence of fibred categories $\overline{\mathcal{L}}_n \cong \mathcal{C}_{\Upsilon(A_{n-1})}$.

Let (C, s_-, s_+, S) be a degree- n -pointed chain of \mathbb{P}^1 over a field K . We look at the closed embedding $C \rightarrow \mathbb{P}_K(H^0(C, \mathcal{O}_C(S))) \cong \mathbb{P}_K^n$ determined by $\mathcal{O}_C(S)$.

First assume that C is irreducible, that is $C \cong \mathbb{P}_K^1$. The vector space $H^0(C, \mathcal{O}_C(S))$ is $(n + 1)$ -dimensional and we have a basis y_0, \dots, y_n such that the divisor of y_i satisfies $\text{div}(y_i) = is_- + (n - i)s_+$. The ideal sheaf $\mathcal{I} = \mathcal{O}_C(-S) \rightarrow \mathcal{O}_C$ defining S is a line bundle. Tensoring with $\mathcal{O}_C(S)$ we have an inclusion $\mathcal{O}_C \rightarrow \mathcal{O}_C(S)$ with cokernel \mathcal{O}_S , and the image of the 1-section of \mathcal{O}_C is a global section $\sum_{i=0}^n a_i y_i \in H^0(C, \mathcal{O}_C(S))$. The subscheme $S \subset C$ is given by the equation $\sum_{i=0}^n a_i y_i = 0$, where $a_0, a_n \neq 0$ as S does not meet s_-, s_+ . We can choose the basis y_0, \dots, y_n such that $a_0 = a_n = 1$.

The embedding defined by $\mathcal{O}_C(S)$, the n -fold Veronese embedding or n -uple embedding, gives an isomorphism of C onto the subscheme in \mathbb{P}_K^n determined by the equations

$$y_i y_{j+1} = b_{i+1} \cdots b_j y_{i+1} y_j$$

for $0 \leq i < j < n$ and certain numbers $b_1, \dots, b_{n-1} \in K^*$. These equations express the condition that the rank of the matrix

$$\begin{pmatrix} y_0 & b_1 y_1 & \cdots & b_1 \cdots b_{n-1} y_{n-1} \\ y_1 & y_2 & \cdots & y_n \end{pmatrix}$$

is less than 2. The subscheme S in the embedded curve is given by the additional linear equation

$$y_n + a_{n-1} y_{n-1} + \cdots + a_1 y_1 + y_0 = 0.$$

Similarly, we have a natural embedding of reducible degree- n -pointed chains of \mathbb{P}^1 into \mathbb{P}_K^n .

PROPOSITION 3.2. — *Let (C, s_-, s_+, S) be a degree- n -pointed chain of \mathbb{P}^1 over a field K . It decomposes into irreducible components $C_1, \dots, C_m \cong \mathbb{P}_K^1$ with poles $(p_1^-, p_1^+), \dots, (p_m^-, p_m^+)$ such that $s_- = p_1^-$, $s_+ = p_m^+$ and C_i intersects C_{i+1} in $p_i^+ = p_{i+1}^-$. Let n_1, \dots, n_m be the degrees of S on the components C_1, \dots, C_m and $N_k = \sum_{i=1}^k n_i$. Then there is a basis y_0, \dots, y_n of $H^0(C, \mathcal{O}_C(S))$ characterised up to nonzero scalars by the following conditions: y_i is nonzero only on the components C_k satisfying $N_{k-1} \leq i \leq N_k$ and in this case $\text{div}(y_i)|_{C_k} = (i - N_{k-1})p_k^- + (N_k - i)p_k^+$. We scale y_0, y_n such that the image of the 1-section under the inclusion $\mathcal{O}_C \rightarrow \mathcal{O}_C(S)$ is $\sum_{i=0}^n a_i y_i \in H^0(C, \mathcal{O}_C(S))$ with $a_0 = a_n = 1$, that is, S is given by an equation*

$$y_n + a_{n-1} y_{n-1} + \cdots + a_1 y_1 + y_0 = 0 \tag{3.1}$$

for some $a_1, \dots, a_{n-1} \in K$. These sections y_0, \dots, y_n satisfy the equations

$$y_i y_{j+1} = b_{i+1} \cdots b_j y_{i+1} y_j \tag{3.2}$$

for $0 \leq i < j < n$ and certain numbers $b_1, \dots, b_{n-1} \in K$ such that $b_j = 0$ exactly if $j \in \{N_1, \dots, N_{m-1}\}$.

The curve C embeds into $\mathbb{P}_K(H^0(C, \mathcal{O}_C(S)))$, the image being the subscheme defined by the equations (3.2). The subscheme S of the embedded curve is given by the additional equation (3.1). The sections s_-, s_+ are $(1:0:\dots:0), (0:\dots:0:1)$.

The numbers a_1, \dots, a_{n-1} and b_1, \dots, b_{n-1} in (3.1) and (3.2) have the property that not both $a_j = 0$ and $b_j = 0$.

We will not work out the proof in detail, but add some remarks.

Remark 3.3. — The component C_k is embedded into the projective subspace of $\mathbb{P}_K(H^0(C, \mathcal{O}_C(S)))$ spanned by the coordinates $y_{N_{k-1}}, \dots, y_{N_k}$ via a Veronese embedding, the image is given by the equations corresponding to the condition that the rank of the matrix

$$\begin{pmatrix} y_{N_{k-1}} & b_{N_{k-1}+1}y_{N_{k-1}+1} & \cdots & b_{N_{k-1}+1} \cdots b_{N_k-1}y_{N_{k-1}} \\ y_{N_{k-1}+1} & y_{N_{k-1}+2} & \cdots & y_{N_k} \end{pmatrix}$$

is less than 2. The equation (3.1) reduces on C_k to $a_{N_k}y_{N_k} + \dots + a_{N_{k-1}}y_{N_{k-1}} = 0$ which defines a finite subscheme S_k of degree n_k in $C_k \subseteq \mathbb{P}_K^{n_k}$. A subscheme S_k of this form does not meet the poles of C_k provided that $a_{N_k}, a_{N_{k-1}} \neq 0$.

We generalise this to degree- n -pointed chains over schemes.

PROPOSITION 3.4. — *Let $(\pi: C \rightarrow Y, s_-, s_+, S)$ be a degree- n -pointed chain of \mathbb{P}^1 over a scheme Y . For any $y \in Y$ there is an open affine neighbourhood $Y' \subseteq Y$ such that there is a decomposition $\pi_*\mathcal{O}_C(S)|_{Y'} \cong \bigoplus_{i=0}^n \mathcal{O}_{Y'}y_i$ characterised on the fibres by the properties of proposition 3.2. The generators $y_0, \dots, y_n \in H^0(Y', \pi_*\mathcal{O}_C(S))$ of the individual summands, after possibly rescaling by a global section of $\mathcal{O}_{Y'}^*$, satisfy:*

(i) *The image of the 1-section under the inclusion $\mathcal{O}_C \rightarrow \mathcal{O}_C(S)$ is of the form*

$$y_n + a_{n-1}y_{n-1} + \dots + a_1y_1 + y_0.$$

over Y' for some $a_1, \dots, a_{n-1} \in \mathcal{O}_{Y'}(Y')$.

(ii) *The kernel of the homomorphism of algebras $\text{Sym } \pi_*(\mathcal{O}_C(S)) \rightarrow \bigoplus_{k=0}^\infty \pi_*\mathcal{O}_C(kS)$ is over Y' generated by the equations*

$$y_iy_{j+1} = b_{i+1} \cdots b_jy_{i+1}y_j$$

for $0 \leq i < j < n$ and some $b_1, \dots, b_{n-1} \in \mathcal{O}_{Y'}(Y')$.

The line bundle $\mathcal{O}_C(S)$ determines a closed embedding $C \rightarrow C' \subset \mathbb{P}_Y(\pi_(\mathcal{O}_C(S)))$ over Y . Over Y' in the coordinates y_0, \dots, y_n the embedded curve $C'_{Y'} \subset \mathbb{P}_{Y'}(\pi_*(\mathcal{O}_C(S))|_{Y'}) \cong \mathbb{P}_{Y'}^n$, is defined by the equations in (ii), the image of $S_{Y'}$ in $C'_{Y'}$, by the additional equation in (i), and the sections s_-, s_+ are $(1:0:\dots:0), (0:\dots:0:1)$.*

Proof. — The decomposition of the fibre C_y over a point $y \in Y$ into irreducible components $C_y = C_1 \cup \dots \cup C_m$ determines over an open affine subscheme $Y' \subseteq Y$ containing y a decomposition of S into divisors S_1, \dots, S_m which are disjoint and such that S_k only meets one component on each fibre and the component C_k over y . Each S_k determines a morphism onto a \mathbb{P}^1 -bundle over Y' , which on the fibres is an isomorphism on the component containing S_k and contracts the other components (similar to the

contraction morphisms in [14], cf. also [2, 3.3]). After possibly shrinking Y' , we have global sections $y_0^{(k)}, \dots, y_n^{(k)}$ of $\mathcal{O}_{\mathbb{P}^1_{Y'}}(S_k)$ that satisfy $\text{div}(y_i^{(k)}) = (i - N_{k-1})s_- + (N_k - i)s_+$ for $N_{k-1} \leq i \leq N_k$, $\text{div}(y_i^{(k)}) = (N_k - N_{k-1})s_+$ for $i < N_{k-1}$ and $\text{div}(y_i^{(k)}) = (N_k - N_{k-1})s_-$ for $i > N_k$ (using the notation S_k, s_-, s_+ also for their images in $\mathbb{P}^1_{Y'}$). Using the pull-backs of these sections to $C_{Y'}$ denoted by the same symbols, let $y_i = y_i^{(1)} \cdots y_i^{(m)} \in H^0(C_{Y'}, \mathcal{O}_C(S)) = H^0(C_{Y'}, \mathcal{O}_{C_{Y'}}(S_1) \otimes \cdots \otimes \mathcal{O}_{C_{Y'}}(S_m))$. Over the open affine neighbourhood Y' of y the sections y_0, \dots, y_n define a decomposition of $\pi_*\mathcal{O}_C(S)$ with the required properties.

The image of the 1-section under the inclusion $\mathcal{O}_C \rightarrow \mathcal{O}_C(S)$ gives a global section of $\mathcal{O}_C(S)$ which over Y' is of the form $\sum_{i=0}^n a_i y_i$ with $a_i \in \mathcal{O}_{Y'}(Y')$. Since $a_0, a_n \in \mathcal{O}_{Y'}^*(Y')$ we can assume that $a_0, a_n = 1$.

Using what is known about the fibres and results from [9, III] (cf. also [14]), we derive that $\pi_*(\mathcal{O}_C(kS))$ for $k > 0$ is locally free of rank $kn + 1$, further that the homomorphism $\pi^*\pi_*\mathcal{O}_C(S) \rightarrow \mathcal{O}_C(S)$ is surjective and defines a closed embedding $C \rightarrow \mathbb{P}_Y(\pi_*\mathcal{O}_C(S))$.

The embedding $C \rightarrow \mathbb{P}_Y^n$ corresponds to the surjection of graded algebras $\text{Sym } \pi_*(\mathcal{O}_C(S)) \rightarrow \bigoplus_{k=0}^\infty \pi_*\mathcal{O}_C(kS)$. Its kernel \mathcal{I} is the graded ideal that defines the embedded curve $C' \subset \mathbb{P}_Y^n$. Each part \mathcal{I}_k of \mathcal{I} is locally free, being the kernel of a surjective homomorphism of locally free sheaves. The graded ideal \mathcal{I} is generated in degree 2 since this is the case on the fibres $\mathcal{I} \otimes \kappa(y)$ for each point $y \in Y$. The part \mathcal{I}_2 of degree 2 is a vector bundle of rank $\frac{1}{2}n(n - 1)$.

Working over Y' , for $i < j$ the subsheaves $\langle y_{i+1}y_j, y_i y_{j+1} \rangle$ and $\langle y_{i+1}y_j \rangle$ of $\pi_*\mathcal{O}_C(S)|_{Y'}$, i.e. the subsheaves generated by the respective sections, coincide as this is true on the fibres. Considering the case $j = i + 1$, the kernel of the surjective homomorphism $\text{Sym}^2 \pi_*(\mathcal{O}_C(S))|_{Y'} \supset \mathcal{O}_{Y'} y_{i+1}^2 \oplus \mathcal{O}_{Y'} y_i y_{i+2} \rightarrow \langle y_{i+1}^2, y_i y_{i+2} \rangle \subset \pi_*\mathcal{O}_C(2S)|_{Y'}$ is generated by an element $y_i y_{i+2} - b_{i+1} y_{i+1}^2$ for some $b_{i+1} \in \mathcal{O}_{Y'}(Y')$. For general $i < j$ we have as kernel $y_i y_{j+1} - b_{i+1,j} y_{i+1} y_j$ for some $b_{i+1,j} \in \mathcal{O}_{Y'}(Y')$ and from the equation $(y_{i+2} \cdots y_j) b_{i+1,j} y_{i+1} y_j = (y_{i+2} \cdots y_j) y_i y_{j+1} = (b_{i+1} \cdots b_j) (y_{i+2} \cdots y_j) y_{i+1} y_j$ in $\pi_*\mathcal{O}_C(S)|_{Y'}$ we conclude that $b_{i+1,j} = b_{i+1} \cdots b_j$. \square

We define morphisms of fibred categories $\Phi: \overline{\mathcal{L}}_n \rightarrow \mathcal{C}_{\mathbf{Y}(A_{n-1})}$ and $\Psi: \mathcal{C}_{\mathbf{Y}(A_{n-1})} \rightarrow \overline{\mathcal{L}}_n$.

CONSTRUCTION 3.5. — Let $\mathcal{C} = (C \rightarrow Y, s_-, s_+, S)$ be a degree- n -pointed chain of \mathbb{P}^1 over a scheme Y . For any point $y \in Y$ we have an open neighbourhood $U \subseteq Y$ over which we have a decomposition $\pi_*\mathcal{O}_C(S)|_U \cong$

$\mathcal{O}_U^{\oplus n+1}$ and a basis y_0, \dots, y_n as in proposition 3.4, and we obtain functions $a_1, \dots, a_{n-1}, b_1, \dots, b_{n-1} \in \mathcal{O}_U(U)$. We define $(\mathcal{L}_{\varrho_i}, a_i) := (\mathcal{O}_U, a_i)$, $(\mathcal{L}_{\tau_i}, b_i) := (\mathcal{O}_U, b_i)$ and have the isomorphisms $c_i: \mathcal{L}_{\tau_i} \otimes \mathcal{L}_{\varrho_{i-1}} \otimes \mathcal{L}_{\varrho_i}^{\otimes -2} \otimes \mathcal{L}_{\varrho_{i+1}} \rightarrow \mathcal{O}_U$ (omit $\mathcal{L}_{\varrho_0}, \mathcal{L}_{\varrho_n}$) given by the identities on \mathcal{O}_U . These data form a $\Upsilon(A_{n-1})$ -collection over U , the nondegeneracy condition that not both $a_i = 0$ and $b_i = 0$ in each point is satisfied by construction and proposition 3.2.

Different choices of bases y_0, \dots, y_n and y'_0, \dots, y'_n over U and U' as above are related over $U'' = U \cap U'$ by $y_i = \kappa'_i y'_i$ for some $\kappa'_i \in \mathcal{O}_{U''}^*(U'')$, where $\kappa'_0 = \kappa'_n$. Let $\kappa_i = \kappa'_i / \kappa'_0$. There is an isomorphism between the corresponding $\Upsilon(A_{n-1})$ -collections over U'' that, with respect to the given trivialisations, is given by the collection $\kappa_1, \dots, \kappa_{n-1}, \lambda_1, \dots, \lambda_{n-1} \in \mathcal{O}_{U''}^*(U'')$ as in remark 2.5.

We cover Y by open subschemes U as above, obtain $\Upsilon(A_{n-1})$ -collections on this covering and glue them to a $\Upsilon(A_{n-1})$ -collection $\Phi\mathcal{C}$ on Y .

For a morphism $\mathcal{C}' \rightarrow \mathcal{C}$ of degree- n -pointed chains over $f: Y' \rightarrow Y$, i.e. a cartesian diagram consisting of f , a morphism $F: C' \rightarrow C$ that maps s'_-, s'_+, S' to s_-, s_+, S and $\pi: C \rightarrow Y$, $\pi': C' \rightarrow Y'$, we have a morphism of $\Upsilon(A_{n-1})$ -collections $\mathcal{L}' = \Phi\mathcal{C}' \rightarrow \mathcal{L} = \Phi\mathcal{C}$ over $f: Y' \rightarrow Y$. Locally over $U'' = f^{-1}(U) \cap U'$, where U' and U are elements of the chosen open coverings of Y' and Y , we have chosen local bases f^*y_0, \dots, f^*y_n and y'_0, \dots, y'_n of $f^*\pi_*\mathcal{O}_C(S) \cong \pi'_*F^*\mathcal{O}_C(S) \cong \pi'_*\mathcal{O}_{C'}(S')$. Comparing these bases gives rise to isomorphisms $f^*\mathcal{L}_{\varrho_i}|_{U''} \rightarrow \mathcal{L}'_{\varrho_i}|_{U''}$, $f^*\mathcal{L}_{\tau_i}|_{U''} \rightarrow \mathcal{L}'_{\tau_i}|_{U''}$ as above, and these can be glued to a morphism $\mathcal{L}' \rightarrow \mathcal{L}$.

One checks that this defines a functor $\Phi: \tilde{\mathcal{L}}_n \rightarrow \mathcal{C}_{\Upsilon(A_{n-1})}$. The functor Φ is base-preserving and sends cartesian arrows to cartesian arrows.

CONSTRUCTION 3.6. — Let $\mathcal{L} = ((\mathcal{L}_{\varrho_i}, a_i)_i, (\mathcal{L}_{\tau_i}, b_i)_i, (c_i)_i)$ be a $\Upsilon(A_{n-1})$ -collection on a scheme Y . For any point $y \in Y$ we have an open neighbourhood $U \subseteq Y$ over which we can choose trivialisations $\mathcal{L}_{\varrho_i}|_U, \mathcal{L}_{\tau_i}|_U \cong \mathcal{O}_U$ such that the isomorphisms $c_i: (\mathcal{L}_{\tau_i} \otimes \mathcal{L}_{\varrho_{i-1}} \otimes \mathcal{L}_{\varrho_i}^{\otimes -2} \otimes \mathcal{L}_{\varrho_{i+1}})|_U \rightarrow \mathcal{O}_U$ for $i = 1, \dots, n - 1$ (omit $\mathcal{L}_{\varrho_0}, \mathcal{L}_{\varrho_n}$) are the identities on \mathcal{O}_U . Let C be the closed subscheme of \mathbb{P}_U^n given by the equations $y_i y_{j+1} = b_{i+1} \cdots b_j y_{i+1} y_j$ for $0 \leq i < j \leq n - 1$, where y_0, \dots, y_n are homogeneous coordinates of \mathbb{P}_U^n and b_1, \dots, b_{n-1} are considered as regular functions on U via the isomorphisms $\mathcal{L}_{\tau_i}|_U \cong \mathcal{O}_U$, and let $\pi: C \rightarrow U$ be induced by $\mathbb{P}_U^n \rightarrow U$. By construction, the subscheme $C \subseteq \mathbb{P}_U^n$ is isomorphic to $\text{Proj}_U \mathcal{S}$ where \mathcal{S} is the graded algebra $\mathcal{O}_U[y_0, \dots, y_n] / \langle y_i y_{j+1} = b_{i+1} \cdots b_j y_{i+1} y_j; i < j \rangle$. The morphism $\pi: C \rightarrow U$ is flat since each graded piece of \mathcal{S} is locally free ([9, III, (7.9.14)]). Indeed, we have $\mathcal{S}_k \cong \mathcal{O}_U^{\oplus kn+1}$ with basis y_n^k and

$y_i^{k-l}y_{i+1}^l$ for $i = 0, \dots, n-1$ and $l = 0, \dots, k-1$. Let s_-, s_+ be the sections $(1 : 0 : \dots : 0), (0 : \dots : 0 : 1)$ with respect to the coordinates y_0, \dots, y_n and let $S \subset C$ be the subscheme given by the additional equation $y_n + a_{n-1}y_{n-1} + \dots + a_1y_1 + y_0 = 0$, where again a_1, \dots, a_{n-1} are considered as regular functions via $\mathcal{L}_{\rho_i}|_U \cong \mathcal{O}_U$. This defines a degree- n -pointed chain $(C \rightarrow U, s_-, s_+, S)$ over U .

Different choices of local trivialisations of the line bundles \mathcal{L}_{ρ_i} over U, U' are related by multiplication by some $\kappa_i \in \mathcal{O}_{U''}^*(U'')$ over $U'' = U \cap U'$. The corresponding degree- n -pointed chains in $\mathbb{P}_{U''}^n$ are connected by the automorphism of $\mathbb{P}_{U''}^n$ given by multiplying the homogeneous coordinates with $\kappa_0, \dots, \kappa_n$ (set $\kappa_0 = \kappa_n = 1$).

We cover Y by open subschemes U as above, obtain degree- n -pointed chains over this covering and glue them to a degree- n -pointed chain $\Psi\mathcal{L}$ over Y .

For a morphism $\mathcal{L}' \rightarrow \mathcal{L}$ of $\mathfrak{Y}(A_{n-1})$ -collections over $f: Y' \rightarrow Y$, i.e. a collection of isomorphisms of line bundles with sections $(f^*\mathcal{L}_{\rho_i}, f^*a_i) \rightarrow (\mathcal{L}'_{\rho_i}, a'_i), (f^*\mathcal{L}_{\tau_i}, f^*b_i) \rightarrow (\mathcal{L}'_{\tau_i}, b'_i)$, we have a morphism $\mathcal{C}' = \Psi\mathcal{L}' \rightarrow \mathcal{C} = \Psi\mathcal{L}$ of degree- n -pointed chains over $f: Y' \rightarrow Y$: locally on $U'' = f^{-1}(U) \cap U'$, where U' and U are elements of the chosen open coverings of Y' and Y , using the given trivialisations $\mathcal{L}'_{\rho_i}|_{U'} \cong \mathcal{O}_{U'}$ and $f^*\mathcal{L}_{\rho_i}|_U \cong f^*\mathcal{O}_U = \mathcal{O}_{f^{-1}(U)}$, the isomorphisms $f^*\mathcal{L}_{\rho_i} \rightarrow \mathcal{L}'_{\rho_i}$ are given by multiplication with elements $\kappa_i \in \mathcal{O}_{U''}^*(U'')$, and the automorphism of $\mathbb{P}_{U''}^n$ such that the coordinates y'_i and f^*y_i are related by multiplication by κ_i (set $\kappa_0 = \kappa_n = 1$) induces an isomorphism of the embedded degree- n -pointed chains over U'' . By glueing we obtain an isomorphism $\mathcal{C}' \rightarrow f^*\mathcal{C}$.

This defines a functor $\Psi: \mathcal{C}_{\mathfrak{Y}(A_{n-1})} \rightarrow \bar{\mathcal{L}}_n$ which is base-preserving and sends cartesian arrows to cartesian arrows.

These two functors give the equivalence of fibred categories stated in the theorem.

Proof of theorem 3.1. — We show that the fibred categories $\bar{\mathcal{L}}_n$ and $\mathcal{C}_{\mathfrak{Y}(A_{n-1})}$ are equivalent using the functors $\Psi: \mathcal{C}_{\mathfrak{Y}(A_{n-1})} \rightarrow \bar{\mathcal{L}}_n$ and $\Phi: \bar{\mathcal{L}}_n \rightarrow \mathcal{C}_{\mathfrak{Y}(A_{n-1})}$.


For an object $\mathcal{C} = (C \rightarrow Y, s_-, s_+, S)$ in $\bar{\mathcal{L}}_n$, after choice of a suitable open covering, the data $\Phi\mathcal{C}$ is given locally by $(\mathcal{L}_{\rho_i}|_U = \mathcal{O}_U, a_i), (\mathcal{L}_{\tau_i}|_U = \mathcal{O}_U, b_i)$ and locally \mathcal{C} is isomorphic to the embedded object in \mathbb{P}_U^n defined by the functions a_i, b_i with respect to coordinates y_0, \dots, y_n of \mathbb{P}_U^n , see construction 3.5. Applying the functor Ψ , we choose a covering and isomorphisms $\mathcal{L}_{\rho_i}|_{U'}, \mathcal{L}_{\tau_i}|_{U'} \rightarrow \mathcal{O}_{U'}$ as in construction 3.6 giving rise to functions \tilde{a}_i, \tilde{b}_i , and these define the object $\Psi\Phi\mathcal{C}$ locally embedded in

$\mathbb{P}_{U'}^n$, with homogeneous coordinates \tilde{y}_i . Comparing the two isomorphisms $\mathcal{L}_{\varrho_i}|_{U''} \rightarrow \mathcal{O}_{U''}$ over open subschemes $U'' = U \cap U'$, we obtain an isomorphism $\Psi\Phi\mathcal{C} \rightarrow \mathcal{C}$ locally over U'' using the two local embeddings of $\Psi\Phi\mathcal{C}$ and \mathcal{C} in $\mathbb{P}_{U''}^n$. One checks that these isomorphisms form an isomorphism of functors $\Psi \circ \Phi \cong \text{Id}$.


Starting with a $\Upsilon(A_{n-1})$ -collection \mathcal{L} on Y , after choice of a covering and isomorphisms $\mathcal{L}|_U \cong ((\mathcal{O}_U, a_i)_i, (\mathcal{O}_U, b_i)_i, (id)_i)$, we construct an object $\Psi\mathcal{L} = (\pi: C \rightarrow Y, s_-, s_+, S)$, locally embedded in \mathbb{P}_U^n with homogeneous coordinates y_0, \dots, y_n using the functions $a_i, b_i \in \mathcal{O}_U(U)$. From $\Psi\mathcal{L}$ we extract data $\Phi\Psi\mathcal{L}$ after choice of a suitable covering and local bases $\tilde{y}_0, \dots, \tilde{y}_n$ of $\pi_*(\mathcal{O}_C(S))$, locally given by $(\tilde{\mathcal{L}}_{\varrho_i}|_{U'} = \mathcal{O}_{U'}, \tilde{a}_i)$, $(\tilde{\mathcal{L}}_{\tau_i}|_{U'} = \mathcal{O}_{U'}, \tilde{b}_i)$ for elements U' of the covering. Comparing the two collections of homogeneous coordinates, satisfying the conditions of proposition 3.4, of $\mathbb{P}_U^n \times_U U'' \xrightarrow{\sim} \mathbb{P}_{U''}^n(\tilde{\pi}_*\mathcal{O}_{\mathbb{P}_U^n}(\tilde{S})|_{U''}) \cong \mathbb{P}_{U''}^n(\pi_*\mathcal{O}_C(S)|_{U''})$ where $\tilde{S} \subset \mathbb{P}_U^n$ is the hyperplane determined by the equation $\sum_i a_i y_i = 0$ (set $a_0, a_{n+1} = 1$) and $\tilde{\pi}: \mathbb{P}_U^n \rightarrow U$, and thus the two ways the object \mathcal{C} is locally embedded in $\mathbb{P}_{U''}^n$ for open sets $U'' = U \cap U'$, we obtain an isomorphism of $\Upsilon(A_{n-1})$ -collections $\Phi\Psi\mathcal{L} \rightarrow \mathcal{L}$ locally given by some $\kappa_1, \dots, \kappa_{n-1}, \lambda_1, \dots, \lambda_{n-1} \in \mathcal{O}_{U''}^*(U'')$ as in remark 2.5. One verifies that this gives an isomorphism of functors $\Phi \circ \Psi \cong \text{Id}$. \square

COROLLARY 3.7. — *The coarse moduli space of $\overline{\mathcal{L}}_n$, which coincides with the quotient $\overline{\mathcal{L}}_n/S_n$, is isomorphic to the toric variety $Y(A_{n-1})$ corresponding to the simplicial fan $\Upsilon(A_{n-1})$ underlying the stacky fan $\Upsilon(A_{n-1})$.*

Example 3.8. — In the case $n = 2$ we have the isomorphism $\overline{\mathcal{L}}_2 \cong \mathcal{Y}(A_1)$. The stacky fan of $\mathcal{Y}(A_1)$ was pictured in example 2.2. We have the following types of pointed chains over $\overline{\mathcal{L}}_2 \cong \mathcal{Y}(A_1)$ (cf. also example 1.8):



Example 3.9. — In the case $n = 3$ we have the isomorphism $\overline{\mathcal{L}}_3 \cong \mathcal{Y}(A_2)$. The stacky fan of $\mathcal{Y}(A_2)$ appeared in example 2.3. Here we picture the types of pointed chains over the torus invariant divisors of the moduli stack $\overline{\mathcal{L}}_3$.



4. The functor of $X(A_{n-1})$, Losev-Manin moduli spaces and the morphism to $\bar{\mathcal{L}}_n$

We start by comparing three descriptions of the functor of the toric variety associated with root systems of type A . We use notations as in [2, Section 2.1], in particular we have the lattice $M(A_{n-1}) = \langle u_i - u_j : i, j \in \{1, \dots, n\} \rangle \subset \bigoplus_{i=1}^n \mathbb{Z}u_i$, generated by the roots $\beta_{ij} = u_i - u_j$ and forming the character lattice for the toric variety $X(A_{n-1})$. Its dual $N(A_{n-1}) = \bigoplus_{i=1}^n \mathbb{Z}v_i / \sum_i v_i$, where $(u_i)_i$ and $(v_i)_i$ are dual bases, is the lattice for the fan $\Sigma(A_{n-1})$ of $X(A_{n-1})$.

The functor of the toric variety $X(A_{n-1})$ was described in [2] in terms of A_{n-1} -data, i.e. families $(\mathcal{L}_{\{\pm\beta_{ij}\}}, \{t_{\beta_{ij}}, t_{-\beta_{ij}}\})_{\{\pm\beta_{ij}\}}$ of line bundles with two generating sections that satisfy $t_\alpha t_\beta t_{-\gamma} = t_{-\alpha} t_{-\beta} t_\gamma$ if $\gamma = \alpha + \beta$, up to isomorphism of line bundles with a pair of sections. With pull-back of line bundles and its sections we have the functor $F_{A_{n-1}}$ of A_{n-1} -data, see [2, Def. 1.17].

On $X(A_{n-1})$ we have the universal A_{n-1} -data, which can be defined using the morphisms $\varphi_{\{\pm\beta_{ij}\}} : X(A_{n-1}) \rightarrow \mathbb{P}^1$ induced by pairs of opposite roots $\{\pm\beta_{ij}\}$ in A_{n-1} (see [2, Ex. 1.5 and 1.13]). We have homogeneous coordinates $z_{\beta_{ij}}, z_{-\beta_{ij}} \in H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$ such that $x^{\beta_{ij}} = \varphi_{\{\pm\beta_{ij}\}}^*(z_{\beta_{ij}}/z_{-\beta_{ij}})$, where x^u for $u \in M(A_{n-1})$ is the rational function corresponding to an element of the root lattice. Let $\mathcal{L}_{\{\pm\beta_{ij}\}} = \varphi_{\{\pm\beta_{ij}\}}^* \mathcal{O}_{\mathbb{P}^1}(1)$ and $t_{\beta_{ij}}, t_{-\beta_{ij}}$ be the pull-back of $z_{\beta_{ij}}, z_{-\beta_{ij}}$.

By [2, Thm. 1.20] the toric variety $X(A_{n-1})$ together with the universal A_{n-1} -data represents the functor $F_{A_{n-1}}$.

We can also apply the description of the functor of a smooth toric variety of Cox [5] to $X(A_{n-1})$. The fan $\Sigma(A_{n-1})$ gives rise to the notion of a $\Sigma(A_{n-1})$ -collection $((\mathcal{L}_I, w_I)_{I, (c_{ij})_{ij}}$ on a scheme Y , consisting of

line bundles on Y with a global section (\mathcal{L}_I, w_I) for $\emptyset \neq I \subsetneq \{1, \dots, n\}$ and isomorphisms $c_{ij} : (\bigotimes_{i \in I, j \notin I} \mathcal{L}_I) \otimes (\bigotimes_{i \notin I, j \in I} \mathcal{L}_I^{\otimes -1}) \rightarrow \mathcal{O}_Y$ for $i, j \in \{1, \dots, n\}$, $i \neq j$, such that identifications of the form $c_{ij} \otimes c_{jk} = c_{ik}$ hold. These data have to satisfy the nondegeneracy condition that for any point $y \in Y$ there are sets $I_1 \subset \dots \subset I_{n-1} \subset \{1, \dots, n\}$ with $|I_i| = i$ such that $w_I(y) \neq 0$ if $I \neq I_1, \dots, I_{n-1}$. We denote the functor of $\Sigma(A_{n-1})$ -collections by $C_{\Sigma(A_{n-1})}$.

On $X(A_{n-1})$ we have the universal $\Sigma(A_{n-1})$ -collection given by the line bundles $\mathcal{L}_I = \mathcal{O}_{X(A_{n-1})}(D_I)$, where D_I is the torus invariant prime divisor corresponding to the ray generated by $\sum_{i \in I} v_i$, with the section w_I arising as the image of the 1-section under the natural inclusion $\mathcal{O}_{X(A_{n-1})} \rightarrow \mathcal{O}_{X(A_{n-1})}(D_I)$ and the isomorphisms $c_{ij} : \mathcal{O}_{X(A_{n-1})}(\sum_{i \in I, j \notin I} D_I - \sum_{i \notin I, j \in I} D_I) \rightarrow \mathcal{O}_{X(A_{n-1})}$ induced by multiplication with the rational functions $x^{\beta_{ij}}$ on $X(A_{n-1})$.

By [5] the toric variety $X(A_{n-1})$ together with the universal $\Sigma(A_{n-1})$ -collection $((\mathcal{O}_{X(A_{n-1})}(D_I), w_I)_I, (c_{ij})_{ij})$ represents the functor $C_{\Sigma(A_{n-1})}$.

As both functors $C_{\Sigma(A_{n-1})}$ and $F_{A_{n-1}}$ are isomorphic to the functor of the toric variety $X(A_{n-1})$, we have an isomorphism of functors $C_{\Sigma(A_{n-1})} \rightarrow F_{A_{n-1}}$, which we describe explicitly.

PROPOSITION 4.1. — *By the following procedure we can construct A_{n-1} -data $(\mathcal{L}_{\{\pm\beta_{ij}\}}, \{t_{\beta_{ij}}, t_{-\beta_{ij}}\})_{\{\pm\beta_{ij}\}}$ out of a $\Sigma(A_{n-1})$ -collection $((\mathcal{L}_I, w_I)_I, (c_{ij})_{ij})$ over a scheme Y : for a pair of opposite roots $\pm\beta_{ij}$ in A_{n-1} we have isomorphisms $\bigotimes_{i \in I, j \notin I} \mathcal{L}_I \xrightarrow{\sim} \bigotimes_{i \notin I, j \in I} \mathcal{L}_I$ of line bundles on Y defined by c_{ij}, c_{ji} inverse to each other, and we let $\mathcal{L}_{\{\pm\beta_{ij}\}}$ be a line bundles in the same isomorphism class, with the sections $t_{\beta_{ij}}, t_{-\beta_{ij}}$ defined as the images of $\prod_{i \in I, j \notin I} w_I, \prod_{i \notin I, j \in I} w_I$ in $\mathcal{L}_{\{\pm\beta_{ij}\}}$ under isomorphisms compatible with the above. This construction defines an isomorphism of functors $C_{\Sigma(A_{n-1})} \rightarrow F_{A_{n-1}}$, mapping the universal $\Sigma(A_{n-1})$ -collection to the universal A_{n-1} -data.*

Proof. — This construction defines a morphism of functors $C_{\Sigma(A_{n-1})} \rightarrow F_{A_{n-1}}$, in particular the requirement that the two sections $t_{\pm\beta_{ij}}$ as defined in the construction generate the line bundle $\mathcal{L}_{\{\pm\beta_{ij}\}}$ follows from the nondegeneracy condition of $\Sigma(A_{n-1})$ -data. One can show that this morphism of functors is an isomorphism by showing that it coincides with the composition of isomorphisms $C_{\Sigma(A_{n-1})} \rightarrow \text{Mor}(\cdot, X(A_{n-1})) \rightarrow F_{A_{n-1}}$. This follows from the fact that the universal $\Sigma(A_{n-1})$ -collection is mapped to the universal A_{n-1} -data, which is easy to verify. \square

Considering the universal data on $X(A_{n-1})$, we have isomorphisms $\bigotimes_{i \in I} \mathcal{L}_I \xrightarrow{\sim} \bigotimes_{j \in I} \mathcal{L}_I$ via multiplication by the rational function $x^{\beta_{ij}}$. For any chosen $j \in \{1, \dots, n\}$ we may define $x_1, \dots, x_n \in H^0(X(A_{n-1}), \bigotimes_{j \in I} \mathcal{L}_I)$ as images of the sections $\prod_{1 \in I} w_I, \dots, \prod_{n \in I} w_I$ under these isomorphisms. We then have $x^{\beta_{ij}} = x_i/x_j$.

DEFINITION 4.2. — Given an ordering i_1, \dots, i_n of the set $\{1, \dots, n\}$, we define line bundles $\mathcal{L}_1, \dots, \mathcal{L}_{n-1}$ on $X(A_{n-1})$ and sections x_J of $\mathcal{L}_{|J|}$. Let

$$\mathcal{L}_j = \bigotimes_I \mathcal{L}_I^{\otimes (|\{i_1, \dots, i_j\} \cap I| - \max\{0, |I| + j - n\})}$$

be defined in terms of the universal $\Sigma(A_{n-1})$ -collection. The line bundles $\bigotimes_I \mathcal{L}_I^{\otimes (|J \cap I| - \max\{0, |I| + j - n\})}$ for any $J \subset \{1, \dots, n\}$ of cardinality j are isomorphic to \mathcal{L}_j via multiplication by $\prod_{i \in J} x_i / \prod_{k=1}^j x_{i_k}$ (can also be expressed in terms of the isomorphisms c_{ij} being part of the universal $\Sigma(A_{n-1})$ -collection). We define $x_J \in H^0(X(A_{n-1}), \mathcal{L}_j)$ as the image of $\prod_I w_I^{|J \cap I| - \max\{0, |I| + j - n\}}$ under this isomorphism.

For these sections $x_J \in H^0(X(A_{n-1}), \mathcal{L}_{|J|})$ we have equations of rational functions

$$\prod_{j \in J} x_j / \prod_{j \in J'} x_j = x_J / x_{J'} = \prod_{j \in J'} x_{\{1, \dots, n\} \setminus \{j\}} / \prod_{j \in J} x_{\{1, \dots, n\} \setminus \{j\}}.$$

Remark 4.3. — The line bundle \mathcal{L}_j was defined as $\mathcal{O}_{X(A_{n-1})}(D)$ in terms of the divisor $D = \sum_I d_I D_i$, where $d_I = |\{i_1, \dots, i_j\} \cap I| - \max\{0, |I| + j - n\}$, which corresponds to the lattice polytope

$$\begin{aligned} \Delta_j(A_{n-1}) &= \bigcap_I \{u \in M(A_{n-1})_{\mathbb{Q}} : \sum_{i \in I} v_i(u) \geq -d_I\} \\ &= \text{conv} \{ \sum_{i \in J} u_i - \sum_{k=1}^j u_{i_k} : |J| = j \} \end{aligned}$$

in $M(A_{n-1})_{\mathbb{Q}}$; the elements $x_J \in H^0(X(A_{n-1}), \mathcal{L}_j)$ for $|J| = j$ form a basis of global sections. Different choices of the ordering fixed in the definition give rise to translated polytopes.

The line bundle \mathcal{L}_1 with its basis of global sections x_1, \dots, x_n defines a morphism $X(A_{n-1}) \rightarrow \mathbb{P}^{n-1}$. This morphism is a composition of toric blow-ups as described in [12, (4.3.13)]. It maps the divisors $D_{\{i\}}$ of $X(A_{n-1})$ to the torus invariant prime divisor $D_i = \{x_i = 0\}$ of \mathbb{P}^{n-1} , and more generally D_I to $\bigcap_{i \in I} D_i$. Similarly, we have a morphism $X(A_{n-1}) \rightarrow \mathbb{P}^{n-1}$ defined by the line bundle \mathcal{L}_{n-1} mapping the divisor $D_{\{1, \dots, n\} \setminus \{i\}}$ of $X(A_{n-1})$ to the torus invariant prime divisor $D_i^* = \{x_{\{1, \dots, n\} \setminus \{i\}} = 0\}$ of \mathbb{P}^{n-1} , and more generally $D_{\{1, \dots, n\} \setminus I}$ to $\bigcap_{i \in I} D_i^*$, see also [12, (4.3.14)]. We also consider the morphisms defined by the other line bundles \mathcal{L}_j :

PROPOSITION 4.4. — *The line bundle \mathcal{L}_j is generated by the basis of global sections $(x_J)_{|J|=j}$, it determines a projective toric morphism*

$$X(A_{n-1}) \rightarrow \mathbb{P}(\langle x_J : |J| = j \rangle) \cong \mathbb{P}^{\binom{n}{j}-1}$$

which is birational onto its image. Together, these morphisms form a closed embedding

$$X(A_{n-1}) \rightarrow \prod_{j=1}^{n-1} \mathbb{P}(\langle x_J : |J| = j \rangle) \cong \prod_{j=1}^{n-1} \mathbb{P}^{\binom{n}{j}-1}. \tag{4.1}$$

The subscheme $X(A_{n-1})$ in this product is defined by homogeneous equations

$$\prod_{i=1}^l x_{J_i} = \prod_{i=1}^l x_{J'_i} \tag{4.2}$$

where $\emptyset \neq J_i, J'_i \subsetneq \{1, \dots, n\}$ such that $|J_i| = |J'_i|$ and the equation for the characteristic functions $\sum_i \chi_{J_i} = \sum_i \chi_{J'_i}$ is satisfied.

Proof. — That \mathcal{L}_j is generated by global sections $(x_J)_{|J|=j}$ and determines a projective toric morphism follows from the fact that \mathcal{L}_j can be reconstructed from the polytope $\Delta_j(A_{n-1})$, see remark 4.3. This morphism is birational onto its image since the polytope is full-dimensional. We describe these morphisms in terms of the corresponding maps of fans.

For the toric variety $\mathbb{P}(\langle x_J : |J| = j \rangle)$ we have the character lattice $M(A_{n-1})_j \subset \bigoplus_{|J|=j} \mathbb{Z}u_J$ generated by differences $u_J - u_{J'}$ and the dual lattice $N(A_{n-1})_j = (\bigoplus_{|J|=j} \mathbb{Z}v_J) / (\sum_J v_J)$. The fan of $\mathbb{P}(\langle x_J : |J| = j \rangle)$ has the one-dimensional cones generated by the v_J . The morphism $X(A_{n-1}) \rightarrow \mathbb{P}(\langle x_J : |J| = j \rangle)$ is determined by the map of lattices $M(A_{n-1})_j \rightarrow M(A_{n-1})$, $u_J \mapsto \sum_{i \in J} u_i$, or dually $N(A_{n-1}) \rightarrow N(A_{n-1})_j$, $v_i \mapsto \sum_{i \in J} v_J$, which defines a map of fans.

The product of these morphisms is given by the map of lattices $\bigoplus_{j=1}^n M(A_{n-1})_j \rightarrow M(A_{n-1})$ with kernel generated by elements of the form $\sum_i n_{J_i} - \sum_i n_{J'_i}$ such that $|J_i| = |J'_i|$ and $\sum_i \chi_{J_i} = \sum_i \chi_{J'_i}$. This gives rise to the homogeneous equations.

The maximal cones of the fan of $\mathbb{P}(\langle x_J : |J| = j \rangle)$ are the cones σ_J generated by $\{v_{J'} \mid J' \neq J, |J'| = j\}$. For an ordering i_1, \dots, i_n of $\{1, \dots, n\}$ the preimage of the maximal cone $\sigma_{\{i_n\}} \times \sigma_{\{i_n, i_{n-1}\}} \times \dots \times \sigma_{\{i_n, \dots, i_2\}}$ of the product fan is the maximal cone of $\Sigma(A_{n-1})$ generated by $v_{i_1}, v_{i_1} + v_{i_2}, \dots, v_{i_1} + \dots + v_{i_{n-1}}$, thus the open sets corresponding to maximal cones of this form cover the image of $X(A_{n-1})$. The corresponding maps of coordinate algebras are surjective, so the morphism is a closed embedding. \square

Remark 4.5. — The $(n-1)$ -dimensional permutohedron, usually defined in an n -dimensional vector space as convex hull of the orbit of $(n, n-1, \dots, 1)$ under the action of the symmetric group S_n permuting the given

basis (cf. for example [12, (4.3.10)]), can be considered as a lattice polytope in $M(A_{n-1})_{\mathbb{Q}} \subset \mathbb{Q}^n$ after a translation moving one of its vertices, specified by fixing an ordering i_1, \dots, i_n of the set $\{1, \dots, n\}$, to the origin:

$$\Delta(A_{n-1}) = \text{conv} \left\{ \sum_{k=1}^{n-1} (n-k)u_{\sigma(k)} - \sum_{k=1}^{n-1} (n-k)u_{i_k} : \sigma \in S_n \right\}.$$

We have Minkowski sum decompositions of the permutohedron, first

$$\Delta(A_{n-1}) = \sum_{k < j} l_{i_j i_k}$$

into line segments $l_{i_j} = \{r \cdot \beta_{i_j} \mid 0 \leq r \leq 1\}$ corresponding to the line bundles $\mathcal{L}_{\{\pm\beta_{i_j}\}}$ forming the universal A_{n-1} -data (choosing $\mathcal{O}_{X(A_{n-1})}(\sum_{i_k \in I, i_j \notin I} D_I)$ in the isomorphism class of $\mathcal{L}_{\{\pm\beta_{i_j i_k}\}}$ if $k < j$), and second

$$\Delta(A_{n-1}) = \Delta_1(A_{n-1}) + \dots + \Delta_{n-1}(A_{n-1})$$

into the polytopes corresponding to the line bundles \mathcal{L}_j .

Remark 4.6. — The closed embedding (4.1) together with the functor of projective spaces gives another description of the functor of the toric variety $X(A_{n-1})$. We have a contravariant functor on the category of schemes: its data on a scheme Y are line bundles with generating sections $(\mathcal{L}_j, (x_J)_{|J|=j})_{j=1, \dots, n-1}$ up to isomorphism such that the sections satisfy the relations (4.2), and for morphisms of schemes we have the pull-back of line bundles with sections. We call the data on $X(A_{n-1})$ introduced in definition 4.2 the universal data on $X(A_{n-1})$. Then, the toric variety $X(A_{n-1})$ together with the universal data represents this functor. Further, the method of definition 4.2 applied to $\Sigma(A_{n-1})$ -data over arbitrary schemes gives a morphism from $C_{\Sigma(A_{n-1})}$ to this functor, mapping the universal $\Sigma(A_{n-1})$ -collection to the universal data. As in the proof of proposition 4.1 this implies that we have an isomorphism of functors.

The following observation can be directly calculated from the definition of the line bundles \mathcal{L}_j .

LEMMA 4.7. — *We have isomorphisms*

$$\mathcal{L}_{j-1}^{\otimes -1} \otimes \mathcal{L}_j^{\otimes 2} \otimes \mathcal{L}_{j+1}^{\otimes -1} \cong \bigotimes_{|J|=n-j} \mathcal{L}_J \tag{4.3}$$

where we set $\mathcal{L}_0 = \mathcal{L}_n = \mathcal{O}_{X(A_{n-1})}$.

DEFINITION 4.8. — *We define the divisors C_1, \dots, C_{n-1} and D_1, \dots, D_{n-1} on $X(A_{n-1})$. Let $D_j = \sum_{|J|=n-j} D_J$ and let C_j be the zero divisor of the section $\sum_{|J|=j} x_J$ of the line bundle \mathcal{L}_j .*

Remark 4.9. — We may write the isomorphism (4.3) as linear equivalence of divisors

$$2C_j - C_{j-1} - C_{j+1} \sim D_j.$$

The rational function

$$\frac{(\sum_{|I|=j-1} x_I)(\sum_{|I|=j+1} x_I)}{(\sum_{|I|=j} x_I)^2} \tag{4.4}$$

has divisor $D_j + C_{j-1} + C_{j+1} - 2C_j$.

LEMMA 4.10. — For $j = 1, \dots, n-1$ we have $C_j \cap D_j = \emptyset$.

Proof. — Can easily be checked locally using the covering of the following remark. □

Remark 4.11. — Given a $\Sigma(A_{n-1})$ -collection $((\mathcal{L}_I, w_I)_I, (c_{ij})_{ij})$ on a scheme Y , by nondegeneracy we have the following covering of Y by open subschemes: for a permutation $\sigma \in S_n$ set $\mathcal{I}_\sigma = \{\{\sigma(n)\}, \{\sigma(n), \sigma(n-1)\}, \dots, \{\sigma(n), \dots, \sigma(2)\}\}$ and let W_σ be the open subscheme of Y where $w_I \neq 0$ for $I \notin \mathcal{I}_\sigma$.

In the case of the universal $\Sigma(A_{n-1})$ -collection on $X(A_{n-1})$ the subscheme $W_\sigma \subset X(A_{n-1})$ corresponds to the maximal cone $\langle v_{\sigma(n)}, \dots, v_{\sigma(n)} + \dots + v_{\sigma(2)} \rangle \subset N(A_{n-1})_{\mathbb{Q}}$ dual to the cone generated by the simple roots $u_{\sigma(n)} - u_{\sigma(n-1)}, \dots, u_{\sigma(2)} - u_{\sigma(1)}$ and has as coordinate algebra the polynomial ring generated by $\frac{x_{\sigma(n)}}{x_{\sigma(n-1)}}, \dots, \frac{x_{\sigma(2)}}{x_{\sigma(1)}}$.

By [2, Thm. 3.19] there is an isomorphism between the functor $F_{A_{n-1}}$ and the moduli functor of n -pointed chains of \mathbb{P}^1 \bar{L}_n mapping the universal A_{n-1} -data to the universal n -pointed chain $(X(A_n) \rightarrow X(A_{n-1}), s_-, s_+, s_1, \dots, s_n)$ defined in [2, Con. 3.6]. This means that the toric variety $X(A_{n-1})$ coincides with the Losev-Manin moduli space \bar{L}_n (we use the same symbol for the functor and the moduli space). The construction uses an embedding of n -pointed chains into $(\mathbb{P}^1)^n$.

This also implies that there is an isomorphism between the functor $C_{\Sigma(A_{n-1})}$ and the moduli functor \bar{L}_n compatible with the other isomorphisms of functors. We make this isomorphism explicit using an embedding of n -pointed chains into \mathbb{P}^n .

CONSTRUCTION 4.12. — Let $((\mathcal{L}_I, w_I)_I, (c_{ij})_{ij})$ be a $\Sigma(A_{n-1})$ -collection over a scheme Y . We construct an n -pointed chain of \mathbb{P}^1 $(C \rightarrow Y, s_-, s_+, s_1, \dots, s_n)$ using the covering of Y by $(W_\sigma)_{\sigma \in S_n}$ (see remark 4.11).

For $\sigma \in S_n$ the restricted $\Sigma(A_{n-1})$ -collection $((\mathcal{L}_I|_{W_\sigma}, w_I|_{W_\sigma})_I, (c_{ij}|_{W_\sigma})_{ij})$ is isomorphic to a $\Sigma(A_{n-1})$ -collection $((\mathcal{L}_I^\sigma, w_I^\sigma)_I, (c_{ij}^\sigma)_{ij})$ on W_σ with the property $(\mathcal{L}_I^\sigma, w_I^\sigma) = (\mathcal{O}_{W_\sigma}, 1)$ for $I \notin \mathcal{I}_\sigma$, and for $i = 1, \dots, n-1$ we have isomorphisms $c_{\sigma(i+1), \sigma(i)}^\sigma : \mathcal{L}_{\{\sigma(n), \dots, \sigma(i+1)\}}^\sigma \rightarrow \mathcal{O}_{W_\sigma}$. Let $w_i^\sigma \in \mathcal{O}_{W_\sigma}(W_\sigma)$ be the image of $w_{\{\sigma(n), \dots, \sigma(i+1)\}}$. Equivalently, we can use the restricted original data $((\mathcal{L}_I|_{W_\sigma}, w_I|_{W_\sigma})_I, (c_{ij}|_{W_\sigma})_{ij})$ and the image of the respective product of the restricted w_I 's under $c_{\sigma(i+1), \sigma(i)}|_{W_\sigma}$.

>From these functions $w_1^\sigma, \dots, w_{n-1}^\sigma$ we construct an n -pointed chain over W_σ embedded in the projective space $\mathbb{P}_{W_\sigma}^n$ with homogeneous coordinates y_0, \dots, y_n . Let C_σ be the subscheme of $\mathbb{P}_{W_\sigma}^n$ defined by the equations $y_i y_{j+1} = w_{i+1}^\sigma \cdots w_j^\sigma y_{i+1} y_j$ for $0 \leq i < j < n$ (cf. construction 3.6), the sections $s_{\sigma(i)}^\sigma$ defined by the additional equation $y_{i-1} = y_i \neq 0$, and let s_-^σ, s_+^σ be the sections $(1:0:\dots:0), (0:\dots:0:1)$.

These n -pointed chains $(C_\sigma \rightarrow W_\sigma, s_-^\sigma, s_+^\sigma, s_1^\sigma, \dots, s_n^\sigma)$ can be glued to an n -pointed chain $(C \rightarrow Y, s_-, s_+, s_1, \dots, s_n)$ over Y .

PROPOSITION 4.13. — *Construction 4.12 is valid and defines an isomorphism between the functor $C_{\Sigma(A_{n-1})}$ and the moduli functor of n -pointed chains of \mathbb{P}^1 mapping the universal $\Sigma(A_{n-1})$ -collection $((\mathcal{O}_{X(A_{n-1})}(D_I), w_I)_I, (c_{ij})_{ij})$ to the universal n -pointed chain $(X(A_n) \rightarrow X(A_{n-1}), s_-, s_+, s_1, \dots, s_n)$.*

Proof. — Given $\Sigma(A_{n-1})$ -data over a scheme Y , it is easy to show that construction 4.12 locally over the open subschemes $W_\sigma \subseteq Y$ defines n -pointed chains of \mathbb{P}^1 (compare also to construction 3.6).

We show that, applying the isomorphism of functors $\bar{L}_n \rightarrow F_{A_{n-1}}$ to these objects over W_σ , we obtain A_{n-1} -data which coincide with the data we get by applying the functor $C_{\Sigma(A_{n-1})} \rightarrow F_{A_{n-1}}$ to the restricted data. According to [2, Section 3.3] we extract A_{n-1} -data from an n -pointed chain $(C_\sigma \rightarrow W_\sigma, s_-^\sigma, s_+^\sigma, s_1^\sigma, \dots, s_n^\sigma)$ via projections to $\mathbb{P}_{W_\sigma}^1$ such that s_-^σ, s_+^σ become the $(1:0), (0:1)$ -section and a given section s_i^σ becomes the section $(1:1)$. In the present case for $i = 1, \dots, n$ the morphism determined by the rational functions $1, y_i/y_{i-1}$ restricted to the component of C_σ containing $s_{\sigma(i)}^\sigma$ after contracting the other components transforms the sections $s_-^\sigma, s_+^\sigma, s_{\sigma(i)}^\sigma$ into the $(1:0), (0:1), (1:1)$ -sections. For $n = 1, \dots, n-1$ the section $s_{\sigma(i+1)}^\sigma$ becomes the section $(w_i^\sigma : 1)$, and this gives $(t_{-\beta_{\sigma(i), \sigma(i+1)}} : t_{\beta_{\sigma(i), \sigma(i+1)}}) = (w_i^\sigma : 1)$ which coincides with the data obtained via proposition 4.1.

Thus, the chains over the open subschemes W_σ can be glued to an n -pointed chain over the scheme Y and construction 4.12 defines a morphism of functors $C_{\Sigma(A_{n-1})} \rightarrow \bar{L}_n$, such that its composition with $\bar{L}_n \rightarrow F_{A_{n-1}}$

coincides with the isomorphism of functors $C_{\Sigma(A_{n-1})} \rightarrow F_{A_{n-1}}$ defined in proposition 4.1. Since the other morphisms of functors are isomorphisms and map the given universal objects to the given universal objects, this is also true for $C_{\Sigma(A_{n-1})} \rightarrow \bar{L}_n$. \square

THEOREM 4.14. — *The morphism $\bar{L}_n \rightarrow \bar{L}_n$ that arises by forgetting the labels of the n sections is given by the following $\Upsilon(A_{n-1})$ -collection on $\bar{L}_n = X(A_{n-1})$: for $i = 1, \dots, n-1$ let $\mathcal{L}_{\varrho_i} = \mathcal{O}_{\bar{L}_n}(C_i)$ and $\mathcal{L}_{\tau_i} = \mathcal{O}_{\bar{L}_n}(D_i)$, let*

$$c_i : \mathcal{L}_{\tau_i} \mathcal{L}_{\varrho_{i-1}} \mathcal{L}_{\varrho_i}^{\otimes -2} \mathcal{L}_{\varrho_{i+1}} = \mathcal{O}_{\bar{L}_n}(D_i + C_{i-1} - 2C_i + C_{i+1}) \rightarrow \mathcal{O}_{\bar{L}_n}$$

be given by multiplication by the rational function (4.4), and let the sections a_i, b_i be defined as the images of the 1-sections under the inclusions $\mathcal{O}_{\bar{L}_n} \rightarrow \mathcal{L}_{\varrho_i}, \mathcal{O}_{\bar{L}_n} \rightarrow \mathcal{L}_{\tau_i}$.

Proof. — The data defined form a $\Upsilon(A_{n-1})$ -collection, nondegeneracy follows from $C_i \cap D_i = \emptyset$, see lemma 4.10.

We use the covering by $W_\sigma, \sigma \in S_n$ (see remark 4.11). We have an isomorphism $((\mathcal{L}_{\varrho_i}, a_i)_i, (\mathcal{L}_{\tau_i}, b_i)_i, (c_i)_i)|_{W_\sigma} \rightarrow ((\mathcal{O}_{W_\sigma}, a_i^\sigma)_i, (\mathcal{O}_{W_\sigma}, b_i^\sigma)_i, (id)_i)$ of $\Upsilon(A_{n-1})$ -collections on W_σ consisting of isomorphisms $\mathcal{L}_{\tau_i}|_{W_\sigma} \rightarrow \mathcal{O}_{W_\sigma}$ given by multiplication with $x_{\sigma(i+1)}/x_{\sigma(i)}$ (compare to construction 4.12) and $\mathcal{L}_{\varrho_i}|_{W_\sigma} \rightarrow \mathcal{O}_{W_\sigma}$ by multiplication with $(\sum_{|I|=i} x_I)/x_{\{\sigma(1), \dots, \sigma(i)\}}$.

We show that the degree- n -pointed chain constructed from these data coincides with the degree- n -pointed chain that arises by forgetting the labels of the universal n -pointed chain coming from the universal $\Sigma(A_{n-1})$ -collection by proposition 4.13.

Applying construction 3.6 to these data, we get a chain of $\mathbb{P}^1 C \subset \mathbb{P}_{W_\sigma}^n$ defined by the functions $b_i^\sigma = x_{\sigma(i+1)}/x_{\sigma(i)}$ and a subscheme $S \subset C$ finite of degree n over W_σ defined by the functions $a_i^\sigma = (\sum_{|I|=i} x_I)/x_{\{\sigma(1), \dots, \sigma(i)\}}$.

Applying construction 4.12 to the universal $\Sigma(A_{n-1})$ -collection on $X(A_{n-1})$, locally over $W_\sigma \subset X(A_{n-1})$ again we get the chain of $\mathbb{P}^1 C \subset \mathbb{P}_{W_\sigma}^n$ defined by the functions $b_i^\sigma = x_{\sigma(i+1)}/x_{\sigma(i)}$. The n sections are

$$s_{\sigma(i)} = (\dots : \frac{x_{\sigma(i)}^2}{x_{\sigma(i-2)}x_{\sigma(i-1)}} : \frac{x_{\sigma(i)}}{x_{\sigma(i-1)}} : 1 : 1 : \frac{x_{\sigma(i+1)}}{x_{\sigma(i)}} : \frac{x_{\sigma(i+1)}x_{\sigma(i+2)}}{x_{\sigma(i)}^2} : \dots)$$

in terms the coordinates y_0, \dots, y_n of $\mathbb{P}_{W_\sigma}^n$, that is, we have $y_{i-1}(s_{\sigma(i)}) = y_i(s_{\sigma(i)})$ which we may set to 1, and then $y_k(s_{\sigma(i)}) = x_{\sigma(i)}^{i-k} \frac{x_{\sigma(1)} \dots x_{\sigma(k)}}{x_{\sigma(1)} \dots x_{\sigma(i)}}$. The sections are contained in the hyperplane defined by $\sum_{k=0}^n \binom{n}{k} a_k^\sigma y_k = 0$ (set $a_0^\sigma = a_n^\sigma = 1$):

$$\begin{aligned}
 \sum_{k=0}^n (-)^k a_k^\sigma y_k(s_{\sigma(i)}) &= \frac{x_{\sigma(i)}^i}{x_{\sigma(1)} \cdots x_{\sigma(i)}} \sum_{k=0}^n (-)^k \sum_{\substack{|I|=k \\ \sigma(i) \notin I}} \frac{x_I x_{\sigma(1)} \cdots x_{\sigma(k)}}{x_{\{\sigma(1), \dots, \sigma(k)\}} x_{\sigma(i)}^k} \\
 &= \frac{x_{\sigma(i)}^i}{x_{\sigma(1)} \cdots x_{\sigma(i)}} \sum_{k=0}^n (-)^k \left(\sum_{\substack{|I|=k \\ \sigma(i) \notin I}} \frac{x_I x_{\sigma(1)} \cdots x_{\sigma(k)}}{x_{\{\sigma(1), \dots, \sigma(k)\}} x_{\sigma(i)}^k} + \sum_{\substack{|I|=k \\ \sigma(i) \in I}} \frac{x_I x_{\sigma(1)} \cdots x_{\sigma(k)}}{x_{\{\sigma(1), \dots, \sigma(k)\}} x_{\sigma(i)}^k} \right) \\
 &= \frac{x_{\sigma(i)}^i}{x_{\sigma(1)} \cdots x_{\sigma(i)}} \sum_{k=0}^n \left((-)^k \sum_{\substack{|I|=k \\ \sigma(i) \notin I}} \frac{x_I x_{\sigma(1)} \cdots x_{\sigma(k)}}{x_{\{\sigma(1), \dots, \sigma(k)\}} x_{\sigma(i)}^k} - \right. \\
 &\qquad \qquad \qquad \left. (-)^{k-1} \sum_{\substack{|I|=k-1 \\ \sigma(i) \notin I}} \frac{x_I x_{\sigma(1)} \cdots x_{\sigma(k-1)}}{x_{\{\sigma(1), \dots, \sigma(k-1)\}} x_{\sigma(i)}^{k-1}} \right) \\
 &= 0
 \end{aligned}$$

The relative effective divisors $\sum_i s_i$ and S in C over W_σ coincide since they coincide over the open dense subscheme of \overline{L}_n parametrising chains with distinct sections. \square

Remark 4.15. — The results of this section imply a construction of a morphism of fibred categories from the functor of $\Sigma(A_{n-1})$ -collections, considering the S_n -operation on this functor, to the category of $\Upsilon(A_{n-1})$ -collections such that the diagram

$$\begin{array}{ccc}
 C_{\Sigma(A_{n-1})} & \xleftarrow{\sim} & \overline{L}_n \\
 \downarrow & & \downarrow \\
 C_{\Upsilon(A_{n-1})} & \xleftarrow{\sim} & \overline{\mathcal{L}}_n
 \end{array}$$

commutes.

Example 4.16. — In the case $n = 2$ the functor $F_{A_1} \cong C_{\Sigma(A_1)} \rightarrow C_{\Upsilon(A_1)}$ maps an object $(\mathcal{L}_{\{\pm\beta_{12}\}}, \{t_{\beta_{12}}, t_{-\beta_{12}}\})$ of F_{A_1} over a scheme Y to the object

$$\begin{aligned}
 &((\mathcal{O}_Y(C_1), a_1), (\mathcal{O}_Y(D_1), b_1), c_1: \mathcal{O}_Y(D_1) \otimes \mathcal{O}_Y(C_1)^{\otimes -2} \xrightarrow{\sim} \mathcal{O}_Y) \cong \\
 &((\mathcal{L}_{\{\pm\beta_{12}\}}, t_{\beta_{12}} + t_{-\beta_{12}}), (\mathcal{L}_{\{\pm\beta_{12}\}}^{\otimes 2}, t_{\beta_{12}} t_{-\beta_{12}}), \mathcal{L}_{\{\pm\beta_{12}\}}^{\otimes 2} \otimes \mathcal{L}_{\{\pm\beta_{12}\}}^{\otimes -2} \xrightarrow{\sim} \mathcal{O}_Y)
 \end{aligned}$$

of $C_{\Upsilon(A_1)}$, cf. example 1.8.

5. Pointed chains with involution and Cartan matrices of type B and C

As a natural variation of $\overline{\mathcal{L}}_n$ we consider moduli stacks $\overline{\mathcal{L}}_n^\pm$ of stable degree- $2n$ -pointed chains of \mathbb{P}^1 with an involution.

DEFINITION 5.1. — We define the fibred category $\overline{\mathcal{L}}_n^\pm$ of stable degree- $2n$ -pointed chains of \mathbb{P}^1 with involution. An object over a scheme Y is a collection $(C \rightarrow Y, I, s_-, s_+, S)$, where $(C \rightarrow Y, s_-, s_+, S)$ is a stable

degree- $2n$ -pointed chain of \mathbb{P}^1 over Y (definition 1.1), I an automorphism of C over Y such that $I^2 = id_C$ and $I(s_-) = s_+$, and S is invariant under I . Morphisms between objects are morphisms of degree- $2n$ -pointed chains which commute with the involution I .

As in the case of $\overline{\mathcal{L}}_n$, see proposition 1.4, the fibred category $\overline{\mathcal{L}}_n^\pm$ is a stack in the fpqc topology with representable finite diagonal.

Considering degree- $2n$ -pointed chains of \mathbb{P}^1 with involution as degree- $2n$ -pointed chains defines a morphism of stacks $\overline{\mathcal{L}}_n^\pm \rightarrow \overline{\mathcal{L}}_{2n}$ which makes $\overline{\mathcal{L}}_n^\pm$ a subcategory of $\overline{\mathcal{L}}_{2n}$ but in general not a substack, because a stable degree- $2n$ -pointed chain may have automorphisms not commuting with an additional involution.

The moduli stack $\overline{\mathcal{L}}_n^\pm$ decomposes, unless we are working in characteristic 2, into two components $\overline{\mathcal{L}}_n^\pm = \overline{\mathcal{L}}_{n,+}^\pm \cup \overline{\mathcal{L}}_{n,-}^\pm$, where the component $\overline{\mathcal{L}}_{n,+}^\pm$ parametrises isomorphism classes of stable degree- $2n$ -pointed chains with involution (C, I, s_-, s_+, S) such that the degree of S in each of the fixed points under the involution is even. We first consider this main component $\overline{\mathcal{L}}_{n,+}^\pm$.

The component $\overline{\mathcal{L}}_{n,+}^\pm$ is related to the moduli space $\overline{L}_n^\pm \cong X(C_n)$ of $2n$ -pointed chains with involution defined in [3, Section 6]. There is a morphism $\overline{L}_n^\pm \rightarrow \overline{\mathcal{L}}_{n,+}^\pm$ forgetting the labels of the sections. This morphism is equivariant with respect to the natural action of the Weyl group $W(C_n) = (\mathbb{Z}/2\mathbb{Z})^n \rtimes S_n$ on \overline{L}_n^\pm , the coarse moduli space of $\overline{\mathcal{L}}_{n,+}^\pm$ is $\overline{L}_n^\pm / W(C_n)$. Similar as in proposition 1.5 one can show that the morphism $\overline{L}_n^\pm \rightarrow \overline{\mathcal{L}}_{n,+}^\pm$ is faithfully flat and finite of degree $|W(C_n)| = 2^n n!$.

These morphisms together with the morphisms $\overline{L}_{2n} \rightarrow \overline{\mathcal{L}}_{2n}$ (see section 1) and $\overline{L}_n^\pm \rightarrow \overline{\mathcal{L}}_{2n}$ (see [3, Rem. 6.16]) form a commutative diagram

$$\begin{array}{ccc} \overline{L}_n^\pm & \longrightarrow & \overline{L}_{2n} \\ \downarrow & & \downarrow \\ \overline{\mathcal{L}}_{n,+}^\pm & \longrightarrow & \overline{\mathcal{L}}_{2n} \end{array}$$

where \overline{L}_n^\pm is a component of the fibred product.

The stack $\overline{\mathcal{L}}_{n,+}^\pm$ compactifies the space of finite subschemes of degree $2n$ in $\mathbb{P}^1 \setminus \{0, \infty\}$ which are invariant under the involution and of even degree in each of the fixed points of the involution. Equivalently, this is the space of polynomials $\sum_{i=0}^{2n} a'_i y^i$ of degree $2n$ with the symmetry $a'_{2n-i} = a'_i$ in the coefficients, up to change of the variable by multiplication by -1 . These

polynomials can contain $y - 1$ and $y + 1$ only with even multiplicity. After dividing by the coefficient $a'_{2n} = a'_0$, we have a polynomial of the form

$$y^{-n} + a_{n-1}y^{-n+1} + \dots + a_1y^{-1} + a_0 + a_1y + \dots + a_{n-1}y^{n-1} + y^n$$

determined by the isomorphism class up to multiplication of y with -1 (together with multiplication of the whole expression by $(-1)^n$).

In general, embedding a chain (C, I, s_-, s_+, S) into the projective space $\mathbb{P}^{2n} = \mathbb{P}(H^0(C, \mathcal{O}_C(S)))$, the image of C is given by equations arising from the 2×2 minors of a matrix of the form (decompose into several matrices if some of the b_i are zero, cf. remark 3.3; symbol $\sqrt{b_0}$ introduced for symmetry reasons)

$$\begin{pmatrix} \cdots & y_{-2} & y_{-1} & \sqrt{b_0}y_0 & \sqrt{b_0}b_1y_1 & \cdots \\ \cdots & \sqrt{b_0}b_1y_{-1} & \sqrt{b_0}y_0 & y_1 & y_2 & \cdots \end{pmatrix}$$

where $y_{-n}, \dots, y_0, \dots, y_n$ is a basis of $H^0(C, \mathcal{O}_C(S))$ defined similar as in proposition 3.2, 3.4 and such that the involution maps $y_{-i} \leftrightarrow y_i$. The sections s_-, s_+ become the sections $(1 : 0 : \dots : 0), (0 : \dots : 0 : 1)$ and the subscheme $S \subset C \subset \mathbb{P}^{2n}$ is determined by an equation

$$y_{-n} + a_{n-1}y_{-(n-1)} + \dots + a_1y_{-1} + a_0y_0 + a_1y_1 + \dots + a_{n-1}y_{n-1} + y_n = 0.$$

For an algebraically closed field K a K -valued point of $\overline{\mathcal{L}}_{n,+}^\pm$ corresponds to a collection $(a_{n-1}, \dots, a_0, b_{n-1}, \dots, b_0) \in K^{2n}$ up to the equivalence

$$(a_{n-1}, \dots, a_0, b_{n-1}, \dots, b_0) \sim (\kappa_{n-1}a_{n-1}, \dots, \kappa_0a_0, \lambda_{n-1}b_{n-1}, \dots, \lambda_0b_0)$$

with $(\kappa_{n-1}, \dots, \kappa_0, \lambda_{n-1}, \dots, \lambda_0) \in (K^*)^{2n}$ satisfying $\lambda_{n-1} = \kappa_{n-1}^2/\kappa_{n-2}$, $\lambda_{n-2} = \kappa_{n-2}^2/(\kappa_{n-3}\kappa_{n-1})$, \dots , $\lambda_1 = \kappa_1^2/(\kappa_0\kappa_2)$, $\lambda_0 = \kappa_0^2/\kappa_1^2$. This gives rise to a toric orbifold whose exact sequence of tori

$$1 \longrightarrow G \cong (\mathbb{G}_m)^n \longrightarrow (\mathbb{G}_m)^{2n} \longrightarrow T_N \cong (\mathbb{G}_m)^n \longrightarrow 1$$

corresponds to the exact sequence of lattices

$$0 \longrightarrow M \cong \mathbb{Z}^n \xrightarrow{\begin{pmatrix} -C \\ I_n \end{pmatrix}} \mathbb{Z}^{2n} \xrightarrow{\begin{pmatrix} I_n & C \end{pmatrix}} \mathbb{Z}^n \longrightarrow 0$$

where $C = C(C_n)^\top$ is the transpose of the Cartan matrix

$$C(C_n) = \begin{pmatrix} 2 & -1 & 0 & \cdots & 0 \\ -1 & 2 & \ddots & \ddots & \vdots \\ 0 & \ddots & \ddots & -1 & 0 \\ \vdots & \ddots & -1 & 2 & -1 \\ 0 & \cdots & 0 & -2 & 2 \end{pmatrix}$$

of the root system C_n .

DEFINITION 5.2. — We define the toric orbifold $\mathcal{Y}(C_n)$ associated to the Cartan matrix of the root system C_n in terms of the stacky fan $\Upsilon(C_n) = (N, \Upsilon(C_n), \beta)$, where $N = \mathbb{Z}^n$ and the linear map $\beta: \mathbb{Z}^{2n} \rightarrow N$ is given by the $n \times 2n$ matrix $(-C(C_n) \ I_n)$. The fan $\Upsilon(C_n)$ has the $2n$ one-dimensional cones $\varrho_{n-1}, \dots, \varrho_0, \tau_{n-1}, \dots, \tau_0$ generated by the columns of the matrix $(-C(C_n) \ I_n)$. A subset of one-dimensional cones generates a higher dimensional cone of $\Upsilon(C_n)$ if it does not contain one of the sets $\{\varrho_0, \tau_0\}, \dots, \{\varrho_{n-1}, \tau_{n-1}\}$. This defines a fan containing 2^n n -dimensional cones σ_I generated by sets $\{\varrho_i\}_{i \notin I} \cup \{\tau_i\}_{i \in I}$ for subsets $I \subseteq \{0, \dots, n-1\}$.

The functor of $\Upsilon(C_n)$ -collections $\mathcal{C}_{\Upsilon(C_n)} \cong \mathcal{Y}(C_n)$ has objects of the form $((\mathcal{L}_{\varrho_i}, a_i)_{i=0, \dots, n-1}, (\mathcal{L}_{\tau_i}, b_i)_{i=0, \dots, n-1}, (c_i)_{i=0, \dots, n-1})$ over a scheme Y , where the c_i are isomorphisms of line bundles on Y

$$\begin{aligned} c_{n-1} &: \mathcal{L}_{\tau_{n-1}} \otimes \mathcal{L}_{\varrho_{n-1}}^{\otimes -2} \otimes \mathcal{L}_{\varrho_{n-2}} \rightarrow \mathcal{O}_Y, \\ c_{n-2} &: \mathcal{L}_{\tau_{n-2}} \otimes \mathcal{L}_{\varrho_{n-1}} \otimes \mathcal{L}_{\varrho_{n-2}}^{\otimes -2} \otimes \mathcal{L}_{\varrho_{n-3}} \rightarrow \mathcal{O}_Y, \\ &\vdots \\ c_1 &: \mathcal{L}_{\tau_1} \otimes \mathcal{L}_{\varrho_2} \otimes \mathcal{L}_{\varrho_1}^{\otimes -2} \otimes \mathcal{L}_{\varrho_0} \rightarrow \mathcal{O}_Y, \\ c_0 &: \mathcal{L}_{\tau_0} \otimes \mathcal{L}_{\varrho_1}^{\otimes 2} \otimes \mathcal{L}_{\varrho_0}^{\otimes -2} \rightarrow \mathcal{O}_Y. \end{aligned}$$

We have a morphism of stacks $\mathcal{C}_{\Upsilon(C_n)} \rightarrow \mathcal{C}_{\Upsilon(A_{2n-1})}$ by considering the collection

$$\begin{aligned} &((\mathcal{L}_{\varrho_{n-1}}, a_{n-1}), \dots, (\mathcal{L}_{\varrho_0}, a_0), \dots, (\mathcal{L}_{\varrho_{n-1}}, a_{n-1}), (\mathcal{L}_{\tau_{n-1}}, b_{n-1}), \\ &\dots, (\mathcal{L}_{\tau_0}, b_0), \dots, (\mathcal{L}_{\tau_{n-1}}, b_{n-1}), c_{n-1}, \dots, c_0, \dots, c_{n-1}), \end{aligned} \tag{5.1}$$

built out of a $\Upsilon(C_n)$ -collection, as a $\Upsilon(A_{2n-1})$ -collection. This morphism can be described by the map of fans $\Upsilon(C_n) \rightarrow \Upsilon(A_{2n-1})$ mapping $e'_{n-1} \mapsto e_{2n-1} + e_1, \dots, e'_1 \mapsto e_{n+1} + e_{n-1}, e'_0 \mapsto e_n$, where e'_{n-1}, \dots, e'_0 are the generators of $\tau_{n-1}, \dots, \tau_0$ of $\Upsilon(C_n)$ and e_{2n-1}, \dots, e_1 are those of $\tau_{2n-1}, \dots, \tau_1$ of $\Upsilon(A_{2n-1})$. It corresponds to a toric morphism $\mathcal{Y}(C_n) \rightarrow \mathcal{Y}(A_{2n-1})$ making $\mathcal{Y}(C_n)$ a subcategory of $\mathcal{Y}(A_{2n-1})$.

THEOREM 5.3. — There is an isomorphism of stacks $\overline{\mathcal{L}}_{n,+}^{\pm} \cong \mathcal{Y}(C_n)$.

Proof. — Applying construction 3.5 to a degree- $2n$ -pointed chain with involution, it is possible to choose y_0, \dots, y_{2n} such that the involution maps $y_i \leftrightarrow y_{2n-i}$, and we obtain a $\Upsilon(A_{2n-1})$ -collection of the form (5.1). Applying construction 3.6 to a $\Upsilon(A_{2n-1})$ -collection of the form (5.1), making symmetric choices, we can introduce an involution on the resulting degree- $2n$ -pointed chain by $y_i \leftrightarrow y_{2n-i}$. A $\Upsilon(A_{2n-1})$ -collection of the form (5.1)

is equivalent to a $\mathfrak{Y}(C_n)$ -collection, and further, morphisms of the corresponding degree- $2n$ -chains with involution that commute with the involution are equivalent to morphisms of $\mathfrak{Y}(C_n)$ -collections. \square

The case of the other component $\overline{\mathcal{L}}_{n,-}^\pm$ is very similar. The stack $\overline{\mathcal{L}}_{n,-}^\pm$ parametrises isomorphism classes of stable degree- $2n$ -pointed chains with involution (C, I, s_-, s_+, S) such that the degree of S in each of the fixed points of the involution is odd if there are two fixed points, and positive if there is only one fixed point. It is related to the moduli stack $\mathcal{X}(C_{n-1})$ defined in [3, Section 6]: there is a morphism $\mathcal{X}(C_{n-1}) \rightarrow \overline{\mathcal{L}}_{n,-}^\pm$ determined by forgetting the labels of the sections and adding the fixed point subscheme of the involution as a subscheme of degree 2 to the $2n - 2$ sections.

The stack $\overline{\mathcal{L}}_{n,-}^\pm$ compactifies the stack of finite subschemes S of degree $2n$ in $\mathbb{P}^1 \setminus \{0, \infty\}$ invariant under the involution such that S has odd degree in $(1:1)$ and $(1:-1)$ (positive degree in $(1:1) = (1:-1)$ in characteristic 2). Equivalently we may consider polynomials $\sum_{i=0}^{2n} a'_i y^i$ of degree $2n$ with the symmetry $a'_{2n-i} = -a'_i$ in the coefficients and $a'_n = 0$. We may represent each isomorphism class by an expression of the form

$$-y^{-n} - a_{n-1}y^{-n+1} - \dots - a_1y^{-1} + 0 + a_1y + \dots + a_{n-1}y^{n-1} + y^n$$

determined up to multiplication of y with -1 (together with multiplication of the whole expression by $(-1)^n$). It has a factor $(y - 1)(y^{-1} + 1)$, occurring with odd multiplicity in characteristic $\neq 2$.

In general, similar as in the case of $\overline{\mathcal{L}}_{n,+}^\pm$, a not necessarily irreducible chain can be naturally embedded into $\mathbb{P}^{2n} = \mathbb{P}(H^0(C, \mathcal{O}_C(S)))$ and described by equations arising from the 2×2 minors of a matrix of the form

$$\begin{pmatrix} \cdots & y_{-2} & y_{-1} & y_0 & b_1y_1 & \cdots \\ \cdots & b_1y_{-1} & y_0 & y_1 & y_2 & \cdots \end{pmatrix}$$

and the subscheme $S \subset C \subset \mathbb{P}^{2n}$ is given by an equation

$$-y_{-n} - a_{n-1}y_{-(n-1)} - \dots - a_1y_{-1} + a_1y_1 + \dots + a_{n-1}y_{n-1} + y_n = 0.$$

Over an algebraically closed field K a K -valued point of $\overline{\mathcal{L}}_{n,-}^\pm$ corresponds to a collection $(a_{n-1}, \dots, a_1, b_{n-1}, \dots, b_1) \in K^{2n-2}$ up to the equivalence

$$(a_{n-1}, \dots, a_1, b_{n-1}, \dots, b_1) \sim (\kappa_{n-1}a_{n-1}, \dots, \kappa_1a_1, \lambda_{n-1}b_{n-1}, \dots, \lambda_1b_1)$$

with $(\kappa_{n-1}, \dots, \kappa_1, \lambda_{n-1}, \dots, \lambda_1) \in (K^*)^{2n-2}$ satisfying $\lambda_{n-1} = \kappa_{n-1}^2 / \kappa_{n-2}$, $\lambda_{n-2} = \kappa_{n-2}^2 / (\kappa_{n-3}\kappa_{n-1})$, \dots , $\lambda_2 = \kappa_2^2 / (\kappa_1\kappa_3)$, $\lambda_1^2 = \kappa_1^2 / \kappa_2^2$.

We will see that this stack can be described by a toric stack that differs from $\mathcal{Y}(C_{n-1})$ by replacing the matrix $(-C(C_{n-1}) \ I_{n-1})$ defining the map β of the stacky fan $\mathbf{Y}(C_{n-1})$ by the matrix

$$\begin{pmatrix} & & & 1 & 0 & & \\ & & & 0 & \ddots & \ddots & \\ & -C(C_{n-1}) & & & \ddots & & \\ & & & & & 1 & 0 \\ & & & & & 0 & 2 \end{pmatrix}.$$

In the case $n = 1$ we define it to be $B\mu_2$. This toric stack corresponds to the category of collections of the form $((\mathcal{L}_{\varrho_i}, a_i)_{i=1, \dots, n-1}, (\mathcal{L}_{\tau_i}, b_i)_{i=1, \dots, n-1}, (c_i)_{i=1, \dots, n-1})$ over a scheme Y , where the c_i are isomorphisms of line bundles

$$\begin{aligned} c_{n-1} &: \mathcal{L}_{\tau_{n-1}} \otimes \mathcal{L}_{\varrho_{n-1}}^{\otimes -2} \otimes \mathcal{L}_{\varrho_{n-2}} \rightarrow \mathcal{O}_Y, \\ c_{n-2} &: \mathcal{L}_{\tau_{n-2}} \otimes \mathcal{L}_{\varrho_{n-1}} \otimes \mathcal{L}_{\varrho_{n-2}}^{\otimes -2} \otimes \mathcal{L}_{\varrho_{n-3}} \rightarrow \mathcal{O}_Y, \\ &\vdots \\ c_2 &: \mathcal{L}_{\tau_2} \otimes \mathcal{L}_{\varrho_3} \otimes \mathcal{L}_{\varrho_2}^{\otimes -2} \otimes \mathcal{L}_{\varrho_1} \rightarrow \mathcal{O}_Y, \\ c_1 &: \mathcal{L}_{\tau_1}^{\otimes 2} \otimes \mathcal{L}_{\varrho_2}^{\otimes 2} \otimes \mathcal{L}_{\varrho_1}^{\otimes -2} \rightarrow \mathcal{O}_Y. \end{aligned}$$

PROPOSITION 5.4. — *The stack $\overline{\mathcal{L}}_{n,-}^{\pm}$ is isomorphic to the above toric stack. It can be embedded into $\overline{\mathcal{L}}_{n,+}^{\pm}$ as the divisor D_{ϱ_0} corresponding to the cone ϱ_0 and defined by $a_0 = 0$.*

Proof. — Similar as in the proof of theorem 5.3 one can show that $\overline{\mathcal{L}}_{n,-}^{\pm}$ is isomorphic to the above toric stack by embedding this toric stack as a subcategory into $\mathcal{Y}(A_{2n-1})$.

An embedding of $\overline{\mathcal{L}}_{n,-}^{\pm}$ into $\overline{\mathcal{L}}_{n,+}^{\pm}$ as divisor D_{ϱ_0} is given by mapping a collection $((\mathcal{L}_{\varrho_i}, a_i)_{i=1, \dots, n-1}, (\mathcal{L}_{\tau_i}, b_i)_{i=1, \dots, n-1}, (c_i)_{i=1, \dots, n-1})$ over a scheme Y to the collection $((\mathcal{L}_{\varrho_i}, a_i)_{i=0, \dots, n-1}, (\mathcal{L}_{\tau_i}, b_i)_{i=0, \dots, n-1}, (c_i)_{i=0, \dots, n-1})$ where $\mathcal{L}_{\varrho_0} = \mathcal{L}_{\tau_1} \otimes \mathcal{L}_{\varrho_2}$, $a_0 = 0$ and $\mathcal{L}_{\tau_0} = \mathcal{O}_Y$, $b_0 = 1$ and c_0 is defined using c_1 . This corresponds to mapping $(C \rightarrow Y, I, s_-, s_+, S_-)$ to $(C \rightarrow Y, I, s_-, s_+, S_+)$ such that S_+ is given by

$$y_n + a_{n-1}y_{-(n-1)} + \dots + a_1y_{-1} + a_1y_1 + \dots + a_{n-1}y_{n-1} + y_n$$

if S_- is given by


$$-y_n - a_{n-1}y_{-(n-1)} - \dots - a_1y_{-1} + a_1y_1 + \dots + a_{n-1}y_{n-1} + y_n.$$

□


Example 5.5. — The toric orbifold $\overline{\mathcal{L}}_{1,+}^{\pm}$ is isomorphic to the weighted projective line $\mathbb{P}(1, 2) \cong \mathcal{Y}(C_1)$. Here the inclusion as subcategory $\overline{\mathcal{L}}_{1,+}^{\pm} \rightarrow$

$\overline{\mathcal{L}}_2$ is an isomorphism of stacks as any degree-2-pointed chain is isomorphic to a symmetric object under an involution whose isomorphisms commute with the involution. So we have the same situation as in examples 1.8, 2.2, 3.8. The component $\overline{\mathcal{L}}_{1,-}^\pm$ is isomorphic to $B\mu_2$.


Example 5.6. — The stacky fan of the toric orbifold $\overline{\mathcal{L}}_{2,+}^\pm \cong \mathcal{Y}(C_2)$ is given by the matrix $\begin{pmatrix} -2 & 1 & 1 & 0 \\ 2 & -2 & 0 & 1 \end{pmatrix}$:



We picture the types of pointed chains over the torus invariant divisors of the moduli stack $\overline{\mathcal{L}}_{2,+}^\pm$.



The toric orbifold $\overline{\mathcal{L}}_{2,-}^{\pm} \cong \mathbb{P}^1/\mu_2$ corresponds to the stacky fan given by the matrix $(-2 \ 2)$. We have the following types of pointed chains over $\overline{\mathcal{L}}_{2,-}^{\pm}$:



One also may consider chains with involution and a subscheme of odd degree.

DEFINITION 5.7. — Let the fibred category $\overline{\mathcal{L}}_n^{0,\pm}$ of stable degree- $(2n + 1)$ -pointed chains of \mathbb{P}^1 with involution be defined analogously to definition 5.1.

The fibred category $\overline{\mathcal{L}}_n^{0,\pm}$ is a stack in the fpqc topology with representable finite diagonal.

The moduli stack $\overline{\mathcal{L}}_n^{0,\pm}$ forms a subcategory of $\overline{\mathcal{L}}_{2n+1}$. It is related to the moduli space $\overline{L}_n^{0,\pm} \cong X(B_n)$ of $(2n + 1)$ -pointed chains with involution defined in [3, Section 1]. We have a morphism $\overline{L}_n^{0,\pm} \rightarrow \overline{\mathcal{L}}_n^{0,\pm}$ forgetting the labels of the sections, which is equivariant with respect to the action of the Weyl group $W(B_n) = (\mathbb{Z}/2\mathbb{Z})^n \rtimes S_n$ on $\overline{L}_n^{0,\pm}$. The coarse moduli space of $\overline{\mathcal{L}}_n^{0,\pm}$ is $\overline{L}_n^{0,\pm}/W(B_n)$. As in the C -case the morphism $\overline{L}_n^{0,\pm} \rightarrow \overline{\mathcal{L}}_n^{0,\pm}$ is faithfully flat and finite of degree $|W(B_n)| = 2^n n!$, and we have a commutative diagram

$$\begin{array}{ccc} \overline{L}_n^{0,\pm} & \longrightarrow & \overline{L}_{2n+1} \\ \downarrow & & \downarrow \\ \overline{\mathcal{L}}_n^{0,\pm} & \longrightarrow & \overline{\mathcal{L}}_{2n+1} \end{array}$$

Embedding a degree- $(2n + 1)$ -pointed chain with involution (C, I, s_-, s_+, S) into the projective space $\mathbb{P}^{2n+1} = \mathbb{P}(H^0(C, \mathcal{O}_C(S)))$, the image of C is given by equations arising from the 2×2 minors of a matrix of the form

$$\begin{pmatrix} \cdots & y_{-5/2} & y_{-3/2} & y_{-1/2} & b_1 y_{1/2} & b_1 b_2 y_{3/2} & \cdots \\ \cdots & b_1 b_2 y_{-3/2} & b_1 y_{-1/2} & y_{1/2} & y_{3/2} & y_{5/2} & \cdots \end{pmatrix}$$

and the subscheme S by

$$y_{-(2n+1)/2} + a_n y_{-(2n-1)/2} + \cdots + a_1 y_{-1/2} + a_1 y_{1/2} + \cdots + a_n y_{(2n-1)/2} + y_{(2n+1)/2}.$$

where $y_{-(2n+1)/2}, \dots, y_{-3/2}, y_{-1/2}, y_{1/2}, y_{3/2}, \dots, y_{(2n+1)/2}$ is a basis of $H^0(C, \mathcal{O}_C(S))$ defined similar as in proposition 3.2, 3.4 and such that the involution maps $y_{-i/2} \leftrightarrow y_{i/2}$.

DEFINITION 5.8. — We define the toric orbifold $\mathcal{Y}(B_n)$ in terms of the stacky fan $\Upsilon(B_n)$ as in definition 5.2 replacing the Cartan matrix $C(C_n)$ of the root system C_n by the Cartan matrix $C(B_n)$ of the root system B_n .

It turns out that $\overline{\mathcal{L}}_n^{0,\pm}$ is not quite $\mathcal{Y}(B_n)$, but coincides with the underlying canonical toric stack $\mathcal{Y}(B_n)^{\text{can}}$ (as defined in [7]). So instead of the Cartan matrix of the root system B_n we have the matrix

$$\begin{pmatrix} 2 & -1 & 0 & \cdots & \cdots & 0 \\ -1 & 2 & \ddots & \ddots & & \vdots \\ 0 & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 2 & -1 & 0 \\ \vdots & & \ddots & -1 & 2 & -1 \\ 0 & \cdots & \cdots & 0 & -1 & 1 \end{pmatrix}$$

where the rightmost column is half of the column of the Cartan matrix. The functor of $\Upsilon(B_n)^{\text{can}}$ -collections $\mathcal{C}_{\Upsilon(B_n)^{\text{can}}} \cong \mathcal{Y}(B_n)^{\text{can}}$ has objects of the form $((\mathcal{L}_{\varrho_i}, a_i)_{i=1, \dots, n}, (\mathcal{L}_{\tau_i}, b_i)_{i=1, \dots, n}, (c_i)_{i=1, \dots, n})$ over a scheme Y , where the c_i are isomorphisms of line bundles

$$c_n: \mathcal{L}_{\tau_n} \otimes \mathcal{L}_{\varrho_n}^{\otimes -2} \otimes \mathcal{L}_{\varrho_{n-1}} \rightarrow \mathcal{O}_Y, c_{n-1}: \mathcal{L}_{\tau_{n-1}} \otimes \mathcal{L}_{\varrho_n} \otimes \mathcal{L}_{\varrho_{n-1}}^{\otimes -2} \otimes \mathcal{L}_{\varrho_{n-2}} \rightarrow \mathcal{O}_Y, \dots, c_2: \mathcal{L}_{\tau_2} \otimes \mathcal{L}_{\varrho_3} \otimes \mathcal{L}_{\varrho_2}^{\otimes -2} \otimes \mathcal{L}_{\varrho_1} \rightarrow \mathcal{O}_Y, c_1: \mathcal{L}_{\tau_1} \otimes \mathcal{L}_{\varrho_2} \otimes \mathcal{L}_{\varrho_1}^{\otimes -1} \rightarrow \mathcal{O}_Y.$$

The inclusion as subcategory $\mathcal{Y}(B_n)^{\text{can}} \rightarrow \mathcal{Y}(A_{2n})$ can be described as $\mathcal{C}_{\Upsilon(B_n)^{\text{can}}} \rightarrow \mathcal{C}_{\Upsilon(A_{2n})}$ by considering the collection

$$((\mathcal{L}_{\varrho_n}, a_n), \dots, (\mathcal{L}_{\varrho_1}, a_1), (\mathcal{L}_{\varrho_1}, a_1), \dots, (\mathcal{L}_{\varrho_n}, a_n), (\mathcal{L}_{\tau_n}, b_n), \dots, (\mathcal{L}_{\tau_1}, b_1), (\mathcal{L}_{\tau_1}, b_1), \dots, (\mathcal{L}_{\tau_n}, b_n), c_n, \dots, c_1, c_1, \dots, c_n),$$

formed out of a $\Upsilon(B_n)^{\text{can}}$ -collection, as a $\Upsilon(A_{2n})$ -collection.


As in the case of degree- $2n$ -pointed chains with involution one can prove:

THEOREM 5.9. — There is an isomorphism of stacks $\overline{\mathcal{L}}_n^{0,\pm} \cong \mathcal{Y}(B_n)^{\text{can}}$.


Example 5.10. — In the case $n = 1$ we have a scheme $\overline{\mathcal{L}}_1^{0,\pm} \cong \mathcal{Y}(B_1)^{\text{can}}$ isomorphic to \mathbb{P}^1 .

Example 5.11. — The toric orbifold $\overline{\mathcal{L}}_2^{0,\pm} \cong \mathcal{Y}(B_2)^{\text{can}}$ is given by the matrix $\begin{pmatrix} -2 & 1 & 1 & 0 \\ 1 & -1 & 0 & 1 \end{pmatrix}$. In the picture of the stacky fan $\Upsilon(B_2)^{\text{can}}$ the

dotted arrow corresponds to the generator of the ray ϱ_1 determined by the stacky fan $\Upsilon(B_2)$. $\overline{\mathcal{L}}_{2,-}^\pm$:



We picture the types of pointed chains over the torus invariant divisors of the moduli stack $\overline{\mathcal{L}}_2^{0,\pm}$. $\overline{\mathcal{L}}_{2,-}^\pm$:



BIBLIOGRAPHY

- [1] D. ABRAMOVICH, M. OLSSON & A. VISTOLI, “Tame stacks in positive characteristic”, *Ann. Inst. Fourier* **58** (2008), p. 1057-1091, arXiv:math/0703310.
- [2] V. BATYREV & M. BLUME, “The functor of toric varieties associated with Weyl chambers and Losev-Manin moduli spaces”, *Tohoku Math. J.* **63** (2011), p. 581-604, arXiv:0911.3607.
- [3] ———, “On generalisations of Losev-Manin moduli spaces for classical root systems”, *Pure and Applied Mathematics Quarterly* **7** (2011), (Special Issue: In memory of Eckart Viehweg), p. 1053-1084, arXiv:0912.2898.
- [4] L. BORISOV, L. CHEN & G. SMITH, “The orbifold Chow ring of toric Deligne-Mumford stacks”, *J. Amer. Math. Soc.* **18** (2005), p. 193-213, arXiv:math/0309229.
- [5] D. COX, “The functor of a smooth toric variety”, *Tohoku Math. J.* **47** (1995), p. 251-262, arXiv:alg-geom/9312001.

- [6] ———, “The homogeneous coordinate ring of a toric variety”, *J. Algebraic Geom.* **4** (1995), p. 17-50, arXiv:alg-geom/9210008.
- [7] B. FANTECHI, E. MANN & F. NIRONI, “Smooth toric Deligne-Mumford stacks”, *J. reine angew. Math.* **648** (2010), p. 201-244, arXiv:0708.1254.
- [8] J. GIRAUD, *Cohomologie non abélienne*, Grundlehren math. Wiss., vol. 179, Springer-Verlag, Berlin - Heidelberg - New York, 1971.
- [9] A. GROTHENDIECK & J. DIEUDONNÉ, *Éléments de Géométrie Algébrique*, vol. 4,8,11,17,20,24,28,32, Publ. Math. IHES, 1960-1967.
- [10] A. GROTHENDIECK et al., *Séminaire de géométrie algébrique, Théorie des topos et cohomologie étale des schémas, Tome 3*, Lecture Notes in Mathematics, vol. 305, Springer-Verlag, Berlin - Heidelberg - New York, 1973.
- [11] I. IWANARI, “Integral Chow rings of toric stacks”, arXiv:0705.3524.
- [12] M. KAPRANOV, “Chow quotients of Grassmannians I”, *Adv. Soviet Math.* **16** (1993), p. 29-110, arXiv:alg-geom/9210002.
- [13] S. KLEIMAN, “The Picard scheme”, in *Fundamental Algebraic Geometry*, Math. Surveys Monogr., vol. 123, Amer. Math. Soc., Providence, RI, 2005, p. 237-321, arXiv:math/0504020.
- [14] F. KNUDSEN, “The projectivity of the moduli space of stable curves II: The stacks $M_{g,n}$ ”, *Math. Scand.* **52** (1983), p. 161-199.
- [15] G. LAUMON & L. MORET-BAILLY, *Champs algébriques*, Springer-Verlag, Berlin - Heidelberg - New York, 2000.
- [16] A. LOSEV & Y. MANIN, “New Moduli Spaces of Pointed Curves and Pencils of Flat Connections”, *Michigan Math. J.* **48** (2000), p. 443-472, arXiv:math/0001003.
- [17] F. PERRONI, “A note on toric Deligne-Mumford stacks”, *Tohoku Math. J.* **60** (2008), p. 441-458, arXiv:0705.3823.
- [18] A. VISTOLI, “Grothendieck topologies, fibered categories and descent theory”, in *Fundamental Algebraic Geometry*, Math. Surveys Monogr., vol. 123, Amer. Math. Soc., Providence, RI, 2005, p. 1-104, arXiv:math/0412512.

Manuscrit reçu le 3 juillet 2012,
accepté le 14 décembre 2012.

Mark BLUME
Mathematisches Institut,
Universität Münster,
Einsteinstrasse 62,
48149 Münster,
Germany
mark.blume@uni-muenster.de