



ANNALES

DE

L'INSTITUT FOURIER

Amaury FRESLON

Permanence of approximation properties for discrete quantum groups

Tome 65, n° 4 (2015), p. 1437-1467.

http://aif.cedram.org/item?id=AIF_2015__65_4_1437_0



© Association des Annales de l'institut Fourier, 2015,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

PERMANENCE OF APPROXIMATION PROPERTIES FOR DISCRETE QUANTUM GROUPS

by Amaury FRESLON

ABSTRACT. — We prove several results on the permanence of weak amenability and the Haagerup property for discrete quantum groups. In particular, we improve known facts on free products by allowing amalgamation over a finite quantum subgroup. We also define a notion of relative amenability for discrete quantum groups and link it with amenable equivalence of von Neumann algebras, giving additional permanence properties.

RÉSUMÉ. — Nous prouvons plusieurs résultats concernant la permanence de la moyennabilité faible et de la propriété de Haagerup pour les groupes quantiques discrets. En particulier, nous améliorons des résultats connus sur les produits libres en autorisant l'amalgamation sur un sous-groupe quantique fini. Nous définissons également une notion de moyennabilité relative pour les groupes quantiques discrets et nous la relierons à l'équivalence moyennable d'algèbres de von Neumann, ce qui donne de nouvelles propriétés de permanence.

1. Introduction

A fruitful way of studying compact quantum groups is through their dual discrete quantum group. Geometric group theory is then a rich source of inspiration, even though results can seldom be straightforwardly transferred from the classical to the quantum setting. A very important property from this point of view is of course *amenability*, the study of which culminated in R. Tomatsu's work [34]. However, some important examples of discrete quantum groups, often called "free quantum groups", are known not to be amenable by [4, Cor 5]. One should then look for weak versions of amenability like the *Haagerup property* or *weak amenability*. Some of these

Keywords: Quantum groups, approximation properties, relative amenability.

Math. classification: 20G42, 46L65.

approximation properties were investigated for unimodular discrete quantum groups by J. Kraus and Z.-J. Ruan in [27], but no genuinely quantum example was found.

There recently was a regain of interest in the subject thanks to M. Brannan's breakthrough paper [10] proving the Haagerup property for the free orthogonal and free unitary quantum groups O_N^+ and U_N^+ . On the one hand, his techniques were extended to free wreath products by F. Lemeux in [28] and to more general classes of unitary easy quantum groups by the author in [24]. On the other hand, weak amenability was also established for O_N^+ and U_N^+ by the author in [22] and later extended, with K. De Commer and M. Yamashita in [16] using monoidal equivalence.

At that point, it seemed reasonable to try to establish a general theory of approximation properties for discrete quantum groups. For the Haagerup property, this was done by M. Daws, P. Fima, A. Skalski and S. White in [14] using the more general setting of locally compact quantum groups. For weak amenability, this was part of the author's PhD thesis [23]. In particular, we gave in the latter several permanence properties, i.e. group-theoretic constructions preserving weak amenability. One of the most important is the following one : if two discrete quantum groups are weakly amenable with Cowling-Haagerup constant equal to 1, then their free product is again weakly amenable with Cowling-Haagerup constant equal to 1, which was proved in [21].

However, there are several other natural constructions to look at like direct products there or inductive limits. Moreover, the techniques of E. Ricard and X. Qu [31], which are crucial in the proof of the aforementioned result on free products, also work for amalgamated free products under certain assumptions. For discrete groups, it is not very difficult to see that these assumptions are satisfied as soon as the amalgam is finite. That this also works for quantum groups will be proved in Section 3. Note that even if it was certainly known to experts, the result for classical groups did not appear yet in the literature. Let us also mention that our proofs are based on multiplier techniques, so that they can also be applied to the Haagerup property, improving [14, Prop 7.13] by removing the unmodularity assumption.

The last permanence property which we will study is *relative amenability*. In fact, this was the first to be used in combination with weak amenability by M. Cowling and U. Haagerup in [13, Thm 6.4] to prove that lattices in different symplectic groups yield non-isomorphic von Neumann algebras. Their proof relied on [13, Prop 6.2] :

THEOREM (Cowling-Haagerup). — *Let G be a locally compact group and let Γ be a lattice in G . Then, $\Lambda_{cb}(\Gamma) = \Lambda_{cb}(G)$, where Λ_{cb} denotes the Cowling-Haagerup constant.*

The core of the proof is the finite covolume assumption, which is however too restrictive in our context. In [17], P. Eymard defined a subgroup $H \subset G$ to be *co-amenable* if there exists a mean on the homogeneous space G/H (i.e. a state on $\ell^\infty(G/H)$) which is invariant with respect to the translation action of G (so lattices in locally compact groups are in particular co-amenable). He investigated this property as a weakening of the notion of amenability since a group is amenable if and only if all its subgroups are co-amenable. It was proved in [2, Paragraphe 4.10] that if H is co-amenable in G , then $\Lambda_{cb}(H) = \Lambda_{cb}(G)$. Since this proof makes use of the machinery of von Neumann algebras and correspondences, it may be used as a starting point for a generalization to quantum groups, which will be done in Section 4.

Let us briefly outline the content of this work. In Section 2, we introduce the necessary material concerning quantum groups and their actions as well as approximation properties. In Section 3 we give several permanence results. The main one is the stability of weak amenability under free products amalgamated over finite quantum subgroups, if the Cowling-Haagerup constant is equal to 1. Finally, Section 4 deals with relative amenability. We define it and prove that it is equivalent to amenable equivalence of the associated von Neumann algebras. We then study the simplest examples, namely finite index quantum subgroups and eventually prove permanence properties for relatively amenable discrete quantum groups.

Acknowledgments

The results of this work were part of the author's PhD thesis and he wishes to thank his advisor E. Blanchard for his supervision. The author is also thankful to S. Vaes for pointing out a mistake in a preliminary version, to C. Anantharaman-Delaroche for discussions on amenable equivalence and to A. Skalski and the anonymous referee for their useful comments.

2. Preliminaries

In this paper, we will denote by \otimes the tensor product of Hilbert spaces and the minimal tensor product of C^* -algebras, and by $\bar{\otimes}$ the tensor product of von Neumann algebras. Once the spaces involved are clear, we will denote tensor products of elements or maps by \otimes .

2.1. Discrete quantum groups

Discrete quantum groups will be seen as duals of compact quantum groups in the sense of S.L. Woronowicz. We briefly present this theory as introduced in [43], i.e. in the C^* -algebraic setting. We will then explain how one passes to the von Neumann algebraic setting, which will prove more convenient when dealing with relative amenability (see Remark 4.15). In the sequel, all tensor products of C^* -algebras are minimal.

DEFINITION 2.1. — *A compact quantum group \mathbb{G} is a pair $(C(\mathbb{G}), \Delta)$ where $C(\mathbb{G})$ is a unital C^* -algebra and $\Delta : C(\mathbb{G}) \rightarrow C(\mathbb{G}) \otimes C(\mathbb{G})$ is a unital $*$ -homomorphism such that*

$$(\Delta \otimes \iota) \circ \Delta = (\iota \otimes \Delta) \circ \Delta$$

and both $\Delta(C(\mathbb{G}))(1 \otimes C(\mathbb{G}))$ and $\Delta(C(\mathbb{G}))(C(\mathbb{G}) \otimes 1)$ span dense subspaces of $C(\mathbb{G}) \otimes C(\mathbb{G})$.

The main feature of compact quantum groups is the existence of a *Haar state*, which is both left and right invariant (see [43, Thm 1.3]).

PROPOSITION 2.2. — *Let \mathbb{G} be a compact quantum group. Then, there exists a unique state h on \mathbb{G} , called the Haar state, such that for all $a \in C(\mathbb{G})$,*

$$(\iota \otimes h) \circ \Delta(a) = h(a).1$$

$$(h \otimes \iota) \circ \Delta(a) = h(a).1$$

Let $(L^2(\mathbb{G}), \xi_h)$ be the associated GNS construction and let $C_{\text{red}}(\mathbb{G})$ be the image of $C(\mathbb{G})$ under the GNS map, called the *reduced form* of \mathbb{G} . Let W be the unique (by [43, Thm 4.1]) unitary operator on $L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$ satisfying, for all for $\xi \in L^2(\mathbb{G})$ and $a \in C(\mathbb{G})$,

$$W^*(\xi \otimes a\xi_h) = \Delta(a)(\xi \otimes \xi_h)$$

and let $\widehat{W} := \Sigma W^* \Sigma$. Then, W is a *multiplicative unitary* in the sense of [3], i.e. $W_{12}W_{13}W_{23} = W_{23}W_{12}$ and we have the following equalities :

$$C_{\text{red}}(\mathbb{G}) = \overline{\text{span}}(\iota \otimes \mathcal{B}(L^2(\mathbb{G}))_*)(W) \text{ and } \Delta(x) = W^*(1 \otimes x)W$$

for all $x \in C_{\text{red}}(\mathbb{G})$. Moreover, we can define the *dual discrete quantum group* $\widehat{\mathbb{G}} = (C_0(\widehat{\mathbb{G}}), \widehat{\Delta})$ by

$$C_0(\widehat{\mathbb{G}}) = \overline{\text{span}}(\mathcal{B}(L^2(\mathbb{G}))_* \otimes \iota)(W) \text{ and } \widehat{\Delta}(x) = \widehat{W}^*(1 \otimes x)\widehat{W}$$

for all $x \in C_0(\mathbb{G})$. The two von Neumann algebras associated to these quantum groups are

$$L^\infty(\mathbb{G}) = C_{\text{red}}(\mathbb{G})'' \text{ and } \ell^\infty(\widehat{\mathbb{G}}) = C_0(\widehat{\mathbb{G}})''$$

where the bicommutants are taken in $\mathcal{B}(L^2(\mathbb{G}))$. The coproducts extend to normal maps on these von Neumann algebras and one can prove that $W \in L^\infty(\mathbb{G}) \overline{\otimes} \ell^\infty(\widehat{\mathbb{G}})$. Moreover, the Haar state of \mathbb{G} extends to a normal state on $L^\infty(\mathbb{G})$. In order to give an alternative description of $\ell^\infty(\widehat{\mathbb{G}})$, we need to define representations of quantum groups.

DEFINITION 2.3. — *Let \mathbb{G} be a compact quantum group.*

- *A representation of \mathbb{G} on a Hilbert space H is an invertible operator $U \in L^\infty(\mathbb{G}) \overline{\otimes} \mathcal{B}(H)$ such that $(\Delta \otimes \iota)(U) = U_{13}U_{23}$.*
- *A representation of $\widehat{\mathbb{G}}$ on a Hilbert space H is an invertible operator $U \in \ell^\infty(\widehat{\mathbb{G}}) \overline{\otimes} \mathcal{B}(H)$ such that $(\widehat{\Delta} \otimes \iota)(U) = U_{13}U_{23}$.*
- *A representation is said to be unitary if the operator U is unitary.*

The linear span of coefficients of unitary representations of \mathbb{G} forms a Hopf- $*$ -algebra $\text{Pol}(\mathbb{G})$ which is dense in $C(\mathbb{G})$. Let $\text{Irr}(\mathbb{G})$ be the set of isomorphism classes of irreducible unitary representations of \mathbb{G} (which are all finite-dimensional by [43, Thm 1.2]). If $\alpha \in \text{Irr}(\mathbb{G})$, we will denote by u^α a representative of α and by H_α the finite-dimensional Hilbert space on which it acts. There are $*$ -isomorphisms

$$C_0(\widehat{\mathbb{G}}) = \bigoplus_{\alpha \in \text{Irr}(\mathbb{G})} \mathcal{B}(H_\alpha) \text{ and } \ell^\infty(\widehat{\mathbb{G}}) = \prod_{\alpha \in \text{Irr}(\mathbb{G})} \mathcal{B}(H_\alpha).$$

The minimal central projection in $\ell^\infty(\widehat{\mathbb{G}})$ corresponding to the identity of $\mathcal{B}(H_\alpha)$ will be denoted by p_α and there exist positive matrices $(Q_\alpha)_{\alpha \in \text{Irr}(\mathbb{G})}$

such that the two normal semi-finite faithful (in short n.s.f.) weights

$$(2.1) \quad h_L : x \mapsto \sum_{\alpha \in \text{Irr}(\mathbb{G})} \text{Tr}(Q_\alpha) \text{Tr}(Q_\alpha(xp_\alpha))$$

$$(2.2) \quad h_R : x \mapsto \sum_{\alpha \in \text{Irr}(\mathbb{G})} \text{Tr}(Q_\alpha^{-1}) \text{Tr}(Q_\alpha^{-1}(xp_\alpha))$$

on $\ell^\infty(\widehat{\mathbb{G}})$ are respectively left and right invariant. These so-called *Haar weights* are both unique up to multiplication by a scalar.

2.2. Actions on von Neumann algebras

Actions of quantum groups can be defined both on C*-algebras and on von Neumann algebras. However, only the latter will be used in this paper. The main feature, which will prove of importance later on, is the existence of a unitary implementation. The standard reference on this subject is [36].

DEFINITION 2.4. — A (left) action of $\widehat{\mathbb{G}}$ on a von Neumann algebra M is a unital normal *-homomorphism $\rho : M \rightarrow \ell^\infty(\widehat{\mathbb{G}}) \overline{\otimes} M$ such that

$$(\widehat{\Delta} \otimes \iota) \circ \rho = (\iota \otimes \rho) \circ \rho.$$

The *fixed point algebra* of the action ρ is the subalgebra $M^\rho = \{x \in M, \rho(x) = 1 \otimes x\}$. A subalgebra N of M is said to be *stable* under the action if $\rho(N) \subset \ell^\infty(\widehat{\mathbb{G}}) \overline{\otimes} N$. In that case, there is a *restricted action* of $\widehat{\mathbb{G}}$ on N which will still be denoted by ρ . The crossed-product construction in this setting generalizes the classical definition.

DEFINITION 2.5. — Let $\widehat{\mathbb{G}}$ be a discrete quantum group acting on a von Neumann algebra M . The crossed-product $\widehat{\mathbb{G}} \rtimes_\rho M$ is the von Neumann subalgebra of $\mathcal{B}(L^2(\mathbb{G})) \overline{\otimes} M$ generated by $\rho(M)$ and $L^\infty(\mathbb{G}) \otimes 1$.

The crossed-product is endowed with a *dual action* $\widehat{\rho}$ of \mathbb{G}^{op} (i.e. with respect to the flipped coproduct) defined by

$$\begin{cases} \widehat{\rho}(\rho(m)) &= 1 \otimes \rho(m) \\ \widehat{\rho}(a \otimes 1) &= [(\sigma \circ \Delta)(a)] \otimes 1 \end{cases}$$

for all $m \in M$ and $a \in L^\infty(\mathbb{G})$. Let $\widehat{\mathbb{G}}$ be a discrete quantum group and let M be a von Neumann algebra together with a fixed n.s.f. weight θ with GNS construction $(K, \iota, \Lambda_\theta)$. It is proven in [36] that any action of $\widehat{\mathbb{G}}$ on M is unitarily implementable, i.e. there exists a unitary

$$U^\rho \in \ell^\infty(\mathbb{G}) \overline{\otimes} \mathcal{B}(K)$$

which is the adjoint of a representation of $\widehat{\mathbb{G}}$ and such that

$$\rho(x) = U^\rho(1 \otimes x)(U^\rho)^*$$

for all $x \in M$.

2.3. Quantum subgroups and quotients

Let us give some details concerning the notions of discrete quantum subgroups and quotients, which will appear in all the constructions of this work. Let \mathbb{G} be a compact quantum group and let \mathbb{H} be another compact quantum group such that $C(\mathbb{H}) \subset C(\mathbb{G})$ and the coproduct of \mathbb{H} is given by the restriction of the coproduct of \mathbb{G} . Then, $\widehat{\mathbb{H}}$ is said to be a *discrete quantum subgroup* of $\widehat{\mathbb{G}}$. The following important fact was proved in [39, Prop 2.2].

PROPOSITION 2.6 (Vergnioux). — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group, let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup and denote the respective Haar states of \mathbb{G} and \mathbb{H} by $h_{\mathbb{G}}$ and $h_{\mathbb{H}}$. Then, there exists a faithful conditional expectation $\mathbb{E}_{\mathbb{H}} : C_{\text{red}}(\mathbb{G}) \rightarrow C_{\text{red}}(\mathbb{H})$ such that $h_{\mathbb{H}} \circ \mathbb{E}_{\mathbb{H}} = h_{\mathbb{G}}$. Moreover, $\mathbb{E}_{\mathbb{H}}$ extends to a faithful conditional expectation from $L^\infty(\mathbb{G})$ to $L^\infty(\mathbb{H})$, still denoted by $\mathbb{E}_{\mathbb{H}}$.*

Note that the inclusion $C(\mathbb{H}) \subset C(\mathbb{G})$ extends to inclusions of matrix algebras $M_n(C(\mathbb{H})) \subset M_n(C(\mathbb{G}))$ so that any finite-dimensional representation of \mathbb{H} can be seen as a finite-dimensional representation of \mathbb{G} . This inclusion obviously preserves intertwiners, so that we have an inclusion $\text{Irr}(\mathbb{H}) \subset \text{Irr}(\mathbb{G})$. Let us define a central projection

$$p_{\mathbb{H}} = \sum_{\alpha \in \text{Irr}(\mathbb{H})} p_\alpha \in \ell^\infty(\widehat{\mathbb{G}}).$$

We can use $p_{\mathbb{H}}$ to describe the structure of $\widehat{\mathbb{H}}$ from the structure of $\widehat{\mathbb{G}}$ (see [18, Prop 2.3] for details).

PROPOSITION 2.7. — *With the notations above, we have*

- (1) $\widehat{\Delta}(p_{\mathbb{H}})(p_{\mathbb{H}} \otimes 1) = p_{\mathbb{H}} \otimes p_{\mathbb{H}}$
- (2) $\ell^\infty(\widehat{\mathbb{H}}) = p_{\mathbb{H}} \ell^\infty(\widehat{\mathbb{G}})$
- (3) $\widehat{\Delta}_{\mathbb{H}}(a) = \widehat{\Delta}(a)(p_{\mathbb{H}} \otimes p_{\mathbb{H}})$
- (4) *If h_L is a left Haar weight for $\widehat{\mathbb{G}}$, then $h_{L,\mathbb{H}} : x \mapsto h_L(xp_{\mathbb{H}})$ is a left Haar weight for $\widehat{\mathbb{H}}$.*

It is easy to see, using the above statements, that the map $a \mapsto \widehat{\Delta}(a)(1 \otimes p_{\mathbb{H}})$ defines a right action (right actions of discrete quantum groups are defined in the same way as left actions with the obvious modifications) of $\widehat{\mathbb{H}}$ on $\ell^\infty(\widehat{\mathbb{G}})$. Let $\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ be the fixed point subalgebra for this action. Using Proposition 2.7 again, we see that the restriction of the coproduct $\widehat{\Delta}$ to $\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ yields a left action of $\widehat{\mathbb{G}}$ on this von Neumann algebra which will be denoted by τ . Let h_L be the left-invariant weight of $\widehat{\mathbb{G}}$ defined by Equation (2.1). It is known from [18, Prop 2.4] that the map

$$T : x \mapsto (\iota \otimes h_{L, \mathbb{H}})[\widehat{\Delta}(x)(1 \otimes p_{\mathbb{H}})]$$

is a normal faithful operator-valued weight from $\ell^\infty(\widehat{\mathbb{G}})$ to $\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ and that there exists a n.s.f. weight θ on $\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ such that $h_L = \theta \circ T$. Let U be the unitary implementation of the action τ with respect to the weight θ . Then, $\mathcal{R} = U^*$ will be called the *quasi-regular representation* of $\widehat{\mathbb{G}}$ modulo $\widehat{\mathbb{H}}$.

Remark 2.8. — By a straightforward calculation, we see that when both sides are well-defined,

$$\tau \circ T(x) = (\iota \otimes \iota \otimes h_{L, \mathbb{H}})[(\widehat{\Delta} \otimes \iota)(\widehat{\Delta}(x)(1 \otimes p_{\mathbb{H}}))] = (\iota \otimes T) \circ \tau(x).$$

The weight θ can therefore be interpreted as an almost invariant measure on the quotient space with respect to the action τ .

2.4. Approximation properties

Two approximation properties will be considered in this paper : weak amenability and the Haagerup property. They have both been defined in earlier works and enjoy various characterizations (see for example [27], [21] and [14]). For our purpose, the point of view of *multipliers* is the best suited. We refer the reader to [11, Ch 12] for an introduction to approximation properties for classical groups, which motivates the following definitions.

DEFINITION 2.9. — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group and $a \in \ell^\infty(\widehat{\mathbb{G}})$. The left multiplier associated to a is the map $m_a : \text{Pol}(\widehat{\mathbb{G}}) \rightarrow \text{Pol}(\widehat{\mathbb{G}})$ defined by*

$$(m_a \otimes \iota)(u^\alpha) = (1 \otimes ap_\alpha)u^\alpha,$$

for every irreducible representation α of \mathbb{G} . A net (a_t) of elements of $\ell^\infty(\widehat{\mathbb{G}})$ is said to converge pointwise to $a \in \ell^\infty(\widehat{\mathbb{G}})$ if for every irreducible representation α of \mathbb{G} , $a_t p_\alpha \rightarrow ap_\alpha$ in $\mathcal{B}(H_\alpha)$. An element $a \in \ell^\infty(\widehat{\mathbb{G}})$ is said to have finite support if $ap_\alpha = 0$ for all but finitely many $\alpha \in \text{Irr}(\mathbb{G})$.

DEFINITION 2.10. — A discrete quantum group $\widehat{\mathbb{G}}$ is said to be weakly amenable if there exists a net (a_t) of elements of $\ell^\infty(\widehat{\mathbb{G}})$ such that

- a_t has finite support for all t .
- (a_t) converges pointwise to 1.
- $K := \limsup_t \|m_{a_t}\|_{cb}$ is finite.

The lower bound of the constants K for all nets satisfying these properties is denoted by $\Lambda_{cb}(\widehat{\mathbb{G}})$ and called the Cowling-Haagerup constant of $\widehat{\mathbb{G}}$. By convention, $\Lambda_{cb}(\widehat{\mathbb{G}}) = \infty$ if $\widehat{\mathbb{G}}$ is not weakly amenable.

DEFINITION 2.11. — A discrete quantum group $\widehat{\mathbb{G}}$ is said to have the Haagerup property if there exists a net (a_t) of elements of $\ell^\infty(\widehat{\mathbb{G}})$ such that

- $a_t \in C_0(\widehat{\mathbb{G}})$ for all t .
- (a_t) converges pointwise to 1.
- m_{a_t} is completely positive for all t .

As in the classical case, these properties are connected to corresponding approximation properties for the associated operator algebras (see for instance [11, Ch 12] for the definitions). However, the link is not fully understood yet when the quantum group is not *unimodular*. Let us therefore only state results in the unimodular case (i.e. when the Haar state is a trace), which were proved in [27, Thm 5.14].

THEOREM 2.12 (Kraus-Ruan). — Let $\widehat{\mathbb{G}}$ be a unimodular discrete quantum group. Then,

- $\widehat{\mathbb{G}}$ has the Haagerup property $\Leftrightarrow C_{red}(\mathbb{G})$ has the Haagerup property relative to $h \Leftrightarrow L^\infty(\mathbb{G})$ has the Haagerup property.
- $\Lambda_{cb}(\widehat{\mathbb{G}}) = \Lambda_{cb}(C_{red}(\mathbb{G})) = \Lambda_{cb}(L^\infty(\mathbb{G}))$.

3. Permanence results

This section is divided into two parts. In the first one, we will prove permanence of approximation properties under several elementary constructions. In the second part, we will extend the result of [21] to free products amalgamated over a finite quantum subgroup. The result also holds for the Haagerup property, thus improving [14, Thm 7.8] by removing the unimodularity assumption.

3.1. First results

We start with the simplest case, namely passing to quantum subgroups. The permanence of the Haagerup property by passing to discrete quantum

subgroups (or more generally to closed quantum subgroups of a locally compact quantum group) was proved in [14, Prop 5.7]. We therefore only consider weak amenability.

PROPOSITION 3.1. — *Let $\widehat{\mathbb{G}}$ be a weakly amenable discrete quantum group and let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup of $\widehat{\mathbb{G}}$. Then, $\widehat{\mathbb{H}}$ is weakly amenable and $\Lambda_{cb}(\widehat{\mathbb{H}}) \leq \Lambda_{cb}(\widehat{\mathbb{G}})$.*

Proof. — Let $\epsilon > 0$ and let (a_t) be a net of finitely supported elements in $\ell^\infty(\widehat{\mathbb{G}})$ converging pointwise to 1 and such that $\limsup \|m_{a_t}\|_{cb} \leq \Lambda_{cb}(\widehat{\mathbb{G}}) + \epsilon$. Then, using the notations of Proposition 2.7, $(a_t p_{\mathbb{H}})$ is a net of finitely supported elements in $\ell^\infty(\widehat{\mathbb{H}})$ which converges pointwise to 1. Using the conditional expectation of Proposition 2.6, we see that $\|m_{a_t p_{\mathbb{H}}}\|_{cb} = \|\mathbb{E}_{\mathbb{H}} \circ m_{a_t}\|_{cb} \leq \|m_{a_t}\|_{cb}$. Thus, $\Lambda_{cb}(\widehat{\mathbb{H}}) \leq \Lambda_{cb}(\widehat{\mathbb{G}}) + \epsilon$. \square

It is proved in [42] that if $\widehat{\mathbb{G}}$ and $\widehat{\mathbb{H}}$ are two discrete quantum groups, then the minimal tensor product $C(\mathbb{G}) \otimes C(\mathbb{H})$ can be turned into a compact quantum group in a natural way. Its dual discrete quantum group is denoted by $\widehat{\mathbb{G}} \times \widehat{\mathbb{H}}$ and called the *direct product* of $\widehat{\mathbb{G}}$ and $\widehat{\mathbb{H}}$.

PROPOSITION 3.2. — *Let $\widehat{\mathbb{G}}$ and $\widehat{\mathbb{H}}$ be two discrete quantum groups. Then, $\widehat{\mathbb{G}} \times \widehat{\mathbb{H}}$ is weakly amenable if and only if both $\widehat{\mathbb{G}}$ and $\widehat{\mathbb{H}}$ are weakly amenable. Moreover,*

$$\max(\Lambda_{cb}(\widehat{\mathbb{G}}), \Lambda_{cb}(\widehat{\mathbb{H}})) \leq \Lambda_{cb}(\widehat{\mathbb{G}} \times \widehat{\mathbb{H}}) \leq \Lambda_{cb}(\widehat{\mathbb{G}})\Lambda_{cb}(\widehat{\mathbb{H}}).$$

Proof. — The "only if" part is a direct consequence of Proposition 3.1, as well as the first inequality. To prove the second inequality, let $\epsilon > 0$ and let (a_t) and (b_s) be nets of finitely supported elements respectively in $\ell^\infty(\widehat{\mathbb{G}})$ and in $\ell^\infty(\widehat{\mathbb{H}})$ converging pointwise to 1 and such that $\limsup \|m_{a_t}\|_{cb} \leq \Lambda_{cb}(\widehat{\mathbb{G}}) + \epsilon$ and $\limsup \|m_{b_s}\|_{cb} \leq \Lambda_{cb}(\widehat{\mathbb{H}}) + \epsilon$. Set

$$c_{(t,s)} = a_t \otimes b_s \in \ell^\infty(\widehat{\mathbb{G}} \times \widehat{\mathbb{H}}).$$

From the description of the representation theory of direct products given in [42, Thm 2.11], we see that $(c_{(t,s)})$ is a net of finitely supported elements converging pointwise to 1. Moreover, since $m_{c_{(t,s)}} = m_{a_t} \otimes m_{b_s}$, we have $\Lambda_{cb}(\widehat{\mathbb{G}} \times \widehat{\mathbb{H}}) \leq (\Lambda_{cb}(\widehat{\mathbb{G}}) + \epsilon)(\Lambda_{cb}(\widehat{\mathbb{H}}) + \epsilon)$, which concludes the proof. \square

Remark 3.3. — It is a general fact that for any two C*-algebras A and B , $\Lambda_{cb}(A \otimes B) = \Lambda_{cb}(A)\Lambda_{cb}(B)$ (see e.g. [11, Thm 12.3.13]). Hence, we always have

$$\Lambda_{cb}(C_{\text{red}}(\mathbb{G} \otimes \mathbb{H})) = \Lambda_{cb}(C_{\text{red}}(\mathbb{G}))\Lambda_{cb}(C_{\text{red}}(\mathbb{H})).$$

Moreover, Theorem 2.12 implies that $\Lambda_{cb}(\widehat{\mathbb{G}} \times \widehat{\mathbb{H}}) = \Lambda_{cb}(\widehat{\mathbb{G}})\Lambda_{cb}(\widehat{\mathbb{H}})$ as soon as the discrete quantum groups are unimodular. It is very likely that this equality holds in general but we were not able to prove it.

A similar statement holds for the Haagerup property.

PROPOSITION 3.4. — *Let $\widehat{\mathbb{G}}$ and $\widehat{\mathbb{H}}$ be two discrete quantum groups. Then, $\widehat{\mathbb{G}} \times \widehat{\mathbb{H}}$ has the Haagerup property if and only if both $\widehat{\mathbb{G}}$ and $\widehat{\mathbb{H}}$ have the Haagerup property.*

Proof. — The "only if" part comes from stability under passing to quantum subgroups. To prove the "if" part, one can build, as in the proof of Proposition 3.2, multipliers on $\widehat{\mathbb{G}} \times \widehat{\mathbb{H}}$ by tensoring multipliers on the two quantum groups. To finish the proof, note that a tensor product of unital completely positive maps is again unital and completely positive and that the tensor product of two elements in $C_0(\widehat{\mathbb{G}})$ and $C_0(\widehat{\mathbb{H}})$ respectively lies in $C_0(\widehat{\mathbb{G}} \times \widehat{\mathbb{H}})$. □

The third construction we will study is inductive limits of discrete quantum groups, or equivalently inverse limits of compact quantum groups. It was proved in [41, Prop 3.1] that given a family of discrete quantum groups $(\widehat{\mathbb{G}}_i)_i$ with connecting maps $\pi_{ij} : C(\widehat{\mathbb{G}}_i) \rightarrow C(\widehat{\mathbb{G}}_j)$ intertwining the coproducts and satisfying $\pi_{jk} \circ \pi_{ij} = \pi_{ik}$, there is a natural compact quantum group structure on the inductive limit C*-algebra. Its dual discrete quantum group is called the *inductive limit* of the system $(\widehat{\mathbb{G}}_i, \pi_{ij})$. In order to study approximation properties, we first need to understand its representation theory. Since we were not able to find a reference for this, we give a statement even though it is certainly well-known to experts.

PROPOSITION 3.5. — *Let $(\widehat{\mathbb{G}}_i, \pi_{ij})$ be an inductive system of discrete quantum groups with inductive limit $\widehat{\mathbb{G}}$ and assume that all the maps π_{ij} are injective. Then, there is a one-to-one correspondence between the irreducible representations of $\widehat{\mathbb{G}}$ and the increasing union of the sets of irreducible representations of the $\widehat{\mathbb{G}}_i$'s.*

Proof. — The maps π_{ij} being injective, we can identify each $\widehat{\mathbb{G}}_i$ with a discrete quantum subgroup of the $\widehat{\mathbb{G}}_j$'s for $j \geq i$. This gives inclusions of the sets of irreducible representations and we denote by \mathcal{S} the increasing union of these sets. We can also identify each $C(\widehat{\mathbb{G}}_i)$ with a C*-subalgebra of $C(\widehat{\mathbb{G}})$ in such a way that

$$\overline{\bigcup C(\widehat{\mathbb{G}}_i)} = C(\widehat{\mathbb{G}}).$$

Under this identification, the discrete quantum groups $\widehat{\mathbb{G}}_i$ are quantum subgroups of $\widehat{\mathbb{G}}$, hence any irreducible representation of some \mathbb{G}_i yields an irreducible representation of \mathbb{G} and we have proved that $\mathcal{S} \subset \text{Irr}(\mathbb{G})$. Moreover, the algebra

$$\mathcal{A} := \bigcup_i \text{Pol}(\mathbb{G}_i)$$

is a dense Hopf- $*$ -subalgebra of $C(\mathbb{G})$ spanned by coefficients of irreducible representations. Because of Schur's orthogonality relations, the density implies that the coefficients of all irreducible representations of \mathbb{G} are in \mathcal{A} , i.e. $\mathcal{A} = \text{Pol}(\mathbb{G})$. This means that any irreducible representation of \mathbb{G} comes from an element of \mathcal{S} and $\text{Irr}(\mathbb{G}) = \mathcal{S}$. \square

We can now prove the permanence of weak amenability under this construction.

PROPOSITION 3.6. — *Let $(\widehat{\mathbb{G}}_i, \pi_{ij})$ be an inductive system of discrete quantum groups with inductive limit $\widehat{\mathbb{G}}$ and limit maps $\pi_i : C(\mathbb{G}_i) \rightarrow C(\mathbb{G})$. Then, if all the maps π_i are injective,*

$$\sup_i \Lambda_{cb}(\widehat{\mathbb{G}}_i) = \Lambda_{cb}(\widehat{\mathbb{G}}).$$

In particular, the inductive limit is weakly amenable if and only if the quantum groups are all weakly amenable with uniformly bounded Cowling-Haagerup constant.

Proof. — The injectivity of the limit maps ensures that each $\widehat{\mathbb{G}}_i$ can be seen as a discrete quantum subgroup of \mathbb{G} . Hence, Corollary 3.1 gives the inequality

$$\sup_i \Lambda_{cb}(\widehat{\mathbb{G}}_i) \leq \Lambda_{cb}(\widehat{\mathbb{G}}).$$

To prove the converse inequality, fix an $\epsilon > 0$ and let $(a_t^i)_t$ be a net of finitely supported elements in $\ell^\infty(\widehat{\mathbb{G}}_i)$ converging pointwise to 1 and such that $\limsup \|m_{a_t^i}\|_{cb} \leq \Lambda_{cb}(\widehat{\mathbb{G}}_i) + \epsilon$. Using the description of the representation theory of inductive limits given by Proposition 3.5, we can see $(a_t^i)_{(i,t)}$ as a net of finitely supported elements of $\ell^\infty(\widehat{\mathbb{G}})$ converging pointwise to 1 by setting $a_t^i p_\alpha = 0$ for any $\alpha \notin \text{Irr}(\mathbb{G}_i)$. The associated multiplier is $m_{a_t^i} \circ \mathbb{E}_{\mathbb{G}_i}$, so that it has the same completely bounded norm as $m_{a_t^i}$. The conditions on the completely bounded norms then gives $\Lambda_{cb}(\widehat{\mathbb{G}}) \leq \sup_i \Lambda_{cb}(\widehat{\mathbb{G}}_i) + \epsilon$. \square

PROPOSITION 3.7. — *Let $(\widehat{\mathbb{G}}_i, \pi_{ij})$ be an inductive system of discrete quantum groups with inductive limit $\widehat{\mathbb{G}}$ and limit maps $\pi_i : C(\mathbb{G}_i) \rightarrow C(\mathbb{G})$. Then, if all the maps π_i are injective, $\widehat{\mathbb{G}}$ has the Haagerup property if and only if all the quantum groups \mathbb{G}_i have the Haagerup property.*

Proof. — The "only if" part comes from stability under passing to quantum subgroups. To prove the "if" part, we simply have to prove that complete positivity is preserved when a multiplier is extended to the inductive limit. This comes from the fact that the multiplier is equal to $m_{a_i} \circ \mathbb{E}_{\mathbb{G}_i}$. \square

3.2. Amalgamated free products

Consider two discrete quantum groups $\widehat{\mathbb{G}}_1$ and $\widehat{\mathbb{G}}_2$ together with a common discrete quantum subgroup $\widehat{\mathbb{H}}$ and let us consider the C^* -algebra A obtained by taking the reduced amalgamated free product $C_{\text{red}}(\mathbb{G}_1) *_{C_{\text{red}}(\mathbb{H})} C_{\text{red}}(\mathbb{G}_2)$ with respect to the conditional expectations given by 2.6. The coproducts of \mathbb{G}_1 and \mathbb{G}_2 induce a map Δ on A which is shown in [41] to be a coproduct turning (A, Δ) into a compact quantum group. In analogy with the classical case, the dual of A will be called the *free product of $\widehat{\mathbb{G}}_1$ and $\widehat{\mathbb{G}}_2$ amalgamated over $\widehat{\mathbb{H}}$* .

It is well-known that a free product of amenable groups need not be amenable. However, it was proved in [14, Thm 7.8] that the Haagerup property passes to free products of discrete quantum groups and it was proved in [21] that a free product of discrete quantum groups with Cowling-Haagerup constant equal to 1 again has Cowling-Haagerup constant equal to 1. Our goal in this subsection is to extend those two results by allowing amalgamation over a finite quantum subgroup. Note that such a statement for the Haagerup property was proved in [14, Prop 7.13] when the quantum groups are unimodular, using the associated von Neumann algebras. Our proof does not require unimodularity, but the price to pay is dealing all the way long with multipliers. For this, we need the following generalization of Gilbert's criterion proved in [15, Prop 4.1 and Thm 4.2].

THEOREM 3.8 (Daws). — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group and let $a \in \ell^\infty(\widehat{\mathbb{G}})$. Then, m_a extends to a completely bounded multiplier on $\mathcal{B}(L^2(\mathbb{G}))$ if and only if there exists a Hilbert space K and two maps $\xi, \eta \in \mathcal{B}(L^2(\mathbb{G}), L^2(\mathbb{G}) \otimes K)$ such that $\|\xi\| \|\eta\| = \|m_a\|_{cb}$ and*

$$(3.1) \quad (1 \otimes \eta)^* \widehat{W}_{12}^* (1 \otimes \xi) \widehat{W} = a \otimes 1.$$

Moreover, we then have $m_a(x) = \eta^*(x \otimes 1)\xi$.

Let us give a proof of the stability of the Haagerup property under free products in the language of multipliers, in order to make the extension to the amalgamated case more clear.

PROPOSITION 3.9. — *Let $(\widehat{\mathbb{G}}_i)_{i \in I}$ be a family of discrete quantum groups with the Haagerup property. Then, $*_i \widehat{\mathbb{G}}_i$ has the Haagerup property.*

Proof. — It is clearly enough to prove the result for two quantum groups \mathbb{G}_1 and \mathbb{G}_2 . First note that according to [15, Thm 5.9], the complete positivity of a multiplier m_a implies that $a \in C_0(\widehat{\mathbb{G}}_i)$ can in fact be chosen to be of the form $(\omega_a \otimes \iota)(W_i)$ for some state ω_a on the envelopping C^* -algebra $C_{\max}(\mathbb{G}_i)$ of $\text{Pol}(\mathbb{G}_i)$. This means that if we consider two elements $a \in C_0(\widehat{\mathbb{G}}_1)$ and $b \in C_0(\widehat{\mathbb{G}}_2)$, then the free product $m_a * m_b$ of completely positive maps corresponds to the multiplier m_c with

$$c = (\omega_c \otimes \iota)(W),$$

where ω_c is the free product of the states ω_a and ω_b . So let us take nets $(\omega_{a_t} \otimes \iota)(W_1)$ and $(\omega_{b_t} \otimes \iota)(W_2)$ implementing the Haagerup property for $\widehat{\mathbb{G}}_1$ and $\widehat{\mathbb{G}}_2$ respectively. Using the remark above, we get a net of elements $c_t = (\omega_{c_t} \otimes \iota)(W)$ converging pointwise to the identity and yielding completely positive multipliers (because ω_{c_t} is again a state). Now, [8, Thm 3.9] asserts that the free product map $m_{c_t} = m_{a_t} * m_{b_t}$ is L^2 -compact as soon as m_{a_t} and m_{b_t} are. If $\widehat{\mathbb{K}} = \widehat{\mathbb{G}}_1 * \widehat{\mathbb{G}}_2$, we have for any $\alpha \in \text{Irr}(\mathbb{K})$,

$$\begin{aligned} (h \otimes \iota)((m_{c_t} \otimes \iota)(u^\alpha)(u^\alpha)^*) &= (h \otimes \iota)((\omega_{c_t} \otimes \iota \otimes \iota)(u_{13}^\alpha u_{23}^\alpha)(u^\alpha)^*) \\ &= (h \otimes \iota)((\omega_{c_t} \otimes \iota \otimes \iota)[u_{13}^\alpha u_{23}^\alpha (u_{23}^\alpha)^*]) \\ &= (h \otimes \iota)((\omega_{c_t} \otimes \iota \otimes \iota)(u_{13}^\alpha)) \\ &= c_t p_\alpha. \end{aligned}$$

This and the fact that m_{c_t} is L^2 -compact prove that $c_t \in C_0(\widehat{\mathbb{G}}_1 * \widehat{\mathbb{G}}_2)$, concluding the proof. □

The strategy to handle the amalgamated case is to produce multipliers which, while implementing the desired approximation property, are the identity on the amalgam. Here finiteness proves crucial through the next Lemma. If $\widehat{\mathbb{H}}$ is a finite quantum group, it is unimodular and we will denote by \widehat{h} the unique Haar weight on $\ell^\infty(\widehat{\mathbb{H}})$ which is both left and right invariant, i.e. $\widehat{h}(a) = \sum_\alpha \text{Tr}(ap_\alpha)$. We first define the averaging process.

DEFINITION 3.10. — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group, $\widehat{\mathbb{H}}$ a finite quantum subgroup of $\widehat{\mathbb{G}}$ and let $a \in \ell^\infty(\widehat{\mathbb{G}})$. The averaging of a over $\widehat{\mathbb{H}}$ is the element $c \in \ell^\infty(\widehat{\mathbb{G}})$ defined by*

$$c = (\widehat{h} \otimes \iota)[(p_{\mathbb{H}} \otimes \iota)\widehat{\Delta}(a)].$$

We now prove that this averaging process is well-behaved with respect to completely bounded norms and that it yields the identity on $\widehat{\mathbb{H}}$.

LEMMA 3.11. — *The averaging of a over $\widehat{\mathbb{H}}$ satisfies the inequality $\|m_c\|_{cb} \leq |\widehat{h}(p_{\mathbb{H}})| \|m_a\|_{cb}$. Moreover, m_c is a multiple of the identity on $C_{red}(\mathbb{H}) \subset C_{red}(\mathbb{G})$.*

Proof. — If $\eta, \xi : L^2(\mathbb{G}) \rightarrow L^2(\mathbb{G}) \otimes K$ are the maps coming from Theorem 3.8, we set

$$(3.2) \quad \xi' = (\widehat{h} \otimes \iota \otimes \iota)[\widehat{W}_{12}^*(p_{\mathbb{H}} \otimes \xi)\widehat{W}].$$

Let us make the meaning of this definition clear : if $x \in \ell^2(\mathbb{G}) \otimes L^2(\mathbb{G})$, then $\widehat{W}x$ belongs to the same space so that we can apply ξ to its second leg, yielding an element $y \in \ell^2(\widehat{\mathbb{G}}) \otimes L^2(\mathbb{G}) \otimes K$ to which we can apply \widehat{W}_{12}^* . Now, taking $p_{\mathbb{H}}$ into account we see that the term inside the brackets is a linear map from $\ell^2(\mathbb{H}) \otimes L^2(\mathbb{G})$ to $\ell^2(\mathbb{H}) \otimes L^2(\mathbb{G}) \otimes K$. Eventually, applying \widehat{h} to the first leg (note that it is defined on all $\mathcal{B}(\ell^2(\widehat{\mathbb{H}}))$ because it is finite-dimensional) gives a map from $L^2(\mathbb{G})$ to $L^2(\mathbb{G}) \otimes K$.

We are going to prove that $c \otimes 1 = (1 \otimes \eta^*)\widehat{W}_{12}^*(1 \otimes \xi')\widehat{W}$, which will imply our claim on the completely bounded norm since $\|\xi'\| \leq |\widehat{h}(p_{\mathbb{H}})| \|\xi\|$. First note that since

$$(\widehat{\Delta} \otimes \iota)(x) = \widehat{W}_{12}^*(1 \otimes x)\widehat{W}_{12},$$

we have

$$\begin{aligned} \widehat{\Delta}(a) \otimes 1 &= \widehat{W}_{12}^*(1 \otimes 1 \otimes \eta^*)(1 \otimes \widehat{W}_{12}^*)(1 \otimes 1 \otimes \xi)(1 \otimes \widehat{W})\widehat{W}_{12} \\ &= (1 \otimes 1 \otimes \eta^*)(\widehat{W}_{12}^* \otimes 1)(1 \otimes \widehat{W}_{12}^*)(\widehat{W}_{12} \otimes 1) \\ &\quad \times (1 \otimes 1 \otimes \xi)\widehat{W}_{12}^*\widehat{W}_{23}\widehat{W}_{12} \\ &= (1 \otimes 1 \otimes \eta^*)(\widehat{W}_{23}^*\widehat{W}_{13}^* \otimes 1)(1 \otimes 1 \otimes \xi)\widehat{W}_{13}\widehat{W}_{23} \end{aligned}$$

where we used twice the pentagonal equation for \widehat{W} . Applying $\widehat{h}(p_{\mathbb{H}} \cdot)$ to the first leg yields the result. Now, if $\alpha \in \text{Irr}(\mathbb{H})$, we get using the invariance of \widehat{h} ,

$$\begin{aligned} cp_{\alpha} &= (\widehat{h} \otimes \iota)[(p_{\mathbb{H}} \otimes \iota)\widehat{\Delta}(a)]p_{\alpha} \\ &= (\widehat{h} \otimes \iota)[(p_{\mathbb{H}} \otimes p_{\mathbb{H}})\widehat{\Delta}(a)]p_{\alpha} \\ &= (\widehat{h} \otimes \iota)[\widehat{\Delta}(p_{\mathbb{H}}a)]p_{\alpha} \\ &= \widehat{h}(p_{\mathbb{H}}a)p_{\alpha} \end{aligned}$$

and $m_c = \widehat{h}(p_{\mathbb{H}}a) \text{Id}$ on $C_{red}(\mathbb{H})$. □

Using this averaging technique, we get the result for the Haagerup property.

THEOREM 3.12. — *Let $(\widehat{\mathbb{G}}_i)_{i \in I}$ be a family of discrete quantum groups with the Haagerup property and let $\widehat{\mathbb{H}}$ be a common finite quantum subgroup. Then, $*_{\widehat{\mathbb{H}}}\widehat{\mathbb{G}}_i$ has the Haagerup property.*

Proof. — We first prove that completely positive multipliers on $C_{\text{red}}(\mathbb{G}_i)$ can be averaged so that they become the identity on $C_{\text{red}}(\mathbb{H})$. Let us first recall that by [27, Prop 2.6], if $c \in \ell^\infty(\widehat{\mathbb{G}}_i)$ satisfies $m_c(x) = \eta^*(x \otimes 1)\xi$ for some maps $\xi, \eta \in \mathcal{B}(L^2(\mathbb{G}_i), L^2(\mathbb{G}_i) \otimes K)$, then

$$m_{\widehat{S}(c)^*}(x) = \xi^*(x \otimes 1)\eta,$$

where \widehat{S} denotes the antipode of $\widehat{\mathbb{G}}_i$. Let (a_t) be a net of elements in $C_0(\widehat{\mathbb{G}}_i)$ converging pointwise to 1 and such that m_{a_t} is unital and completely positive. Let b_t be the averaging of a_t over $\widehat{\mathbb{H}}$. Average again $\widehat{S}(b_t)^*$ to produce a third element b'_t . Then, by Lemma 3.11, (b'_t) is a net of elements in $C_0(\widehat{\mathbb{G}}_i)$ converging pointwise to 1 and such that $m_{b'_t}$ is the identity on $C_{\text{red}}(\mathbb{H})$. In the case of a unital completely positive multiplier, the proof of Theorem 3.8 yields a map $\xi : L^2(\mathbb{G}_i) \rightarrow L^2(\mathbb{G}_i) \otimes K$ such that $m_{a_t}(x) = \xi^*(x \otimes \iota)\xi$. This implies that $m_{b'_t}(x) = (\xi')^*(x \otimes 1)\xi'$ is unital and completely positive. Moreover, the conditional expectation $\mathbb{E}_{\widehat{\mathbb{H}}} \circ m_{b'_t}$ is invariant with respect to the Haar state, because $m_{b'_t}$ is invariant. Since, by [39, Prop 2.2], $\mathbb{E}_{\widehat{\mathbb{H}}}$ is the unique conditional expectation satisfying this invariance, we have $\mathbb{E}_{\widehat{\mathbb{H}}} \circ m_{b'_t} = \mathbb{E}_{\widehat{\mathbb{H}}}$.

Now that all the multipliers are the identity on $C_{\text{red}}(\mathbb{H})$ and preserve the conditional expectation $\mathbb{E}_{\widehat{\mathbb{H}}}$, their amalgamated free product makes sense (see for example [11, Thm 4.8.5] for details on the construction of the free product of u.c.p. maps). We can then follow the proof of Proposition 3.9 to conclude. □

Let us turn to weak amenability, which is more involved, and let us check that our averaging technique preserves weak amenability.

LEMMA 3.13. — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group, $\widehat{\mathbb{H}}$ a finite quantum subgroup of $\widehat{\mathbb{G}}$ and let (a_t) be a net of finitely supported elements in $\ell^\infty(\widehat{\mathbb{G}})$ converging pointwise to 1. Then, there exists a net (b_t) of finite-rank elements in $\ell^\infty(\widehat{\mathbb{G}})$ converging pointwise to 1, such that*

$$\limsup_t \|m_{b_t}\|_{cb} \leq \limsup_t \|m_{a_t}\|_{cb}$$

and m_{b_t} is the identity on $C_{\text{red}}(\mathbb{H}) \subset C_{\text{red}}(\mathbb{G})$.

Proof. — Let c_t be the averaging of a_t over $\widehat{\mathbb{H}}$ and let $\text{Supp}(a_t)$ be the set of equivalence classes of irreducible representations α of \mathbb{G} such that

$a_t p_\alpha \neq 0$. Then,

$$\begin{aligned} c_t p_\alpha &= (\widehat{h} \otimes \iota) \left[(p_{\mathbb{H}} \otimes p_\alpha) \widehat{\Delta}(a) \right] \\ &= (\widehat{h} \otimes \iota) \left[\sum_{\beta \in \text{Irr}(\mathbb{H})} \sum_{\gamma \in \text{Supp}(a)} \widehat{\Delta}(p_\gamma) (p_\beta \otimes p_\alpha) \widehat{\Delta}(a) \right] \end{aligned}$$

By definition, $\widehat{\Delta}(p_\gamma) (p_\beta \otimes p_\alpha) \neq 0$ if and only if $\gamma \subset \beta \otimes \alpha$, which by Frobenius reciprocity (see e.g. [33, Prop 3.1.11]) is equivalent to $\bar{\alpha} \subset \bar{\gamma} \otimes \beta$. Hence, $c_t p_\alpha$ is non-zero only if α belongs to the finite set

$$\bigcup_{\beta \in \text{Irr}(\mathbb{H})} \bigcup_{\gamma \in \text{Supp}(a)} \{ \alpha \in \text{Irr}(\mathbb{G}), \bar{\alpha} \subset \bar{\gamma} \otimes \beta \}$$

and c_t has finite support. The same holds for

$$b_t = \frac{1}{\widehat{h}(p_{\mathbb{H}} a_t)} c_t,$$

which induces multipliers m_{b_t} which are the identity on $C_{\text{red}}(\mathbb{H})$.

Assume now that a_t converges pointwise to 1, and note that the inequality of the completely bounded norms is obvious from Lemma 3.11. Fix $\beta \in \text{Irr}(\mathbb{G})$, $\epsilon > 0$ and let D be the (finite) set of $\gamma \in \text{Irr}(\mathbb{G})$ which are contained in $\alpha \otimes \beta$ for some $\alpha \in \text{Irr}(\mathbb{H})$. We denote by p_D the sum of the projections p_γ for $\gamma \in D$. Let t be such that :

- $\|a_t p_D - p_D\| \leq \frac{\epsilon}{4}$
- $\|a_t p_{\mathbb{H}} - p_{\mathbb{H}}\| \leq \frac{\epsilon}{4}$
- $|\widehat{h}(a_t p_{\mathbb{H}})| > \frac{1}{2}$

Then,

$$\begin{aligned} \|b_t p_\beta - p_\beta\| &= \| |\widehat{h}(a_t p_{\mathbb{H}})|^{-1} (\widehat{h} \otimes \iota) [(p_{\mathbb{H}} \otimes p_\beta) \widehat{\Delta}(a_t)] - p_\beta \| \\ &= |\widehat{h}(a_t p_{\mathbb{H}})|^{-1} \| (\widehat{h} \otimes \iota) [(p_{\mathbb{H}} \otimes p_\beta) \widehat{\Delta}(a_t) - (a_t p_{\mathbb{H}} \otimes p_\beta)] \| \\ &= |\widehat{h}(a_t p_{\mathbb{H}})|^{-1} \| (\widehat{h} \otimes \iota) [(p_{\mathbb{H}} \otimes p_\beta) \widehat{\Delta}(a_t p_D) - (a_t p_{\mathbb{H}} \otimes p_\beta)] \| \\ &\leq |\widehat{h}(a_t p_{\mathbb{H}})|^{-1} \| (\widehat{h} \otimes \iota) [(p_{\mathbb{H}} \otimes p_\beta) \widehat{\Delta}(a_t p_D - p_D)] \| \\ &\quad + |\widehat{h}(a_t p_{\mathbb{H}})|^{-1} \| (\widehat{h} \otimes \iota) [(p_{\mathbb{H}} \otimes p_\beta) \widehat{\Delta}(p_D) - p_{\mathbb{H}} \otimes p_\beta] \| \\ &\quad + |\widehat{h}(a_t p_{\mathbb{H}})|^{-1} \| (\widehat{h} \otimes \iota) [(p_{\mathbb{H}} - a_t p_{\mathbb{H}}) \otimes p_\beta] \| \\ &\leq |\widehat{h}(a_t p_{\mathbb{H}})|^{-1} (\|a_t p_D - p_D\| + \|a_t p_{\mathbb{H}} - p_{\mathbb{H}}\|) \\ &\leq 2 \left(\frac{\epsilon}{4} + \frac{\epsilon}{4} \right) = \epsilon \end{aligned}$$

using the fact that $(p_{\mathbb{H}} \otimes p_{\beta})\widehat{\Delta}(p_D) = p_{\mathbb{H}} \otimes p_{\beta}$. □

From this it is easy to prove using the amalgamated version of [31, Thm 4.3] that if the quantum groups \mathbb{G}_i are amenable, then the free product amalgamated over a finite quantum subgroup is weakly amenable with Cowling-Haagerup constant equal to 1, thus generalizing a result of M. Bożejko and M.A. Picardello [9] (see [23, Thm 2.3.15] for a proof). When the quantum groups are not amenable but are weakly amenable with Cowling-Haagerup constant equal to 1, we need the full power of the work of E. Ricard and X. Qu. Let us introduce some notations : if A is a C^* -algebra with a conditional expectation \mathbb{E} , we denote by $L^2(A, \mathbb{E})$ (resp. $L^2(A, \mathbb{E})^{op}$) the Hilbert module obtained by the GNS constructions using the inner product $\langle a, b \rangle = \mathbb{E}(ab^*)$ (resp. $\langle a, b \rangle = \mathbb{E}(a^*b)$).

THEOREM 3.14 (Ricard, Xu). — *Let C be a C^* -algebra, let $(B_i)_{i \in I}$ be unital C^* -algebras together with GNS-faithful conditional expectations $\mathbb{E}_{B_i} : B_i \rightarrow C$. Let $A_i \subset B_i$ be unital C^* -subalgebras with GNS-faithful conditional expectations $\mathbb{E}_{A_i} : A_i \rightarrow C$ which are the restrictions of \mathbb{E}_{B_i} . Assume that for each i , there is a net of finite-rank maps $(V_{i,j})_j$ on A_i converging to the identity pointwise, satisfying $\mathbb{E}_{A_i} \circ V_{i,j} = \mathbb{E}_{A_i}$ and such that $\limsup_j \|V_{i,j}\|_{cb} = 1$. Assume moreover that for each pair (i, j) , there is a completely positive unital map $U_{i,j} : A_i \rightarrow B_i$ satisfying $\mathbb{E}_{B_i} \circ U_{i,j} = \mathbb{E}_{A_i}$ and such that*

$$\begin{aligned} & \|V_{i,j} - U_{i,j}\|_{cb} + \|V_{i,j} - U_{i,j}\|_{\mathcal{B}(L^2(A_i, \mathbb{E}_{A_i}), L^2(B_i, \mathbb{E}_{B_i}))} \\ & + \|V_{i,j} - U_{i,j}\|_{\mathcal{B}(L^2(A_i, \mathbb{E}_{A_i})^{op}, L^2(B_i, \mathbb{E}_{B_i})^{op})} \xrightarrow{j} 0. \end{aligned}$$

*Assume moreover that the maps $V_{i,j}$ and $U_{i,j}$ are the identity on C for all i, j . Then, the reduced amalgamated free product $*_C(A_i, \mathbb{E}_{A_i})$ has Cowling-Haagerup constant equal to 1.*

Remark 3.15. — This statement is the same as [31, Prop 4.11] with states replaced by conditional expectations. This amalgamated version does not appear explicitly in [31] but is a straightforward consequence of the amalgamated version of the Khintchine inequality proved in [31, Sec 5]. The idea of the proof is that the "free product" of the maps $V_{i,j}$, which does not make sense *a priori* can be defined using the free product of the u.c.p. maps $U_{i,j}$ and the approximation assumption. Let us note that it is necessary to have bigger C^* -algebras B_i as ranges for the completely positive maps $U_{i,j}$, otherwise we would be assuming that the C^* -algebras A_i are nuclear.

Based on the non-amalgamated case [21], one can try the following strategy : if (a_t) is a net of elements implementing weak amenability, first average a_t and then $S(a_t)^*$ over $\widehat{\mathbb{H}}$ to produce an element b'_t in $\ell^\infty(\widehat{\mathbb{G}})$ and two maps η'_t and ξ'_t in $\mathcal{B}(L^2(\mathbb{G}), L^2(\mathbb{G}) \otimes K)$ satisfying

$$m_{b'_t}(x) = (\eta'_t)^*(x \otimes 1)\xi'_t$$

and such that the multiplier $m_{b'_t}$ is the identity on $C_{\text{red}}(\mathbb{H})$. Then, mimicking [21, Lem 4.3], setting $\gamma'_t = (\eta'_t + \xi'_t)/2$ and $\tilde{\gamma}'_t = \gamma'_t|\gamma'_t|^{-1}$, the unital completely positive approximation we are looking for should be

$$M_{\tilde{\gamma}'_t}(x) = (\tilde{\gamma}'_t)^*(x \otimes 1)\tilde{\gamma}'_t.$$

Note that $\|1 - (\gamma'_t)^*\gamma'_t\| \leq \|m_{b'_t}\|_{cb} - 1$ so that $(\gamma'_t)^*\gamma'_t$ is invertible if $m_{b'_t}$ is sufficiently close to the identity in completely bounded norm. The problem is then to prove that this operator is the identity on $C_{\text{red}}(\mathbb{H})$. We do not know whether this fact holds or not. However, we can use again an averaging trick, this time at the level of C^* -algebras, to build a new unital completely positive approximation which will be the identity on $C_{\text{red}}(\mathbb{H})$. If $T : C_{\text{red}}(\mathbb{G}) \rightarrow \mathcal{B}(L^2(\mathbb{G}))$ is a linear map, we define a linear map $R_{\widehat{\mathbb{H}}}(T)$ by

$$R_{\widehat{\mathbb{H}}}(T) : x \mapsto \int_{\mathcal{U}(C_{\text{red}}(\mathbb{H}))} T(xv^*)v dv,$$

where the integration is done with respect to the normalized Haar measure of the compact group $\mathcal{U}(C_{\text{red}}(\mathbb{H}))$ (recall that $C_{\text{red}}(\mathbb{H})$ is finite-dimensional). Similarly, we define a linear map $L_{\widehat{\mathbb{H}}}(T)$ by

$$L_{\widehat{\mathbb{H}}}(T) : x \mapsto \int_{\mathcal{U}(C_{\text{red}}(\mathbb{H}))} u^*T(ux)du.$$

Let us give some elementary properties of these two operations.

LEMMA 3.16. — *If T is completely bounded, then $R_{\widehat{\mathbb{H}}}(T)$ (resp. $L_{\widehat{\mathbb{H}}}(T)$) is also completely bounded with $\|R_{\widehat{\mathbb{H}}}(T)\|_{cb} \leq \|T\|_{cb}$ (resp. $\|L_{\widehat{\mathbb{H}}}(T)\|_{cb} \leq \|T\|_{cb}$). Moreover, for any $a, b \in C_{\text{red}}(\mathbb{H})$,*

$$(3.3) \quad R_{\widehat{\mathbb{H}}} \circ L_{\widehat{\mathbb{H}}}(T)(axb) = a[R_{\widehat{\mathbb{H}}} \circ L_{\widehat{\mathbb{H}}}(T)(x)]b.$$

Proof. — Let us prove the first part for $R_{\mathbb{H}}^{\wedge}(T)$ (the computation is similar for $L_{\mathbb{H}}^{\wedge}(T)$). For any integer n and any $x \in C_{\text{red}}(\mathbb{G}) \otimes M_n(\mathbb{C})$, we have

$$\begin{aligned} \|(R_{\mathbb{H}}^{\wedge}(T) \otimes \text{Id}_{M_n(\mathbb{C})})(x)\| &\leq \int_{\mathcal{U}(C_{\text{red}}(\mathbb{H}))} \|(T \otimes \text{Id}_{M_n(\mathbb{C})})(x(v^* \otimes 1))(v \otimes 1)\| dv \\ &\leq \int_{\mathcal{U}(C_{\text{red}}(\mathbb{H}))} \|T\|_{cb} \|x\| dv \\ &= \|T\|_{cb} \|x\|. \end{aligned}$$

Writing any element in $C_{\text{red}}(\mathbb{H})$ as a linear combination of four unitaries (up to a scalar), we can restrict ourselves to prove Equation (3.3) when a and b are unitaries. In that case, the changes of variables $u = u'a$ and $v = v'b$ yield

$$\begin{aligned} R_{\mathbb{H}}^{\wedge} \circ L_{\mathbb{H}}^{\wedge}(T)(a^*xb) &= \iint_{\mathcal{U}(C_{\text{red}}(\mathbb{H})) \times \mathcal{U}(C_{\text{red}}(\mathbb{H}))} u^*T(ua^*xbv^*)vdudv \\ &= \iint_{\mathcal{U}(C_{\text{red}}(\mathbb{H})) \times \mathcal{U}(C_{\text{red}}(\mathbb{H}))} a^*(u')^*T(u'x(v')^*)v'bdu'dv' \\ &= a^*[R_{\mathbb{H}}^{\wedge} \circ L_{\mathbb{H}}^{\wedge}(T)(x)]b. \end{aligned}$$

□

With this in hand, we will be able to average the completely positive maps approximating the multipliers. Let us check that this averaging behaves nicely on the multipliers $m_{b'(t)}$.

LEMMA 3.17. — *The maps $A_t = R_{\mathbb{H}}^{\wedge} \circ L_{\mathbb{H}}^{\wedge}(m_{b'(t)})$ have finite rank, converge pointwise to the identity, are equal to the identity on $C_{\text{red}}(\mathbb{H})$ and satisfy $\limsup_t \|A_t\|_{cb} = \limsup_t \|m_{b'(t)}\|_{cb}$.*

Proof. — The pointwise convergence, the identity property and the bound on the completely bounded norms follow from the construction and Lemma 3.16. To prove that the rank is finite, first note that if $\alpha \in \text{Irr}(\mathbb{G})$ and $u, v \in C_{\text{red}}(\mathbb{H})$, then $u(u_{i,j}^\alpha)v$ belongs to the linear span of coefficients of irreducible subrepresentations γ of $\beta_1 \otimes \alpha \otimes \beta_2$ for $\beta_1, \beta_2 \in \text{Irr}(\mathbb{H})$. Thus, by Frobenius reciprocity,

$$A_t(u_{i,j}^\alpha) = \iint_{\mathcal{U}(C_{\text{red}}(\mathbb{H})) \times \mathcal{U}(C_{\text{red}}(\mathbb{H}))} u^*m_{b'(t)}(u(u_{i,j}^\alpha)v^*)vdudv$$

is equal to 0 as soon as α is not in the finite the set

$$\bigcup_{\beta_1, \beta_2 \in \text{Irr}(\mathbb{H})} \bigcup_{\gamma \in \text{Supp}(b'(t))} \{\alpha \in \text{Irr}(\mathbb{G}), \bar{\alpha} \in \beta_2 \otimes \bar{\gamma} \otimes \beta_1\}.$$

□

The following theorem is the the best known statement on stability of weak amenability with respect to free products for discrete quantum groups. Note that, to our knowledge, it is also new for classical groups, even though it may be well-known to experts.

THEOREM 3.18. — *Let $(\widehat{\mathbb{G}}_i)_{i \in I}$ be a family of weakly amenable discrete quantum groups such that $\Lambda_{cb}(\widehat{\mathbb{G}}_i) = 1$ for every $i \in I$ and let $\widehat{\mathbb{H}}$ be a common finite quantum subgroup. Then,*

$$\Lambda_{cb}(*\widehat{\mathbb{G}}_i) = 1.$$

Proof. — Using the notations of Theorem 3.14, we set :

- $A_i = C_{\text{red}}(\mathbb{G}_i)$
- $B_i = \mathcal{B}(L^2(\mathbb{G}_i))$
- $C = C_{\text{red}}(\mathbb{H})$
- $\mathbb{E}_{A_i} = \mathbb{E}_{\mathbb{H}}$

To define the conditional expectations \mathbb{E}_{B_i} , first consider the orthogonal projection

$$P_{\mathbb{H}}^i : L^2(\mathbb{G}_i) \rightarrow L^2(\mathbb{H}).$$

Then, $\mathbb{E}'_i : x \mapsto P_{\mathbb{H}}^i x P_{\mathbb{H}}^i$ is a conditional expectation from $\mathcal{B}(L^2(\mathbb{G}_i))$ to $\mathcal{B}(L^2(\mathbb{H}))$ with the property that for any coefficient x of an irreducible representation in $\text{Irr}(\mathbb{G}_i) \setminus \text{Irr}(\mathbb{H})$, $\mathbb{E}'_i(x) = 0$. In fact, the restriction of \mathbb{E}'_i to $C_{\text{red}}(\mathbb{G}_i)$ is precisely the conditional expectation $\mathbb{E}_{\mathbb{H}}$ of Proposition 2.6. Because $C_{\text{red}}(\mathbb{H})$ is finite-dimensional, there is also a conditional expectation

$$\mathbb{E}''_i : \mathcal{B}(L^2(\mathbb{H})) \rightarrow C_{\text{red}}(\mathbb{H}).$$

We set $\mathbb{E}_{B_i} = \mathbb{E}''_i \circ \mathbb{E}'_i$. Since \mathbb{E}_{A_i} is the restriction of \mathbb{E}'_i to A_i it is also the restriction of \mathbb{E}_{B_i} .

Let us fix an index i , let $(a_j)_j$ be a net of elements in $\ell^\infty(\mathbb{G}_i)$ implementing weak amenability. Applying the construction of Lemma 3.13 to a_j we get an element b_j . Applying it again to $S(b_j)^*$ yields another element b'_j and we can assume that $b'_j = S(b'_j)^*$ as in the proof of [21, Thm 4.2]. We therefore set $V_{i,j} = R_{\widehat{\mathbb{H}}} \circ L_{\widehat{\mathbb{H}}}(m_{b'_j})$. By Lemma 3.17, these maps satisfy all the required properties. Recall that $\gamma'_j = (\xi'_j + \eta'_j)/2$ and $\tilde{\gamma}'_j = \gamma'_j |\gamma'_j|^{-1}$. Set

$$\zeta_j = \int_{\mathcal{U}(C_{\text{red}}(\mathbb{H}))} (1 \otimes u^*) \tilde{\gamma}'_j u du$$

and observe that

$$R_{\widehat{\mathbb{H}}} \circ L_{\widehat{\mathbb{H}}}(M_{\tilde{\gamma}'_j}) = M_{\zeta_j} : x \mapsto \zeta_j^* (1 \otimes x) \zeta_j$$

is a completely positive map which is the identity on $C_{\text{red}}(\mathbb{H})$ by Equation (3.3). We therefore set $U_{i,j} = M_{\zeta_j}$. Then, Lemma 3.17 yields

$$\|V_{i,j} - U_{i,j}\|_{cb} = \|R_{\mathbb{H}} \circ L_{\mathbb{H}}(m_{b'_j} - M_{\tilde{\gamma}_j})\|_{cb} \leq \|m_{b'_j} - M_{\tilde{\gamma}_j}\|_{cb},$$

so that, by [21, Lem 4.3] the convergence in completely bounded norm holds. We still have to check the compatibility of the maps with the conditional expectations and the L^2 -norm convergence.

Let us consider the conditional expectation

$$\mathbb{E} = \mathbb{E}_{B_i} \circ U_{i,j} : C_{\text{red}}(\mathbb{G}_i) \rightarrow C_{\text{red}}(\mathbb{H}).$$

We claim that $\mathbb{E}(x) = 0$ whenever x is a coefficient of a representation in $\text{Irr}(\mathbb{G}) \setminus \text{Irr}(\mathbb{H})$. In fact, it follows directly from the explicit expression of ξ'_j given in Lemma 3.11 that there is a vector $v_{\xi'_j} \in K$ such that for any $y \in C_{\text{red}}(\mathbb{H})$, $\xi'_j(y) = y \otimes v_{\xi'_j}$. The same holds for η'_j and γ'_j (with different vectors), hence also for $\tilde{\gamma}'_j$. From this, a straightforward calculation yields

$$(3.4) \quad P_{\mathbb{H}} M_{\tilde{\gamma}'_j}(x) P_{\mathbb{H}} = 0.$$

Since elements in $\mathcal{U}(C_{\text{red}}(\mathbb{H}))$ respect the decomposition $L^2(\mathbb{G}) = L^2(\mathbb{H}) \oplus L^2(\mathbb{H})^\perp$, Equation (3.4) also holds for M_{ζ_j} , proving the claim. Since $\mathbb{E}(x) = x$ for any $x \in C_{\text{red}}(\mathbb{H})$, we see that $\mathbb{E} = \mathbb{E}_{\mathbb{H}}$. The same argument shows that $\mathbb{E}_{A_i} \circ V_{i,j} = \mathbb{E}_{A_i}$.

Using this compatibility, the same argument as in [21, Lem 4.5] shows that $M_{\tilde{\gamma}'_j}$ also approximates $m_{b'_j}$ in both L^2 -norms (here L^2 -norms means norms as operator between the Hilbert modules associated to the conditional expectations). The proof of Lemma 3.16 also works for the L^2 -norms so that we can conclude that $U_{i,j}$ approximates $V_{i,j}$ in both L^2 -norms, concluding the proof. \square

In the unimodular case, one can use a free product trick to deduce results on HNN extensions (as defined in [19]) from what has been done. The optimal result for unimodular quantum groups, mixing amalgamated free products and HNN extensions, was stated in the language of graphs of quantum groups in [20, Thm 4.6].

It is known that $\mathbb{Z}^2 \rtimes SL(2, \mathbb{Z})$ neither has the Haagerup property nor is weakly amenable, whereas both \mathbb{Z}^2 and $SL(2, \mathbb{Z})$ do. Thus, these properties are not preserved under extensions in general. Moreover, recall that since an extension of amenable groups is again amenable, the groups $\mathbb{Z}^2 \rtimes \mathbb{Z}/4\mathbb{Z}$ and $\mathbb{Z}^2 \rtimes \mathbb{Z}/6\mathbb{Z}$ are amenable. Thus, they have the Haagerup property and are weakly amenable with Cowling-Haagerup constant equal to 1. However,

noticing that

$$\mathbb{Z}^2 \rtimes SL(2, \mathbb{Z}) = (\mathbb{Z}^2 \rtimes \mathbb{Z}/4\mathbb{Z}) \underset{\mathbb{Z}^2 \rtimes \mathbb{Z}/2\mathbb{Z}}{*} (\mathbb{Z}^2 \rtimes \mathbb{Z}/6\mathbb{Z}),$$

we see that the finiteness condition in Theorem 3.18 cannot be removed.

4. Relative amenability

We now turn to the study of the notion of relative amenability in the context of discrete quantum groups. The definition is straightforward, recalling that τ is the action of $\widehat{\mathbb{G}}$ on the quotient.

DEFINITION 4.1. — *Let $\widehat{\mathbb{G}}$ be discrete quantum group and let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup of $\widehat{\mathbb{G}}$. We say that $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$ if the quotient space has an invariant mean for the action τ , i.e. if there exists a state m on $\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ such that for all $x \in \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$,*

$$(\iota \otimes m) \circ \tau(x) = m(x).1$$

Remark 4.2. — One could define an action of a discrete (or even locally compact) quantum group on a von Neumann algebra to be *amenable* if such an invariant mean exists. This is the notion of "amenable homogeneous space" introduced by P. Eymard in [17]. However, there is another notion of amenable action, due to R.J. Zimmer [45]. This notion was generalized to locally compact groups acting on von Neumann algebras by C. Anantharaman-Delaroche in [1] and to Kac algebras by M. Joita and S. Petrescu in [25]. As one can expect, this notion is dual to ours in the following sense : if \mathbb{G} is a Kac algebra together with an amenable action (in the sense of [25, Def 3.1]) on a von Neumann algebra M and if M admits an invariant state, then \mathbb{G} is amenable (see [25, Thm 3.5] for a proof). Let us also mention that a C*-algebraic version of amenable actions of discrete quantum groups in the sense of R.J. Zimmer was introduced by S. Vaes and R. Vergnioux in [38].

Remark 4.3. — Relative amenability does not pass to subgroups, even in the classical case. Examples of triples of discrete groups $K < H < G$ with G amenable relative to K but H not amenable relative to K were constructed in [29] and [30].

It follows from [7, Thm 7.8] that if the quasi-regular representation of $\widehat{\mathbb{G}}$ modulo $\widehat{\mathbb{H}}$ weakly contains the trivial one, then it has a left-invariant mean, implying in turn that $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$. In the classical

case, all three properties are known to be equivalent. Such an equivalence is not known yet for discrete quantum groups, but a weak converse involving correspondences will be given in Proposition 4.9.

4.1. Amenable equivalence

We are going to use ideas from the work of C. Anantharaman-Delaroche on group actions on von Neumann algebras [1] and follow the path of [2] to prove a general statement on von Neumann algebras associated to relatively amenable discrete quantum groups, involving the notion of *amenable equivalence* of von Neumann algebras which was introduced in [2, Def 4.1]. Recall from [32] that two von Neumann algebras M and N are said to be *Morita equivalent* if there exists an $M - N$ correspondence H such that M is isomorphic to $\mathcal{L}_N(H)$.

DEFINITION 4.4. — *Let M and N be two von Neumann algebras. We say that M is amenablely dominated by N if there exists a von Neumann algebra N_1 which is Morita equivalent to N and contains M in such a way that there is a norm-one projection from N_1 to M . We then write $M \prec_a N$. We say that M and N are amenablely equivalent if $M \prec_a N$ and $N \prec_a M$.*

The following theorem is a generalization of a classical result (see Paragraphe 4.10 in [2]).

THEOREM 4.5. — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group and let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup such that $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$. Then, $L^\infty(\mathbb{H})$ is amenablely equivalent to $L^\infty(\mathbb{G})$.*

In order to prove this theorem, we have to build a norm-one projection from a well-chosen von Neumann algebra to $L^\infty(\mathbb{G})$. We will do this by adapting some ideas of [1] to the setting of discrete quantum groups.

LEMMA 4.6. — *Let M_1, M_2 be von Neumann algebras and let ρ be an action of a discrete quantum group $\widehat{\mathbb{G}}$ on M_2 . Assume that we have a von Neumann subalgebra N_2 of M_2 which is stable under the action ρ and a norm-one (non-necessarily normal) equivariant surjective projection $P : M_2 \rightarrow N_2$ (i.e. $\rho \circ P = (\iota \otimes P) \circ \rho$). Then, there exists a norm-one surjective projection $Q : M_1 \overline{\otimes} M_2 \rightarrow M_1 \overline{\otimes} N_2$ such that*

$$Q(x_1 \otimes x_2) = x_1 \otimes P(x_2)$$

for all $x_1 \in M_1$ and $x_2 \in M_2$. Moreover, Q is equivariant with respect to $\mu = (\sigma \otimes \iota) \circ (\iota \otimes \rho)$.

Proof. — The existence of the projection Q is proved in [35, Thm 4] (we thank the referee for pointing out to us this reference). It is proved in [1, Lem 2.1] that J. Tomiyama’s construction preserves equivariance under a classical group action and the proof for a quantum group action is exactly the same. □

PROPOSITION 4.7. — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group and let (M, N, P) be a triple consisting in a von Neumann algebra M endowed with an action ρ of $\widehat{\mathbb{G}}$, a von Neumann subalgebra N of M which is stable under the action ρ and a norm-one equivariant projection $P : M \rightarrow N$. Then, there exists a norm-one projection $\widetilde{Q} : \widehat{\mathbb{G}} \times M \rightarrow \widehat{\mathbb{G}} \times N$.*

Proof. — By Lemma 4.6, there exists a norm-one projection

$$Q : \mathcal{B}(L^2(\mathbb{G})) \overline{\otimes} M \rightarrow \mathcal{B}(L^2(\mathbb{G})) \overline{\otimes} N$$

which is equivariant with respect to $\mu = (\sigma \otimes \iota) \circ (\iota \otimes \rho)$ and such that

$$Q(x_1 \otimes x_2) = x_1 \otimes P(x_2).$$

Let us consider the explicit $*$ -isomorphisms of [36, Thm 2.6]

$$\begin{aligned} \Phi_M : \mathcal{B}(L^2(\mathbb{G})) \overline{\otimes} M &\rightarrow \mathbb{G}^{op} \times (\widehat{\mathbb{G}} \times M) \\ \Phi_N : \mathcal{B}(L^2(\mathbb{G})) \overline{\otimes} N &\rightarrow \mathbb{G}^{op} \times (\widehat{\mathbb{G}} \times N) \end{aligned}$$

Since any von Neumann algebra can be recovered in a crossed-product as the fixed points algebra under the dual action by [36, Thm 2.7], we only have to prove that the norm-one projection

$$\widetilde{Q} = \Phi_N \circ Q \circ \Phi_M^{-1} : \mathbb{G}^{op} \times (\widehat{\mathbb{G}} \times M) \rightarrow \mathbb{G}^{op} \times (\widehat{\mathbb{G}} \times N)$$

is equivariant with respect to the the bidual action to conclude.

Let us use the notations of [36]. The bidual action $\widehat{\rho}$ on the double crossed product can be transported to $\mathcal{B}(L^2(\mathbb{G})) \overline{\otimes} M$ in the following way : there is an operator

$$\mathcal{J} : L^\infty(\mathbb{G}) \rightarrow L^\infty(\mathbb{G}'^{op})$$

and a map $\gamma = \text{Ad}_{\Sigma V^* \Sigma \otimes 1} \circ \mu$, for some operator V , such that

$$\widehat{\rho} \circ \Phi = (\mathcal{J} \otimes \Phi) \circ \gamma.$$

It is clear that the $\widehat{\rho}$ -equivariance of \widetilde{Q} is equivalent to the γ -equivariance of Q . We already know by Lemma 4.6 that

$$\mu \circ Q = (\iota \otimes Q) \circ \mu$$

and, using the approximating projections P'_j , we see that $\iota \otimes Q$ also commutes to $\text{Ad}_{\Sigma V^* \Sigma \otimes 1}$. Hence,

$$\gamma \circ Q = (\iota \otimes Q) \circ \gamma.$$

□

We can now prove our main result.

Proof of Theorem 4.5. — The fact that the mean is invariant precisely means that m is an equivariant norm-one projection to the von Neumann subalgebra $\mathbb{C}.1$ of M . Consequently, proposition 4.7 applied to the triple $(\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}}), \mathbb{C}, m)$ yields a norm-one projection from $\widehat{\mathbb{G}} \rtimes \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ to $\widehat{\mathbb{G}} \rtimes \mathbb{C} = L^\infty(\mathbb{G})$. Since according e.g. to [37, Rmk 4.3], $\widehat{\mathbb{G}} \rtimes \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ is Morita equivalent to $L^\infty(\mathbb{H})$, we have proven that $L^\infty(\mathbb{G}) \prec_a L^\infty(\mathbb{H})$. The conditional expectation from $L^\infty(\mathbb{G})$ to $L^\infty(\mathbb{H})$ defined in [39, Prop 2.2] then gives $L^\infty(\mathbb{H}) \prec_a L^\infty(\mathbb{G})$, concluding the proof. □

Remark 4.8. — The basic construction $\langle L^\infty(\mathbb{G}), L^\infty(\mathbb{H}) \rangle$ is naturally isomorphic to $\widehat{\mathbb{G}} \rtimes \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ (see e.g. the beginning of Section 4 of [37]). This means that if $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$, then $L^\infty(\mathbb{G})$ is amenable relative to $L^\infty(\mathbb{H})$ in the sense of [29].

Using the machinery of correspondences, we can also give a partial converse to the fact that if \mathcal{R} weakly contains the trivial representations, then $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$.

PROPOSITION 4.9. — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group and let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup of $\widehat{\mathbb{G}}$ such that $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$. Then, the correspondence associated to the quasi-regular representation of $\widehat{\mathbb{G}}$ modulo $\widehat{\mathbb{H}}$ weakly contains the identity correspondence.*

Proof. — Let us denote by H the Hilbert space $L^2(\mathbb{G})$ seen as the standard correspondence between $L^\infty(\mathbb{G})$ and $L^\infty(\mathbb{H})$. By Theorem 4.5, H is a *left injective correspondence* in the sense of [2, Def 3.1]. By [2, Prop 3.6], it is also a *left amenable correspondence* in the sense of [2, Def 2.1], i.e. the correspondence $H \otimes \overline{H}$ weakly contains the identity correspondence of $L^\infty(\mathbb{G})$. Moreover, $H \otimes \overline{H}$ is precisely the Hilbert space $\ell^2(\widehat{\mathbb{G}} \rtimes \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}}))$ associated to the GNS construction for the dual weight $\tilde{\theta}$ on the crossed-product. Thanks to [36, Prop 3.10], the GNS construction for the dual weight may be explicitly described. In fact, the map

$$\mathcal{I} : (a \otimes 1)\alpha(x) \mapsto a \otimes x$$

for $a \in L^\infty(\mathbb{G})$ and $x \in \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ extends to an isomorphism between $\ell^2(\widehat{\mathbb{G}} \rtimes \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}}))$ and $L^2(\mathbb{G}) \otimes \ell^2(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$. Recall that \mathcal{R} denotes the adjoint of the unitary implementation of τ . Let us endow the latter Hilbert space with the structure of a correspondence from $L^\infty(\mathbb{G})$ to itself induced by the quasi-regular representation, i.e. the left action π_l and the right action

π_r are given, for every $a \in L^\infty(\mathbb{G})$, by

$$\pi_l(a) = \mathcal{R}^*(a \otimes 1)\mathcal{R} \text{ and } \pi_r(a) = (Ja^*J) \otimes 1.$$

Then, the previous isomorphism intertwines these actions with the natural left and right actions of $L^\infty(\mathbb{G})$ on $\ell^2(\widehat{\mathbb{G}} \times \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}}))$ (inherited from its identification with $H \otimes \overline{H}$). Thus, the correspondence associated with the quasi-regular representation weakly contains the identity correspondence. □

4.2. Finite index quantum subgroups

The simplest source of examples of relatively amenable discrete quantum subgroups is of course *finite index* quantum subgroups.

DEFINITION 4.10. — *A discrete quantum subgroup $\widehat{\mathbb{H}}$ of a discrete quantum group $\widehat{\mathbb{G}}$ is said to have finite index if the quotient von Neumann algebra $\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ is finite-dimensional.*

PROPOSITION 4.11. — *Let $\widehat{\mathbb{H}}$ be a finite index quantum subgroup of a discrete quantum group $\widehat{\mathbb{G}}$. Then, $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$.*

Proof. — Let us prove that the weight θ yields an invariant state. Indeed, the operator-valued weight $T : \ell^\infty(\widehat{\mathbb{G}}) \rightarrow \ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ is finite in that case and we can therefore only consider elements of the form $T(x)$. Using the equality $h_L = \theta \circ T$, we have

$$(\iota \otimes \theta) \circ \tau(T(x)) = (\iota \otimes (\theta \circ T)) \circ \tau(x) = (\iota \otimes h_L) \circ \widehat{\Delta}(x) = h_L(x).1 = \theta(T(x)).1$$

Thus, $\theta(1)^{-1}\theta$ is an invariant mean for τ . □

Example 4.12. — Let $N \in \mathbb{N}$, let \mathbb{G} be the free orthogonal quantum group O_N^+ and let $u = (u_{ij})_{1 \leq i, j \leq N}$ be its fundamental representation. If $N = 2$, its dual discrete quantum group $\widehat{\mathbb{G}}$ is amenable, hence it is amenable relative to any quantum subgroup. If $N \geq 3$, consider the subalgebra of $L^\infty(O_N^+)$ generated by the elements $u_{ij}u_{kl}$ for all i, j, k, l . This subalgebra is stable under the coproduct and thus defines a discrete quantum subgroup $\widehat{\mathbb{H}}$ of $\widehat{\mathbb{G}}$ called its *even part*. It is clear that under the usual identification $\text{Irr}(\mathbb{G}) = \mathbb{N}$, $\text{Irr}(\mathbb{H})$ corresponds to the even integers and it is not very difficult to see that $\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}}) = \mathbb{C} \oplus \mathbb{C}$. Thus $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$. Note that this quantum group is isomorphic to the quantum automorphism group of $M_N(\mathbb{C})$ (with respect to a suitably chosen trace) according to [5].

As for classical groups, relative amenability becomes equivalent to finite index in the presence of Kazhdan's property (T).

PROPOSITION 4.13. — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group with Kazhdan's property (T) as defined in [18, Def 3.1] and let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup such that $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$. Then, $\widehat{\mathbb{H}}$ has finite index in $\widehat{\mathbb{G}}$.*

Proof. — Recall from Proposition 4.9 that the correspondence associated with the quasi-regular representation weakly contains the identity correspondence. Since $L^\infty(\widehat{\mathbb{G}})$ has property (T) in the sense of [12] by [18, Thm 3.1], it actually contains the identity correspondence. Since any property (T) discrete quantum group is unimodular by [18, Prop 3.2], we can apply [26, Lem 7.1] to conclude that \mathcal{R} has a fixed vector. This implies by [18, Lem 2.3] that the quotient $\ell^\infty(\widehat{\mathbb{G}}/\widehat{\mathbb{H}})$ is finite-dimensional. \square

4.3. Applications

Theorem 4.5 links relative amenability for discrete quantum groups and amenable equivalence of von Neumann algebras. We can thus now use the work of C. Anantharaman-Delaroche [2] on von Neumann algebras to derive permanence results. However, going back to the quantum group is not always possible at the present state of our knowledge, hence the restriction to unimodular quantum groups in the sequel.

COROLLARY 4.14. — *Let $\widehat{\mathbb{G}}$ be a discrete quantum group and let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup such that $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$, then $\Lambda_{cb}(L^\infty(\mathbb{H})) = \Lambda_{cb}(L^\infty(\mathbb{G}))$. If moreover $\widehat{\mathbb{G}}$ (and consequently $\widehat{\mathbb{H}}$) is unimodular, then $\Lambda_{cb}(\widehat{\mathbb{G}}) = \Lambda_{cb}(\widehat{\mathbb{H}})$.*

Proof. — It was proved in [2, Thm 4.9] that amenable equivalent von Neumann algebras have equal Cowling-Haagerup constant. We conclude by [27, Thm 5.14]. \square

Remark 4.15. — A more direct (and C*-algebraic) proof of Corollary 4.14 for groups is given in [11, Prop 12.3.11]. However, it is quite ill-suited to the setting of quantum groups since it is based on the use of a section of the quotient which may fail to exist in a reasonable sense in the quantum case, for example if the subgroup is not *divisible* in the sense of [40, Def 4.1].

A similar statement holds for the Haagerup property.

COROLLARY 4.16. — Let $\widehat{\mathbb{G}}$ be a unimodular discrete quantum group and let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup such that $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$. Then, if $L^\infty(\mathbb{H})$ has the Haagerup property, $L^\infty(\mathbb{G})$ also has the Haagerup property.

Proof. — This is a combination of Theorem 4.5 and [6, Thm 5.1]. \square

We end with an application to hyperlinearity. A unimodular discrete quantum group $\widehat{\mathbb{G}}$ is said to be *hyperlinear* if the von Neumann algebra $L^\infty(\mathbb{G})$ tracially embeds into an ultraproduct of the hyperfinite II_1 factor.

COROLLARY 4.17. — Let $\widehat{\mathbb{G}}$ be a unimodular discrete quantum group and let $\widehat{\mathbb{H}}$ be a discrete quantum subgroup such that $\widehat{\mathbb{G}}$ is amenable relative to $\widehat{\mathbb{H}}$. Then, $\widehat{\mathbb{G}}$ is hyperlinear if and only if $\widehat{\mathbb{H}}$ is hyperlinear.

Proof. — This is a direct consequence of [44, Thm 3.1] since, by Remark 4.8, $L^\infty(\mathbb{G})$ is amenable relative to $L^\infty(\mathbb{H})$. \square

BIBLIOGRAPHY

- [1] C. ANANTHARAMAN-DELAROCHE, “Action moyennable d’un groupe localement compact sur une algèbre de von Neumann”, *Math. Scand* **45** (1979), p. 289-304.
- [2] ———, “Amenable correspondences and approximation properties for von Neumann algebras”, *Pacific J. Math.* **171** (1995), no. 2, p. 309-341.
- [3] S. BAAJ & G. SKANDALIS, “Unitaires multiplicatifs et dualité pour les produits croisés de C^* -algèbres”, *Ann. Sci. École Norm. Sup.* **26** (1993), no. 4, p. 425-488.
- [4] T. BANICA, “Le groupe quantique compact libre $U(n)$ ”, *Comm. Math. Phys.* **190** (1997), no. 1, p. 143-172.
- [5] ———, “Symmetries of a generic coaction”, *Math. Ann.* **314** (1999), no. 4, p. 763-780.
- [6] J. BANNON & J. FANG, “Some remarks on Haagerup’s approximation property”, *J. Operator Theory* **65** (2011), no. 2, p. 403-417.
- [7] E. BÉDOS, R. CONTI & L. TUSET, “On amenability and co-amenability of algebraic quantum groups and their corepresentations”, *Canad. J. Math.* **57** (2005), no. 1, p. 17-60.
- [8] F. BOCA, “On the method of constructing irreducible finite index subfactors of Popa”, *Pacific J. Math.* **161** (1993), no. 2, p. 201-231.
- [9] M. BOŽEJKO & M. PICARDELLO, “Weakly amenable groups and amalgamated products”, *Proc. Amer. Math. Soc.* **117** (1993), no. 4, p. 1039-1046.
- [10] M. BRANNAN, “Approximation properties for free orthogonal and free unitary quantum groups”, *J. Reine Angew. Math.* **672** (2012), p. 223-251.
- [11] N. P. BROWN & N. OZAWA, *C^* -algebras and finite-dimensional approximations*, Graduate Studies in Mathematics, vol. 88, American Mathematical Society, Providence, RI, 2008, xvi+509 pages.
- [12] A. CONNES & V. JONES, “Property T for von Neumann algebras”, *Bull. London Math. Soc.* **17** (1985), no. 1, p. 57-62.
- [13] M. COWLING & U. HAAGERUP, “Completely bounded multipliers of the Fourier algebra of a simple Lie group of real rank one”, *Invent. Math.* **96** (1989), no. 3, p. 507-549.

- [14] M. DAWS, P. FIMA, A. SKALSKI & S. WHITE, “The Haagerup property for locally compact quantum groups”, <http://arxiv.org/abs/1303.3261>, to appear in *J. Reine Angew. Math.*, 2014.
- [15] M. DAWS, “Completely positive multipliers of quantum groups”, *Internat. J. Math.* **23** (2012), no. 12, p. 1250132, 23.
- [16] K. DE COMMER, A. FRESLON & M. YAMASHITA, “CCAP for universal discrete quantum groups”, *Comm. Math. Phys.* **331** (2014), no. 2, p. 677-701.
- [17] P. EYMARD, *Moyennes invariantes et représentations unitaires*, Lecture notes in mathematics, vol. 300, Springer, 1972.
- [18] P. FIMA, “Kazhdan’s property T for discrete quantum groups”, *Internat. J. Math.* **21** (2010), no. 1, p. 47-65.
- [19] ———, “K-amenability of HNN extensions of amenable discrete quantum groups”, *J. Funct. Anal.* **265** (2013), no. 4, p. 507-519.
- [20] P. FIMA & A. FRESLON, “Graphs of quantum groups and K-amenability”, *Adv. Math.* **260** (2014), p. 233-280.
- [21] A. FRESLON, “A note on weak amenability for free products of discrete quantum groups”, *C. R. Acad. Sci. Paris Sér. I Math.* **350** (2012), p. 403-406.
- [22] ———, “Examples of weakly amenable discrete quantum groups”, *J. Funct. Anal.* **265** (2013), no. 9, p. 2164-2187.
- [23] ———, “Propriétés d’approximation pour les groupes quantiques discrets”, PhD Thesis, Université Paris VII (France), 2013.
- [24] ———, “Fusion (semi)rings arising from quantum groups”, *J. Algebra* **417** (2014), p. 161-197.
- [25] M. JOITA & S. PETRESCU, “Amenable actions of Katz algebras on von Neumann algebras”, *Rev. Roumaine Math. Pures Appl.* **35** (1990), no. 2, p. 151-160.
- [26] ———, “Property (T) for Kac algebras”, *Rev. Roumaine Math. Pures Appl.* **37** (1992), no. 2, p. 163-178.
- [27] J. KRAUS & Z.-J. RUAN, “Approximation properties for Kac algebras”, *Indiana Univ. Math. J.* **48** (1999), no. 2, p. 469-535.
- [28] F. LEMEUX, “Fusion rules for some free wreath product quantum groups and applications”, *J. Funct. Anal.* **267** (2014), no. 7, p. 2507-2550.
- [29] N. MONOD & S. POPA, “On co-amenability for groups and von Neumann algebras”, *C. R. Math. Acad. Sci. Soc. R. Can.* **25** (2003), no. 3, p. 82-87.
- [30] V. PESTOV, “On some questions of Eymard and Bekka concerning amenability of homogeneous spaces and induced representations”, *C. R. Math. Acad. Sci. Soc. R. Can.* **25** (2003), no. 3, p. 76-81.
- [31] E. RICARD & Q. XU, “Khinchine type inequalities for reduced free products and applications”, *J. Reine Angew. Math.* **599** (2006), p. 27-59.
- [32] M. RIEFFEL, “Morita equivalence for C*-algebras and W*-algebras”, *J. Pure Appl. Algebra* **5** (1974), no. 1, p. 51-96.
- [33] T. TIMMERMANN, *An invitation to quantum groups and duality. From Hopf algebras to multiplicative unitaries and beyond*, EMS, 2008.
- [34] R. TOMATSU, “Amenable discrete quantum groups”, *J. Math. Soc. Japan* **58** (2006), no. 4, p. 949-964.
- [35] J. TOMIYAMA, “On tensor products of von Neumann algebras”, *Pacific J. Math.* **30** (1969), no. 1, p. 263-270.
- [36] S. VAES, “The unitary implementation of a locally compact quantum group action”, *J. Funct. Anal.* **180** (2001), no. 2, p. 426-480.
- [37] ———, “A new approach to induction and imprimitivity results”, *J. Funct. Anal.* **229** (2005), no. 2, p. 317-374.

- [38] S. VAES & R. VERGNIoux, “The boundary of universal discrete quantum groups, exactness and factoriality”, *Duke Math. J.* **140** (2007), no. 1, p. 35-84.
- [39] R. VERGNIoux, “K-amenability for amalgamated free products of amenable discrete quantum groups”, *J. Funct. Anal.* **212** (2004), no. 1, p. 206-221.
- [40] R. VERGNIoux & C. VOIGT, “The K-theory of free quantum groups”, *Math. Ann.* **357** (2013), no. 1, p. 355-400.
- [41] S. WANG, “Free products of compact quantum groups”, *Comm. Math. Phys.* **167** (1995), no. 3, p. 671-692.
- [42] ———, “Tensor products and crossed-products of compact quantum groups”, *Proc. London Math. Soc.* **71** (1995), no. 3, p. 695-720.
- [43] S. WORONOWICZ, “Compact quantum groups”, *Symétries quantiques (Les Houches, 1995)* (1998), p. 845-884.
- [44] J. WU, “Co-amenability and Connes’s embedding problem”, *Sci. China Math.* **55** (2012), no. 5, p. 977-984.
- [45] R. ZIMMER, “Amenable ergodic group actions and an application to Poisson boundaries of random walks”, *J. Funct. Anal.* **27** (1978), no. 3, p. 350-372.

Manuscrit reçu le 4 août 2014,
révisé le 12 octobre 2014,
accepté le 23 octobre 2014.

Amaury FRESLON
Univ. Paris Diderot, Sorbonne Paris Cité
IMJ-PRG, UMR 7586 CNRS
Sorbonne Universités UPMC Univ. Paris 06
F-75013, Paris (France)
amaury.freslon@imj-prg.fr