



ANNALES

DE

L'INSTITUT FOURIER

Phillip S. HARRINGTON & Andrew S. RAICH

Closed Range for $\bar{\partial}$ and $\bar{\partial}_b$ on Bounded Hypersurfaces in Stein Manifolds

Tome 65, n° 4 (2015), p. 1711-1754.

http://aif.cedram.org/item?id=AIF_2015__65_4_1711_0



© Association des Annales de l'institut Fourier, 2015,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

CLOSED RANGE FOR $\bar{\partial}$ AND $\bar{\partial}_b$ ON BOUNDED HYPERSURFACES IN STEIN MANIFOLDS

by Phillip S. HARRINGTON & Andrew S. RAICH (*)

ABSTRACT. — We define weak $Z(q)$, a generalization of $Z(q)$ on bounded domains Ω in a Stein manifold M^n that suffices to prove closed range of $\bar{\partial}$. Under the hypothesis of weak $Z(q)$, we also show (i) that harmonic $(0, q)$ -forms are trivial and (ii) if $\partial\Omega$ satisfies weak $Z(q)$ and weak $Z(n - 1 - q)$, then $\bar{\partial}_b$ has closed range on $(0, q)$ -forms on $\partial\Omega$. We provide examples to show that our condition contains examples that are excluded from $(q - 1)$ -pseudoconvexity and the authors' previous notion of weak $Z(q)$.

RÉSUMÉ. — Nous définissons $Z(q)$ faible, une généralisation de $Z(q)$ sur les domaines bornés Ω dans une variété de Stein M^n qui suffit à prouver que l'image de $\bar{\partial}$ est fermée. Sous l'hypothèse d'une $Z(q)$ faible, nous montrons également que (i) les $(0, q)$ -formes harmoniques sont triviales et (ii) si $\partial\Omega$ satisfait une $Z(q)$ faible et une $Z(n - 1 - q)$ faible, alors $\bar{\partial}_b$ a une image fermée sur les $(0, q)$ -formes sur $\partial\Omega$. Nous fournissons des exemples pour montrer que notre condition contient des exemples qui sont exclus de la $(q - 1)$ -pseudoconvexité et la notion précédente des auteurs de $Z(q)$ faible.

1. Introduction

The purpose of this article is to establish sufficient conditions for the closed range of $\bar{\partial}$ (and $\bar{\partial}_b$) on not necessarily pseudoconvex domains (and their boundaries) in Stein manifolds. We pay particular attention to keeping the boundary regularity at a minimum; our results hold for C^3 boundaries. In [9], we develop a notion of weak $Z(q)$ for which we can prove closed range of $\bar{\partial}_b$ for smooth bounded CR manifolds of hypersurface type in \mathbb{C}^n . In this

Keywords: Stein manifold, $\bar{\partial}_b$, tangential Cauchy-Riemann operator, closed range, $\bar{\partial}$ -Neumann, weak $Z(q)$, q -pseudoconvexity.

Math. classification: 32W05, 32W10, 32Q28, 35N15.

(*) The first author was partially supported by NSF grant DMS-1002332 and the second author was partially supported by NSF grant DMS-0855822 and is partially supported by a grant from the Simons Foundation (#280164 to Andrew Raich).

paper, we generalize our notion of weak $Z(q)$ and relax the smoothness assumption. The microlocal analysis technique of [9] requires significant boundary smoothness, and to replace the microlocal analysis, we assume that our CR manifold is the boundary of a bounded domain in a Stein manifold and attain the closed range of $\bar{\partial}_b$ as a consequence of the $\bar{\partial}$ -theory which we prove. Our analysis of $\bar{\partial}_b$ is in the spirit of [22]. Additionally, we show that the weak $Z(q)$ -hypothesis is sufficient to show that harmonic forms vanish at level $(0, q)$. Finally, we provide examples to show that our new condition is more general than either $(q - 1)$ -pseudoconvexity [27] or the weak $Z(q)$ -hypothesis of [9].

Let $\Omega \subset \mathbb{C}^n$ be a bounded domain with a C^2 defining function ρ . The Levi form of $\partial\Omega$ is the form

$$\mathcal{L}_\rho(t) = \sum_{j,k=1}^n \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_k} t_j \bar{t}_k, \text{ where } \sum_{j=1}^n \frac{\partial \rho}{\partial z_j} t_j = 0.$$

If the Levi form is positive semi-definite for all boundary points, we say Ω is *pseudoconvex*. Suppose that f is a $(0, q)$ -form on Ω with components in L^2 . If $\bar{\partial}f = 0$, where $\bar{\partial}$ is the Cauchy-Riemann operator, we wish to know whether there exists a $(0, q-1)$ -form u with L^2 components such that $\bar{\partial}u = f$. When Ω is a pseudoconvex domain, this question was answered in the affirmative for all $1 \leq q \leq n$ by Hörmander in [13]. In fact, pseudoconvexity is a necessary condition to solve $\bar{\partial}$ in L^2 for all $1 \leq q \leq n$.

If we only wish to solve $\bar{\partial}$ for a fixed value of q , pseudoconvexity is no longer necessary. If the Levi form has at least $n - q$ positive eigenvalues or at least $q + 1$ negative eigenvalues, we say that $\partial\Omega$ satisfies $Z(q)$. It is known that the $\bar{\partial}$ problem can be solved in L^2 if $\partial\Omega$ satisfies $Z(q)$. In fact, $Z(q)$ is equivalent to the solvability of $\bar{\partial}u = f$ if the components of u are required to be elements of the L^2 Sobolev space $W^{1/2}$ (see [13], Theorem 3.2.2 in [7], or [1]).

If we allow the Levi form to degenerate as in the pseudoconvex case, solvability is less well understood. The most natural condition would replace “positive” and “negative” in the definition of $Z(q)$ with “nonnegative” and “nonpositive.” From a function theoretic perspective, this is indeed natural (see for example [3] or [6]). Building on work of Andreotti and Hill [2], Brinkschulte [4] is able to show local smooth solvability of $\bar{\partial}$ at appropriate form levels for such domains. However, global closed range for $\bar{\partial}$ in $L^2_{(0,q)}(\Omega)$ and $L^2_{(0,q+1)}(\Omega)$ remains open on these domains.

In [12], Ho considers domains Ω where the sum of any q -eigenvalues of the Levi-form for $\partial\Omega$ are nonnegative (q -convex domains), and shows that this

suffices for closed range of $\bar{\partial}$ in $L^2_{(0,r)}(\Omega)$ with $r \geq q$. Zampieri has further generalized this as q -pseudoconvexity (see [27]). In q -pseudoconvexity, a subbundle of the tangent bundle of at most dimension q exists locally with the property that the sum of the $q + 1$ smallest eigenvalues of the Levi form is greater than or equal to the trace of the Levi form with respect to the subbundle. As shown in Theorem 1.9.9 in [27], this implies that for $L^2(0, q + 1)$ -forms f in the kernel of $\bar{\partial}$, there exists an $L^2(0, q)$ -form u solving $\bar{\partial}u = f$. To be consistent with the notation convention in $Z(q)$ (and q -convexity in [12]), we will typically refer to this as $(q - 1)$ -pseudoconvexity.

In this paper, we will generalize $(q - 1)$ -pseudoconvexity as follows. Taking the trace of the Levi form with respect to a vector bundle can be thought of in local coordinates as taking the trace of the Levi form with respect to a projection matrix (i.e., a hermitian matrix with eigenvalues of 0 or 1). We will relax this condition by allowing eigenvalues that are between 0 and 1. This gives us needed flexibility, as demonstrated by an example in Proposition 6.1 in which the rank of the identity minus our matrix is forced to be locally nonconstant. By analogy with the nondegenerate case, we will call such domains weak $Z(q)$ domains (see Definition 2.1 for a formal definition). The example in Proposition 6.1 thus illustrates that weak $Z(q)$ is a strictly weaker condition than any previously known condition.

We also allow for the boundary to be disconnected, using techniques developed for the annulus between two pseudoconvex domains in [20]. These techniques allow us to solve $\bar{\partial}$ modulo the space of harmonic $(0, q)$ -forms in some weighted L^2 space. By adapting recent arguments of Shaw [23], we are able to show that the space of harmonic $(0, q)$ -forms in fact vanishes, which allows us to use Hörmander's methods to obtain results in unweighted L^2 spaces. Our main L^2 -result is thus the following:

THEOREM 1.1. — *Let M be an n -dimensional Stein manifold, and let Ω be a bounded subset of M with C^3 boundary satisfying weak $Z(q)$ for some $1 \leq q \leq n - 1$. Then we have*

- (1) *The space of harmonic $(0, q)$ -forms $\mathcal{H}^q(\Omega)$ is trivial.*
- (2) *The $\bar{\partial}$ -Laplacian \square^q has closed range in $L^2_{(0,q)}(\Omega)$.*
- (3) *The $\bar{\partial}$ -Neumann operator N^q exists and is continuous in $L^2_{(0,q)}(\Omega)$.*
- (4) *The operator $\bar{\partial}$ has closed range in $L^2_{(0,q)}(\Omega)$ and $L^2_{(0,q+1)}(\Omega)$.*
- (5) *The operator $\bar{\partial}^*$ has closed range in $L^2_{(0,q)}(\Omega)$ and $L^2_{(0,q-1)}(\Omega)$.*

We work in Stein manifolds motivated by results in [10]. The C^3 boundary is needed for our method of proof because of additional integration by parts that are carried out to handle the nonpositive eigenvalues of the Levi

form. Unfortunately, while the results of Theorem 1.1 do not depend on the metric, our condition appears to depend on the metric (see Proposition 6.7). See [26] for discussion of analogous difficulties surrounding the apparent metric dependence of Property (P_q) . This is another benefit of working in generic Stein manifolds, since our example in Section 6 requires a non-Euclidean metric. Our condition also implies stronger results which do depend on the metric. Let φ be a global plurisubharmonic exhaustion function for M , and let the metric for M be given by the Kähler form $\omega = i\partial\bar{\partial}\varphi$. For a weight function ϕ which is chosen to equal $\pm\varphi$ in a neighborhood of each connected component of $\partial\Omega$, we can define the weighted L^2 -norm $\|f\|_t^2 = \int_{\Omega} e^{-t\phi} |f|^2 dV$. For sufficiently large $t > 0$, we will be able to obtain Sobolev space estimates for the weighted $\bar{\partial}$ -Neumann operator defined with respect to this norm.

THEOREM 1.2. — *Let M be an n -dimensional Stein manifold, and let Ω be a bounded subset of M with C^3 boundary satisfying weak $Z(q)$ for some $1 \leq q \leq n - 1$. Then there exists a constant $\tilde{t} > 0$ such that for all $t > \tilde{t}$ and $-\frac{1}{2} \leq s \leq 1$ we have*

- (1) *The weighted $\bar{\partial}$ -Neumann operator N_t^q exists and is continuous in $L^2_{(0,q)}(\Omega)$.*
- (2) *The canonical solution operators to $\bar{\partial}$ given by $\bar{\partial}_t^* N_t^q : W^s_{(0,q)}(\Omega) \rightarrow W^s_{(0,q-1)}(\Omega)$ and $N_t^q \bar{\partial}_t^* : W^s_{(0,q+1)}(\Omega) \rightarrow W^s_{(0,q)}(\Omega)$ are continuous.*
- (3) *The canonical solution operators to $\bar{\partial}_t^*$ given by $\bar{\partial} N_t^q : W^s_{(0,q)}(\Omega) \rightarrow W^s_{(0,q+1)}(\Omega)$ and $N_t^q \bar{\partial} : W^s_{(0,q-1)}(\Omega) \rightarrow W^s_{(0,q)}(\Omega)$ are continuous.*
- (4) *For every $f \in W^s_{(0,q)}(\Omega) \cap \ker \bar{\partial}$ there exists a $u \in W^s_{(0,q-1)}(\Omega)$ such that $\bar{\partial}u = f$.*

Interestingly, Theorem 1.2 is actually needed to prove Theorem 1.1 when the boundary is disconnected; see Section 5 for details. The estimates of Theorem 1.2 were carried out by the first author in [8] for pseudoconvex domains with C^2 boundary; again, the additional integration by parts needed for weak $Z(q)$ seem to require an additional degree of smoothness. Furthermore, the elliptic regularization carried out in Section 4 seems to require a C^3 boundary.

When $\partial\Omega$ satisfies weak $Z(q)$ and weak $Z(n - q - 1)$, we say that $\partial\Omega$ satisfies weak $Y(q)$. In [9], the authors continued work of [18] and [19] to understand solvability for the boundary operator $\bar{\partial}_b$ on CR -manifolds of hypersurface type. The definition given for weak $Y(q)$ in that paper is completely superseded by the definition in the present paper. When our

bounded weak $Y(q)$ manifold is an actual hypersurface in a Stein manifold, we now have the following result:

THEOREM 1.3. — *Let M be an n -dimensional Stein manifold, and let Ω be a bounded subset of M with connected C^3 boundary satisfying weak $Y(q)$ for some $1 \leq q < n - 1$. For every $f \in L^2_{(0,q)}(\partial\Omega) \cap \ker \bar{\partial}_b$ there exists a $u \in L^2_{(0,q-1)}(\partial\Omega)$ satisfying $\bar{\partial}_b u = f$. Hence, $\bar{\partial}_b$ has closed range in $L^2_{(0,q)}(\partial\Omega)$.*

Additional assumptions seem necessary for $q = n - 1$, even in the pseudoconvex case (see [5] for details).

The authors would like to thank the referee for several helpful comments on Section 6, which led to substantial improvements to Proposition 6.6.

2. Basic Properties and Notation

Let M be an n -dimensional Stein manifold, $n \geq 2$, and fix a smooth, strictly plurisubharmonic exhaustion function φ for M . We endow M with the Kähler metric given by the Kähler form $\omega = i\partial\bar{\partial}\varphi$. In local coordinates z_1, \dots, z_n , we will write

$$\omega = i \sum_{j,k=1}^n g_{j\bar{k}} dz_j \wedge d\bar{z}_k = i \sum_{j,k=1}^n \frac{\partial^2 \varphi}{\partial z_j \partial \bar{z}_k} dz_j \wedge d\bar{z}_k.$$

As usual, $g^{\bar{k}j}$ will denote the inverse matrix to $g_{j\bar{k}}$. By the usual convention, we will use the metric to raise and lower indices, so that, for example,

$$\sum_{\ell=1}^n g_{j\bar{\ell}} b^{\bar{\ell}k} = b_j^{\cdot k} \text{ and } \sum_{\ell=1}^n c_{j\bar{\ell}} g^{\bar{\ell}k} = c_j^{\cdot k}.$$

Additionally, we use the bracket $\langle \cdot, \cdot \rangle$ notation for the metric pairing, i.e., if $L, L' \in T^{1,0}(M)$, then $\langle L, L' \rangle = \omega(i\bar{L}' \wedge L)$ whereas if $\alpha = \sum_{j=1}^n a_j dz_j$ and $\alpha' = \sum_{j=1}^n a'_j dz_j$ in local coordinates, then $\langle \alpha, \alpha' \rangle = \sum_{j,k=1}^n \bar{a}'_k g^{\bar{k}j} a_j$. By multilinearity, $\langle \cdot, \cdot \rangle$ now extends to (p, q) -forms.

Let $\Omega \subset M$ be a bounded domain with C^m boundary. By definition, this means that there exists a C^m function ρ on M such that $\Omega = \{z \in M \mid \rho(z) < 0\}$ and $d\rho \neq 0$ on $\partial\Omega$. Such a ρ is called a C^m defining function for Ω . For $z \in \partial\Omega$, we define the induced CR-structure on $\partial\Omega$ at z by

$$T_z^{1,0}(\partial\Omega) = \{L \in T_z^{1,0}(M) : \partial\rho(L) = 0\}.$$

Let $T^{1,0}(\partial\Omega)$ denote the space of C^{m-1} sections of $T_z^{1,0}(\partial\Omega)$. We will also need $T^{0,1}(\partial\Omega) = \overline{T^{1,0}(\partial\Omega)}$ and the exterior algebra generated by

these: $T^{p,q}(\partial\Omega)$. Let $\Lambda^{p,q}(\partial\Omega)$ denote the bundle of C^{m-1} (p, q) -forms on $T^{p,q}(\partial\Omega)$. We use τ to denote the orthogonal projection and restriction:

$$(2.1) \quad \tau : \Lambda^{p,q}(M) \rightarrow \Lambda^{p,q}(\partial\Omega).$$

For each element X of $T^{p,q}$ (resp. $\Lambda^{p,q}$), we denote the metric dual element of $\Lambda^{p,q}$ (resp. $T^{p,q}$) by X^\sharp . This is defined to satisfy the relationships $X^\sharp(Y) = \langle Y, X \rangle$ for all $Y \in T^{p,q}$ (resp. $Y(X^\sharp) = \langle Y, X \rangle$ for all $Y \in \Lambda^{p,q}$). For example, the dual of the Kähler form is given in local coordinates by

$$\omega^\sharp = i \sum_{j,k=1}^n g^{\bar{k}j} \frac{\partial}{\partial \bar{z}_k} \wedge \frac{\partial}{\partial z_j}.$$

For any C^2 defining function ρ , the *Levi form* \mathcal{L}_ρ is the real element of $\Lambda^{1,1}(\partial\Omega)$ defined by

$$\mathcal{L}_\rho(i\bar{L} \wedge L') = i\partial\bar{\partial}\rho(i\bar{L} \wedge L')$$

for any $L, L' \in T^{1,0}(\partial\Omega)$. As usual, if $\tilde{\rho}$ is another C^2 defining function for Ω , then $\tilde{\rho} = \rho h$ for some nonvanishing C^1 function h , and $\mathcal{L}_{\tilde{\rho}} = h\mathcal{L}_\rho$. We will typically suppress the subscript ρ when the choice of defining function is not relevant.

DEFINITION 2.1. — For $1 \leq q \leq n - 1$, we say $\partial\Omega$ satisfies $Z(q)$ weakly if there exists a real $\Upsilon \in T^{1,1}(\partial\Omega)$ satisfying

- (1) $|\theta|^2 \geq (i\theta \wedge \bar{\theta})(\Upsilon) \geq 0$ for all $\theta \in \Lambda^{1,0}(\partial\Omega)$.
- (2) $\mu_1 + \dots + \mu_q - \mathcal{L}(\Upsilon) \geq 0$ where μ_1, \dots, μ_{n-1} are the eigenvalues of \mathcal{L} in increasing order.
- (3) $\omega(\Upsilon) \neq q$.

Remark 2.2. — Note that this is an intrinsic definition, so it can also be applied to abstract CR-manifolds of hypersurface type. This replaces the definition given in [9], and the main results of that paper still follow with this more general definition.

The fact that $\partial\Omega$ is the boundary of a domain induces a natural orientation on $\partial\Omega$. It is sometimes useful to reverse the orientation and think of $\partial\Omega$ as the boundary of the complement instead. The following observation is trivial for $Z(q)$, and motivates the definition of $Y(q)$, so it is of interest to confirm the corresponding fact for weak $Z(q)$.

PROPOSITION 2.3. — For $1 \leq q \leq n - 2$, let $\Omega \subset M$ be a bounded domain and let $B \subset M$ be a sufficiently large bounded pseudoconvex domain so that $\bar{\Omega} \subset B$. Then $\partial\Omega$ satisfies $Z(q)$ weakly if and only if $\partial(B/\bar{\Omega})$ satisfies $Z(n - q - 1)$ weakly.

Proof. — Suppose $\partial\Omega$ satisfies $Z(q)$ weakly, and let $\tilde{\Upsilon}$ be the element of $T^{1,1}(\partial\Omega)$ given by Definition 2.1. On ∂B , we can define $\Upsilon = 0$, and on $\partial\Omega$ we define $\Upsilon = (\tau\omega)^\sharp - \tilde{\Upsilon}$. If $\tilde{\mu}_1, \dots, \tilde{\mu}_{n-1}$ are the eigenvalues of the Levi form of $\partial\Omega$ in increasing order, then $\mu_1 = -\tilde{\mu}_{n-1}, \dots, \mu_{n-1} = -\tilde{\mu}_1$ are the eigenvalues of the Levi form of $\partial(B/\overline{\Omega})$ (on $\partial\Omega$) in increasing order, so since $\mathcal{L}((\tau\omega)^\sharp) = \mu_1 + \dots + \mu_{n-1}$, we have

$$\mu_1 + \dots + \mu_{n-q-1} - \mathcal{L}(\Upsilon) = \tilde{\mu}_1 + \dots + \tilde{\mu}_q - \tilde{\mathcal{L}}(\tilde{\Upsilon}).$$

Furthermore, $\omega(\Upsilon) = n - 1 - \omega(\tilde{\Upsilon}) \neq n - q - 1$.

Similar computations prove the converse. □

Remark 2.4. — We can replace B with any bounded domain such that ∂B satisfies $Z(n - q - 1)$ weakly.

Motivated by this, we define

DEFINITION 2.5. — For $1 \leq q \leq n - 2$, we say $\partial\Omega$ satisfies $Y(q)$ weakly if $\partial\Omega$ satisfies $Z(q)$ weakly and $Z(n - q - 1)$ weakly.

We note that Definition 2.1 is essentially a local property (modulo connected boundary components).

LEMMA 2.6. — For $1 \leq q \leq n - 1$, let $\sigma : \partial\Omega \rightarrow \{-1, 1\}$ be continuous and suppose that for every $p \in \partial\Omega$ there exists an open neighborhood U_p of p such that $U_p \cap \partial\Omega$ is connected and a real $\Upsilon_p \in T^{1,1}(U_p)$ satisfying

- (1) $|\theta|^2 \geq (i\theta \wedge \bar{\theta})(\Upsilon_p) \geq 0$ for all $\theta \in \Lambda^{1,0}(U_p)$.
- (2) $\mu_1 + \dots + \mu_q - \mathcal{L}(\Upsilon_p) \geq 0$ on U_p where μ_1, \dots, μ_{n-1} are the eigenvalues of the Levi form in increasing order.
- (3) $\sigma(p)(\omega(\Upsilon_p) - q) > 0$ on U_p .

Then $\partial\Omega$ satisfies $Z(q)$ weakly.

Remark 2.7. — The function σ represents a choice of orientation for each connected boundary component of Ω .

Proof. — Choose a finite cover $\{U_p\}_{p \in \mathcal{P}}$ of $\partial\Omega$ and let χ_p be a subordinate partition of unity. If we let $\Upsilon = \sum_{p \in \mathcal{P}} \chi_p \Upsilon_p$, then the necessary properties are satisfied by linearity. Since σ is constant on each connected component of $\partial\Omega$, $\omega(\Upsilon_p) - q$ will have a constant sign on each connected component of $\partial\Omega$, so there is no possibility of cancellation in the corresponding sum. □

Fix $p \in \partial\Omega$, and choose local coordinates that are orthonormal at p and satisfy $\frac{\partial \rho}{\partial z_j}(p) = 0$ for all $1 \leq j \leq n - 1$. At p we can write

$$\Upsilon = i \sum_{j,k=1}^{n-1} b^{\bar{k}j} \frac{\partial}{\partial \bar{z}_k} \wedge \frac{\partial}{\partial z_j} \quad \text{and} \quad \mathcal{L} = i \sum_{j,k=1}^{n-1} c_{j\bar{k}} dz_j \wedge d\bar{z}_k$$

where $b^{\bar{k}j}$ and $c_{j\bar{k}}$ are hermitian $(n - 1) \times (n - 1)$ matrices. Suppose that the local coordinates are chosen to diagonalize $b^{\bar{k}j}$ at p , and write $b^{\bar{k}j} = \delta_{jk} \lambda_j$. When restricted to p , the defining characteristics of weak $Z(q)$ will take the form

- (1) $0 \leq \lambda_j \leq 1$ for all $1 \leq j \leq n - 1$.
- (2) $\mu_1 + \dots + \mu_q - (\lambda_1 c_{1\bar{1}} + \dots + \lambda_{n-1} c_{n-1\overline{n-1}}) \geq 0$.
- (3) $\lambda_1 + \dots + \lambda_{n-1} \neq q$.

If there is an orthonormal local coordinate frame that diagonalizes $b^{\bar{k}j}$ such that $\lambda_1 = \dots = \lambda_m = 1$ and $\lambda_{m+1} = \dots = \lambda_{n-1} = 0$ for some $m \neq q$, then this is the condition studied in [9] which was shown to generalize $Z(q)$ and $(q - 1)$ -pseudoconvexity [27] (with the weight $\varphi(z) = |z|^2$ in [9]).

Alternatively, we can choose orthonormal coordinates that diagonalize the Levi form at a point, so that $c_{j\bar{k}} = \delta_{jk} \mu_j$. The second condition then translates into

$$\mu_1(1 - b^{\bar{1}1}) + \dots + \mu_q(1 - b^{\bar{q}q}) - (\mu_{q+1} b^{\overline{q+1}q+1} + \dots + \mu_{n-1} b^{\overline{n-1}n-1}) \geq 0.$$

Since $(1 - b^{\bar{j}j}) \geq 0$, $b^{\bar{j}j} \geq 0$, and $\mu_j \leq \mu_q \leq \mu_{q+1} \leq \mu_k$ for all $j \leq q < q + 1 \leq k$, it follows that $\mu_q(q - \omega(\Upsilon)) \geq 0$ and $\mu_{q+1}(q - \omega(\Upsilon)) \geq 0$. Hence, if $\mu_q < 0$, then $\omega(\Upsilon) > q$, and if $\mu_{q+1} > 0$, then $\omega(\Upsilon) < q$. Equivalently, we have

LEMMA 2.8. — For $1 \leq q \leq n - 1$ let $\Omega \subset M$ be a bounded domain and suppose that $\partial\Omega$ satisfies $Z(q)$ weakly. Let Υ be as in Definition 2.1. For any fixed boundary point, if $\omega(\Upsilon) < q$ then the Levi form has at least $n - q$ nonnegative eigenvalues, and if $\omega(\Upsilon) > q$, then the Levi form has at least $q + 1$ nonpositive eigenvalues.

Many results will be easier to work with when the boundary is connected. The fact that we are only working with bounded domains induces a natural decomposition into domains with connected boundaries.

LEMMA 2.9. — For any Stein manifold M , supposed that $\Omega \subset M$ is a connected bounded domain with C^3 boundary satisfying $Z(q)$ weakly for some $1 \leq q \leq n - 1$. Then $\Omega = \Omega_1 / \bigcup_{j=2}^m \bar{\Omega}_j$ where Ω_j has connected boundary for each $1 \leq j \leq m$, Ω_1 satisfies $Z(q)$ weakly, and Ω_j satisfies

$Z(n - q - 1)$ weakly for each $2 \leq j \leq m$. The $(1, 1)$ -vector Υ in Definition 2.1 will satisfy $\omega(\Upsilon) < q$ on $\partial\Omega_1$ and $\omega(\Upsilon) > q$ on $\partial\Omega_j$ for $2 \leq j \leq m$.

Proof. — Let $\psi : M \rightarrow \mathbb{C}^{2n+1}$ be an embedding (see Theorem 5.3.9 in [14]). Since Ω is bounded, there exists a minimal radius $R > 0$ such that $\psi[\Omega]$ is contained in a ball centered at zero with radius R . Denote the pullback of this ball under ψ by B . Then B is a strictly pseudoconvex domain in M containing Ω and since R is minimal there exists at least one point $p \in \partial\Omega \cap \partial B$. At p , $\partial\Omega$ must also be strictly pseudoconvex, so $\omega(\Upsilon)(p) < q$ by the contrapositive of Lemma 2.8. By continuity, $\omega(\Upsilon) < q$ on the connected boundary component containing p , so we define this to be $\partial\Omega_1$. Since Ω is connected, the remaining boundary components (finitely many, since Ω is relatively compact with C^3 boundary) can be thought of as boundaries of $Z(n - q - 1)$ domains by Proposition 2.3 and Remark 2.4. Using the argument with a ball in \mathbb{C}^{2n+1} , we again see that each of these subdomains has a strictly pseudoconvex point, and hence the Levi form is negative definite (when viewed as part of $\partial\Omega$). When the Levi form is negative-definite, we must have $\omega(\Upsilon) > q$ by Lemma 2.8. \square

Remark 2.10. — One consequence of this proof is that there are no bounded weak $Z(0)$ domains, since $\omega(\Upsilon) < 0$ is impossible. On the other hand, bounded weak $Z(n - 1)$ domains can exist (e.g., pseudoconvex domains), but they must have connected boundaries (otherwise some boundary components would bound weak $Z(0)$ domains). For analysis of the $q = n - 1$ case on domains with disconnected boundaries, see [15] and [23].

To prove our basic estimates, we will need extensions of Υ to M .

LEMMA 2.11. — *Suppose that $\partial\Omega$ satisfies $Z(q)$ weakly, and let Υ be as in Definition 2.1. Let ρ be any C^m defining function for Ω . There exist relatively compact open sets U^+ , U^- , and U^0 covering $\bar{\Omega}$ such that $\partial\Omega \cap \bar{U}^0 = \emptyset$ and $\bar{U}^+ \cap \bar{U}^- = \emptyset$, along with real Υ^+ , $\Upsilon^- \in T^{1,1}(M)$ satisfying*

- (1) $|\theta|^2 \geq (i\theta \wedge \bar{\theta})(\Upsilon^\pm) \geq 0$ for all $\theta \in \Lambda^{1,0}(M)$.
- (2) $\omega(\Upsilon^+) < q$ and $\omega(\Upsilon^-) > q$ on M .
- (3) For any $\theta \in \Lambda^{1,0}(M)$ we have

$$(i\theta \wedge \bar{\theta})(\Upsilon^\pm) = (i\tau\theta \wedge \tau\bar{\theta})(\Upsilon)$$

on $\partial\Omega \cap U^\pm$.

- (4) On a neighborhood of $\partial\Omega \cap U^\pm$, we have $(\theta \wedge \bar{\partial}\rho)(\Upsilon^\pm) = 0$ for all $\theta \in \Lambda^{1,0}(M)$.

Remark 2.12. — Note that $U^+ \neq \emptyset$ by Lemma 2.9. On the other hand, if $\partial\Omega$ is connected, we can set $U^- = \emptyset$ and $U^0 = \emptyset$.

Proof. — Let $K^+ = \{z \in \partial\Omega : \omega(\Upsilon) < q\}$ and $K^- = \{z \in \partial\Omega : \omega(\Upsilon) > q\}$. By the continuity of $\omega(\Upsilon)$ these are disconnected from each other, so there exist open neighborhoods U^+ and U^- such that $K^\pm \subset U^\pm$ and $K^\pm \cap \bar{U}^\mp = \emptyset$. Choose U^0 such that $\partial\Omega \cap \bar{U}^0 = \emptyset$ and $\bar{\Omega} \subset U^0 \cup U^+ \cup U^-$.

Let U be a neighborhood of $\partial\Omega$ on which $d\rho \neq 0$ for some C^m defining function ρ . Let $\psi(w, t) : U \times [0, |\rho(w)|] \rightarrow U$ solve the initial value problem $\psi(w, 0) = w$ and $\frac{\partial}{\partial t}\psi(w, t) = -\left((\text{sgn}\rho) |d\rho|^{-2} (d\rho)^\sharp\right) (\psi(w, t))$. By construction, this will satisfy

$$\frac{\partial}{\partial t}\rho(\psi(w, t)) = -(\text{sgn}\rho) |d\rho|^{-2} \langle d\rho, d\rho \rangle (\psi(w, t)) = -(\text{sgn}\rho) (\psi(w, t)),$$

so $\psi(w, |\rho(w)|) \in \partial\Omega$.

We denote parallel translation along $\psi(w, t)$ by

$$P_{a,w}^b : T_{\psi(w,a)}^{p,q}(M) \rightarrow T_{\psi(w,b)}^{p,q}(M).$$

Choose $\chi \in C_0^\infty(U)$ such that $\chi \equiv 1$ on a neighborhood of $\partial\Omega$. We define Υ^\pm on $\partial\Omega \cap U^\pm$ by

$$(i\theta \wedge \bar{\theta})(\Upsilon^\pm) = (i\tau\theta \wedge \tau\bar{\theta})(\Upsilon)$$

for any $\theta \in \Lambda^{1,0}(M)$. On $U \cap U^\pm$, we parallel translate Υ^\pm along ψ , as follows. We define

$$\Upsilon^+(w) = \chi(w) P_{|\rho(w)|,w}^0 \Upsilon^+(\psi(w, |\rho(w)|))$$

and

$$\Upsilon^-(w) = \chi(w) P_{|\rho(w)|,w}^0 \Upsilon^-(\psi(w, |\rho(w)|)) + (1 - \chi(w))\omega^\sharp.$$

We can now define $\Upsilon^+ = 0$ on $M/(U \cap U^+)$ and $\Upsilon^- = \omega^\sharp$ on $M/(U \cap U^-)$. □

We are now ready to define our weight function. Let U^\pm, U^0 , and Υ^\pm be as in Lemma 2.11. Fix $\chi \in C_0^\infty(M/\bar{U}^-)$ such that $\chi \equiv 1$ on \bar{U}^+ . Set

$$\phi = \chi\varphi - (1 - \chi)\varphi.$$

While the complex Hessian of ϕ on U^0 will involve derivatives of χ , we still have

$$(2.2) \quad i\partial\bar{\partial}\phi = \begin{cases} \omega & \text{on } U^+ \\ -\omega & \text{on } U^- \end{cases}.$$

We next define the usual weighted L^2 -inner products. For $f, h \in L^2_{(0,q)}(\Omega)$, define

$$(f, h)_t = \int_{\Omega} e^{-t\phi} \langle f, h \rangle dV.$$

and $\|f\|_t = \sqrt{(f, f)_t}$. Since $e^{-t\phi}$ is uniformly bounded on Ω , the space of $(0, q)$ -forms bounded in $\|\cdot\|_t$ is equal to $L^2_{(0,q)}(\Omega)$. The operator

$$\bar{\partial} : L^2_{(0,q)}(\Omega, e^{-t\phi}) \rightarrow L^2_{(0,q+1)}(\Omega, e^{-t\phi})$$

is given its L^2 -maximal definition, and the adjoint

$$\bar{\partial}_t^* : L^2_{(0,q+1)}(\Omega, e^{-t\phi}) \rightarrow L^2_{(0,q)}(\Omega, e^{-t\phi})$$

is defined with respect to the weighted inner product $(\cdot, \cdot)_t$. We also have $\square_t^q = \bar{\partial}\bar{\partial}_t^* + \bar{\partial}_t^*\bar{\partial}$ with the induced domain. The space of harmonic forms is given by

$$\mathcal{H}_t^q(\Omega) = L^2_{(0,q)}(\Omega, e^{-t\phi}) \cap \ker \bar{\partial} \cap \ker \bar{\partial}_t^*,$$

with the projection onto these denoted $H_t^q : L^2_{(0,q)}(\Omega, e^{-t\phi}) \rightarrow \mathcal{H}_t^q(\Omega, e^{-t\phi})$.

When it exists, the weighted $\bar{\partial}$ -Neumann operator

$$N_t^q : L^2_{(0,q)}(\Omega, e^{-t\phi}) \rightarrow \text{Dom}(\square_t^q)$$

satisfies $\square_t^q N_t^q = I - H_t^q$.

Let \mathcal{I}_q denote the set of increasing multi-indices over $\{1, \dots, n\}$ of length q . For an open set $U \subset M$ with local coordinates $\{z_1^U, \dots, z_n^U\}$, we let ∇_j^U denote the covariant derivative with respect to $\frac{\partial}{\partial z_j^U}$. We also use $\nabla_{j,t}^{U,*} = -\bar{\nabla}_j^U + t \frac{\partial \phi}{\partial \bar{z}_j^U}$. This satisfies the adjoint relationship

$$\sum_{j,k=1}^n (g_U^{\bar{k}j} \nabla_j^U f, h_k)_t = \sum_{j,k=1}^n (g_U^{\bar{k}j} f, \nabla_{j,t}^{U,*} h_k)_t$$

assuming f and h_k are compactly supported. If \mathcal{U} is a finite open cover of Ω by such sets, we let $\{\chi^U\}_{U \in \mathcal{U}}$ denote a partition of unity subordinate to this cover and define the following gradient terms on $(0, q)$ -forms f for any $\Upsilon \in T^{1,1}(M)$:

$$(2.3) \quad \|\bar{\nabla} f\|_t^2 = \sum_{U \in \mathcal{U}} \sum_{j,k=1}^n (\chi^U g_U^{\bar{k}j} \bar{\nabla}_k^U f, \bar{\nabla}_j^U f)_t$$

$$(2.4) \quad \|\bar{\nabla}_\Upsilon f\|_t^2 = \sum_{U \in \mathcal{U}} \sum_{j,k=1}^n (\chi^U b_U^{\bar{k}j} \bar{\nabla}_k^U f, \bar{\nabla}_j^U f)_t$$

$$(2.5) \quad \|\nabla_\Upsilon f\|_t^2 = \sum_{U \in \mathcal{U}} \sum_{j,k=1}^n (\chi^U b_U^{\bar{k}j} \bar{\nabla}_{j,t}^{U,*} f, \bar{\nabla}_{k,t}^{U,*} f)_t,$$

where

$$\Upsilon = i \sum_{j,k=1}^n b_U^{\bar{k}j} \frac{\partial}{\partial \bar{z}_k^U} \wedge \frac{\partial}{\partial z_j^U}$$

on U . We also introduce vector fields which will figure prominently in our error terms:

$$E = \sum_{U \in \mathcal{U}} \sum_{j,k,\ell} \chi^U g_U^{\bar{k}\ell} \left(\frac{\partial}{\partial \bar{z}_k^U} b_{\ell,U}^j \right) \frac{\partial}{\partial z_j^U}$$

$$E_\Upsilon = \sum_{U \in \mathcal{U}} \sum_{j,k,\ell,r} \chi^U g_U^{\bar{k}\ell} \left(\frac{\partial}{\partial \bar{z}_k^U} b_{\ell,U}^r \right) b_{r,U}^j \frac{\partial}{\partial z_j^U}.$$

Note that if we change coordinates, $b_{\ell,U}^r$ will be multiplied by matrices of holomorphic functions, which will be annihilated by $\frac{\partial}{\partial \bar{z}_k^U}$, so the vector fields remain invariant under changes of coordinates. At any point $p \in \Omega$, choose orthonormal coordinates at p that diagonalize $b^{\bar{k}j}$, with eigenvalues λ_j corresponding to the eigenvector $\frac{\partial}{\partial z_j}$ at p . If Υ satisfies property (1) in Lemma 2.11, then $0 \leq \lambda_j \leq 1$. If Υ is C^1 , then at p we can write

$$E = \sum_{j=1}^n A^j \frac{\partial}{\partial z_j} \text{ and } E_\Upsilon = \sum_{j=1}^n A^j \lambda_j \frac{\partial}{\partial z_j}$$

where A^j are continuous functions on our local coordinate patch. Hence, at p , since $\lambda_j^2 \leq \lambda_j$ we have

$$\left| \bar{\nabla}_{E_\Upsilon, t}^* f \right|^2 = \left| \sum_{j=1}^n A^j \lambda_j \bar{\nabla}_{j,t}^* f \right|^2 \leq C \sum_{j=1}^n \lambda_j \left| \bar{\nabla}_{j,t}^* f \right|^2$$

for some constant $C > 0$. Integrating, this gives us

$$(2.6) \quad \left\| \bar{\nabla}_{E_\Upsilon, t}^* f \right\|_t^2 \leq C \left\| \nabla_\Upsilon f \right\|_t^2.$$

On the other hand, since $(1 - \lambda_j)^2 \leq (1 - \lambda_j)$, we also have at p

$$\left| \bar{\nabla}_E f - \bar{\nabla}_{E_\Upsilon} f \right|^2 = \left| \sum_{j=1}^n A_j (1 - \lambda_j) \bar{\nabla}_j f \right|^2 \leq C \sum_{j=1}^n (1 - \lambda_j) \left| \bar{\nabla}_j f \right|^2$$

for some constant $C > 0$. Integration gives us

$$(2.7) \quad \left\| \bar{\nabla}_E f - \bar{\nabla}_{E_\Upsilon} f \right\|_t^2 \leq C \left(\left\| \bar{\nabla} f \right\|_t^2 - \left\| \bar{\nabla}_\Upsilon f \right\|_t^2 \right).$$

We also abuse notation and define the action of $(1, 1)$ -forms on $(0, q)$ -forms. Let $f \in C^1_{(0,q)}(\bar{\Omega}) \cap \text{Dom}(\bar{\partial}_t^*)$. For any point $p \in \Omega$, choose local coordinates that are orthonormal at p and define

$$i\bar{\partial}\bar{\partial}\phi(f, f)(p) = \sum_{J \in \mathcal{I}_{q-1}} \sum_{j,k=1}^n \frac{\partial^2 \phi}{\partial z_j \partial \bar{z}_k} f_{kJ} \bar{f}_j$$

where $f_{kJ} = (-1)^\sigma f_K$ for $K \in \mathcal{I}_q$ if $\{k\} \cup J = K$ as sets and σ is the length of the permutation that changes kJ into K . Due to (2.2), we have

$$(2.8) \quad i\partial\bar{\partial}\phi(f, f) = \begin{cases} q|f|^2 & \text{on } U^+ \\ -q|f|^2 & \text{on } U^- \end{cases}.$$

For any point $p \in \partial\Omega$, choose local coordinates that are orthonormal at p such that $\frac{\partial\rho}{\partial z_j}(p) = 0$ for all $1 \leq j \leq n - 1$. We define

$$\mathcal{L}(f, f)(p) = \sum_{J \in \mathcal{I}_{q-1}} \sum_{j,k=1}^{n-1} \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_k} f_{kJ} \bar{f}_{jJ}.$$

We note for future reference that if μ_1, \dots, μ_{n-1} are the eigenvalues of \mathcal{L} arranged in increasing order, then (adapting the proof of Lemma 4.7 in [25]), we have

$$(2.9) \quad \mathcal{L}(f, f) \geq (\mu_1 + \dots + \mu_q) |f|^2.$$

3. The Basic Estimate

Let ρ be a C^m defining function for Ω with $|\rho| = 1$ on Ω . For $f \in C^1_{(0,q)}(\bar{\Omega}) \cap \text{Dom}(\bar{\partial}_t^*)$ with $1 \leq q \leq n$, we have the Morrey-Kohn-Hörmander equality (see for example [5], [7], [14], or [25]):

$$(3.1) \quad \begin{aligned} \|\bar{\partial}f\|_t^2 + \|\bar{\partial}_t^* f\|_t^2 &= \|\bar{\nabla}f\|_t^2 + t \int_{\Omega} i\partial\bar{\partial}\phi(f, f)e^{-t\phi} dV \\ &\quad + \int_{\partial\Omega} \mathcal{L}(f, f)e^{-t\phi} dS + O(\|f\|_t^2). \end{aligned}$$

The error term involves the curvature of the Kähler metric, and can be computed explicitly using the Bochner-Kodaira technique [24]. Since this term can be controlled by choosing t large enough, we will not need the precise value.

We wish to understand integration by parts in the gradient term. Note that in (2.3), (2.4), and (2.5), the integrated terms are invariant under changes of coordinate, so derivatives of the partition of unity χ^U that arise from integration by parts will cancel. Hence, for clarity of notation, we can suppress the partition of unity without losing information.

Let U^\pm , U^0 , and Υ^\pm be as in Lemma 2.11. Note that property (3) in this lemma guarantees that Υ^\pm has no normal component on $\partial\Omega \cap U^\pm$, so $\Upsilon^\pm|_{\partial\Omega \cap U^\pm}$ is made up of tangential derivatives which can be integrated

by parts without introducing a boundary term. Hence, working in local coordinates with $f \in C^2_{(0,q)}(\bar{\Omega}) \cap C^2_0(U^\pm)$ we have

$$\begin{aligned}
 \|\bar{\nabla}_{\Upsilon^\pm} f\|_t^2 &= \sum_{j,k=1}^n (b^{\bar{k}j} \bar{\nabla}_k f, \bar{\nabla}_j f)_t \\
 (3.2) \qquad &= \sum_{j,k,\ell=1}^n (g^{\bar{k}\ell} f, \bar{\nabla}_{k,t}^* (b_{\bar{\ell}}^j \bar{\nabla}_j f))_t \\
 &= \sum_{j,k=1}^n (b^{\bar{k}j} f, \bar{\nabla}_{k,t}^* \bar{\nabla}_j f)_t - (f, \bar{\nabla}_E f)_t.
 \end{aligned}$$

To continue, we will need the commutator

$$\sum_{j,k=1}^n b^{\bar{k}j} [\bar{\nabla}_{k,t}^*, \bar{\nabla}_j] f = -t \sum_{j,k=1}^n b^{\bar{k}j} \frac{\partial^2 \phi}{\partial z_k \partial \bar{z}_j} f + O(f) = \mp t \omega(\Upsilon^\pm) f + O(f)$$

where the error terms are independent of t and (2.2) has been used. Substituting in (3.2) we have

$$\begin{aligned}
 \|\bar{\nabla}_{\Upsilon^\pm} f\|_t^2 &= \mp t (\omega(\Upsilon^\pm) f, f)_t + \sum_{j,k=1}^n (b^{\bar{k}j} f, \bar{\nabla}_j \bar{\nabla}_{k,t}^* f)_t \\
 &\quad - (f, \bar{\nabla}_E f)_t + O(\|f\|_t^2) \\
 (3.3) \qquad &= \mp t (\omega(\Upsilon^\pm) f, f)_t + \sum_{j,k,\ell=1}^n (g^{\bar{\ell}j} \bar{\nabla}_{j,t}^* (b_{\bar{\ell}}^k f), \bar{\nabla}_{k,t}^* f)_t \\
 &\quad - (f, \bar{\nabla}_E f)_t + O(\|f\|_t^2) \\
 &= \mp t (\omega(\Upsilon^\pm) f, f)_t + \|\nabla_{\Upsilon^\pm} f\|_t^2 - (f, \bar{\nabla}_{E,t}^* f)_t \\
 &\quad - (f, \bar{\nabla}_E f)_t + O(\|f\|_t^2).
 \end{aligned}$$

When we integrate the error terms by parts, it will be helpful to note that on $\partial\Omega$ we have $E_{\Upsilon^\pm} \rho = 0$ but

$$E\rho = \sum_{j,k,\ell} g^{\bar{k}\ell} \left(\frac{\partial}{\partial \bar{z}_k} b_{\bar{\ell}}^j \right) \frac{\partial \rho}{\partial z_j} = - \sum_{j,k} b^{\bar{k}j} \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_k} = -\mathcal{L}(\Upsilon).$$

It is also important to note that Υ^\pm must be C^2 if integration by parts with respect to $\nabla_{E_{\Upsilon^\pm}}$ is going to be well-defined. Considering the error terms in (3.3), we have

$$\begin{aligned}
 (f, \bar{\nabla}_{E,t}^* f)_t &= (f, \bar{\nabla}_{E_{\Upsilon^\pm},t}^* f)_t + ((\bar{\nabla}_E - \bar{\nabla}_{E_{\Upsilon^\pm}}) f, f)_t \\
 (3.4) \qquad &\quad + \int_{\partial\Omega} \mathcal{L}(\Upsilon) |f|^2 e^{-t\phi} dS + O(\|f\|_t^2)
 \end{aligned}$$

and

$$(3.5) \quad (f, \bar{\nabla}_E f)_t = (\bar{\nabla}_{E_{\Upsilon^\pm}}^* f, f)_t + (f, (\bar{\nabla}_E - \bar{\nabla}_{E_{\Upsilon^\pm}})f)_t + O(\|f\|_t^2).$$

Substituting (3.4) and (3.5) into (3.3), we have

$$(3.6) \quad \begin{aligned} \|\bar{\nabla}_{\Upsilon^\pm} f\|_t^2 &= \mp t(\omega(\Upsilon^\pm)f, f)_t - 2\operatorname{Re}\left((\bar{\nabla}_{E_{\Upsilon^\pm}}^* f, f)_t \right. \\ &\quad \left. + (f, (\bar{\nabla}_E - \bar{\nabla}_{E_{\Upsilon^\pm}})f)_t\right) + \|\nabla_{\Upsilon^\pm} f\|_t^2 \\ &\quad - \int_{\partial\Omega} \mathcal{L}(\Upsilon) |f|^2 e^{-t\phi} dS + O(\|f\|_t^2). \end{aligned}$$

Since we have property (1) in Lemma 2.11, we can now write

$$\|\bar{\nabla} f\|_t^2 = \left(\|\bar{\nabla} f\|_t^2 - \|\bar{\nabla}_{\Upsilon^\pm} f\|_t^2\right) + \|\bar{\nabla}_{\Upsilon^\pm} f\|_t^2$$

and use the Schwarz inequality, the small constant/large constant inequality, (2.7) and (2.6) to control the error terms in (3.6). We conclude

$$(3.7) \quad \|\bar{\nabla} f\|_t^2 \geq \mp t(\omega(\Upsilon^\pm)f, f)_t - \int_{\partial\Omega} \mathcal{L}(\Upsilon) |f|^2 dS + O(\|f\|_t^2)$$

Substituting (3.7) and (2.8) into (3.1), we have

$$(3.8) \quad \begin{aligned} \|\bar{\partial} f\|_t^2 + \|\bar{\partial}_t^* f\|_t^2 &\geq \pm t((q - \omega(\Upsilon^\pm))f, f)_t \\ &\quad + \int_{\partial\Omega} (\mathcal{L}(f, f) - \mathcal{L}(\Upsilon) |f|^2) e^{-t\phi} dS + O(\|f\|_t^2). \end{aligned}$$

We are now ready to prove the basic estimate (see Proposition 3.1 in [20] for the case where Ω is the annuli between two weakly-pseudoconvex domains).

PROPOSITION 3.1. — *Let M be an n -dimensional Stein manifold, $n \geq 2$, and let Ω be a bounded subset of M with C^3 boundary satisfying weak $Z(q)$ for some $1 \leq q \leq n - 1$.*

- (1) *For any constant $\varepsilon > 0$ there exists $t_\varepsilon > 0$ and a $C_\varepsilon > 0$ such that for any $t \geq t_\varepsilon$ and $f \in L^2_{(0,q)}(\Omega, e^{-t\phi}) \cap \operatorname{Dom}(\bar{\partial}) \cap \operatorname{Dom}(\bar{\partial}_t^*)$ we have*

$$\varepsilon \left(\|\bar{\partial} f\|_t^2 + \|\bar{\partial}_t^* f\|_t^2 \right) + C_\varepsilon \|f\|_{t,W^{-1}}^2 \geq \|f\|_t^2$$

where $\|\cdot\|_{t,W^{-1}}$ is the dual norm to $\|\cdot\|_{t,W^1}$.

- (2) *There exist constants $C > 0$ and $\tilde{t} > 0$ such that for all $t \geq \tilde{t}$ and $f \in L^2_{(0,q)}(\Omega, e^{-t\phi}) \cap \operatorname{Dom}(\bar{\partial}) \cap \operatorname{Dom}(\bar{\partial}_t^*) \cap (\mathcal{H}_t^q(\Omega))^\perp$ we have*

$$C \left(\|\bar{\partial} f\|_t^2 + \|\bar{\partial}_t^* f\|_t^2 \right) \geq \|f\|_t^2.$$

(3) If $\partial\Omega$ is connected, then for any constant $\varepsilon > 0$ there exists $t_\varepsilon > 0$ such that for all $t \geq t_\varepsilon$ and $f \in L^2_{(0,q)}(\Omega, e^{-t\phi}) \cap \text{Dom}(\bar{\partial}) \cap \text{Dom}(\bar{\partial}_t^*)$ we have

$$\varepsilon \left(\|\bar{\partial}f\|_t^2 + \|\bar{\partial}_t^* f\|_t^2 \right) \geq \|f\|_t^2.$$

Proof. — Let U^\pm, U^0 , and Υ^\pm be as in Lemma (2.11). Let χ^\pm and χ^0 form a partition of unity subordinate to U^\pm and U^0 . Given $f \in C^2_{(0,q)}(\bar{\Omega}) \cap \text{Dom}(\bar{\partial}_t^*)$, we define $f^\pm = \chi^\pm f$ and $f^0 = \chi^0 f$. Since $\partial\Omega$ is C^3 , Υ^\pm are C^2 , and hence (3.8) holds for f^\pm . By (2.9) and property (2) of Definition 2.1, the boundary term in (3.8) is positive, so we have

$$\|\bar{\partial}f^\pm\|_t^2 + \|\bar{\partial}_t^* f^\pm\|_t^2 \geq \pm t((q - \omega(\Upsilon^\pm))f^\pm, f^\pm)_t + O(\|f^\pm\|_t^2).$$

By property (2) of Lemma 2.11, we have $\pm(q - \omega(\Upsilon^\pm)) > 0$. Since Ω is bounded we know that $\pm(q - \omega(\tilde{\Upsilon})) \geq C_0$ for some constant $C_0 > 0$. Furthermore,

$$\|\bar{\partial}f^\pm\|_t^2 + \|\bar{\partial}_t^* f^\pm\|_t^2 \leq 2\|\bar{\partial}f\|_t^2 + 2\|\bar{\partial}_t^* f\|_t^2 + O(\|f\|_t^2),$$

so

$$2\|\bar{\partial}f\|_t^2 + 2\|\bar{\partial}_t^* f\|_t^2 \geq tC_0\|f^\pm\|_t^2 + O(\|f\|_t^2).$$

Since f_0 is compactly supported in Ω , we have Gårding’s inequality

$$\|f^0\|_{t,W^1}^2 \leq C_t(\|\bar{\partial}f^0\|_t^2 + \|\bar{\partial}_t^* f^0\|_t^2 + \|f^0\|_t^2)$$

for some constant $C_t > 0$. Using the duality between W_0^1 and W^{-1} we have

$$\|f^0\|_t^2 \leq \|f^0\|_{t,W^{-1}} \sqrt{C_t(\|\bar{\partial}f^0\|_t^2 + \|\bar{\partial}_t^* f^0\|_t^2 + \|f^0\|_t^2)}.$$

Associating $\sqrt{C_t}$ with the $\|f^0\|_{t,W^{-1}}$ and applying the standard small constant/large constant inequality, we have for any $s > 0$

$$\|f^0\|_t^2 \leq \frac{s}{2}C_t\|f^0\|_{t,W^{-1}}^2 + \frac{1}{2s}(\|\bar{\partial}f^0\|_t^2 + \|\bar{\partial}_t^* f^0\|_t^2 + \|f^0\|_t^2).$$

Subtracting $\frac{1}{2s}\|f^0\|_t^2$ from both sides and multiplying by $2s$ we have

$$(2s - 1)\|f^0\|_t^2 \leq s^2C_t\|f^0\|_{t,W^{-1}}^2 + \|\bar{\partial}f^0\|_t^2 + \|\bar{\partial}_t^* f^0\|_t^2.$$

Letting $s = \frac{1}{2}(1 + tC_0)$, we have

$$tC_0\|f^0\|_t^2 \leq \frac{1}{4}(1 + tC_0)^2C_t\|f\|_{t,W^{-1}}^2 + 2\|\bar{\partial}f\|_t^2 + 2\|\bar{\partial}_t^* f\|_t^2 + O(\|f\|_t^2).$$

Combining the estimates for f^0 and f^\pm , we conclude

$$\frac{tC_0}{3}\|f\|_t^2 + O(\|f\|_t^2) \leq \frac{1}{4}(1 + tC_0)^2C_t\|f\|_{t,W^{-1}}^2 + 2\|\bar{\partial}f\|_t^2 + 2\|\bar{\partial}_t^* f\|_t^2.$$

We can now choose t sufficiently large so that

$$\frac{tC_0}{6} \|f\|_t^2 + O(\|f\|_t^2) \geq \frac{1}{\varepsilon} \|f\|_t^2$$

and the estimate is complete. Standard density results (see for example Lemma 4.3.2 in [5]) complete the proof of part (1). The proof of part (2) is completed in the same manner as Lemma 3.1 in [20], after setting $\tilde{t} = \inf_{\varepsilon > 0} t_\varepsilon$.

When the boundary is connected, we note that $U^0 = U^- = \emptyset$ (see Remark 2.12), so there is no need to estimate f_0 . Hence the W^{-1} terms are not necessary, and part (3) follows. \square

We immediately have the standard consequences of such L^2 estimates.

THEOREM 3.2. — *Let M be an n -dimensional Stein manifold, $n \geq 2$, and let Ω be a bounded subset of M with C^3 boundary satisfying weak $Z(q)$ for some $1 \leq q \leq n - 1$. Then there exists a constant $\tilde{t} > 0$ such that for all $t > \tilde{t}$ we have*

- (1) $\mathcal{H}_t^q(\Omega)$ is finite dimensional. If $\partial\Omega$ is connected, then $\mathcal{H}_t^q(\Omega) = \{0\}$.
- (2) The weighted $\bar{\partial}$ -Laplacian \square_t^q has closed range in $L^2_{(0,q)}(\Omega, e^{-t\phi})$.
- (3) The weighted $\bar{\partial}$ -Neumann operator N_t^q exists and is continuous.
- (4) The operator $\bar{\partial}$ has closed range in $L^2_{(0,q)}(\Omega, e^{-t\phi})$ and $L^2_{(0,q+1)}(\Omega, e^{-t\phi})$.
- (5) The operator $\bar{\partial}_t^*$ has closed range in $L^2_{(0,q)}(\Omega, e^{-t\phi})$ and $L^2_{(0,q-1)}(\Omega, e^{-t\phi})$.
- (6) The canonical solution operators to $\bar{\partial}$ given by $\bar{\partial}_t^* N_t^q : L^2_{(0,q)}(\Omega, e^{-t\phi}) \rightarrow L^2_{(0,q-1)}(\Omega, e^{-t\phi})$ and $N_t^q \bar{\partial}_t^* : L^2_{(0,q+1)}(\Omega, e^{-t\phi}) \rightarrow L^2_{(0,q)}(\Omega, e^{-t\phi})$ are continuous.
- (7) The canonical solution operators to $\bar{\partial}_t^*$ given by $\bar{\partial} N_t^q : L^2_{(0,q)}(\Omega, e^{-t\phi}) \rightarrow L^2_{(0,q+1)}(\Omega, e^{-t\phi})$ and $N_t^q \bar{\partial} : L^2_{(0,q-1)}(\Omega, e^{-t\phi}) \rightarrow L^2_{(0,q)}(\Omega, e^{-t\phi})$ are continuous.
- (8) For every $f \in L^2_{(0,q)}(\Omega) \cap \ker \bar{\partial} \cap (\mathcal{H}_t^q(\Omega))^\perp$ there exists a $u \in L^2_{(0,q-1)}(\Omega)$ such that $\bar{\partial}u = f$.

4. Sobolev Estimates

In this section we will use elliptic regularization to obtain estimates in the L^2 -Sobolev space W^1 when $\partial\Omega$ is connected. The first author obtained such estimates for C^2 -pseudoconvex domains in [8]. In that paper, he used an exhaustion by smooth strictly pseudoconvex domains. Although smooth $Z(q)$ domains can exhaust bounded weakly $Z(q)$ domains with connected boundaries, constructing Υ on the exhaustion domains in such a way that

the estimates are uniform may not be possible. Hence, we will use elliptic regularization in the present paper. Our discussion follows the argument in Section 3.3 of [25], focusing on steps where the reduced boundary regularity requires more careful estimates.

We will need two equivalent norms on $W^1(\Omega)$: the standard norm $\|u\|_{W^1}^2 = \|u\|^2 + \|\nabla u\|^2$ and the weighted norm $\|u\|_{t,W^1}^2 = \|u\|_t^2 + \|\nabla u\|_t^2$. Although these are equivalent, the constant involved will depend on t , so for estimates where the dependency on t is significant we will need to use the weighted norm. Only at the end of the proof will we be able to pass to estimates for the standard norm, which is more suitable for interpolation. For $u \in W^1_{(0,q)}(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$ and $\delta > 0$ we define

$$Q_{t,\delta}(u, u) = \|\bar{\partial}u\|_t^2 + \|\bar{\partial}_t^*u\|_t^2 + \delta \|\nabla u\|_t^2.$$

As in [25], we have a unique self-adjoint operator $\square_{t,\delta}^q$ on $L^2_{(0,q)}(\Omega)$ satisfying $(\square_{t,\delta}^q u, v)_t = Q_{t,\delta}(u, v)$ for all $u \in \text{Dom}(\square_{t,\delta}^q)$ and $v \in W^1_{(0,q)}(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$, where $\text{Dom}(\square_{t,\delta}^q)$ is the subspace of $W^1_{(0,q)}(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$ on which $\tilde{\square}_{t,\delta}^q u \in L^2_{(0,q)}(\Omega)$ and $\tilde{\square}_{t,\delta}^q$ is the canonical identification between $W^1_{(0,q)}(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$ and its conjugate dual. We also obtain a unique solution operator $N_{t,\delta}^q$ mapping $L^2_{(0,q)}(\Omega)$ onto $\text{Dom}(\square_{t,\delta}^q)$ satisfying $(u, v)_t = Q_{t,\delta}(N_{t,\delta}^q u, v)$.

By Proposition 3.5 in [25], $N_{t,\delta}^q$ maps $L^2_{(0,q)}(\Omega)$ continuously to $W^2_{(0,q)}(\Omega)$. Although this proposition is stated for smooth domains, the proof for the $s = 0$ case holds on C^3 domains and we now outline the key step to illustrate the role of boundary smoothness. Let ρ be the signed distance function for Ω , so that ρ is a C^3 defining function [16], and let (x_1, \dots, x_{2n-1}) be coordinates on $\partial\Omega$ with ρ as the transverse coordinate. Similarly, we choose an orthonormal basis for $(1, 0)$ -forms consisting of $\omega_1, \dots, \omega_n$, where $\omega_n = \partial\rho$. If we express u in this basis, then the components will involve first derivatives of ρ . If we let D_j^h denote a difference quotient with respect to x_j , we can define $D_j^h u$ by considering difference quotients of components of u in our special basis. This will preserve $\text{Dom}(\bar{\partial}^*)$, but uniform bounds on $D_j^h u$ will now involve the C^2 norm of ρ . Finally we wish to estimate $Q_{t,\delta}(D_j^h u, v)$. The details for this estimate are contained in (3.38) through (3.41) in [25], but we will simply observe that they involve uniform bounds for $[\bar{\partial}, D_j^h]u$, $[\bar{\partial}_t^*, D_j^h]u$, and $[\nabla, D_j^h]u$, which will all involve the C^3 norm of ρ . Working with the smooth cutoff functions necessary to work locally will not involve additional derivative of ρ , so Straube’s argument will allow us to bound tangential derivatives of u in the W^1 norm. As usual, the structure of $\square_{t,\delta}^q$ as a second-order elliptic operator will allow us to estimate the second

derivatives in the normal direction (which only involve second derivatives of ρ).

Our goal is to show that the W^1 norm of $N_{t,\delta}^q f$ is bounded by the W^1 norm of f with a constant that is independent of δ , so that we may use a limiting argument to show that this estimate also holds for $N_t^q f$. Since $\bar{\partial} \oplus \bar{\partial}_t^*$ is an elliptic system, it will suffice to estimate tangential derivatives, but first we must clarify how a differential operator acts globally on a $(0, q)$ -form. Let $\psi \in W_{(0,q)}^2(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$ and let T be a differential operator defined on $\bar{\Omega}$ that is tangential on the boundary. Since $\partial\Omega$ is C^3 , we may assume T has C^2 coefficients. For $U \subset M$ with local coordinates $\{z_1, \dots, z_n\}$ we can write

$$\psi = \sum_{I \in \mathcal{I}_q} \psi_I d\bar{z}_I,$$

where the components ψ_I are all in $W^2(\Omega)$. The covariant derivative $\nabla_T \psi$ is globally defined and in local coordinates we can write

$$\nabla_T \psi = \sum_{I \in \mathcal{I}_q} T\psi_I d\bar{z}_I + O(\psi),$$

where the coefficients in the zero order term involve coefficients of T and are hence C^2 . However, $\nabla_T \psi$ is probably not in the domain of $\bar{\partial}_t^*$. On the other hand, for U sufficiently small, we can also choose a C^2 orthonormal basis for the space of $(1, 0)$ -forms $\omega_1, \dots, \omega_n$ where $\omega_n = \partial\rho$. In this basis we write

$$\psi = \sum_{I \in \mathcal{I}_q} \tilde{\psi}_I \bar{\omega}_I.$$

Since the transition matrices between dz and ω have C^2 entries, $\tilde{\psi}_I$ can be obtained by applying a linear operator with C^2 coefficients to ψ_I . We define

$$D_T^U \psi = \sum_{I \in \mathcal{I}_q} T\tilde{\psi}_I \omega_I,$$

and sum over a partition of unity to obtain $D_T \psi$. Note that D_T preserves tangential and normal components of ψ , so it will also preserve the domain of $\bar{\partial}_t^*$. However, returning to local coordinates,

$$D_T^U \psi = \sum_{I \in \mathcal{I}_q} T\psi_I d\bar{z}_I + O(\psi),$$

where the coefficients of the zero order terms are obtained by differentiating the C^2 transitions matrices between dz and ω , so they are only C^1 . Hence, $D_T - \nabla_T$ is a zero-order operator with C^1 coefficients. This requires some caution. For example, if ∇_D^2 is a second-order differential

operator with C^1 coefficients, then $[\nabla_D^2, \nabla_T]$ is a second-order differential operator with continuous coefficients, and hence $[\nabla_D^2, \nabla_T]\psi$ is a form in $L^2_{(0,q)}$. However, $[\nabla_D^2, D_T - \nabla_T]$ may not be a differential operator with continuous coefficients, so we cannot make use of $[\nabla_D^2, D_T]\psi$. On the other hand, commutators between D_T and first-order differential operators will still have continuous (hence bounded) coefficients.

For $\varepsilon > 0$, let t_ε and t be as in (3) of Proposition 3.1. Then for $u \in W^1_{(0,q)}(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$, when $\partial\Omega$ is connected we have $\varepsilon Q_{t,\delta}(u, u) \geq \|u\|_t^2$. Let $f \in W^1_{(0,q)}(\Omega)$. We immediately obtain

$$(4.1) \quad \left\| N_{t,\delta}^q f \right\|_t \leq \varepsilon \|f\|_t.$$

Since $N_{t,\delta}^q f \in W^2_{(0,q)}$, we can set $u = D_T N_{t,\delta}^q f$ and obtain $u \in W^1_{(0,q)}(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$. Hence,

$$(4.2) \quad \left\| D_T N_{t,\delta}^q f \right\|_t^2 \leq \varepsilon Q_{t,\delta}(D_T N_{t,\delta}^q f, D_T N_{t,\delta}^q f).$$

To estimate $Q_{t,\delta}(D_T N_{t,\delta}^q f, D_T N_{t,\delta}^q f)$, we will need to work with slightly smoother forms. To that end, we introduce the following density lemma:

LEMMA 4.1. — *Let $\Omega \subset M$ be a bounded domain with C^3 boundary, and let $u \in W^1_{(0,q)}(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$. Then there exists a sequence $u_\ell \in C^2_{(0,q)}(\bar{\Omega}) \cap \text{Dom}(\bar{\partial}_t^*)$ converging to u in the W^1 norm.*

Proof. — Let ρ be a C^3 defining function for Ω , and let

$$I_q = \left\{ f \in L^2_{(0,q)}(\Omega) : f = \bar{\partial}\rho \wedge g, g \in L^2_{(0,q-1)}(\Omega) \right\}.$$

Choose $\chi \in C^\infty_0(M)$ such that $\chi \equiv 1$ in a neighborhood of $\partial\Omega$ and $\bar{\partial}\rho \neq 0$ on the support of χ . If we let ν denote the orthogonal projection onto I_q where defined, then $\chi\nu$ is a linear operator with $C^2(\bar{\Omega})$ coefficients. Since $u \in W^1_{(0,q)}(\Omega)$, u has a boundary trace in L^2 . By the usual density lemma (e.g., Lemma 4.3.2 in [5]) and the usual characterization of $\text{Dom}(\bar{\partial}_t^*)$ (e.g., Lemma 4.2.1 in [5]), we have that the boundary trace of $\chi\nu u$ is zero a.e. Since components of $\chi\nu u$ are in $W^1_0(\Omega)$, $\chi\nu u$ is the limit in W^1 of a sequence $u'_\ell \in C^\infty_{0,(0,q)}(\Omega)$, so we can write $u'_\ell \rightarrow \chi\nu u$ in $W^1_{(0,q)}(\Omega)$. We can also write u as a limit in $W^1_{(0,q)}(\Omega)$ of forms $\tilde{u}_\ell \in C^\infty_{(0,q)}(\bar{\Omega})$. If we set $u_\ell = u'_\ell + (1 - \chi\nu)\tilde{u}_\ell$, then $u_\ell \rightarrow u$ in $W^1_{(0,q)}(\Omega)$ since

$$\|u_\ell - u\|_{W^1} \leq \|u'_\ell - \chi\nu u\|_{W^1} + \|(1 - \chi\nu)(\tilde{u}_\ell - u)\|_{W^1}.$$

Since ν has $C^2(\bar{\Omega})$ coefficients, u_ℓ is in $C^2_{(0,q)}(\bar{\Omega})$. Furthermore, since $\nu(1 - \chi\nu) = (1 - \chi)\nu$, we have $\nu u_\ell = 0$ on the boundary of Ω , so $u_\ell \in \text{Dom}(\bar{\partial}_t^*)$. \square

With this density lemma in place, we are ready to prove the key lemma for our estimate (analogous to (3.50) in [25]).

LEMMA 4.2. — *Let $\Omega \subset M$ be a bounded domain with connected C^3 boundary satisfying weak $Z(q)$. There exist constants $C > 0$ independent of t and $C_t > 0$ depending on t such that for any $f \in W^1_{(0,q)}(\Omega)$ and differential operator T with $C^2(\bar{\Omega})$ coefficients that is tangential on the boundary of Ω , we have*

$$(4.3) \quad Q_{t,\delta}(D_T N^q_{t,\delta} f, D_T N^q_{t,\delta} f) \leq C(D_T f, D_T N^q_{t,\delta} f)_t + C \left\| N^q_{t,\delta} f \right\|^2_{t,W^1} + C_t \|f\|^2_t.$$

Proof. — We will adopt the convention that the values of $C > 0$ and $C_t > 0$ may increase from line to line. Let $u = D_T N^q_{t,\delta} f$ and let $u_\ell \in C^2_{(0,q)}(\bar{\Omega}) \cap \text{Dom}(\bar{\partial}_t^*)$ be the sequence converging to $D_T N^q_{t,\delta} f$ given by Lemma 4.1. Note that the principal part of $\bar{\partial}_t^*$ is the same as the principal part of $\bar{\partial}^*$, so only the low order terms depend on t . Hence, $[D_T, \bar{\partial}_t^*]$ has a first order component which is independent of t and a lower order component which depends on t , so

$$\begin{aligned} \left\| [D_T, \bar{\partial}_t^*] N^q_{t,\delta} f \right\|_t &\leq C \left\| N^q_{t,\delta} f \right\|_{t,W^1} + C_t \left\| N^q_{t,\delta} f \right\|_t \\ &\leq C \left\| N^q_{t,\delta} f \right\|_{t,W^1} + C_t \|f\|_t, \end{aligned}$$

where the second inequality follows from (4.1). Estimating commutators in this fashion gives us

$$\begin{aligned} Q_{t,\delta}(u, u_\ell) &\leq (D_T \bar{\partial} N^q_{t,\delta} f, \bar{\partial} u_\ell)_t + (D_T \bar{\partial}_t^* N^q_{t,\delta} f, \bar{\partial}_t^* u_\ell)_t \\ &\quad + \delta(D_T \nabla N^q_{t,\delta} f, \nabla u_\ell)_t \\ &\quad + \left(C \left\| N^q_{t,\delta} f \right\|_{t,W^1} + C_t \|f\|_t \right) \sqrt{Q_{t,\delta}(u_\ell, u_\ell)}. \end{aligned}$$

Since T is tangential, we can integrate by parts and commute again to obtain

$$\begin{aligned} Q_{t,\delta}(u, u_\ell) &\leq Q_{t,\delta}(N_{t,\delta}^q f, (D_T)^* u_\ell) + (\bar{\partial} N_{t,\delta}^q f, [(D_T)^*, \bar{\partial}] u_\ell)_t \\ &\quad + (\bar{\partial}_t^* N_{t,\delta}^q f, [(D_T)_t^*, \bar{\partial}_t^*] u_\ell)_t \\ &\quad + \delta(\nabla N_{t,\delta}^q f, [(D_T)_t^*, \nabla] u_\ell)_t \\ &\quad + \left(C \left\| N_{t,\delta}^q f \right\|_{t,W^1} + C_t \|f\|_t \right) \sqrt{Q_{t,\delta}(u_\ell, u_\ell)}. \end{aligned}$$

By definition,

$$Q_{t,\delta}(N_{t,\delta}^q f, (D_T)^* u_\ell) = (f, (D_T)^* u_\ell)_t = (D_T f, u_\ell)_t.$$

Since all terms with u_ℓ can now be estimated by the W^1 norm of u_ℓ , we can take limits and obtain

$$\begin{aligned} Q_{t,\delta}(u, u) &\leq (D_T f, u)_t + (\bar{\partial} N_{t,\delta}^q f, [(D_T)^*, \bar{\partial}] u)_t \\ &\quad + (\bar{\partial}_t^* N_{t,\delta}^q f, [(D_T)_t^*, \bar{\partial}_t^*] u)_t \\ &\quad + \delta(\nabla N_{t,\delta}^q f, [(D_T)_t^*, \nabla] u)_t \\ &\quad + \left(C \left\| N_{t,\delta}^q f \right\|_{t,W^1} + C_t \|f\|_t \right) \sqrt{Q_{t,\delta}(u, u)}. \end{aligned}$$

Now, we may also use the estimate

$$\begin{aligned} \left(C \left\| N_{t,\delta}^q f \right\|_{t,W^1} + C_t \|f\|_t \right) \sqrt{Q_{t,\delta}(u, u)} \\ \leq \frac{1}{2} \left(C \left\| N_{t,\delta}^q f \right\|_{t,W^1} + C_t \|f\|_t \right)^2 + \frac{1}{2} Q_{t,\delta}(u, u) \end{aligned}$$

and absorb the last term in the left-hand side to obtain

$$\begin{aligned} Q_{t,\delta}(u, u) &\leq 2(D_T f, u)_t + 2(\bar{\partial} N_{t,\delta}^q f, [(D_T)^*, \bar{\partial}] u)_t \\ &\quad + 2(\bar{\partial}_t^* N_{t,\delta}^q f, [(D_T)_t^*, \bar{\partial}_t^*] u)_t \\ &\quad + 2\delta(\nabla N_{t,\delta}^q f, [(D_T)_t^*, \nabla] u)_t \\ &\quad + C \left\| N_{t,\delta}^q f \right\|_{t,W^1}^2 + C_t \|f\|_t^2. \end{aligned}$$

The remaining commutators will each be estimated using the same technique; we will illustrate the method with $(\bar{\partial}_t^* N_{t,\delta}^q f, [(D_T)_t^*, \bar{\partial}_t^*] u)_t$ since it could potentially involve additional factors of t . Recall that ∇_T has C^2 coefficients, while $D_T - \nabla_T$ is a zero-order operator with C^1 coefficients.

Since adjoints of zero-order operators won't introduce additional factors of t , we can break down the commutator as follows:

$$(4.4) \quad [(D_T)^*, \bar{\partial}_t^*] = [(D_T - \nabla_T)^*, \bar{\partial}^*] + [(D_T - \nabla_T)^*, (\bar{\partial}_t^* - \bar{\partial}^*)] + [(\nabla_T)^*, \bar{\partial}_t^*],$$

The zero-order component of $[(D_T - \nabla_T)^*, \bar{\partial}^*]$ is independent of t but only has continuous coefficients; we denote this A . If we let $B_t = [(D_T)^*, \bar{\partial}_t^*] - A$, then B_t is a first-order operator depending on t (in the zero-order component) with C^1 coefficients. Thus, if we commute D_T with B_t but not A we obtain

$$\begin{aligned} (\bar{\partial}_t^* N_{t,\delta}^q f, [(D_T)^*, \bar{\partial}_t^*] u)_t &= (\bar{\partial}_t^* N_{t,\delta}^q f, (A + B_t) D_T N_{t,\delta}^q f)_t \\ &\leq (\bar{\partial}_t^* N_{t,\delta}^q f, D_T B_t N_{t,\delta}^q f)_t + C \left\| N_{t,\delta}^q f \right\|_{t,W^1}^2 \\ &\quad + C_t \|f\|_t^2. \end{aligned}$$

Integrating D_T by parts, commuting with $\bar{\partial}_t^*$, and using (4.4), we have

$$\begin{aligned} (\bar{\partial}_t^* N_{t,\delta}^q f, [(D_T)^*, \bar{\partial}_t^*] u)_t &\leq (\bar{\partial}_t^* (D_T)^* N_{t,\delta}^q f, B_t N_{t,\delta}^q f)_t + C \left\| N_{t,\delta}^q f \right\|_{t,W^1}^2 \\ &\quad + C_t \|f\|_t^2. \end{aligned}$$

However, this can be bounded by

$$\begin{aligned} (\bar{\partial}_t^* N_{t,\delta}^q f, [(D_T)^*, \bar{\partial}_t^*] u)_t &\leq \sqrt{Q_{t,\delta}((D_T)^* N_{t,\delta}^q f, (D_T)^* N_{t,\delta}^q f)} \\ &\quad \times \left(C \left\| N_{t,\delta}^q f \right\|_{t,W^1} + C_t \|f\|_t \right) \\ &\quad + C \left\| N_{t,\delta}^q f \right\|_{t,W^1}^2 + C_t \|f\|_t^2. \end{aligned}$$

The same upper bound will be obtained if $\bar{\partial}_t^*$ is replaced with $\bar{\partial}$ or ∇ , so we have

$$(4.5) \quad \begin{aligned} Q_{t,\delta}(u, u) &\leq \sqrt{Q_{t,\delta}((D_T)^* N_{t,\delta}^q f, (D_T)^* N_{t,\delta}^q f)} \left(C \left\| N_{t,\delta}^q f \right\|_{t,W^1} + C_t \|f\|_t \right) \\ &\quad + 2(D_T f, u)_t + C \left\| N_{t,\delta}^q f \right\|_{t,W^1}^2 + C_t \|f\|_t^2. \end{aligned}$$

Note that $(D_T)^*_t = -D_T + a_t$, where a_t is a C^1 function depending on t . In particular,

$$\begin{aligned} Q_{t,\delta}((D_T)^*_t N_{t,\delta}^q f + u, (D_T)^*_t N_{t,\delta}^q f + u) &= Q_{t,\delta}(a_t N_{t,\delta}^q f, a_t N_{t,\delta}^q f) \\ &\leq (f, |a_t|^2 N_{t,\delta}^q f)_t + C_t \|f\|_t \sqrt{Q_{t,\delta}(a_t N_{t,\delta}^q f, a_t N_{t,\delta}^q f)}. \end{aligned}$$

After using the small constant/large constant inequality to absorb the $Q_{t,\delta}$ term on the right-hand side, we have

$$Q_{t,\delta}((D_T)_t^* N_{t,\delta}^q f + u, (D_T)_t^* N_{t,\delta}^q f + u) \leq C_t \|f\|_t^2,$$

and hence

$$Q_{t,\delta}((D_T)_t^* N_{t,\delta}^q f, (D_T)_t^* N_{t,\delta}^q f) \leq C Q_{t,\delta}(u, u) + C_t \|f\|_t^2.$$

Substituting this into (4.5) and again using the small constant/large constant estimate to absorb the $Q_{t,\delta}$ term on the left-hand side will imply (4.3). \square

Combining Lemma 4.2 and (4.2), we have

$$\|D_T N_{t,\delta}^q f\|_t^2 \leq \varepsilon C \left(|(D_T f, D_T N_{t,\delta}^q f)_t| + \|N_{t,\delta}^q f\|_{t,W^1}^2 \right) + C_t \|f\|_t^2.$$

Another application of the small constant/large constant inequality to the first term on the right-hand side gives us

$$\|D_T N_{t,\delta}^q f\|_t^2 \leq \varepsilon C \left(\|f\|_{t,W^1}^2 + \|N_{t,\delta}^q f\|_{t,W^1}^2 \right) + C_t \|f\|_t^2.$$

By Lemma 2.2 in [25], the normal derivatives of $N_{t,\delta}^q f$ consist of linear combinations of $\bar{\partial} N_{t,\delta}^q f$, $\bar{\partial}^* N_{t,\delta}^q f$, and tangential derivatives of $N_{t,\delta}^q f$. We can convert $\bar{\partial}^*$ to $\bar{\partial}_t^*$ by adding a zeroth order term, so summing over all tangential derivatives and the normal derivative gives us

$$\|N_{t,\delta}^q f\|_{t,W^1}^2 \leq \varepsilon C \left(\|f\|_{t,W^1}^2 + \|N_{t,\delta}^q f\|_{t,W^1}^2 \right) + C_t \|f\|_t^2.$$

When ε is sufficiently small (and hence t is sufficiently large), we can absorb the W^1 norm of $N_{t,\delta}^q f$ on the left-hand side, obtaining

$$(4.6) \quad \|N_{t,\delta}^q f\|_{t,W^1}^2 \leq \varepsilon C \|f\|_{t,W^1}^2 + C_t \|f\|_t^2.$$

Since all constants have been chosen independently of δ , the usual limiting argument will give us W^1 estimates for N_t^q . Standard arguments now give us

THEOREM 4.3. — *Let M be an n -dimensional Stein manifold, and let Ω be a bounded subset of M with connected C^3 boundary satisfying weak $Z(q)$ for some $1 \leq q \leq n - 1$. Then there exists a constant $\tilde{t} > 0$ such that for all $t > \tilde{t}$ and $-\frac{1}{2} \leq s \leq 1$ we have*

- (1) *The weighted $\bar{\partial}$ -Neumann operator N_t^q is continuous in $W_{(0,q)}^s(\Omega)$.*

- (2) The canonical solution operators to $\bar{\partial}$ given by $\bar{\partial}_t^* N_t^q : W_{(0,q)}^s(\Omega) \rightarrow W_{(0,q-1)}^s(\Omega)$ and $N_t^q \bar{\partial}_t^* : W_{(0,q+1)}^s(\Omega) \cap \text{Dom}(\bar{\partial}_t^*) \rightarrow W_{(0,q)}^s(\Omega)$ are continuous.
- (3) The canonical solution operators to $\bar{\partial}_t^*$ given by $\bar{\partial} N_t^q : W_{(0,q)}^s(\Omega) \rightarrow W_{(0,q+1)}^s(\Omega)$ and $N_t^q \bar{\partial} : W_{(0,q-1)}^s(\Omega) \cap \text{Dom}(\bar{\partial}) \rightarrow W_{(0,q)}^s(\Omega)$ are continuous.
- (4) For every $f \in W_{(0,q)}^s(\Omega) \cap \ker \bar{\partial}$ there exists a $u \in W_{(0,q-1)}^s(\Omega)$ such that $\bar{\partial} u = f$.

Remark 4.4. — Theorem 4.3 is identical to Theorem 1.2 except for the additional hypothesis that $\partial\Omega$ is connected. Also, $N_t^q \bar{\partial}_t^*$ is a problematic solution operator to $\bar{\partial}$ since it places a boundary condition on the data. However, estimates for this operator are needed to obtain estimates for the dual operator $\bar{\partial} N_t^q$ in the dual Sobolev space.

Proof. — We have already completed the proof of (1) when $s = 0$ and $s = 1$, so interpolation will give us the result for $0 < s < 1$. We emphasize that the weighted Sobolev spaces used in (4.6) are equivalent to standard Sobolev spaces which are amenable to interpolation. When $-\frac{1}{2} \leq s < 0$ we use the duality between W^s and W^{-s} and the fact that N_t^q is self-adjoint:

$$\begin{aligned} \|N_t^q f\|_{W^s} &\leq C_t \sup_{h \in W^{-s}(\Omega), \|h\|_{W^{-s}=1}} |(N_t^q f, h)_t| \\ &\leq C_t \sup_{h \in W^{-s}(\Omega), \|h\|_{W^{-s}=1}} |(f, N_t^q h)_t| \\ &\leq C_t \|f\|_{W^s}. \end{aligned}$$

For $\bar{\partial}_t^* N_t^q$ and $\bar{\partial} N_t^q$, we use Lemma 3.2 in [25]. Although this Lemma assumes $N_t^q f$ is smooth, we can use elliptic regularization and Lemma 4.1 as before to obtain sufficient regularity. Interpolation will give us estimates for $0 \leq s \leq 1$.

For $N_t^q \bar{\partial}$ and $N_t^q \bar{\partial}_t^*$, we first note that since $f \in W_{(0,q)}^1(\Omega)$ implies $N_t^q f$, $\bar{\partial} N_t^q f$, and $\bar{\partial}_t^* N_t^q f$ are all in $W^1(\Omega)$, we can conclude that

$$\bar{\partial} D_T N_t^q f = [\bar{\partial}, D_T] N_t^q f + D_T \bar{\partial} N_t^q f \in L_{(0,q+1)}^2(\Omega)$$

and

$$\bar{\partial}_t^* D_T N_t^q f = [\bar{\partial}_t^*, D_T] N_t^q f + D_T \bar{\partial}_t^* N_t^q f \in L_{(0,q-1)}^2(\Omega).$$

Thus we can substitute into (3) of Proposition 3.1 and estimate commutators to obtain

$$\begin{aligned} & \|D_T N_t^q f\|_t^2 \\ & \leq \varepsilon \left(\|D_T \bar{\partial} N_t^q f\|_t^2 + \|D_T \bar{\partial}_t^* N_t^q f\|_t^2 + C \|N_t^q f\|_{t,W^1}^2 \right) + C_t \|N_t^q f\|_t^2. \end{aligned}$$

Estimating the normal derivatives using Lemma 2.2 in [25] we can estimate the W^1 norm of $N_t^q f$, and hence the error term on the right hand side can be absorbed into the left for sufficiently small $\varepsilon > 0$. Hence we have (after possibly increasing t_ε)

$$\|N_t^q f\|_{t,W^1}^2 \leq \varepsilon \left(\|\bar{\partial} N_t^q f\|_{t,W^1}^2 + \|\bar{\partial}_t^* N_t^q f\|_{t,W^1}^2 \right) + C_t \|N_t^q f\|_t^2.$$

For $f_1 \in W_{(0,q-1)}^2(\Omega)$, let $f = \bar{\partial} f_1$ to obtain

$$\|N_t^q \bar{\partial} f_1\|_{t,W^1}^2 \leq \varepsilon \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1}^2 + C_t \|N_t^q \bar{\partial} f_1\|_t^2.$$

To estimate the projection, we use integration by parts:

$$\begin{aligned} \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1}^2 & \leq (N_t^q \bar{\partial} f_1, \bar{\partial} f_1)_{t,W^1} + C \|N_t^q \bar{\partial} f_1\|_{t,W^1} \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1} \\ & \quad + C_t \|N_t^q \bar{\partial} f_1\|_t \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1} \\ & \quad + C_t \|N_t^q \bar{\partial} f_1\|_{t,W^1} \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_t. \end{aligned}$$

Repeated use of the small constant/large constant inequality allows us to absorb the $\|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1}^2$ terms on the left-hand side and obtain

$$\begin{aligned} \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1}^2 & \leq C (N_t^q \bar{\partial} f_1, \bar{\partial} f_1)_{t,W^1} + C \|N_t^q \bar{\partial} f_1\|_{t,W^1}^2 \\ & \quad + C_t \|N_t^q \bar{\partial} f_1\|_t^2 + C_t \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_t^2. \end{aligned}$$

A second integration by parts gives us

$$\begin{aligned} \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1}^2 & \leq C (\bar{\partial}_t^* N_t^q \bar{\partial} f_1, f_1)_{t,W^1} + C \|N_t^q \bar{\partial} f_1\|_{t,W^1} \|f_1\|_{t,W^1} \\ & \quad + C_t \|N_t^q \bar{\partial} f_1\|_t \|f_1\|_{t,W^1} + C \|N_t^q \bar{\partial} f_1\|_{t,W^1}^2 \\ & \quad + C_t \|N_t^q \bar{\partial} f_1\|_t^2 + C_t \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_t^2, \end{aligned}$$

and more small constant/large constant inequalities yield

$$\begin{aligned} \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1}^2 & \leq C \|f_1\|_{t,W^1}^2 + C \|N_t^q \bar{\partial} f_1\|_{t,W^1}^2 \\ & \quad + C_t \|f_1\|_t^2 + C_t \|N_t^q \bar{\partial} f_1\|_t^2 + C_t \|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_t^2. \end{aligned}$$

All of the terms with C_t in front are bounded by $\|f_1\|_t^2$, so

$$\|\bar{\partial}_t^* N_t^q \bar{\partial} f_1\|_{t,W^1}^2 \leq C \|f_1\|_{t,W^1}^2 + C \|N_t^q \bar{\partial} f_1\|_{t,W^1}^2 + C_t \|f_1\|_t^2.$$

Now we can substitute into the original estimate for $\|N_t^q \bar{\partial} f_1\|_{t,W^1}^2$ and, by making ε sufficiently small and adjusting t_ε we obtain

$$\|N_t^q \bar{\partial} f_1\|_{t,W^1}^2 \leq \varepsilon \|f_1\|_{t,W^1}^2 + C_t \|f_1\|_t^2.$$

For $f_2 \in W_{(0,q+1)}^2(\Omega) \cap \text{Dom}(\bar{\partial}_t^*)$, let $f = \bar{\partial}_t^* f_2$ and use similar techniques to estimate $N_t^q \bar{\partial}_t^* f_2$. Density lets us generalize to f_1 and f_2 in W^1 .

As before, we interpolate to obtain estimates for $0 < s < 1$ and use duality to obtain estimates for $-\frac{1}{2} \leq s < 0$ (since we now have estimates for the adjoint of each operator). □

5. Proofs of Main Theorems

When $s < \frac{1}{2}$, we immediately obtain a solution operator for the $\bar{\partial}$ -Cauchy Problem (see Section 9.1 in [5]).

COROLLARY 5.1. — *Let M be an n -dimensional Stein manifold, let Ω be a bounded subset of M with connected C^3 boundary satisfying weak $Z(n - q - 1)$ for some $1 \leq q < n - 1$, and let $-\frac{1}{2} \leq s < \frac{1}{2}$. For every $f \in W_{(0,q+1)}^s(M) \cap \ker \bar{\partial}$ supported in $\bar{\Omega}$ there exists a $u \in W_{(0,q)}^s(M)$ supported in $\bar{\Omega}$ such that $\bar{\partial}u = f$, and the solution operator is continuous.*

Proof. — The proof is identical to that given for Proposition 3.4 in [22]. It suffices to have estimates for $\bar{\partial}N^{(n,n-q-1)}$. On $Z(n - q - 1)$ domains we have estimates for the operator $\bar{\partial}N^{(0,n-q-1)}$, but these are equivalent since the holomorphic component of (p,q) -forms has no impact on our estimates (except in the curvature terms, but these are all dominated when t is large). □

Furthermore, we can now use techniques of Shaw [23] to remove the requirement that the boundary of Ω be connected.

COROLLARY 5.2. — *Let M be an n -dimensional Stein manifold, and let Ω be a bounded subset of M with C^3 boundary satisfying weak $Z(q)$ for some $1 \leq q < n - 1$. For every $f \in L_{(0,q)}^2(\Omega) \cap \ker \bar{\partial}$ there exists a $u \in L_{(0,q-1)}^2(\Omega)$ such that $\bar{\partial}u = f$. In particular, $\mathcal{H}^q(\Omega) = \{0\}$.*

Remark 5.3. — This is also known for bounded weak $Z(n - 1)$ domains, by Remark 2.10 and Theorem 3.2.

Proof. — Without loss of generality, assume that Ω is connected. By Lemma 2.9 we have the decomposition $\Omega = \Omega_1 \setminus \bigcup_{j=2}^m \overline{\Omega}_j$ where Ω_1 is a weak $Z(q)$ domain with connected boundary and each Ω_j with $2 \leq j \leq m$ is a weak $Z(n - q - 1)$ domain with connected boundary and $\overline{\Omega}_j \subset \Omega_1$.

The proof now follows like the proof of Theorem 3.2 in [23], but we must make a few clarifying remarks. Shaw's proof initially finds a solution U in $W^{-1}(\Omega_1)$, and then proposes two techniques for regularizing this to a solution in $L^2(\Omega_1)$. The first technique requires approximating Ω_1 from within by smooth strongly pseudoconvex domains. While we can approximate from within by smooth $Z(q)$ domains, the role of Υ on these domains is unpredictable, so the constants in our estimates may not be uniform. Hence, we use the second technique, which decomposes U into a component supported near the boundary of Ω_1 and a compactly supported component. This technique requires solving $\bar{\partial}$ for compactly supported forms in Ω_1 , but as Shaw points out, this can be accomplished by solving $\bar{\partial}$ on a ball containing Ω_1 .

This gives us vanishing of $\mathcal{H}_t^q(\Omega)$. However, this is isomorphic to a Dolbeault cohomology group which is independent of t , so the dimension of $\mathcal{H}_t^q(\Omega)$ must also be independent of t . Hence, $\mathcal{H}^q(\Omega) = \{0\}$ as well. \square

Now that the vanishing of the space of harmonic $(0, q)$ -forms has been established, Theorem 1.1 will follow for the unweighted operators and spaces (see Theorem 4.4.1 in [5]) from Theorem 3.2.

In the proof of Theorem 4.3 we required a connected boundary only because it gave us vanishing for the space of harmonic $(0, q)$ -forms. Now that we've established this for disconnected boundaries, the proof of Theorem 4.3 can be carried through for all bounded domains, as in the statement of Theorem 1.2.

In [21], Shaw uses the $\bar{\partial}$ operator to solve $\bar{\partial}_b$ extrinsically on smooth pseudoconvex domains. Since we are working on C^3 domains, we will more closely follow the methods of [22], which are adapted to work on non-smooth domains. We now prove Theorem 1.3.

Proof of Theorem 1.3. — We outline the proof, following the construction in [22]. By embedding M in \mathbb{C}^{2n+1} , we can pullback a ball containing the image of $\partial\Omega$ to obtain a strictly pseudoconvex set B such that $\overline{\Omega} \subset B$. Let $\Omega^+ = B \setminus \overline{\Omega}$ and $\Omega^- = \Omega$. In [11] a Martinelli-Bochner-Koppelman type kernel is constructed for Stein manifolds, and in [17] it is shown that the transformation induced by this kernel satisfies a jump formula. As a result, there exists an integral kernel $K_q(\zeta, z)$ of type $(0, q)$ in z and $(n, n - q - 1)$ in ζ satisfying a Martinelli-Bochner-Koppelman formula such that we can

define

$$\int_{\partial\Omega} K_q(\zeta, z) \wedge f(\zeta) = \begin{cases} f^+(z) & z \in \Omega^+ \\ f^-(z) & z \in \Omega^- \end{cases}$$

(see section 2.4 in [11]). If, by an abuse of notation, we extend $f^+(z)$ and $f^-(z)$ to $\partial\Omega$ by considering non-tangential limits, we have the jump formula

$$\tau f^+(z) - \tau f^-(z) = (-1)^q f(z)$$

on $\partial\Omega$ where τ is defined in (2.1) (see Proposition 2.3.1 in [17]). Since $\bar{\partial}_b f = 0$, we have $\bar{\partial} f^+ = 0$ and $\bar{\partial} f^- = 0$. Since f^+ and f^- have non-tangential boundary values in L^2 , they have components in $W^{1/2}$.

Let E denote a linear extension operator from Ω^+ to B that is continuous in the Sobolev spaces $W^s(\Omega^+)$ for $0 \leq s \leq 1/2$. Applying this componentwise to f^+ , we obtain a $(0, q)$ -form $E f^+$ that is in $W^{1/2}(B)$. Since $\bar{\partial} f^+ = 0$ on Ω^+ , we have $\bar{\partial} E f^+$ supported in Ω^- . Furthermore, $\bar{\partial} E f^+$ has components in $W^{-1/2}(M)$, so by Corollary 5.1 there exists $V \in W_{(0,q)}^{-1/2}(M)$ supported in Ω^- satisfying $\bar{\partial} V = \bar{\partial} E f^+$. Thus $\tilde{f}^+ = E f^+ - V \in W_{(0,q)}^{-1/2}(B)$ satisfies $\bar{\partial} \tilde{f}^+ = 0$ on B and $\tilde{f}^+ = f^+$ on Ω^+ .

We may now use the canonical solution operator to define $u^+ = \bar{\partial}^* N_B^q \tilde{f}^+$ and $u^- = \bar{\partial}_t^* N_{t,\Omega}^q f^-$. By interior regularity for N_B^q , u^+ gains a derivative on compact subsets of B , so u^+ has components in $W^{1/2}$ on a neighborhood of $\bar{\Omega}$. By Theorem 4.3, u^- also has components in $W^{1/2}$. Since each of these forms satisfies an elliptic system, they have trace values in $L^2(\partial\Omega)$, so abusing notation we can write $u = (-1)^q(u^+ - u^-)$ on $\partial\Omega$. □

6. Examples

Our first goal in this section is to motivate Definition 2.1 by constructing an example where this definition holds but earlier definitions fail. In [9] we proposed a more restrictive definition for weak $Z(q)$. In the context of bounded domains in \mathbb{C}^n with connected boundaries, this was equivalent to $(q - 1)$ -pseudoconvexity in the sense of [27]. The key difference is that in $(q - 1)$ -pseudoconvexity the eigenvalues of Υ must all be zero or one. This can often be achieved locally by changing the metric. The critical issue seems to be that if Υ is continuous then $(q - 1)$ -pseudoconvexity forces the rank of Υ and the rank of $I - \Upsilon$ to be locally constant. We will construct an example where this is impossible, but the added flexibility of Definition 2.1 still allows us establish closed range. We summarize the first result of this section as follows:

PROPOSITION 6.1. — *There exists a smooth bounded domain $\Omega \subset \mathbb{C}^3$ with $0 \in \partial\Omega$ such that:*

- (1) *There does not exist a hermitian metric in a neighborhood of the origin for which $\partial\Omega$ is 1-pseudoconvex at the origin.*
- (2) *In the Euclidean metric, $\partial\Omega$ is weakly $Z(2)$.*

In what follows, we assume $M = \mathbb{C}^n$ is equipped with a strictly plurisubharmonic exhaustion function φ . After adding a pluriharmonic polynomial to φ , we may assume $\varphi(0) = 0$, $d\varphi(0) = 0$, and $\frac{\partial^2 \varphi}{\partial z_j \partial \bar{z}_k}(0) = 0$ for all $1 \leq j, k \leq n$. Under these assumptions, φ is strictly convex in a neighborhood of the origin, so $\varphi(z) - R^2$ is a defining function for a strictly convex domain when R is sufficiently small.

DEFINITION 6.2. — *Let $\Omega \subset \mathbb{C}^n$ be a domain with C^3 boundary. For $p \in \partial\Omega$, let the Kähler form ω in a neighborhood of p be given by $\omega = i\partial\bar{\partial}\varphi$ for some smooth strictly plurisubharmonic function φ satisfying $\varphi(p) = 0$, $d\varphi(p) = 0$, and $\frac{\partial^2 \varphi}{\partial z_j \partial \bar{z}_k}(p) = 0$ for all $1 \leq j, k \leq n$. For $1 \leq q \leq n - 1$, we say the weak $Z(q)$ property for Ω is radially stable at p if there exists a neighborhood U of p and a real $\Upsilon \in T^{1,1}(\partial\Omega \cap U)$ satisfying*

- (1) $|\theta|^2 \geq (i\theta \wedge \bar{\theta})(\Upsilon) \geq 0$ for all $\theta \in \Lambda^{1,0}(\partial\Omega \cap U)$.
- (2) *For every neighborhood \tilde{U} of p relatively compact in U , there exists $\tilde{\varepsilon} > 0$ such that $\mu_1^\varepsilon + \dots + \mu_q^\varepsilon - \mathcal{L}^\varepsilon(\Upsilon) \geq 0$ on $(U \setminus \tilde{U}) \cap \partial\Omega$ for all $0 \leq \varepsilon < \tilde{\varepsilon}$ where $\mathcal{L}^\varepsilon(i\bar{L} \wedge L) = (i\partial\bar{\partial}\rho + i\varepsilon\partial\varphi \wedge \bar{\partial}\varphi)(i\bar{L} \wedge L)$ for $L \in T^{1,0}(\partial\Omega)$ and $\mu_1^\varepsilon, \dots, \mu_{n-1}^\varepsilon$ are the eigenvalues of \mathcal{L}^ε in increasing order.*
- (3) $\omega(\Upsilon) \neq q$ on $\partial\Omega \cap U$.

Remark 6.3. — The term “radially stable” reflects the observation that in orthonormal coordinates $\varphi(z) = |z - p|^2 + O(|z - p|^3)$, so $d\varphi$ points in the radial direction near p and \mathcal{L}^ε represents a perturbation of \mathcal{L} in the radial direction.

Remark 6.4. — A simple obstruction to radial stability would be the situation where $\mu_q = \mu_{q+1} = 0$, $i\partial\varphi \wedge \bar{\partial}\varphi(\Upsilon) > 0$ is necessary for condition (2) of Definition 6.2 to hold (because of local behavior of the eigenspaces), and $\partial\varphi$ lies in the kernel of \mathcal{L} , i.e., whenever L is in the kernel of (the matrix) \mathcal{L} , $\partial\varphi(L) = 0$. In this case $\mu_1^\varepsilon + \dots + \mu_q^\varepsilon = \mu_1 + \dots + \mu_q$, so $\mu_1^\varepsilon + \dots + \mu_q^\varepsilon - \mathcal{L}^\varepsilon(\Upsilon) < 0$ for all $\varepsilon > 0$.

Remark 6.5. — If $p \in \partial\Omega \subset \mathbb{C}^n$ and U is a neighborhood of p so that $U \cap \partial\Omega$ satisfies weak $Z(n - 1)$, then weak $Z(n - 1)$ is radially stable at p .

As in the proof of Proposition 2.3, we know $\mu_1 + \dots + \mu_{n-1} = \mathcal{L}((\tau\omega)^\sharp)$. Thus, we have

$$\begin{aligned} \mu_1^\varepsilon + \dots + \mu_{n-1}^\varepsilon - \mathcal{L}^\varepsilon(\Upsilon) &= \mathcal{L}^\varepsilon((\tau\omega)^\sharp - \Upsilon) \geq \\ \mathcal{L}((\tau\omega)^\sharp - \Upsilon) &= \mu_1 + \dots + \mu_{n-1} - \mathcal{L}(\Upsilon) \geq 0. \end{aligned}$$

Our motivation for studying radial stability is that it allows us to construct global examples from local data, as per the following proposition.

PROPOSITION 6.6. — *Let Ω_1 be an unbounded C^k domain in \mathbb{C}^n , $k \geq 3$, with $0 \in \partial\Omega_1$. Moreover, suppose that for some $1 \leq q \leq n - 1$, the weak $Z(q)$ property for Ω_1 is radially stable at the origin. There exists a bounded domain $\Omega \subset \Omega_1$ defined by a C^k defining function ρ such that $\Omega \cap U = \Omega_1 \cap U$ on a neighborhood U of the origin and Ω satisfies weak $Z(q)$.*

Proof. — Let ρ_1 be a defining function for Ω_1 . After a rotation and rescaling, we may assume $\partial\rho_1(0) = \frac{i}{2}dz_n$. We write $z_n = x_n + iy_n$. Using the Implicit Function Theorem, we may find a C^k function $P(z_1, \dots, z_{n-1}, x_n)$ such that in some relatively compact neighborhood V of the origin, $\rho_1(z) < 0$ if and only if $-y_n + P(z_1, \dots, z_{n-1}, x_n) < 0$ and we have $|P| \leq O(|z|^2)$. Without loss of generality, we may assume $\rho_1|_V = -y_n + P(z_1, \dots, z_{n-1}, x_n)$. We have the projection $\pi : V \rightarrow \partial\Omega_1$ defined by $\pi(z) = (z_1, \dots, z_{n-1}, x_n + iP(z_1, \dots, z_{n-1}, x_n))$. We will frequently make use of the fact that our chosen defining function ρ_1 satisfies $\partial\rho_1(z) = \partial\rho_1(\pi(z))$ and $\partial\bar{\partial}\rho_1(z) = \partial\bar{\partial}\rho_1(\pi(z))$, where this composition is interpreted componentwise (we do not use the pullback since π is not holomorphic). Since π is not holomorphic, we define the modified pushforward $\tilde{\pi}_* : T^{1,0}(V) \rightarrow T^{1,0}(\partial\Omega_1)$ by $\tilde{\pi}_*(L) = L - \frac{\partial\rho_1(L)}{\partial\rho_1/\partial z_n} \frac{\partial}{\partial z_n}$. Note that $\frac{\partial\rho_1}{\partial z_n} = \frac{1}{2} \left(\frac{\partial P}{\partial x_n} + i \right)$, so $\left| \frac{\partial\rho_1}{\partial z_n} \right| \geq \frac{1}{2}$. Thus

$$|L - \tilde{\pi}_*(L)| \leq \sqrt{2} |\partial\rho_1(L)|.$$

On V , we have

$$(6.1) \quad \left| i\partial\bar{\partial}\rho_1(i\bar{L} \wedge L) - \mathcal{L}_{\rho_1}|_{\pi(z)}(i\overline{\tilde{\pi}_*(L)} \wedge \tilde{\pi}_*(L)) \right| \leq O(\|P\|_{C^2(V)} |L| |\partial\rho_1(L)|).$$

Since $\|P\|_{C^2(V)}$ will not depend on any of our parameters, we will suppress it in future error terms.

Let $\rho_2(z) = \varphi(z) - R^2$ for some $0 < R < 1$. Observe that in coordinates (w_1, \dots, w_n) that are orthonormal at 0, $\varphi(w) = |w|^2 + O(|w|^3)$, so ρ_2 defines a bounded strictly convex domain for $R > 0$ sufficiently small. We choose $R > 0$ sufficiently small so that this domain is contained in V . Since $d\varphi(0) =$

0, on V we have

$$(6.2) \quad |\partial\varphi| \leq O(\|\varphi\|_{C^2(V)} R).$$

As with $\|P\|_{C^2(V)}$, we will not continue to track $\|\varphi\|_{C^2(V)}$ in our error terms. We will assume henceforth that $R > 0$ is chosen sufficiently small so that $\left| \frac{\partial\varphi}{\partial y_n} \right|$ is bounded by $\frac{1}{2}$, and hence

$$(6.3) \quad \left| 1 + \frac{\partial\varphi}{\partial y_n} \right| \geq \frac{1}{2}$$

Our goal is to construct a C^k approximation to $\max\{\rho_1, \rho_2\}$. Constructing such an approximation is relatively easy, but showing that weak $Z(q)$ is satisfied for points where $\rho_1(z) \approx \rho_2(z)$ (i.e., those points which lie near the smoothed corner of our new domain) will take some work. Let $\chi \in C_0^\infty(\mathbb{R})$ be a smooth, even, nonnegative function satisfying $\text{supp } \chi = [-1, 1]$ and $\int_{\mathbb{R}} \chi = 1$. For $0 < r < 1$, define

$$\psi_r(x) = \int_{-\infty}^x r^{-1}(x-t)\chi(r^{-1}t) dt.$$

We can check that $\psi_r \in C^\infty(\mathbb{R})$ is a convex increasing function satisfying $\psi_r(x) = 0$ when $x \leq -r$ and $\psi_r(x) = x$ for $x \geq r$ (note that we use $\int_{-1}^1 t\chi(t) dt = 0$ since χ is even). Furthermore,

$$(6.4) \quad \psi_r(x) \leq (x+r) \int_{-\infty}^x r^{-1}\chi(r^{-1}t) dt = (x+r)\psi'_r(x).$$

We can now define

$$\rho(z) = \rho_1(z) + \psi_r(\rho_2(z) - \rho_1(z)).$$

For $|\rho_2 - \rho_1| \geq r$, $\rho(z) = \max\{\rho_1(z), \rho_2(z)\}$, so this is a candidate for our C^k defining function. At the origin $\rho_1(0) - \rho_2(0) = R^2$, so for $R^2 > r$, $\rho(z) = \rho_1(z)$ in a neighborhood of the origin. Thus, by taking $R^2 > r > 0$ sufficiently small, $\rho(z)$ defines a bounded domain Ω and $\Omega \cap U = \Omega_1 \cap U$ for a sufficiently small neighborhood $U \subset V$ of the origin.

We compute

$$(6.5) \quad \partial\rho = (1 - \psi'_r(\rho_2 - \rho_1))\partial\rho_1 + \psi'_r(\rho_2 - \rho_1)\partial\varphi$$

and

$$(6.6) \quad \begin{aligned} i\partial\bar{\partial}\rho &= (1 - \psi'_r(\rho_2 - \rho_1))i\partial\bar{\partial}\rho_1 + \psi'_r(\rho_2 - \rho_1)\omega \\ &\quad + i\psi''_r(\rho_2 - \rho_1)(\partial\varphi - \partial\rho_1) \wedge (\bar{\partial}\varphi - \bar{\partial}\rho_1). \end{aligned}$$

Since ρ_2 is strictly convex, $\partial\Omega$ is strictly convex when $\rho_2 - \rho_1 \geq r$. Thus, there must exist some $0 < r_0 < r$ such that $\partial\Omega$ is strictly convex when

$\rho_2 - \rho_1 > r_0$. For $z \in \partial\Omega$ satisfying $\rho_2(z) - \rho_1(z) \leq r_0$ and $L \in T^{1,0}(\partial\Omega)$, we can use $\partial\rho(L) = 0$ at z , (6.2), and (6.5) to show

$$(6.7) \quad |\partial\rho_1(L)| \leq O\left(\frac{R\psi'_r(\rho_2 - \rho_1)}{1 - \psi'_r(\rho_2 - \rho_1)}|L|\right).$$

By construction, $\frac{1}{1 - \psi'_r(\rho_2 - \rho_1)} \leq \frac{1}{1 - \psi'_r(r_0)} < \infty$ when $\rho_2 - \rho_1 \leq r_0 < r$. Using this with (6.6) and (6.1), we have

$$\begin{aligned} \mathcal{L}_\rho(i\bar{L} \wedge L) \geq & (1 - \psi'_r(\rho_2 - \rho_1))\mathcal{L}_{\rho_1}|_{\pi(z)}(i\overline{\tilde{\pi}_*(L)} \wedge \tilde{\pi}_*(L)) + \psi'_r(\rho_2 - \rho_1)|L|^2 \\ & + \psi''_r(\rho_2 - \rho_1)|\partial\varphi(L) - \partial\rho_1(L)|^2 - O\left(\frac{R\psi'_r(\rho_2 - \rho_1)}{1 - \psi'_r(r_0)}|L|^2\right). \end{aligned}$$

By choosing $R > 0$ sufficiently small, we may assume $\psi'_r(\rho_2 - \rho_1)|L|^2$ is strictly larger than the error term, so when $\rho_2 - \rho_1 > -r$ we are adding a positive definite form to the Levi-form of Ω_1 . Hence, Ω satisfies $Z(q)$ whenever $\rho_2 - \rho_1 > -r$. Since we know Ω inherits weak $Z(q)$ from Ω_1 when $\rho_2 - \rho_1 < -r$, we conclude that we only need to show that weak $Z(q)$ is satisfied on a neighborhood of the set of boundary points where $\rho_2 - \rho_1 = -r$. To that end, we define

$$K = \{z \in \partial\Omega : \rho_2(z) - \rho_1(z) = -r\} \subset \partial\Omega \cap \partial\Omega_1.$$

For the delicate estimates near K , we will need an estimate for $|z - \pi(z)|$. Since $\frac{\partial\rho_1}{\partial y_n} = -1$, we can use (6.5) to show

$$\frac{\partial\rho}{\partial y_n} + 1 = \psi'_r(\rho_2 - \rho_1) \left(1 + \frac{\partial\varphi}{\partial y_n}\right).$$

Since $\rho_1(\pi(z)) = 0$, we have $\rho(\pi(z)) = \psi_r(\rho_2(\pi(z)))$ by definition. If $z \in \partial\Omega$, we also have $\rho(z) = 0$ by definition, so we can integrate $\frac{\partial\rho}{\partial y_n} + 1$ in y_n to obtain

$$(6.8) \quad \begin{aligned} & \psi_r(\rho_2(\pi(z))) + (P(z_1, \dots, z_{n-1}, x_n) - y_n) \\ & = \int_{y_n}^{P(z_1, \dots, z_{n-1}, x_n)} \left(\psi'_r(\rho_2 - \rho_1) \left(1 + \frac{\partial\varphi}{\partial y_n}\right)\right)(z_1, \dots, z_{n-1}, x_n + it) dt. \end{aligned}$$

This equality motivates the following estimates. Note that each of the following conditions is open and trivial on K , so for every $\frac{1}{4} > \eta > 0$ there exists a neighborhood $W_\eta \subset V$ of K satisfying

- (1) $\rho_2(z) < 0$ on W_η .
- (2) If $z \in \partial\Omega \cap W_\eta$, then the line segment $[z, \pi(z)] \subset W_\eta$.
- (3) $\left|\psi'_r(\rho_2 - \rho_1) \left(1 + \frac{\partial\varphi}{\partial y_n}\right)\right| < \eta$ on W_η .
- (4) $\left|\psi''_r(\rho_2 - \rho_1) \left(1 + \frac{\partial\varphi}{\partial y_n}\right)\right| < \eta$ on W_η .

Note that (2) can be accomplished if W_η is the cross product of a ball in $\mathbb{C}^{n-1} \times \mathbb{R}$ with coordinates $(z_1, \dots, z_{n-1}, x_n)$ and an interval in \mathbb{R} with coordinate y_n . Since (4) tells us that $\left| \frac{\partial}{\partial y_n} \psi'_r(\rho_2 - \rho_1) \right| < \eta$ on W_η , the Mean Value Theorem implies

$$|\psi'_r(\rho_2(\pi(z))) - \psi'_r(\rho_2(z) - \rho_1(z))| < \eta |\pi(z) - z|.$$

Using (6.4) and (1), we have

$$\begin{aligned} \psi_r(\rho_2(\pi(z))) &\leq (\rho_2(\pi(z)) + r)\psi'_r(\rho_2(\pi(z))) \\ &\leq r(\psi'_r(\rho_2(z) - \rho_1(z)) + \eta |\pi(z) - z|) \end{aligned}$$

Since $|\pi(z) - z| = |P(z_1, \dots, z_{n-1}, x_n) - y_n|$, we can substitute this into (6.8) along with property (3) to show

$$|\pi(z) - z| < r(\psi'_r(\rho_2(z) - \rho_1(z)) + \eta |\pi(z) - z|) + \eta |\pi(z) - z|.$$

Since $0 < \eta < \frac{1}{4}$ and $0 < r < 1$ imply $0 < \frac{1}{1-(r+1)\eta} < 2$, we have on W_η

$$(6.9) \quad |\pi(z) - z| < \frac{r}{2} \psi'_r(\rho_2 - \rho_1).$$

Using (6.3) with (3), we have $\psi'_r(\rho_2 - \rho_1) < 2\eta < \frac{1}{2}$ on W_η , so $\frac{1}{1-\psi'_r(\rho_2-\rho_1)} < 2$, and we may neglect the corresponding coefficient in (6.7) on W_η . With (6.2), (6.7), (6.9), we can use our assumption that $r < R^2 < R$ to prove for $z \in W_\eta$ and $L \in T^{1,0}(\partial\Omega \cap W_\eta)$:

$$(6.10) \quad |\partial\varphi|_{\pi(z)}(\tilde{\pi}_*(L)) - \partial\varphi(L) < O(R\psi'_r(\rho_2 - \rho_1)|L|)$$

Note that (6.3) and (4) imply that $\psi''_r(\rho_2 - \rho_1) \leq 2\eta < \frac{1}{2}$ on W_η , so this will not impact our error terms. Given $L \in T^{1,0}(\partial\Omega \cap W_\eta)$, we can use (6.1), (6.6), (6.7), and (6.10), to show

$$\begin{aligned} \mathcal{L}_\rho(i\bar{L} \wedge L) &\geq (1 - \psi'_r(\rho_2 - \rho_1))\mathcal{L}_{\rho_1}|_{\pi(z)}(i\overline{\tilde{\pi}_*(L)} \wedge \tilde{\pi}_*(L)) + \psi'_r(\rho_2 - \rho_1)|L|^2 \\ &\quad + \psi''_r(\rho_2 - \rho_1) |\partial\varphi|_{\pi(z)}(\tilde{\pi}_*(L))|^2 - O(|L|^2 R\psi'_r(\rho_2 - \rho_1)) \end{aligned}$$

If we set $\varepsilon(z) = \left(\frac{\psi''_r(\rho_2-\rho_1)}{1-\psi'_r(\rho_2-\rho_1)} \circ \pi^{-1} \right) (z)$, where $\pi^{-1} : \partial\Omega_1 \rightarrow \partial\Omega$ is well-defined on W_η , we can define $\mathcal{L}^\varepsilon_{\rho_1} = \mathcal{L}_{\rho_1} + i\varepsilon(z)\partial\varphi \wedge \bar{\partial}\varphi$. If the eigenvalues of $\mathcal{L}^\varepsilon_{\rho_1}$ in increasing order are given by $\mu_1^\varepsilon, \dots, \mu_{n-1}^\varepsilon$, then we have

$$(6.11) \quad \begin{aligned} \mu_1 + \dots + \mu_q &\geq (1 - \psi'_r(\rho_2 - \rho_1)) ((\mu_1^\varepsilon + \dots + \mu_q^\varepsilon)|_{\pi(z)}) \\ &\quad + q\psi'_r(\rho_2 - \rho_1) - O(R\psi'_r(\rho_2 - \rho_1)). \end{aligned}$$

Let Υ_1 satisfy the conditions of Definition 6.2. We may assume that $\omega(\Upsilon_1) < q$. Otherwise, we define $\tilde{\rho}_1 = -\rho_1$ and $\tilde{\Upsilon}_1 = (\tau\omega)^\sharp - \Upsilon_1$. As in the proof of Proposition 2.3, we now have a domain $\tilde{\Omega}_1$ such that weak $Z(n - 1 - q)$ is radially stable at the origin and $\omega(\tilde{\Upsilon}_1) < n - 1 - q$ (note

that $\omega(\Upsilon_1) > q$ necessarily implies $q < n - 1$, so $1 \leq n - 1 - q$). The proof proceeds as follows (the change of sign in $\text{Im}z_n$ is irrelevant, as this can be fixed by reflection in z_n), and we use Proposition 2.3 to revert to our original domain. In such cases, our domain will have a disconnected boundary.

Next, we use Υ_1 to construct Υ on $\partial\Omega$, and use our estimates to compare $\mathcal{L}_\rho(\Upsilon)$ with $\mathcal{L}_{\rho_1}^\varepsilon(\Upsilon)$. We extend Υ_1 off of $\partial\Omega_1$ by setting $\Upsilon_1|_z = \Upsilon_1|_{\pi_1(z)}$, translating each coefficient with respect to our local coordinates (we cannot use the pushforward, since this will not respect the complex structure). We can define Υ to be the orthogonal projection of Υ_1 with respect to $\partial\rho$ and $\bar{\partial}\rho$ via the formula

$$\begin{aligned} \Upsilon = \Upsilon_1 - \sum_{j,k=1}^n 2|\partial\rho|^{-2} \text{Re} \left((dz_j \wedge \bar{\partial}\rho)(\Upsilon_1) \frac{\partial\rho}{\partial z_k} \frac{\partial}{\partial z_j} \wedge \frac{\partial}{\partial \bar{z}_k} \right) \\ + \sum_{j,k=1}^n |\partial\rho|^{-4} (\partial\rho \wedge \bar{\partial}\rho)(\Upsilon_1) \frac{\partial\rho}{\partial \bar{z}_j} \frac{\partial\rho}{\partial z_k} \frac{\partial}{\partial z_j} \wedge \frac{\partial}{\partial \bar{z}_k}. \end{aligned}$$

Then $\Upsilon \in T^{1,1}(W_\eta)$ since $(\theta \wedge \bar{\partial}\rho)(\Upsilon) = 0$ for all $\theta \in \Lambda^{1,0}(W_\eta)$. Furthermore for all $\theta \in \Lambda^{1,0}(W_\eta)$

$$(i\theta \wedge \bar{\theta})(\Upsilon) = i((\theta - |\partial\rho|^{-2} \langle \theta, \partial\rho \rangle \partial\rho) \wedge (\bar{\theta} - |\partial\rho|^{-2} \langle \partial\rho, \theta \rangle \bar{\partial}\rho))(\Upsilon_1),$$

so by assumption

$$0 \leq (i\theta \wedge \bar{\theta})(\Upsilon) \leq \left| \theta - |\partial\rho|^{-2} \langle \theta, \partial\rho \rangle \partial\rho \right|^2 = |\theta|^2 - |\partial\rho|^{-2} |\langle \theta, \partial\rho \rangle|^2 \leq |\theta|^2.$$

Note that for any $\theta \in \Lambda^{1,0}(W_\eta)$ we have $\theta \wedge \bar{\partial}\rho_1(\Upsilon_1) = 0$, so by (6.5) $\theta \wedge \bar{\partial}\rho(\Upsilon_1) = \psi'_r(\rho_2 - \rho_1)\theta \wedge \bar{\partial}\rho(\Upsilon_1)$. By (6.2), we have $|\Upsilon - \Upsilon_1| \leq O(R\psi'_r(\rho_2 - \rho_1))$, so for $R > 0$ sufficiently small we have $\omega(\Upsilon) < q$ on $\partial\Omega$. In addition, adapting the proof of (6.11), we have

$$(6.12) \quad \begin{aligned} \mathcal{L}_\rho(\Upsilon) \leq (1 - \psi'_r(\rho_2 - \rho_1)) (\mathcal{L}_{\rho_1}^\varepsilon(\Upsilon_1)|_{\pi(z)}) \\ + \psi'_r(\rho_2 - \rho_1)\omega(\Upsilon) + O(R\psi'_r(\rho_2 - \rho_1)). \end{aligned}$$

Now that we have (6.11) and (6.12), we carefully choose R , r , and η (in that order) to show that radial stability of weak $Z(q)$ on $\partial\Omega_1$ implies weak $Z(q)$ on $\partial\Omega$. On W_η we have

$$\begin{aligned} \mu_1 + \dots + \mu_q - \mathcal{L}_\rho(\Upsilon) \geq (1 - \psi'_r(\rho_2 - \rho_1)) \left((\mu_1^\varepsilon + \dots + \mu_q^\varepsilon - \mathcal{L}_{\rho_1}^\varepsilon(\Upsilon_1))|_{\pi(z)} \right) \\ + \psi'_r(\rho_2 - \rho_1)(q - \omega(\Upsilon)) - O(R\psi'_r(\rho_2 - \rho_1)). \end{aligned}$$

Since $q > \omega(\Upsilon)$, we can choose $R > 0$ sufficiently small so that the error term is dominated by $\psi'_r(\rho_2 - \rho_1)(q - \omega(\Upsilon))$. Hence

$$\mu_1 + \dots + \mu_q - \mathcal{L}_\rho(\Upsilon) \geq (1 - \psi'_r(\rho_2 - \rho_1)) ((\mu_1^\varepsilon + \dots + \mu_q^\varepsilon - \mathcal{L}_{\rho_1}^\varepsilon(\Upsilon_1))|_{\pi(z)}).$$

After choosing $0 < r < R^2$, we can choose $\eta > 0$ sufficiently small to guarantee that $\varepsilon(z)$ is sufficiently small so that Definition 6.2 applies. Thus,

$$\mu_1 + \dots + \mu_q - \mathcal{L}_\rho(\Upsilon) \geq 0,$$

and hence weak $Z(q)$ holds on W_η .

By Lemma 2.6, we can patch to obtain a global Υ . □

Now, we are ready to introduce our unbounded domain.

Proof of Proposition 6.1. — For convenience, we set $z_1 = x + iy$ and define $P(z_1, z_2) = 2x|z_2|^2 - xy^4$. Let $\Omega_1 \subset \mathbb{C}^3$ be defined by $\rho_1(z) = -\text{Im}z_3 + P(z_1, z_2)$. Since $\bar{\partial}x = \frac{1}{2}d\bar{z}_1$ and $\bar{\partial}y = \frac{i}{2}d\bar{z}_1$ we compute:

$$\bar{\partial}\rho_1 = \left(|z_2|^2 - \frac{1}{2}y^4 - 2ixy^3 \right) d\bar{z}_1 + 2xz_2d\bar{z}_2 - \frac{i}{2}d\bar{z}_3$$

and

$$(6.13) \quad \bar{\partial}\bar{\partial}\rho_1 = -3xy^2dz_1 \wedge d\bar{z}_1 + z_2dz_1 \wedge d\bar{z}_2 + \bar{z}_2dz_2 \wedge d\bar{z}_1 + 2xdz_2 \wedge d\bar{z}_2.$$

We choose a basis for $T^{1,0}(\partial\Omega_1)$ by setting $L_j = \frac{\partial}{\partial z_j} + 2i\frac{\partial P}{\partial z_j}\frac{\partial}{\partial z_3}$ for $1 \leq j \leq 2$. Under this basis we can represent the Levi form by the matrix $c_{j\bar{k}} = \mathcal{L}_{\rho_1}(i\bar{L}_k \wedge L_j) = i\bar{\partial}\bar{\partial}\rho_1 \left(i\frac{\partial}{\partial \bar{z}_k} \wedge \frac{\partial}{\partial z_j} \right)$. One can easily check that the Levi-form has one positive and one negative eigenvalue when either $z_2 \neq 0$ or both $x \neq 0$ and $y \neq 0$, so $Z(2)$ is satisfied on a dense subset of the boundary.

Both 1-pseudoconvexity and the definition of weak $Z(2)$ given in [9] require orthonormal coordinates. For an arbitrary hermitian metric, let u_1 and u_2 be an orthonormal basis for $T^{1,0}(\partial\Omega_1)$. Each of these can be written in the form

$$u_j = \sum_{k=1}^2 a_j^k L_k$$

for smooth functions a_j^k . We use $c_{j\bar{k}}^u$ to denote the Levi form with respect to these new coordinates, and note that

$$(6.14) \quad c_{j\bar{k}}^u = \sum_{\ell, m=1}^2 a_j^\ell c_{\ell\bar{m}} \bar{a}_k^m.$$

The eigenvalues of c^u will be denoted $\mu_1^u \leq \mu_2^u$. Computing the trace, we have

$$\mu_1^u + \mu_2^u = \sum_{k,\ell,m=1}^2 a_k^\ell c_{\ell\bar{m}} \bar{a}_k^m = \sum_{\ell,m=1}^2 g^{\bar{m}\ell} c_{\ell\bar{m}}$$

where $g^{\bar{m}\ell}$ is a positive definite 2×2 hermitian matrix. Substituting (6.13), we have

$$(6.15) \quad \mu_1^u + \mu_2^u = -3xy^2g^{\bar{1}1} + 2\text{Re}(z_2g^{\bar{2}1}) + 2xg^{\bar{2}2}.$$

For $\partial\Omega$ to be 1-pseudoconvex, we need either $\mu_1^u + \mu_2^u \geq 0$ or $\mu_1^u + \mu_2^u - c_{\bar{1}1}^u \geq 0$. We will show that each of these leads to contradictions.

First, assume that $\mu_1^u + \mu_2^u \geq 0$. Set $y = 0$, so that by (6.15), $2\text{Re}(z_2g^{\bar{2}1}) + 2xg^{\bar{2}2} \geq 0$. Since the left hand side of this inequality equals zero when $x = z_2 = 0$, this must be a critical point and hence all first derivatives in x or z_2 will also vanish when $x = z_2 = 0$. Hence, when $x = z_2 = 0$ we have $g^{\bar{2}1} = g^{\bar{2}2} = 0$. However, this implies that $g^{\bar{m}\ell}$ has rank 1, contradicting the fact that $g^{\bar{m}\ell}$ is nondegenerate. Thus $\mu_1^u + \mu_2^u$ can not be nonnegative in a neighborhood of the origin.

Now, we assume that $\mu_1^u + \mu_2^u - c_{\bar{1}1}^u \geq 0$. Combining our assumption with (6.15) yields

$$-3xy^2(g^{\bar{1}1} - |a_1^1|^2) + 2\text{Re}(z_2(g^{\bar{2}1} - a_1^1\bar{a}_1^2)) + 2x(g^{\bar{2}2} - |a_1^2|^2) \geq 0.$$

As before, derivatives in x and z_2 must vanish when $x = z_2 = 0$, so at these points we have $g^{\bar{2}1} - a_1^1\bar{a}_1^2 = 0$ and

$$-3y^2(g^{\bar{1}1} - |a_1^1|^2) + 2(g^{\bar{2}2} - |a_1^2|^2) = 0$$

When $y \neq 0$, this means that the rank of $g^{\bar{k}j} - \bar{a}_1^k a_1^j$ is either zero or two. However,

$$g^{\bar{m}\ell} - \bar{a}_1^m a_1^\ell = \sum_{j,k=1}^2 \bar{a}_k^m (\delta_{\bar{k}j} - \delta_{j1}\delta_{k1}) a_j^\ell.$$

Hence $g^{\bar{k}j} - \bar{a}_1^k a_1^j$ must have rank one, contradicting the fact that it must have a rank of zero or two. We conclude that $\mu_1^u + \mu_2^u - c_{\bar{1}1}^u$ can not be nonnegative in a neighborhood of the origin.

Since the above construction was carried out for an arbitrary metric, we conclude that there is no metric in a neighborhood of the origin in which $\partial\Omega$ is 1-pseudoconvex.

We must now show that Definition 2.1 holds for this domain under the Euclidean metric (i.e., $\varphi = |z|^2$). If u_1 and u_2 are orthonormal vectors in

the span of L_1 and L_2 then the sum of the two smallest eigenvalues of the Levi form are

$$\mu_1 + \mu_2 = \mathcal{L}(i\bar{u}_1 \wedge u_1 + i\bar{u}_2 \wedge u_2).$$

If

$$\Upsilon_t = i\bar{u}_1 \wedge u_1 + i\bar{u}_2 \wedge u_2 - t(2i\bar{L}_1 \wedge L_1 + 3y^2 i\bar{L}_2 \wedge L_2),$$

then

$$\mu_1 + \mu_2 - \mathcal{L}(\Upsilon_t) = 0,$$

so (2) of Definition 2.1 is satisfied. Since L_1 and L_2 are also in the span of u_1 and u_2 , condition (1) will follow for $t \geq 0$ sufficiently small. Finally, $\omega(\Upsilon_t) = 2 - t(2|L_1|^2 + 3y^2|L_2|^2)$, so condition (3) holds for any $t \neq 0$. Hence Ω_1 satisfies weak $Z(2)$.

By Remark 6.5, we may now use Proposition 6.6 to turn our example into a bounded domain. □

We will also construct an example demonstrating that our condition is not invariant under changes of metric. Roughly speaking, this example contains a direction which is poorly behaved (at some points in a neighborhood of the origin this direction is an eigenvector of the Levi form with a negative eigenvalue, while at other points it can be an eigenvector corresponding to the largest positive eigenvalue). In order for weak $Z(2)$ to be satisfied, the metric must be chosen to minimize the size of this direction (and prevent it from corresponding to the largest positive eigenvalue).

PROPOSITION 6.7. — *There exists a bounded domain $\Omega \subset \mathbb{C}^4$ with smooth boundary such that*

- (1) $\partial\Omega$ does not satisfy weak $Z(2)$ under the Euclidean metric.
- (2) There exists a strictly plurisubharmonic exhaustion function φ for \mathbb{C}^4 such that $\partial\Omega$ satisfies weak $Z(2)$ with respect to the metric $\omega = i\partial\bar{\partial}\varphi$.

Proof. — As before, we can construct an unbounded domain and use Proposition 6.6 to make this bounded. To minimize the number of subscripts, we will write ρ in place of the ρ_1 used in the statement of Proposition 6.6. As above, let $z_1 = x + iy$ and $\rho(z) = -\text{Im}z_4 + P(z_1, z_2, z_3)$ where $P(z_1, z_2, z_3) = -9|z_1|^4 + 6(x^2|z_2|^2 + y^2|z_3|^2) + |z_2|^2|z_3|^2 + \frac{1}{4}(|z_2|^4 + |z_3|^4)$. We choose a basis for $T^{1,0}(\partial\Omega)$ by setting $L_j = \frac{\partial}{\partial z_j} + 2i\frac{\partial P}{\partial z_j}\frac{\partial}{\partial z_4}$ for $1 \leq j \leq 3$. Let $P_j = \frac{\partial P}{\partial z_j}$.

We compute

$$\begin{aligned} \bar{\partial}\rho &= \left(-18|z_1|^2 z_1 + 6x|z_2|^2 + 6iy|z_3|^2\right) d\bar{z}_1 \\ &\quad + \left(6x^2 z_2 + |z_3|^2 z_2 + \frac{1}{2}|z_2|^2 z_2\right) d\bar{z}_2 \\ &\quad + \left(6y^2 z_3 + |z_2|^2 z_3 + \frac{1}{2}|z_3|^2 z_3\right) d\bar{z}_3 - \frac{i}{2} d\bar{z}_4 \end{aligned}$$

and

$$\begin{aligned} i\partial\bar{\partial}\rho &= i(-36|z_1|^2 + 3|z_2|^2 + 3|z_3|^2) dz_1 \wedge d\bar{z}_1 \\ &\quad + i(6x^2 + |z_2|^2 + |z_3|^2) dz_2 \wedge d\bar{z}_2 \\ &\quad + i(6y^2 + |z_2|^2 + |z_3|^2) dz_3 \wedge d\bar{z}_3 \\ &\quad + 6ix(z_2 dz_1 \wedge d\bar{z}_2 + \bar{z}_2 dz_2 \wedge d\bar{z}_1) \\ &\quad + 6iy(-iz_3 dz_1 \wedge d\bar{z}_3 + i\bar{z}_3 dz_3 \wedge d\bar{z}_1) \\ &\quad + i(\bar{z}_2 z_3 dz_2 \wedge d\bar{z}_3 + z_2 \bar{z}_3 dz_3 \wedge d\bar{z}_2). \end{aligned}$$

If we consider the 2×2 block spanned by dz_2 and dz_3 , we can see that this is positive definite unless $z_2 = z_3 = 0$ and either $x = 0$ or $y = 0$. Hence, the Levi form has at least two positive eigenvalues and satisfies $Z(2)$ except on this set.

To define our metric, we let $\varphi_t(z) = t|z_1|^2 + \sum_{j=2}^4 |z_j|^2$ for some fixed $t \geq 1$ and use the Kähler form $\omega_t = i\partial\bar{\partial}\varphi_t$. We use $|\cdot|_t$ and $\langle \cdot, \cdot \rangle_t$ to denote norms and inner products with respect to this metric. Note that ω_1 is the Euclidean metric.

Observe that $\langle L_j, L_k \rangle_t = \delta_{jk} + 4P_j \bar{P}_k$ when $j \neq 1$ or $k \neq 1$, and $|L_1|_t^2 = t + 4|P_1|^2$. Therefore, by carrying out a Gram-Schmidt process we can construct orthonormal vectors L_2^t and L_3^t satisfying $|L_2 - L_2^t|_t \leq O(|z|^4(|z_2|^2 + |z_3|^2))$ and $|L_3 - L_3^t|_t \leq O(|z|^4(|z_2|^2 + |z_3|^2))$. We assume $3 \geq t \geq 1$, so that we have uniform bounds on constants involving t , and we can safely neglect t in our error terms (although any upper bound larger than 2 will suffice). We can complete our orthonormal basis with L_1^t satisfying $|\frac{1}{\sqrt{t}}L_1 - L_1^t|_t \leq O(|z|^4(|z_1|^2 + |z_2|^2 + |z_3|^2))$. Let $\{\theta_1^t, \theta_2^t, \theta_3^t\}$ be the orthonormal dual basis for $\{L_1^t, L_2^t, L_3^t\}$. We define the non-isotropic error form

$$\Theta = (|z_1|^2 + |z_2|^2 + |z_3|^2)i\theta_1^t \wedge \bar{\theta}_1^t + (|z_2|^2 + |z_3|^2)i(\theta_2^t \wedge \bar{\theta}_2^t + \theta_3^t \wedge \bar{\theta}_3^t),$$

and observe that off-diagonal terms can be estimated by

$$2\text{Re}((ax + by)z_j\theta_1^t \wedge \bar{\theta}_j^t) \leq \sqrt{a^2 + b^2}\Theta$$

for $j = 2$ or $j = 3$. Our Levi-form with respect to our orthonormal basis has the form

$$\begin{aligned} \mathcal{L} = & \frac{(-36|z_1|^2 + 3|z_2|^2 + 3|z_3|^2)}{t} i\theta_1^t \wedge \bar{\theta}_1^t + (6x^2 + |z_2|^2 + |z_3|^2) i\theta_2^t \wedge \bar{\theta}_2^t \\ & + (6y^2 + |z_2|^2 + |z_3|^2) i\theta_3^t \wedge \bar{\theta}_3^t + \operatorname{Re} \left(\frac{12xz_2}{\sqrt{t}} i\theta_1^t \wedge \bar{\theta}_2^t \right) \\ & - \operatorname{Re} \left(\frac{12iyz_3}{\sqrt{t}} i\theta_1^t \wedge \bar{\theta}_3^t \right) + 2\operatorname{Re}(\bar{z}_2 z_3 i\theta_2^t \wedge \bar{\theta}_3^t) + O(|z|^6 \Theta) \end{aligned}$$

We write

$$\Upsilon_t = i \sum_{j,k=1}^{n-1} b_t^{\bar{k}j} \bar{L}_k^t \wedge L_j^t.$$

When $z_2 = z_3 = 0$, the Levi-form is diagonalized with eigenvalues $\mu_1 = \frac{-36|z_1|^2}{t} + O(|z|^6|z_1|^2)$, $\mu_2 = \min\{6x^2, 6y^2\}$, and $\mu_3 = \max\{6x^2, 6y^2\}$. To check condition (2) of Definition 2.1, we compute

$$\begin{aligned} \mu_1 + \mu_2 - \mathcal{L}(\Upsilon_t) = & \min\{6x^2, 6y^2\} + \frac{-36|z_1|^2}{t} (1 - b_t^{\bar{1}1}) \\ & - 6x^2 b_t^{\bar{2}2} - 6y^2 b_t^{\bar{3}3} + O(|z|^6|z_1|^2). \end{aligned}$$

Since $b_t^{\bar{2}2} \geq 0$ and $1 \geq b_t^{\bar{1}1}$, (by condition (1) of Definition 2.1) nonnegativity when $y = 0$ requires $b_t^{\bar{2}2} \leq O(|z|^6)$ and $b_t^{\bar{1}1} \geq 1 - O(|z|^6)$. Similar computations when $x = 0$ require $b_t^{\bar{3}3} \leq O(|z|^6)$ and $b_t^{\bar{1}1} \geq 1 - O(|z|^6)$ on this set. At the origin, Υ_t is now represented by a matrix whose eigenvalues are bounded between 0 and 1 (by condition (1) of Definition 2.1 again) with diagonal entries of 0 and 1, so the off-diagonal entries must vanish. Hence, $\Upsilon_t = i\bar{L}_1^t \wedge L_1^t$ at the origin. Since nonvanishing terms of order $O(|z|)$ would cause the eigenvalues of Υ_t to grow larger than 1 or smaller than 0 in some direction, we conclude that $\Upsilon_t = i\bar{L}_1^t \wedge L_1^t + O(|z|^2)$ near the origin.

We note that $\mu_1 + \mu_2 - \mathcal{L}(\Upsilon_t) \geq 0$ (condition (2) in Definition 2.1) is equivalent to $\operatorname{Tr} \mathcal{L} - \mathcal{L}(\Upsilon_t) \geq \mu_3$, which is in turn equivalent to $(\operatorname{Tr} \mathcal{L} - \mathcal{L}(\Upsilon_t))\omega_t - \mathcal{L} \geq 0$ on $T^{1,1}(\partial\Omega)$. We compute

$$\operatorname{Tr} \mathcal{L} - \mathcal{L}(\Upsilon_t) = 6|z_1|^2 + 2(|z_2|^2 + |z_3|^2) + O(|z|^4).$$

If $(\operatorname{Tr} \mathcal{L} - \mathcal{L}(\Upsilon_t))\omega_t - \mathcal{L} \geq 0$, then every diagonal entry of this form must be nonnegative. If we test nonnegativity of $(\operatorname{Tr} \mathcal{L} - \mathcal{L}(\Upsilon_t)) - \mathcal{L}_{1\bar{1}}$, we see that this is equivalent to

$$6|z_1|^2 + 2(|z_2|^2 + |z_3|^2) - \frac{(-36|z_1|^2 + 3|z_2|^2 + 3|z_3|^2)}{t} + O(|z|^4) \geq 0.$$

Considering coefficients of $|z_2|^2 + |z_3|^2$ when $z_1 = 0$, it is necessary that $2 \geq \frac{3}{t}$, or $t \geq \frac{3}{2}$. Since the Euclidean metric corresponds to $t = 1$, weak $Z(2)$ will fail for the Euclidean metric.

For the positive result, we set $\Upsilon_t = i\bar{L}_1^t \wedge L_1^t$, so that conditions (1) and (3) are immediately satisfied. To check radial stability, we compute

$$\begin{aligned} i\partial\varphi_t \wedge \bar{\partial}\varphi_t &\equiv t|z_1|^2 i\theta_1^t \wedge \bar{\theta}_1^t + |z_2|^2 i\theta_2^t \wedge \bar{\theta}_2^t + |z_3|^2 i\theta_3^t \wedge \bar{\theta}_3^t \\ &\quad + 2\sqrt{t}\operatorname{Re}(\bar{z}_1 z_2 i\theta_1^t \wedge \bar{\theta}_2^t) + 2\sqrt{t}\operatorname{Re}(\bar{z}_1 z_3 i\theta_1^t \wedge \bar{\theta}_3^t) \\ &\quad + 2\operatorname{Re}(\bar{z}_2 z_3 i\theta_2^t \wedge \bar{\theta}_3^t) + O(|z|^3 \Theta) \pmod{\partial\rho, \bar{\partial}\rho}. \end{aligned}$$

If we set $\mathcal{L}^\varepsilon = i\partial\bar{\partial}\rho + i\varepsilon\partial\varphi_t \wedge \bar{\partial}\varphi_t$, we can show

$$\begin{aligned} \operatorname{Tr} \mathcal{L}^\varepsilon - \mathcal{L}^\varepsilon(\Upsilon_t) &= \mathcal{L}_{2\bar{2}}^\varepsilon + \mathcal{L}_{3\bar{3}}^\varepsilon \\ &= 6|z_1|^2 + (2 + \varepsilon)(|z_2|^2 + |z_3|^2) + O(|z|^3(|z_2|^2 + |z_3|^2)). \end{aligned}$$

As before, showing $\mu_1 + \mu_2 - \mathcal{L}^\varepsilon(\Upsilon_t) \geq 0$ is equivalent to showing that the form $\Phi = (\operatorname{Tr} \mathcal{L}^\varepsilon - \mathcal{L}^\varepsilon(\Upsilon_t))\omega_t - \mathcal{L}^\varepsilon$ is positive semi-definite. We compute

$$\begin{aligned} \Phi &= \frac{1}{t} \left((6t - \varepsilon t^2 + 36)|z_1|^2 + ((2 + \varepsilon)t - 3)(|z_2|^2 + |z_3|^2) \right) i\theta_1^t \wedge \bar{\theta}_1^t \\ &\quad + \left(6y^2 + |z_2|^2 + (1 + \varepsilon)|z_3|^2 \right) i\theta_2^t \wedge \bar{\theta}_2^t \\ &\quad + \left(6x^2 + (1 + \varepsilon)|z_2|^2 + |z_3|^2 \right) i\theta_3^t \wedge \bar{\theta}_3^t \\ &\quad - \frac{2}{\sqrt{t}} \operatorname{Re} \left((6x + \varepsilon t \bar{z}_1) z_2 i\theta_1^t \wedge \bar{\theta}_2^t \right) \\ &\quad - \frac{2}{\sqrt{t}} \operatorname{Re} \left((-6iy + \varepsilon t \bar{z}_1) z_3 i\theta_1^t \wedge \bar{\theta}_3^t \right) \\ &\quad - 2(1 + \varepsilon) \operatorname{Re} \left(\bar{z}_2 z_3 i\theta_2^t \wedge \bar{\theta}_3^t \right) + O(|z|^3 \Theta). \end{aligned}$$

For any $1 > \eta > 0$, we may choose $|z|$ sufficiently small so that $O(|z|^3 \Theta) \geq -\eta\Theta$. To control terms off the diagonal, we can estimate

$$\begin{aligned} 2(1 + \varepsilon) \operatorname{Re}(\bar{z}_2 z_3 i\theta_2^t \wedge \bar{\theta}_3^t) &\leq i(1 + \varepsilon)|z_2||z_3|(\theta_2^t \wedge \bar{\theta}_2^t + i\theta_3^t \wedge \bar{\theta}_3^t) \\ &\leq \left(\frac{(1 + \varepsilon)^2}{4(1 + \varepsilon - \eta)} |z_2|^2 + (1 + \varepsilon - \eta)|z_3|^2 \right) i\theta_2^t \wedge \bar{\theta}_2^t \\ &\quad + \left((1 + \varepsilon - \eta)|z_2|^2 + \frac{(1 + \varepsilon)^2}{4(1 + \varepsilon - \eta)} |z_3|^2 \right) i\theta_3^t \wedge \bar{\theta}_3^t, \end{aligned}$$

$$2\operatorname{Re} \left(\frac{6x + \varepsilon t \bar{z}_1}{\sqrt{t}} z_2 i\theta_1^t \wedge \bar{\theta}_2^t \right) \leq \frac{|6x + \varepsilon \bar{z}_1 t|^2}{tC} i\theta_1^t \wedge \bar{\theta}_1^t + C|z_2|^2 i\theta_2^t \wedge \bar{\theta}_2^t,$$

and

$$2\operatorname{Re} \left(\frac{-6iy + \varepsilon t \bar{z}_1}{\sqrt{t}} z_3 i \theta_1^t \wedge \bar{\theta}_3^t \right) \leq \frac{|-6iy + \varepsilon \bar{z}_1 t|^2}{tC} i \theta_1^t \wedge \bar{\theta}_1^t + C |z_3|^2 i \theta_3^t \wedge \bar{\theta}_3^t,$$

for some $C > 0$. Suppose that $0 < \eta < \frac{1}{2}$, so that for ε sufficiently small we know that $C(\eta, \varepsilon) = \frac{4(1+\varepsilon-\eta)(1-\eta)-(1+\varepsilon)^2}{4(1+\varepsilon-\eta)}$ is strictly positive. Combining the above estimates for this value of C , we obtain

$$\begin{aligned} \Phi \geq & \frac{1}{t} ((6t - \varepsilon t^2 + 36 - t\eta) |z_1|^2 + ((2 + \varepsilon - \eta)t - 3)(|z_2|^2 + |z_3|^2)) i \theta_1^t \wedge \bar{\theta}_1^t \\ & - \frac{2 |z_1|^2 (18 + 6\varepsilon t + \varepsilon^2 t^2)}{tC} i \theta_1^t \wedge \bar{\theta}_1^t + 6y^2 i \theta_2 \wedge \bar{\theta}_2 + 6x^2 i \theta_3^t \wedge \bar{\theta}_3^t \end{aligned}$$

If $t > 2$, then we can fix $0 < \eta < \frac{1}{4}$ sufficiently small so that $(6t + 36 - t\eta) > \frac{36}{C(\eta, 0)}$ (note that $C(0, 0) = \frac{3}{4}$). For such η , we know that for all $\varepsilon > 0$ sufficiently small we have $(6t - \varepsilon t^2 + 36 - t\eta) > \frac{2(18 + 6\varepsilon t + \varepsilon^2 t^2)}{C(\eta, \varepsilon)}$. Thus each term has a strictly positive coefficient for sufficiently small ε , so the resulting form is positive for sufficiently small ε and $|z|$. We conclude that weak $Z(2)$ is radially stable, and Proposition 6.6 can now be used to create a bounded domain. \square

BIBLIOGRAPHY

- [1] A. ANDREOTTI & H. GRAUERT, “Théorème de finitude pour la cohomologie des espaces complexes”, *Bull. Soc. Math. France* **90** (1962), p. 193-259.
- [2] A. ANDREOTTI & C. D. HILL, “E. E. Levi convexity and the Hans Lewy problem. I. Reduction to vanishing theorems”, *Ann. Scuola Norm. Sup. Pisa* (3) **26** (1972), p. 325-363.
- [3] A. ANDREOTTI & F. NORGUET, “Problème de Levi pour les classes de cohomologie”, *C. R. Acad. Sci. Paris* **258** (1964), p. 778-781.
- [4] J. BRINKSCHULTE, “Local solvability of the $\bar{\partial}$ -equation with boundary regularity on weakly q -convex domains”, *Math. Ann.* **334** (2006), no. 1, p. 143-152.
- [5] S.-C. CHEN & M.-C. SHAW, *Partial differential equations in several complex variables*, AMS/IP Studies in Advanced Mathematics, vol. 19, American Mathematical Society, Providence, RI; International Press, Boston, MA, 2001, xii+380 pages.
- [6] M. G. EASTWOOD & G. V. SURIA, “Cohomologically complete and pseudoconvex domains”, *Comment. Math. Helv.* **55** (1980), no. 3, p. 413-426.
- [7] G. B. FOLLAND & J. J. KOHN, *The Neumann problem for the Cauchy-Riemann complex*, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1972, Annals of Mathematics Studies, No. 75, viii+146 pages.
- [8] P. S. HARRINGTON, “Sobolev estimates for the Cauchy-Riemann complex on C^1 pseudoconvex domains”, *Math. Z.* **262** (2009), no. 1, p. 199-217.
- [9] P. S. HARRINGTON & A. RAICH, “Regularity results for $\bar{\partial}_b$ on CR-manifolds of hypersurface type”, *Comm. Partial Differential Equations* **36** (2011), no. 1, p. 134-161.

- [10] F. R. HARVEY & H. B. LAWSON, JR., “On boundaries of complex analytic varieties. I”, *Ann. of Math. (2)* **102** (1975), no. 2, p. 223-290.
- [11] G. M. HENKIN & J. LEITERER, “Global integral formulas for solving the $\bar{\partial}$ -equation on Stein manifolds”, *Ann. Polon. Math.* **39** (1981), p. 93-116.
- [12] L.-H. HO, “ $\bar{\partial}$ -problem on weakly q -convex domains”, *Math. Ann.* **290** (1991), no. 1, p. 3-18.
- [13] L. HÖRMANDER, “ L^2 estimates and existence theorems for the $\bar{\partial}$ operator”, *Acta Math.* **113** (1965), p. 89-152.
- [14] ———, *An introduction to complex analysis in several variables*, third ed., North-Holland Mathematical Library, vol. 7, North-Holland Publishing Co., Amsterdam, 1990, xii+254 pages.
- [15] ———, “The null space of the $\bar{\partial}$ -Neumann operator”, *Ann. Inst. Fourier (Grenoble)* **54** (2004), no. 5, p. 1305-1369, xiv, xx.
- [16] S. G. KRANTZ & H. R. PARKS, “Distance to C^k hypersurfaces”, *J. Differential Equations* **40** (1981), no. 1, p. 116-120.
- [17] C. LAURENT-THIÉBAUT, “Transformation de Bochner-Martinelli dans une variété de Stein”, in *Séminaire d'Analyse P. Lelong–P. Dolbeault–H. Skoda, Années 1985/1986*, Lecture Notes in Math., vol. 1295, Springer, Berlin, 1987, p. 96-131.
- [18] A. C. NICOARA, “Global regularity for $\bar{\partial}_b$ on weakly pseudoconvex CR manifolds”, *Adv. Math.* **199** (2006), no. 2, p. 356-447.
- [19] A. RAICH, “Compactness of the complex Green operator on CR-manifolds of hypersurface type”, *Math. Ann.* **348** (2010), no. 1, p. 81-117.
- [20] M.-C. SHAW, “Global solvability and regularity for $\bar{\partial}$ on an annulus between two weakly pseudoconvex domains”, *Trans. Amer. Math. Soc.* **291** (1985), no. 1, p. 255-267.
- [21] ———, “ L^2 -estimates and existence theorems for the tangential Cauchy-Riemann complex”, *Invent. Math.* **82** (1985), no. 1, p. 133-150.
- [22] ———, “ L^2 estimates and existence theorems for $\bar{\partial}_b$ on Lipschitz boundaries”, *Math. Z.* **244** (2003), no. 1, p. 91-123.
- [23] ———, “The closed range property for $\bar{\partial}$ on domains with pseudoconcave boundary”, in *Complex analysis*, Trends Math., Birkhäuser/Springer Basel AG, Basel, 2010, p. 307-320.
- [24] Y. T. SIU, “Complex-analyticity of harmonic maps, vanishing and Lefschetz theorems”, *J. Differential Geom.* **17** (1982), no. 1, p. 55-138.
- [25] E. J. STRAUBE, *Lectures on the L^2 -Sobolev theory of the $\bar{\partial}$ -Neumann problem*, ESI Lectures in Mathematics and Physics, European Mathematical Society (EMS), Zürich, 2010, viii+206 pages.
- [26] ———, “The complex Green operator on CR-submanifolds of \mathbb{C}^n of hypersurface type: compactness”, *Trans. Amer. Math. Soc.* **364** (2012), no. 8, p. 4107-4125.
- [27] G. ZAMPIERI, *Complex analysis and CR geometry*, University Lecture Series, vol. 43, American Mathematical Society, Providence, RI, 2008, viii+200 pages.

Manuscrit reçu le 10 janvier 2012,
 accepté le 13 janvier 2015.

Phillip S. HARRINGTON
 Andrew S. RAICH
 Department of Mathematical Sciences
 1 University of Arkansas
 SCEN 309

Fayetteville, AR 7201 (USA)
psharrin@uark.edu
araich@urak.edu