



ANNALES DE L'INSTITUT FOURIER


Sebastian HERR & Shinya KINOSHITA

**Subcritical well-posedness results for the Zakharov–Kuznetsov
equation in dimension three and higher**

Tome 73, n° 3 (2023), p. 1203–1267.

<https://doi.org/10.5802/aif.3547>

Article mis à disposition par ses auteurs selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE



<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les *Annales de l'Institut Fourier* sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org e-ISSN : 1777-5310

SUBCRITICAL WELL-POSEDNESS RESULTS FOR THE ZAKHAROV–KUZNETSOV EQUATION IN DIMENSION THREE AND HIGHER

by Sebastian HERR & Shinya KINOSHITA (*)

ABSTRACT. — The Zakharov–Kuznetsov equation in space dimension $d \geq 3$ is considered. It is proved that the Cauchy problem is locally well-posed in $H^s(\mathbb{R}^d)$ in the full subcritical range $s > (d-4)/2$, which is optimal up to the endpoint. As a corollary, global well-posedness in $L^2(\mathbb{R}^3)$ and, under a smallness condition, in $H^1(\mathbb{R}^4)$, follow.

RÉSUMÉ. — On considère l'équation de Zakharov–Kuznetsov en dimension $d \geq 3$. On établit que le problème de Cauchy est localement bien posé dans H^s pour tout exposant sous-critique $s > (d-4)/2$. Ceci est optimal jusqu'au cas limite. Comme corollaire, il s'ensuit que l'équation est globalement bien posée dans $L^2(\mathbb{R}^3)$ et, sous une hypothèse de petitesse, dans $H^1(\mathbb{R}^4)$.

1. Introduction

We consider the Zakharov–Kuznetsov equation with the quadratic nonlinearity

$$(1.1) \quad \begin{aligned} \partial_t u + \partial_x \Delta u &= \partial_x u^2 \text{ in } (-T, T) \times \mathbb{R}^d \\ u(0, \cdot) &= u_0 \in H^s(\mathbb{R}^d) \end{aligned}$$

where $u = u(t, x, \mathbf{y})$ is real-valued and Δ denotes the Laplacian with respect to $(x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{d-1}$. The equation (1.1) arises as an asymptotic model wave propagation in a magnetized plasma [4, 32]. It was introduced in [33] in $d = 2, 3$, see also [23] for a formal derivation. More recently, it

Keywords: Well-posedness, Cauchy problem, Zakharov–Kuznetsov equation, bilinear estimate, nonlinear Loomis–Whitney inequality.

2010 Mathematics Subject Classification: 35Q55.

(*) Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – SFB 1283/2 2021 – 317210226.

was rigorously derived from the Euler–Poisson system as a long-wave and small-amplitude limit, see [24, Section 10.3.2.6]. The Zakharov–Kuznetsov equation (1.1) generalizes the Korteweg-de Vries equation (which is the case $d = 1$). In particular, it has solitary wave solutions. Recently, their asymptotic stability has been proven in [10].

Real-valued solutions of (1.1) conserve the L^2 -norm and the energy

$$\frac{1}{2} \int_{\mathbb{R}^d} |\nabla_{x,y} u(t, x, y)|^2 dx dy + \frac{1}{3} \int_{\mathbb{R}^d} u(t, x, y)^3 dx dy.$$

If u is a solution, then for any $\lambda > 0$ the function

$$u_\lambda(t, x, y) = \lambda^2 u(\lambda^3 t, \lambda x, \lambda y)$$

also solves (1.1). This implies that $s_c := (d-4)/2$ is the critical Sobolev regularity for (1.1) in the sense that the corresponding (homogeneous) Sobolev norm is invariant under the rescaling described above.

In this paper, we will focus on the case of spatial dimensions $d \geq 3$ and prove local well-posedness of the Cauchy problem associated with (1.1) in the full sub-critical range. Let $H^s(\mathbb{R}^d)$ denote the Sobolev space of tempered distributions on \mathbb{R}^d all derivatives up to order s in $L^2(\mathbb{R}^d)$, see Subsection 2.1 for a precise definition.

THEOREM 1.1. — *Let $d \geq 3$. For any $s > (d-4)/2$, the Cauchy problem for (1.1) is locally well-posed in $H^s(\mathbb{R}^d)$.*

Note that the energy-subcritical dimensions are $d \leq 5$, and the L^2 -subcritical dimensions are $d \leq 3$. From the conservation laws mentioned above and the Gagliardo–Nirenberg inequality we deduce

COROLLARY 1.2. — *If $d = 3$, the Cauchy problem for (1.1) is globally well-posed for real-valued initial data in $L^2(\mathbb{R}^3)$. If $d = 4$, the Cauchy problem for (1.1) is globally well-posed for real-valued initial data in $H^1(\mathbb{R}^4)$ with sufficiently small $L^2(\mathbb{R}^4)$ -norm.*

Previous results

The Cauchy problem for the Zakharov–Kuznetsov equation and the so-called generalized Zakharov–Kuznetsov equation

$$\partial_t u + \partial_x \Delta u = \partial_x u^{k+1}, \quad (k \in \mathbb{N}),$$

have been studied extensively. In dimension $d = 2$ and for the quadratic nonlinearity ($k = 1$), Faminskii [11] proved global well-posedness in $H^1(\mathbb{R}^2)$,

Linares and Pastor [25] proved local well-posedness for $s > 3/4$, Grünrock and Herr [16] and Molinet and Pilod [29], proved local well-posedness for $s > 1/2$, and recently the second named author [21] proved local well-posedness for $s > -1/4$, which is optimal up to the end-point. In dimension $d = 2$ and for the cubic nonlinearity ($k = 2$) Biagioni and Linares [7] proved global well-posedness in $H^1(\mathbb{R}^2)$, Linares and Pastor [25] proved local well-posedness for $s > 3/4$ and [26] global well-posedness for $s > 53/63$, Ribaud and Vento [30] proved local well-posedness for $s > 1/4$, and recently Bhattacharya, Farah and Roudenko [6] proved global well-posedness for $s > 3/4$, and the second named author [20] proved local well-posedness for $s = 1/4$. In dimension $d = 3$ and for the quadratic nonlinearity ($k = 1$) Linares and Saut [27] proved local well-posedness for $s > 9/8$, Ribaud and Vento [31] proved local well-posedness for $s > 1$ and in $B_2^{1,1}(\mathbb{R}^3)$, and Molinet and Pilod [29] proved global well-posedness for $s > 1$. In dimension $d \geq 3$ and for the cubic nonlinearity ($k = 2$) Grünrock [15] proved local well-posedness in the full sub-critical range $s > d/2 - 1$ and the second named author [20] proved small data global well-posedness at the scaling critical regularity $s = d/2 - 1$. For more results in the case $k \geq 3$, we refer to the papers [12, 14, 26, 30].

Strategy of proof

Theorem 1.1 will be proved by a contraction argument in Fourier restriction spaces $X^{s,b}$, which is based on the bilinear estimate provided in Theorem 3.1 below. Since the general strategy of proof is standard, we refer to [13, Section 2] for details and precise formulations and focus on the key bilinear estimate.

In the low-regularity analysis of the Korteweg-de Vries equation, the set of time-resonances plays a crucial role, see [8, 19]. It is the set of spatial frequencies allowing the product of two solutions to the homogeneous equations to form another solution to the homogeneous equation. In the case of the KdV equation, it is the set

$$\{(\xi, \xi_1) \in \mathbb{R}^2 : 3\xi(\xi - \xi_1)\xi_1 = 0\},$$

while in the case of the ZK equation it is significantly more complex. More precisely, it is the set

$$\{(\xi, \eta, \xi_1, \eta_1) \in \mathbb{R}^{2d} : 3\xi(\xi - \xi_1)\xi_1 + \xi|\eta|^2 - \xi_1|\eta_1|^2 - (\xi - \xi_1)|\eta - \eta_1|^2 = 0\}.$$

First well-posedness results [16, 29] did not rely on the structure of this set. Recently, the second author [21] established well-posedness of the Zakharov–Kuznetsov equation in $d = 2$ for $s > -1/4$, which turned out to be optimal within the purely perturbative regime (up to the endpoint). The key observation is that the bulk of frequencies close to the resonant set are transversal. This allows to invoke the Loomis–Whitney inequality [28], more precisely a nonlinear generalization thereof [3, 5, 22]. This strategy has been used previously in the case of the low-dimensional Zakharov system, see [1, 2], in which case the resonant set is easier to understand.

Outline of the paper

In Section 2 we will discuss preliminaries concerning notation, Strichartz and transversal estimates. The key bilinear estimate is stated as Theorem 3.1 in Section 3 and first reductions are performed. The proof of Theorem 3.1 is then split into cases, which are treated in the following Sections 4–6. In an appendix we include a proof of the well-known transversal L^2 estimate from Section 2.

2. Preliminaries

2.1. Notation

We write $A \lesssim B$ if there exists $C > 0$ such that $A \leq CB$ and $A \ll B$ if $A \leq cB$ for some small enough $c < 1$. Also, $A \sim B$ means $A \lesssim B$ and $B \lesssim A$.

Let $N, L \geq 1$ be dyadic numbers, i.e. there exist $n_1, n_2 \in \mathbb{N}_0$ such that $N = 2^{n_1}$ and $L = 2^{n_2}$, and $\psi \in C_0^\infty((-2, 2))$ be an even, non-negative function which satisfies $\psi(t) = 1$ for $|t| \leq 1$ and letting $\psi_N(t) := \psi(tN^{-1}) - \psi(2tN^{-1})$, $\psi_1(t) := \psi(t)$, the equality $\sum_N \psi_N(t) = 1$ holds. Here and in the sequel we use the convention that capitalized summation indices run over $2^{\mathbb{N}_0}$. Let $d \geq 3$. We distinguish the first and the other spatial variables. To be precise, $(x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{d-1}$ denotes the space variables. Similarly, $(\xi, \boldsymbol{\eta}) \in \mathbb{R} \times \mathbb{R}^{d-1}$ denotes the frequency variables of (x, \mathbf{y}) .

Let $u = u(t, x, \mathbf{y})$. $\mathcal{F}_t u$, $\mathcal{F}_{x, \mathbf{y}} u$ denote the Fourier transform of u in time, space, respectively. $\mathcal{F}_{t, x, \mathbf{y}} u = \hat{u}$ denotes the Fourier transform of u in

space and time. For $s \in \mathbb{R}$ define $H^s(\mathbb{R}^d)$ to be the space of all tempered distributions $f = f(x, \mathbf{y})$ on \mathbb{R}^d satisfying

$$\|f\|_{H^s} := \left(\int_{\mathbb{R}^d} (1 + |\xi|^2 + |\boldsymbol{\eta}|^2)^s |\mathcal{F}_{x,\mathbf{y}} f(\xi, \boldsymbol{\eta})|^2 d\xi d\boldsymbol{\eta} \right)^{\frac{1}{2}} < +\infty.$$

We define frequency and modulation projections P_N, Q_L as

$$\begin{aligned} (\mathcal{F}_{x,\mathbf{y}} P_N u) &:= \psi_N(|\cdot|)(\mathcal{F}_{x,\mathbf{y}} u), \\ \widehat{Q_L u}(\tau, \xi, \boldsymbol{\eta}) &:= \psi_L(\tau - \xi(\xi^2 + |\boldsymbol{\eta}|^2)) \widehat{u}(\tau, \xi, \boldsymbol{\eta}). \end{aligned}$$

Let $B_r(p) \subset \mathbb{R}^d$ denote the open ball with radius $r > 0$ and center $p \in \mathbb{R}^d$, and define the spatial Fourier multiplier $P_{B_r(p)} f = \mathcal{F}_{x,\mathbf{y}}^{-1} \mathbf{1}_{B_r(p)} \mathcal{F}_{x,\mathbf{y}} f$, where $\mathbf{1}_{B_r(p)}$ denotes the characteristic function of $B_r(p)$.

We now define $X^{s,b}(\mathbb{R}^{d+1})$ spaces. Let $s, b \in \mathbb{R}$. Define $X^{s,b}(\mathbb{R}^{d+1})$ as the space of all $f \in \mathcal{S}'(\mathbb{R}^{d+1})$ such that

$$\|f\|_{X^{s,b}} := \left(\sum_{N,L} N^{2s} L^{2b} \|P_N Q_L f\|_{L^2_{t,x,\mathbf{y}}(\mathbb{R}^{d+1})}^2 \right)^{1/2} < +\infty.$$

For convenience, we define the set in frequency as

$$G_{N,L} := \{(\tau, \xi, \boldsymbol{\eta}) \in \mathbb{R}^{d+1} \mid \psi_L(\tau - \xi(\xi^2 + |\boldsymbol{\eta}|^2)) \psi_N(|(\xi, \boldsymbol{\eta})|) \neq 0\}.$$

A simple calculation shows $\overline{X^{s,b}} = X^{s,b}$ and $(X^{s,b})^* = X^{-s,-b}$, for $s, b \in \mathbb{R}$.

Define the propagator for the linear Zakharov–Kuznetsov equation

$$U(t) = e^{-t\partial_x \Delta}$$

and the one-dimensional Fourier multiplier

$$|\partial_x|^s = \mathcal{F}_x^{-1} |\xi|^s \mathcal{F}_x.$$

2.2. Strichartz type estimates and transversal estimates

We start with a Strichartz or (dual) restriction type estimate, where curvature properties of the characteristic set of the differential operator are used.

PROPOSITION 2.1. — *Let $d \geq 3$ and $0 \leq s \leq 1/4$. Then, for all $p \in \mathbb{R}^d$ and $r > 0$ we have*

$$\| |\partial_x|^s U(t) P_{B_r(p)} f \|_{L^4_{t,x,\mathbf{y}}} \lesssim r^{\frac{d-3}{4} + s} \| P_{B_r(p)} f \|_{L^2_{x,\mathbf{y}}}.$$

Proof. — It suffices to show the endpoint cases $s = 0$ and $s = 1/4$. We follow the proof of the Strichartz estimates of the KP-II type equations on cylinders, see Theorem 2 in [17] and [15]. The Littlewood–Paley theorem implies that it suffices to show the claim under the condition

$$(2.1) \quad \text{supp } \mathcal{F}_{x,y} f \subset \{(\xi, \eta) \mid 2^k \leq |\xi| \leq 2^{k+1}\} \cap B_r(p),$$

where k is an arbitrary integer. Let $\psi : \mathbb{R}^{d-1} \rightarrow \mathbb{C}$. We recall the classical (non-endpoint) Strichartz estimate of the Schrödinger equations on \mathbb{R}^{d-1} , i.e.

$$\|\mathrm{e}^{\mathrm{i}t\Delta_y} \psi\|_{L_t^p L_y^q} \lesssim \|\psi\|_{L_y^2},$$

where (p, q) is an admissible pair satisfying $2 < p \leq \infty$ and $2/p = (d-1)(1/2 - 1/q)$. Let $(4, q_0)$ be admissible. Since f satisfies (2.1), if we fix $\xi \in \mathbb{R}$, by using the Sobolev inequality and the above Strichartz estimate, we easily get

$$(2.2) \quad \begin{aligned} \|\mathrm{e}^{\mathrm{i}t\xi\Delta_y} \mathcal{F}_x f(\xi, \cdot)\|_{L_t^4 L_y^4} &\leq r^{\frac{d-3}{4}} |\xi|^{-\frac{1}{4}} \|\mathrm{e}^{\mathrm{i}t\Delta_y} \mathcal{F}_x f(\xi, \cdot)\|_{L_t^4 L_y^{q_0}} \\ &\leq r^{\frac{d-3}{4}} |\xi|^{-\frac{1}{4}} \|\mathcal{F}_x f(\xi, \cdot)\|_{L_y^2}. \end{aligned}$$

Therefore, it follows from Plancherel's theorem in x , Hölder's inequality, we get

$$\begin{aligned} &\|U(t)P_{B_r(p)}f\|_{L_{t,x,y}^4}^2 \\ &= \|(U(t)P_{B_r(p)}f)(U(t)P_{B_r(p)}f)\|_{L_{t,x,y}^2} \\ &\leq \left\| \int_{\mathbb{R}} \left\| (\mathrm{e}^{\mathrm{i}t(\xi-\xi')\Delta_y} \mathcal{F}_x f(\xi - \xi', \cdot)) (\mathrm{e}^{\mathrm{i}t\xi'\Delta_y} \mathcal{F}_x f(\xi', \cdot)) \right\|_{L_{ty}^2} \mathrm{d}\xi' \right\|_{L_{\xi}^2} \\ &\leq \left\| \int_{\mathbb{R}} \left\| \mathrm{e}^{\mathrm{i}t(\xi-\xi')\Delta_y} \mathcal{F}_x f(\xi - \xi', \cdot) \|_{L_{ty}^4} \|\mathrm{e}^{\mathrm{i}t\xi'\Delta_y} \mathcal{F}_x f(\xi', \cdot) \|_{L_{ty}^4} \right\|_{L_{\xi}^2} \mathrm{d}\xi' \right\|_{L_{\xi}^2} \end{aligned}$$

Now, we use (2.2) and continue with

$$\begin{aligned} &\lesssim r^{\frac{d-3}{2}} \left\| \int_{\mathbb{R}} (|\xi - \xi'| |\xi'|)^{-\frac{1}{4}} \|\mathcal{F}_x f(\xi - \xi', \cdot)\|_{L_y^2} \|\mathcal{F}_x f(\xi', \cdot)\|_{L_y^2} \mathrm{d}\xi' \right\|_{L_{\xi}^2} \\ &\lesssim r^{\frac{d-3}{2}} \left\| \int_{\mathbb{R}} \|\mathcal{F}_x f(\xi - \xi', \cdot)\|_{L_y^2} \|\mathcal{F}_x f(\xi', \cdot)\|_{L_y^2} \mathrm{d}\xi' \right\|_{L_{\xi}^{\infty}} \\ &\leq r^{\frac{d-3}{2}} \|f\|_{L_x^2}^2. \end{aligned}$$

This completes the proof for $s = 0$. Here we used $2^k \leq |\xi'| \leq 2^{k+1}$ and $2^k \leq |\xi - \xi'| \leq 2^{k+1}$. Similarly, we have

$$\begin{aligned} & \| |\partial_x|^{1/4} U(t) P_{B_r(p)} f \|_{L_{t,x,y}^4}^2 \\ &= \| (|\partial_x|^{1/4} U(t) P_{B_r(p)} f) (|\partial_x|^{1/4} U(t) P_{B_r(p)} f) \|_{L_{t,x,y}^2} \\ &\lesssim r^{\frac{d-3}{2}} \left\| \int_{\mathbb{R}} \|\mathcal{F}_x f(\xi - \xi', \cdot)\|_{L_y^2} \|\mathcal{F}_x f(\xi', \cdot)\|_{L_y^2} d\xi' \right\|_{L_\xi^2} \\ &\lesssim r^{\frac{d-2}{2}} \left\| \int_{\mathbb{R}} \|\mathcal{F}_x f(\xi - \xi', \cdot)\|_{L_y^2} \|\mathcal{F}_x f(\xi', \cdot)\|_{L_y^2} d\xi' \right\|_{L_\xi^\infty} \\ &\lesssim r^{\frac{d-2}{2}} \|f\|_{L_{x,y}^2}^2, \end{aligned}$$

where we used (2.1) again. □

Next, we recall the standard bilinear estimate which exploits transversality, see e.g. [9, Lemma 2.6] for a proof with a general phase function and for references. We also provide a proof in the appendix.

PROPOSITION 2.2. — *Let $d \geq 2$, $N_2 \leq N_1$, $\varphi(\xi, \eta) = \xi(|\xi|^2 + |\eta|^2)$. Suppose that*

$$\text{supp } \widehat{u}_{N_1, L_1} \subset G_{N_1, L_1} \cap (\mathbb{R} \times B_r(p)), \quad \text{supp } \widehat{v}_{N_2, L_2} \subset G_{N_2, L_2},$$

and there exists K which satisfies $K \gtrsim rN_1$ and

$$|\nabla \varphi(\xi_1, \eta_1) - \nabla \varphi(\xi_2, \eta_2)| \gtrsim K,$$

for all $(\xi_1, \eta_1), (\xi_2, \eta_2)$ in the spatial Fourier support of u_{N_1, L_1} and v_{N_2, L_2} , respectively. Then, we have

$$\begin{aligned} (2.3) \quad & \|u_{N_1, L_1} v_{N_2, L_2}\|_{L_{t,x,y}^2} \\ &\lesssim r^{\frac{d-1}{2}} K^{-\frac{1}{2}} (L_1 L_2)^{\frac{1}{2}} \|u_{N_1, L_1}\|_{L_{t,x,y}^2} \|v_{N_2, L_2}\|_{L_{t,x,y}^2}. \end{aligned}$$

In particular, if $N_2 \leq 2^{-3} N_1$ and

$$\text{supp } \widehat{u}_{N_1, L_1} \subset G_{N_1, L_1}, \quad \text{supp } \widehat{v}_{N_2, L_2} \subset G_{N_2, L_2},$$

we have

$$\begin{aligned} (2.4) \quad & \|u_{N_1, L_1} v_{N_2, L_2}\|_{L_{t,x,y}^2} \\ &\lesssim N_1^{-1} N_2^{\frac{d-1}{2}} (L_1 L_2)^{\frac{1}{2}} \|u_{N_1, L_1}\|_{L_{t,x,y}^2} \|v_{N_2, L_2}\|_{L_{t,x,y}^2}. \end{aligned}$$

A trilinear estimate based on transversality is the following generalization of the classical Loomis–Whitney inequality, which is Corollary 1.5 in [3], see also [5, 22].

PROPOSITION 2.3. — Assume that for $i \in \{1, 2, 3\}$ the surface $S_i \subset \mathbb{R}^3$ is an open and bounded subset of S_i^* which satisfies the following three conditions:

(1) For a convex $U_i \subset \mathbb{R}^3$ such that $\text{dist}(S_i, U_i^c) \geq \text{diam}(S_i)$ we have

$$S_i^* = \{\lambda_i \in U_i \mid \Phi_i(\lambda_i) = 0, \nabla \Phi_i \neq 0, \Phi_i \in C^{1,1}(U_i)\}.$$

(2) The unit normal vector field \mathbf{n}_i on S_i^* satisfies the Hölder condition

$$\sup_{\lambda, \lambda' \in S_i^*} \frac{|\mathbf{n}_i(\lambda) - \mathbf{n}_i(\lambda')|}{|\lambda - \lambda'|} + \frac{|\mathbf{n}_i(\lambda)(\lambda - \lambda')|}{|\lambda - \lambda'|^2} \lesssim 1.$$

(3) There exists $\delta > 0$ such that $\text{diam}(S_i) \lesssim \delta$ and the matrix $N(\lambda_1, \lambda_2, \lambda_3) = (\mathbf{n}_1(\lambda_1), \mathbf{n}_2(\lambda_2), \mathbf{n}_3(\lambda_3))$ satisfies the transversality condition

$$\delta \leq |\det N(\lambda_1, \lambda_2, \lambda_3)| \leq 1, \quad \text{for all } (\lambda_1, \lambda_2, \lambda_3) \in S_1^* \times S_2^* \times S_3^*.$$

Then, for functions $f \in L^2(S_1)$ and $g \in L^2(S_2)$, the restriction of the convolution $f * g$ to S_3 is a well-defined $L^2(S_3)$ -function which satisfies

$$\|f * g\|_{L^2(S_3)} \lesssim \delta^{-\frac{1}{2}} \|f\|_{L^2(S_1)} \|g\|_{L^2(S_2)}.$$

3. The key bilinear estimate

The main contribution of this paper is the following:

THEOREM 3.1. — For any $s > (d - 4)/2$, there exist $b \in (\frac{1}{2}, 1)$ and $b' \in (b - 1, 0)$, such that

$$(3.1) \quad \|\partial_x(uv)\|_{X^{s, b'}} \lesssim \|u\|_{X^{s, b}} \|v\|_{X^{s, b}}.$$

The remainder of the paper will be devoted to its proof. By a duality argument and dyadic decompositions, we observe that

$$(3.2) \quad (3.1) \iff \left| \int w \partial_x(uv) dt dx dy \right| \lesssim \|u\|_{X^{s, b}} \|v\|_{X^{s, b}} \|w\|_{X^{-s, -b'}}.$$

We will use the shorthand notations

$$w_{N_0, L_0} := Q_{L_0} P_{N_0} w, \quad u_{N_1, L_1} := Q_{L_1} P_{N_1} u, \quad v_{N_2, L_2} := Q_{L_2} P_{N_2} v.$$

Obviously, (3.2) follows from

$$(3.3) \quad \sum_{\substack{N_j, L_j \\ (j=0,1,2)}} \left| \int (\partial_x w_{N_0, L_0}) u_{N_1, L_1} v_{N_2, L_2} dt dx dy \right| \lesssim \|u\|_{X^{s, b}} \|v\|_{X^{s, b}} \|w\|_{X^{-s, -b'}}.$$

For brevity, we write

$$\begin{aligned} L_{012}^{\max} &:= \max(L_0, L_1, L_2), \\ N_{012}^{\max} &:= \max(N_0, N_1, N_2), \\ N_{012}^{\min} &:= \min(N_0, N_1, N_2). \end{aligned}$$

3.1. Reductions

Here, we prove (3.3) in the following relatively simple cases:

- (1) $L_{012}^{\max} \gtrsim (N_{012}^{\max})^3$,
- (2) $N_{012}^{\min} \sim 1$ and $L_{012}^{\max} \ll (N_{012}^{\max})^3$

We will use Propositions 2.1 and 2.2, respectively.

We first assume $L_{012}^{\max} \gtrsim (N_{012}^{\max})^3$ and show

$$(3.4) \quad \left| \int w_{N_0, L_0} u_{N_1, L_1} v_{N_2, L_2} dt dx dy \right| \lesssim (N_{012}^{\min})^{\frac{d-3}{2}} (N_{012}^{\max})^{-\frac{3}{2}+3\varepsilon} L_0^{\frac{1}{2}-\varepsilon} (L_1 L_2)^{\frac{1}{2}} \|u_{N_1, L_1}\|_{L^2} \|v_{N_2, L_2}\|_{L^2} \|w_{N_0, L_0}\|_{L^2}$$

for some small $\varepsilon > 0$. Clearly, this inequality gives (3.3). Here we only consider the case $L_0 \gtrsim (N_{012}^{\max})^3$. The other two cases $L_1 \gtrsim (N_{012}^{\max})^3$ and $L_2 \gtrsim (N_{012}^{\max})^3$ can be treated similarly. By the almost orthogonality, we may replace u_{N_1, L_1} and v_{N_2, L_2} by $P_B u_{N_1, L_1}$ and $P_{B'} v_{N_2, L_2}$ where P_B and $P_{B'}$ denote the spatial frequency localization operators for some fixed balls B and B' with radius N_{012}^{\min} , respectively. It follows from the Hölder's inequality and Proposition 2.1 that

$$\begin{aligned} &\left| \int w_{N_0, L_0} (P_B u_{N_1, L_1}) (P_{B'} v_{N_2, L_2}) dt dx dy \right| \\ &\lesssim \|w_{N_0, L_0}\|_{L^2} \|P_B u_{N_1, L_1}\|_{L^4} \|P_{B'} v_{N_2, L_2}\|_{L^4} \\ &\lesssim (N_{012}^{\min})^{\frac{d-3}{2}} (N_{012}^{\max})^{-\frac{3}{2}+3\varepsilon} L_0^{\frac{1}{2}-\varepsilon} (L_1 L_2)^{\frac{1}{2}} \|u_{N_1, L_1}\|_{L^2} \|v_{N_2, L_2}\|_{L^2} \|w_{N_0, L_0}\|_{L^2}, \end{aligned}$$

which completes the proof of (3.4).

Next we deal with the case $N_{012}^{\min} \sim 1$ and $L_{012}^{\max} \ll (N_{012}^{\max})^3$. If $1 \sim N_0 \sim N_1 \sim N_2$, by using the L^4 Strichartz estimate, we get

$$\begin{aligned} &\left| \int w_{N_0, L_0} u_{N_1, L_1} v_{N_2, L_2} dt dx dy \right| \\ &\lesssim \|w_{N_0, L_0}\|_{L^2} \|u_{N_1, L_1}\|_{L^4} \|v_{N_2, L_2}\|_{L^4} \\ &\lesssim (L_1 L_2)^{\frac{1}{2}} \|u_{N_1, L_1}\|_{L^2} \|v_{N_2, L_2}\|_{L^2} \|w_{N_0, L_0}\|_{L^2}, \end{aligned}$$

which implies (3.3). Thus, by symmetry, we only need to consider $1 \sim N_0 \ll N_1 \sim N_2$ and $1 \sim N_2 \ll N_0 \sim N_1$. The both cases are treated by Proposition 2.2. First we assume $1 \sim N_0 \ll N_1 \sim N_2$ and show the following.

$$(3.5) \quad \left| \int w_{N_0, L_0} u_{N_1, L_1} v_{N_2, L_2} dt dx dy \right| \\ \lesssim N_1^{-1} (L_0 L_1)^{\frac{1}{2}} \|u_{N_1, L_1}\|_{L^2} \|v_{N_2, L_2}\|_{L^2} \|w_{N_0, L_0}\|_{L^2},$$

which immediately yields (3.3) since $-2s < 4 - d \leq 1$ and $L_0 \ll N_1^3$. We deduce from $N_0 \sim 1$ and the almost orthogonality, we can replace u_{N_1, L_1} by $P_B u_{N_1, L_1}$ with a fixed ball of spatial frequency B whose radius is 1. Thus, by the Hölder's inequality and Proposition 2.2, we observe

$$\left| \int w_{N_0, L_0} (P_B u_{N_1, L_1}) v_{N_2, L_2} dt dx dy \right| \\ \leq \|w_{N_0, L_0} (P_B u_{N_1, L_1})\|_{L^2} \|v_{N_2, L_2}\|_{L^2} \\ \lesssim N_1^{-1} (L_0 L_1)^{\frac{1}{2}} \|u_{N_1, L_1}\|_{L^2} \|v_{N_2, L_2}\|_{L^2} \|w_{N_0, L_0}\|_{L^2},$$

which completes the proof of (3.5). Similarly, if $1 \sim N_2 \ll N_0 \sim N_1$, by replacing u_{N_1, L_1} with $P_B u_{N_1, L_1}$, it follows from the Hölder's inequality and Proposition 2.2 that

$$\left| \int w_{N_0, L_0} (P_B u_{N_1, L_1}) v_{N_2, L_2} dt dx dy \right| \\ \leq \|(P_B u_{N_1, L_1}) v_{N_2, L_2}\|_{L^2} \|w_{N_0, L_0}\|_{L^2} \\ \lesssim N_1^{-1} (L_1 L_2)^{\frac{1}{2}} \|u_{N_1, L_1}\|_{L^2} \|v_{N_2, L_2}\|_{L^2} \|w_{N_0, L_0}\|_{L^2},$$

which verifies (3.3).

As a consequence, we can assume $L_{012}^{\max} \ll (N_{012}^{\max})^3$ and $1 \ll N_{012}^{\min}$ in the sequel.

4. Proof of the key bilinear estimate: Case 1

The goal of this section is to establish (3.3) under the following assumptions.

Assumption 4.1.

- (1) $L_{012}^{\max} \ll (N_{012}^{\max})^3$,
- (2) $1 \ll N_0 \lesssim N_1 \sim N_2$,
- (3) $\max(|\xi_1|, |\xi_2|) \geq 2^{-5} N_1$.

PROPOSITION 4.2. — Assume Assumption 4.1. Then we get

$$(4.1) \quad \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim N_0^{\frac{d-4}{2}+2\varepsilon} N_1^{-1-\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

where $d\sigma_j = d\tau_j d\xi_j d\eta_j$ and $*$ denotes $(\tau, \xi, \eta) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2)$.

Remarks 4.3.

- (i) Since $L_{012}^{\max} \ll (N_{012}^{\max})^3$, it is easily observed that (4.1) yields (3.3).
- (ii) By replacing the role of \widehat{w}_{N_0, L_0} with that of \widehat{v}_{N_2, L_2} , we can show

$$(4.2) \quad \left| \int_* \widehat{v}_{N_2, L_2}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \eta_1) \widehat{w}_{N_0, L_0}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim N_2^{\frac{d-4}{2}+2\varepsilon} N_1^{-1-\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

under the assumptions

- (1) $L_{012}^{\max} \ll (N_{012}^{\max})^3$,
- (2') $1 \ll N_2 \lesssim N_0 \sim N_1$,
- (3) $\max(|\xi_1|, |\xi_2|) \geq 2^{-5} N_1$.

Clearly, (4.2) gives (3.3) under the same assumptions as above.

We divide the proof of Proposition 4.2 into the three cases.

- (Ia) $\max(|\eta_1|, |\eta_2|) \ll N_1$,
- (Ib) $\min(|\eta_1|, |\eta_2|) \ll N_1$, $\max(|\eta_1|, |\eta_2|) \sim N_1$,
- (Ic) $|\eta_1| \sim |\eta_2| \sim N_1$.

First, we consider the case (Ia): Note that the assumptions $1 \ll N_0 \lesssim N_1 \sim N_2$ and $\max(|\eta_1|, |\eta_2|) \ll N_1$ imply $|\xi_1| \sim |\xi_2| \sim N_1$.

Following [2], for $A \in \mathbb{N}$ we choose a maximally separated set $\{\omega_A^j\}_{j \in \Omega_A}$ of spherical caps of \mathbb{S}^{d-1} of aperture A , i.e. the angle $\angle(\theta_1, \theta_2)$ between any two vectors in $\theta_1, \theta_2 \in \omega_A^j$ satisfies $|\angle(\theta_1, \theta_2)| \leq A^{-1}$ and the characteristic functions $\{\mathbf{1}_{\omega_A^j}\}$ satisfy $1 \leq \sum_{j \in \Omega_A} \mathbf{1}_{\omega_A^j}(\theta) \leq 2^d$, for all $\theta \in \mathbb{S}^{d-1}$.

Further, we define the function

$$\alpha(j_1, j_2) = \inf \left\{ |\angle(\pm \theta_1, \theta_2)| : \theta_1 \in \omega_A^{j_1}, \theta_2 \in \omega_A^{j_2} \right\}$$

which measures the minimal angle between any two straight lines through the spherical caps $\omega_A^{j_1}$ and $\omega_A^{j_2}$, respectively. It is easily observed that for any fixed $j_1 \in \Omega_A$ there exist only a finite number of $j_2 \in \Omega_A$ which satisfies $\alpha(j_1, j_2) \sim A^{-1}$. Based on the above construction, for each $j \in \Omega_A$ we define

$$\mathcal{S}_j^A = \left\{ (\tau, \xi, \eta) \in \mathbb{R} \times (\mathbb{R}^d \setminus \{0\}) : \frac{(\xi, \eta)}{|(\xi, \eta)|} \in \omega_A^j \right\}$$

and the corresponding localization operator

$$\mathcal{F}(R_j^A u)(\tau, \xi, \boldsymbol{\eta}) = \mathbf{1}_{\omega_j^A} \left(\frac{(\xi, \boldsymbol{\eta})}{|(\xi, \boldsymbol{\eta})|} \right) \mathcal{F}u(\tau, \xi, \boldsymbol{\eta}).$$

In addition, we define

$$C_{(Ia)} = \{(\xi, \boldsymbol{\eta}) \in \mathbb{S}^{d-1} \mid |\boldsymbol{\eta}| \ll 1.\}, \quad \Omega_{A,(Ia)} = \{j \in \Omega_A \mid \omega_A^j \cap C_{(Ia)} \neq \emptyset\}.$$

PROPOSITION 4.4. — Assume Assumption 4.1. Let $A \gg 1$ be dyadic, $j_1, j_2 \in \Omega_{A,(Ia)}$ and $\alpha(j_1, j_2) \lesssim A^{-1}$. Then we have

$$(4.3) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_1, L_1} \cap \mathcal{S}_{j_1}^A} \int \widehat{v}_{N_2, L_2} |_{\mathcal{S}_{j_2}^A} (\tau_2, \xi_2, \boldsymbol{\eta}_2) \widehat{w}_{N_0, L_0} (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) d\sigma_2 \right\|_{L^2} \\ & \lesssim A^{-\frac{d-1}{2}} N_1^{\frac{d-3}{2}} (L_0 L_2)^{\frac{1}{2}} \|\widehat{v}_{N_2, L_2} |_{\mathcal{S}_{j_2}^A}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}, \end{aligned}$$

$$(4.4) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_2, L_2} \cap \mathcal{S}_{j_2}^A} \int \widehat{w}_{N_0, L_0} (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) \widehat{u}_{N_1, L_1} |_{\mathcal{S}_{j_1}^A} (\tau_1, \xi_1, \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ & \lesssim A^{-\frac{d-1}{2}} N_1^{\frac{d-3}{2}} (L_0 L_1)^{\frac{1}{2}} \|\widehat{w}_{N_0, L_0}\|_{L^2} \|\widehat{u}_{N_1, L_1} |_{\mathcal{S}_{j_1}^A}\|_{L^2}. \end{aligned}$$

In addition, if $|\xi| \gg A^{-1} N_1$, we get

$$(4.5) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int \widehat{u}_{N_1, L_1} |_{\mathcal{S}_{j_1}^A} (\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2} |_{\mathcal{S}_{j_2}^A} (\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ & \lesssim A^{-\frac{d-2}{2}} N_1^{\frac{d-3}{2}} (L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1} |_{\mathcal{S}_{j_1}^A}\|_{L^2} \|\widehat{v}_{N_2, L_2} |_{\mathcal{S}_{j_2}^A}\|_{L^2}. \end{aligned}$$

Proof. — First, we show (4.5). We observe that $L_{012}^{\max} \ll N_1^3$ yields $|\xi| \ll N_1$. Indeed, we calculate that

$$\begin{aligned} 3L_{012}^{\max} & \geqslant |(\tau - \xi(\xi^2 + |\boldsymbol{\eta}|^2)) - (\tau_1 - \xi_1(\xi_1^2 + |\boldsymbol{\eta}_1|^2)) \\ & \quad - ((\tau - \tau_1) - (\xi - \xi_1)((\xi - \xi_1)^2 + |\boldsymbol{\eta} - \boldsymbol{\eta}_1|^2))| \\ & = |3\xi\xi_1(\xi - \xi_1) + \xi|\boldsymbol{\eta}|^2 - \xi_1|\boldsymbol{\eta}_1|^2 - (\xi - \xi_1)|\boldsymbol{\eta} - \boldsymbol{\eta}_1|^2| \\ & =: |\Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)|. \end{aligned}$$

Since $j_1, j_2 \in \Omega_{A,(Ia)}$ and $L_{012}^{\max} \ll N_1^3$, $|\xi_1| \sim |\xi - \xi_1| \sim N_1$ and $|\boldsymbol{\eta}_1| \ll N_1$, $|\boldsymbol{\eta} - \boldsymbol{\eta}_1| \ll N_1$, the above inequality implies $|\xi| \ll N_1$. By following the

standard Cauchy–Schwarz argument, we get

$$\begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ & \leqslant \sup_{(\tau, \xi, \boldsymbol{\eta}) \in G_{N_0, L_0}} |E(\tau, \xi, \boldsymbol{\eta})|^{1/2} \left\| \left| \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A} \right|^2 * \left| \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A} \right|^2 \right\|_{L^1}^{1/2} \\ & \leqslant \sup_{(\tau, \xi, \boldsymbol{\eta}) \in G_{N_0, L_0}} |E(\tau, \xi, \boldsymbol{\eta})|^{1/2} \| \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A} \|_{L^2} \| \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A} \|_{L^2}, \end{aligned}$$

where $E(\tau, \xi, \boldsymbol{\eta}) \subset \mathbb{R}^{d+1}$ is defined by

$$E(\tau, \xi, \boldsymbol{\eta}) = \{(\tau_1, \xi_1, \boldsymbol{\eta}_1) \in G_{N_1, L_1} \cap \mathcal{S}_{j_1}^A \mid (\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) \in G_{N_2, L_2} \cap \mathcal{S}_{j_2}^A\}.$$

Thus, it suffices to show

$$(4.6) \quad \sup_{(\tau, \xi, \boldsymbol{\eta}) \in G_{N_0, L_0}} |E(\tau, \xi, \boldsymbol{\eta})| \lesssim A^{-(d-2)} N_1^{d-3} L_1 L_2.$$

Clearly, for fixed $(\xi_1, \boldsymbol{\eta}_1)$, it holds

$$(4.7) \quad \sup_{(\tau, \xi, \boldsymbol{\eta}) \in G_{N_0, L_0}} |\{\tau_1 \mid (\tau_1, \xi_1, \boldsymbol{\eta}_1) \in E(\tau, \xi, \boldsymbol{\eta})\}| \lesssim \min(L_1, L_2).$$

A simple calculation yields

$$(4.8) \quad |\partial_{\xi_1} \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)| = |3\xi(\xi - 2\xi_1) + \boldsymbol{\eta} \cdot (\boldsymbol{\eta} - 2\boldsymbol{\eta}_1)|,$$

$$(4.9) \quad |\nabla_{\boldsymbol{\eta}_1} \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)| = 2|(\xi - \xi_1)\boldsymbol{\eta} - \xi\boldsymbol{\eta}_1|.$$

Let $r_1 = |(\xi_1, \boldsymbol{\eta}_1)|$, $r_2 = |(\xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1)|$ and $(\theta_1, \boldsymbol{\theta}'_1) = (\xi_1/r_1, \boldsymbol{\eta}_1/r_1)$, $(\theta_2, \boldsymbol{\theta}'_2) = ((\xi - \xi_1)/r_2, (\boldsymbol{\eta} - \boldsymbol{\eta}_1)/r_2) \in \mathbb{S}^{d-1}$. Since $(\theta_1, \boldsymbol{\theta}'_1) \times (\theta_2, \boldsymbol{\theta}'_2) \in \omega_A^{j_1} \times \omega_A^{j_2}$ with $\alpha(j_1, j_2) \lesssim A^{-1}$ and $|\xi| \ll N_1$, we have $|(\theta_1, \boldsymbol{\theta}'_1) + (\theta_2, \boldsymbol{\theta}'_2)| \lesssim A^{-1}$. Furthermore, the assumption $(\theta_j, \boldsymbol{\theta}'_j) \in C_{(Ia)}$ implies $|\theta_1| \sim |\theta_2| \sim 1$ and $\max(|\boldsymbol{\theta}'_1|, |\boldsymbol{\theta}'_2|) \ll 1$. Therefore, we deduce from the assumption $A^{-1}N_1 \ll |\xi|$ that

$$\begin{aligned} |\boldsymbol{\eta}| &= |r_1 \boldsymbol{\theta}'_1 + r_2 \boldsymbol{\theta}'_2| \leqslant |r_1 - r_2| |\boldsymbol{\theta}'_1| + r_2 |\boldsymbol{\theta}'_1 + \boldsymbol{\theta}'_2| \\ &\leqslant |r_1 \theta_1 - r_2 \theta_1| + r_2 |\boldsymbol{\theta}'_1 + \boldsymbol{\theta}'_2| \\ &\leqslant |r_1 \theta_1 + r_2 \theta_2| + 2r_2 |(\theta_1, \boldsymbol{\theta}'_1) + (\theta_2, \boldsymbol{\theta}'_2)| \leqslant 2|\xi|. \end{aligned}$$

Hence, if $(\tau_1, \xi_1, \boldsymbol{\eta}_1) \in E(\tau, \xi, \boldsymbol{\eta})$, the above inequality and (4.8), (4.9) yield

$$\begin{aligned} |\partial_{r_1} \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)| &= |(\theta_1 \partial_{\xi_1} + \boldsymbol{\theta}'_1 \cdot \nabla_{\boldsymbol{\eta}_1}) \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)| \\ &\geqslant |\theta_1| |\partial_{\xi_1} \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)| - |\boldsymbol{\theta}'_1| |\nabla_{\boldsymbol{\eta}_1} \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)| \\ &\gtrsim |\xi| N_1 \geqslant A^{-1} N_1^2, \end{aligned}$$

which implies that r_1 is confined to a set of measure $\sim A \max(L_1, L_2)/N_1^2$ for fixed $\theta_1, \boldsymbol{\theta}'_1$ since, as we saw above, it holds that

$$\begin{aligned} & \max(L_1, L_2) \\ & \gtrsim |(\tau_1 - \xi_1(\xi_1^2 + |\boldsymbol{\eta}_1|^2)) + ((\tau - \tau_1) - (\xi - \xi_1)((\xi - \xi_1)^2 + |\boldsymbol{\eta} - \boldsymbol{\eta}_1|^2))| \\ & = |(\tau - \xi(\xi^2 + |\boldsymbol{\eta}|^2)) + \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)|. \end{aligned}$$

Therefore, we get

$$\begin{aligned} & |\{(\xi_1, \boldsymbol{\eta}_1) \mid (\tau_1, \xi_1, \boldsymbol{\eta}_1) \in E(\tau, \xi, \boldsymbol{\eta})\}| \\ & = \int_{\boldsymbol{\theta}'_1} \int_{\theta_1} \int_{r_1} \mathbf{1}_{E(\tau, \xi, \boldsymbol{\eta})}(r_1, \theta_1, \boldsymbol{\theta}'_1) r_1^{d-1} dr_1 d\theta_1 d\boldsymbol{\theta}'_1 \\ & \lesssim A^{-(d-2)} N_1^{d-3} \max(L_1, L_2). \end{aligned}$$

This and (4.7) give (4.6).

Next, we consider (4.3). By the same argument it suffices to prove

$$(4.10) \quad \sup_{(\tau_1, \xi_1, \boldsymbol{\eta}_1) \in G_{N_1, L_1}} |E_1(\tau_1, \xi_1, \boldsymbol{\eta}_1)| \lesssim A^{-(d-1)} N_1^{d-3} L_0 L_2,$$

where

$$\begin{aligned} & E_1(\tau_1, \xi_1, \boldsymbol{\eta}_1) \\ & = \{(\tau_2, \xi_2, \boldsymbol{\eta}_2) \in G_{N_2, L_2} \cap \mathcal{S}_{j_2}^A \mid (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) \in G_{N_0, L_0}\}. \end{aligned}$$

As above, we have

$$\begin{aligned} 3L_{012}^{\max} & \geqslant |3(\xi_1 + \xi_2)\xi_1\xi_2 + (\xi_1 + \xi_2)|\boldsymbol{\eta}_1 + \boldsymbol{\eta}_2|^2 - \xi_1|\boldsymbol{\eta}_1|^2 - \xi_2|\boldsymbol{\eta}_2|^2| \\ & =: |\Phi_{\xi_1, \boldsymbol{\eta}_1}^{(1)}(\xi_2, \boldsymbol{\eta}_2)|. \end{aligned}$$

We compute $|\partial_{\xi_2} \Phi_{\xi_1, \boldsymbol{\eta}_1}^{(1)}(\xi_2, \boldsymbol{\eta}_2)| \gtrsim N_1^2$, $|\nabla_{\boldsymbol{\eta}_1} \Phi_{\xi_1, \boldsymbol{\eta}_1}^{(1)}(\xi_2, \boldsymbol{\eta}_2)| \ll N_1^2$, and with the same notation for polar coordinates as above, we obtain

$$|\partial_{r_2} \Phi_{\xi_1, \boldsymbol{\eta}_1}^{(1)}(\xi_2, \boldsymbol{\eta}_2)| \gtrsim N_1^2.$$

If $(\tau_2, \xi_2, \boldsymbol{\eta}_2) \in E_1(\tau_1, \xi_1, \boldsymbol{\eta}_1)$, then for a fixed angular part $(\theta_2, \boldsymbol{\theta}'_2)$ of $(\xi_2, \boldsymbol{\eta}_2)$, the radial direction r_2 is confined to an interval of length $\lesssim \max(L_0, L_2)/N_1^2$. By the analogue of (4.7) we conclude (4.10) and the proof of (4.3) is complete.

Finally, (4.4) follows by symmetry. \square

PROPOSITION 4.5. — Assume Assumption 4.1. Let $A \gg 1$ be dyadic, $j_1, j_2 \in \Omega_{A,(\text{Ia})}$, $\alpha(j_1, j_2) \sim A^{-1}$ and $|\xi_1 + \xi_2| \lesssim A^{-1}N_1$. Then we get

$$(4.11) \quad \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}.$$

Proof. — In the proof of Proposition 4.4 we proved $|\eta| \lesssim |\xi| + A^{-1}N_1 \lesssim A^{-1}N_1$. Thus, by almost orthogonality, we may assume that η_j is confined to a ball whose radius is comparable to $A^{-1}N_1$. We write $\eta_j = (\eta_j, \eta'_j)$. Further, without loss of generality, we can assume $\max(|\eta'_1|, |\eta'_2|) \lesssim A^{-1}N_1$. Indeed, we can apply a rotation in the η -subspace, since the phase function is invariant under such rotations.

Since $\alpha(j_1, j_2) \sim A^{-1}$, we have $\max(|\xi_1 \eta_2 - \xi_2 \eta_1|, |\xi_1 \eta'_2 - \xi_2 \eta'_1|) \sim A^{-1}N_1^2$. We first consider the case $|\xi_1 \eta_2 - \xi_2 \eta_1| \sim A^{-1}N_1^2$. For fixed η'_1, η'_2 , we will show that

$$(4.12) \quad \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\tau_2, \xi_2, \eta_2) d\widehat{\sigma}_1 d\widehat{\sigma}_2 \right| \\ \lesssim A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}(\eta'_1)\|_{L_{\tau\xi\eta}^2} \|\widehat{v}_{N_2, L_2}(\eta'_2)\|_{L_{\tau\xi\eta}^2} \|\widehat{w}_{N_0, L_0}(\eta')\|_{L_{\tau\xi\eta}^2},$$

where $d\widehat{\sigma}_j = d\tau_j d\xi_j d\eta_j$ and $\widehat{*}$ denotes $(\tau, \xi, \eta) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2)$. (4.12) implies (4.11) because

$$\left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim \int \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\tau_1, \xi_1, \eta_1) \right. \\ \times \left. \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\tau_2, \xi_2, \eta_2) d\widehat{\sigma}_1 d\widehat{\sigma}_2 \right| d\eta'_1 d\eta'_2 \\ \lesssim A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \int \|\widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\eta'_1)\|_{L_{\tau\xi\eta}^2} \\ \times \|\widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\eta'_2)\|_{L_{\tau\xi\eta}^2} \|\widehat{w}_{N_0, L_0}(\eta'_1 + \eta'_2)\|_{L_{\tau\xi\eta}^2} d\eta'_1 d\eta'_2 \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

where we used the support conditions in the last step.

Now, we prove (4.12). We use the shorthand notation

$$\begin{aligned} f_{\eta'_1}(\tau_1, \xi_1, \eta_1) &= \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\tau_1, \xi_1, \eta_1), \\ g_{\eta'_2}(\tau_2, \xi_2, \eta_2) &= \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\tau_2, \xi_2, \eta_2), \\ h_{\eta'}(\tau, \xi, \eta) &= \widehat{w}_{N_0, L_0}(\tau, \xi, \eta), \end{aligned}$$

and show

$$\begin{aligned} (4.13) \quad & \left| \int_* h_{\eta'}(\tau, \xi, \eta) f_{\eta'_1}(\tau_1, \xi_1, \eta_1) g_{\eta'_2}(\tau_2, \xi_2, \eta_2) d\hat{\sigma}_1 d\hat{\sigma}_2 \right| \\ & \lesssim A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{\eta'_1}\|_{L_{\tau\xi\eta}^2} \|g_{\eta'_2}\|_{L_{\tau\xi\eta}^2} \|h_{\eta'}\|_{L_{\tau\xi\eta}^2}. \end{aligned}$$

Applying the transformation $\tau_1 = \xi_1(\xi_1^2 + \eta_1^2 + |\eta'_1|^2) + c_1$ and $\tau_2 = \xi_2(\xi_2^2 + \eta_2^2 + |\eta'_2|^2) + c_2$ and Fubini's theorem, we find that it suffices to prove

$$\begin{aligned} (4.14) \quad & \left| \int h_{\eta'}(\phi_{\eta'_1, c_1}(\xi_1, \eta_1) + \phi_{\eta'_2, c_2}(\xi_2, \eta_2)) f_{\eta'_1}(\phi_{\eta'_1, c_1}(\xi_1, \eta_1)) \right. \\ & \quad \times g_{\eta'_2}(\phi_{\eta'_2, c_2}(\xi_2, \eta_2)) d\xi_1 d\eta_1 d\xi_2 d\eta_2 \Big| \\ & \lesssim A^{\frac{1}{2}} N_1^{-2} \|f_{\eta'_1} \circ \phi_{\eta'_1, c_1}\|_{L_{\xi\eta}^2} \|g_{\eta'_2} \circ \phi_{\eta'_2, c_2}\|_{L_{\xi\eta}^2} \|h_{\eta'}\|_{L_{\tau\xi\eta}^2}, \end{aligned}$$

where $h_{\eta'}(\tau, \xi, \eta)$ is supported in $c_0 \leq \tau - \xi(\xi^2 + \eta^2 + |\eta'|^2) \leq c_0 + 1$ and

$$\phi_{\eta'_j, c_j}(\xi_j, \eta_j) = (\xi_j(\xi_j^2 + \eta_j^2 + |\eta'_j|^2) + c_j, \xi_j, \eta_j) \quad \text{for } j = 1, 2.$$

We use the scaling $(\tau, \xi, \eta) \rightarrow (N_1^3 \tau, N_1 \xi, N_1 \eta)$ to define

$$\begin{aligned} \tilde{f}_{\eta'_1}(\tau_1, \xi_1, \eta_1) &= f_{\eta'_1}(N_1^3 \tau_1, N_1 \xi_1, N_1 \eta_1), \\ \tilde{g}_{\eta'_2}(\tau_2, \xi_2, \eta_2) &= g_{\eta'_2}(N_1^3 \tau_2, N_1 \xi_2, N_1 \eta_2), \\ \tilde{h}_{\eta'}(\tau, \xi, \eta) &= h_{\eta'}(N_1^3 \tau, N_1 \xi, N_1 \eta). \end{aligned}$$

Let $\widetilde{\eta_j} = N_1^{-1} \eta'_j$, $\widetilde{c_j} = N_1^{-3} c_j$. The inequality (4.14) reduces to

$$\begin{aligned} & \left| \int \tilde{h}_{\eta'}(\phi_{\widetilde{\eta_1}, \widetilde{c_1}}(\xi_1, \eta_1) + \phi_{\widetilde{\eta_2}, \widetilde{c_2}}(\xi_2, \eta_2)) \right. \\ & \quad \times \tilde{f}_{\eta'_1}(\phi_{\widetilde{\eta_1}, \widetilde{c_1}}(\xi_1, \eta_1)) \tilde{g}_{\eta'_2}(\phi_{\widetilde{\eta_2}, \widetilde{c_2}}(\xi_2, \eta_2)) d\xi_1 d\eta_1 d\xi_2 d\eta_2 \Big| \\ & \lesssim A^{\frac{1}{2}} N_1^{-\frac{3}{2}} \|\tilde{f}_{\eta'_1} \circ \phi_{\widetilde{\eta_1}, \widetilde{c_1}}\|_{L_{\xi\eta}^2} \|\tilde{g}_{\eta'_2} \circ \phi_{\widetilde{\eta_2}, \widetilde{c_2}}\|_{L_{\xi\eta}^2} \|\tilde{h}_{\eta'}\|_{L_{\tau\xi\eta}^2}, \end{aligned}$$

Note that $|\widetilde{\eta_j}| \lesssim A^{-1}$ and we easily see $|\xi_j| \sim 1$, $|\eta_j| \ll 1$ and $|\xi_1 \eta_2 - \xi_2 \eta_1| \sim A^{-1}$ if $(\xi_1, \eta_1) \in \text{supp}(\tilde{f}_{\eta'_1} \circ \phi_{\widetilde{\eta_1}, \widetilde{c_1}})$, $(\xi_2, \eta_2) \in \text{supp}(\tilde{g}_{\eta'_2} \circ \phi_{\widetilde{\eta_2}, \widetilde{c_2}})$. Therefore,

letting $\tilde{\eta} = N_1^{-1}\eta'$, we can assume that $\tilde{h}_{\eta'}$ is supported in $S_3(N_1^{-3})$ where

$$S_3(N_1^{-3}) = \left\{ (\tau, \xi, \eta) \left| \begin{array}{l} |(\xi, \eta)| \lesssim A^{-1}, \\ \frac{c_0}{N_1^3} \leq \tau - \xi(\xi^2 + \eta^2 + |\tilde{\eta}|^2) \leq \frac{c_0 + 1}{N_1^3} \end{array} \right. \right\}.$$

By density and duality, it suffices to show that for continuous $\tilde{f}_{\eta'_1}$ and $\tilde{g}_{\eta'_2}$ it holds that

$$(4.15) \quad \|\tilde{f}_{\eta'_1}|_{S_1} * \tilde{g}_{\eta'_2}|_{S_2}\|_{L^2(S_3(N_1^{-3}))} \lesssim A^{\frac{1}{2}} N_1^{-\frac{3}{2}} \|\tilde{f}_{\eta'_1}\|_{L^2(S_1)} \|\tilde{g}_{\eta'_2}\|_{L^2(S_2)}$$

where S_1, S_2 denote the following surfaces

$$\begin{aligned} S_1 &= \{\phi_{\tilde{\eta}_1, c_1}(\xi_1, \eta_1) \in \mathbb{R}^3 \mid (\xi_1, \eta_1) \in \text{supp}(\tilde{f}_{\eta'_1} \circ \phi_{\tilde{\eta}_1, c_1})\}, \\ S_2 &= \{\phi_{\tilde{\eta}_2, c_2}(\xi_2, \eta_2) \in \mathbb{R}^3 \mid (\xi_2, \eta_2) \in \text{supp}(\tilde{g}_{\eta'_2} \circ \phi_{\tilde{\eta}_2, c_2})\}. \end{aligned}$$

(4.15) is immediately obtained by the following.

$$(4.16) \quad \|\tilde{f}_{\eta'_1}|_{S_1} * \tilde{g}_{\eta'_2}|_{S_2}\|_{L^2(S_3)} \lesssim A^{\frac{1}{2}} \|\tilde{f}_{\eta'_1}\|_{L^2(S_1)} \|\tilde{g}_{\eta'_2}\|_{L^2(S_2)}$$

where

$$S_3 = \left\{ (\psi_{\tilde{\eta}}(\xi, \eta), \xi, \eta) \in \mathbb{R}^3 \left| \begin{array}{l} |(\xi, \eta)| \lesssim A^{-1}, \\ \psi_{\tilde{\eta}}(\xi, \eta) = \xi(\xi^2 + \eta^2 + |\tilde{\eta}|^2) + \frac{c'_0}{N_1^3} \end{array} \right. \right\},$$

for any fixed $c'_0 \in [c_0, c_0 + 1]$. Since $\text{diam}(S_3) \lesssim A^{-1}$, by the almost orthogonality and harmless decompositions, we may assume

$$(4.17) \quad \text{diam}(S_i) \ll A^{-1} \quad \text{for } i = 1, 2, 3.$$

For any $\lambda_i \in S_i$, there exist (ξ_1, η_1) , (ξ_2, η_2) , (ξ, η) such that

$$\lambda_1 = \phi_{\tilde{\eta}_1, c_1}(\xi_1, \eta_1), \quad \lambda_2 = \phi_{\tilde{\eta}_2, c_2}(\xi_2, \eta_2), \quad \lambda_3 = (\psi_{\tilde{\eta}}(\xi, \eta), \xi, \eta),$$

and the unit normals n_i on λ_i are written as

$$n_i(\lambda_i) = \frac{1}{\sqrt{1 + (3\xi_i^2 + \eta_i^2 + |\tilde{\eta}_i|^2)^2 + 4\xi_i^2\eta_i^2}} (-1, 3\xi_i^2 + \eta_i^2 + |\tilde{\eta}_i|^2, 2\xi_i\eta_i)$$

for $i = 1, 2$, and the same for $n_3(\lambda_3)$. Clearly, the surfaces S_1, S_2, S_3 satisfy the following Hölder condition.

$$(4.18) \quad \sup_{\lambda_i, \hat{\lambda}_i \in S_i} \frac{|n_i(\lambda_i) - n_i(\hat{\lambda}_i)|}{|\lambda_i - \hat{\lambda}_i|} + \frac{|n_i(\lambda_i)(\lambda_i - \hat{\lambda}_i)|}{|\lambda_i - \hat{\lambda}_i|^2} \lesssim 1.$$

We may assume that there exist $(\hat{\xi}_1, \hat{\eta}_1)$, $(\hat{\xi}_2, \hat{\eta}_2)$, $(\hat{\xi}, \hat{\eta})$ such that

$$(\hat{\xi}_1, \hat{\eta}_1) + (\hat{\xi}_2, \hat{\eta}_2) = (\hat{\xi}, \hat{\eta}),$$

$$\phi_{\tilde{\eta}_1, c_1}(\hat{\xi}_1, \hat{\eta}_1) \in S_1, \quad \phi_{\tilde{\eta}_2, c_2}(\hat{\xi}_2, \hat{\eta}_2) \in S_2, \quad (\psi_{\tilde{\eta}}(\hat{\xi}, \hat{\eta}), \hat{\xi}, \hat{\eta}) \in S_3,$$

otherwise the left-hand side of (4.16) vanishes. Let $\widehat{\lambda}_1 = \phi_{\widetilde{\eta}_1, c_1}(\widehat{\xi}_1, \widehat{\eta}_1)$, $\widehat{\lambda}_2 = \phi_{\widetilde{\eta}_2, c_2}(\widehat{\xi}_2, \widehat{\eta}_2)$, $\widehat{\lambda}_3 = (\psi_{\widetilde{\eta}}(\widehat{\xi}, \widehat{\eta}), \widehat{\xi}, \widehat{\eta})$. For any $i = 1, 2, 3$ and $\lambda_i, \widehat{\lambda}_i \in S_i$ (4.17) implies that

$$(4.19) \quad |\mathbf{n}_i(\lambda_i) - \mathbf{n}_i(\widehat{\lambda}_i)| \ll A^{-1}.$$

From (4.17) and (4.18), once the following transversality condition

$$(4.20) \quad A^{-1} \lesssim |\det N(\lambda_1, \lambda_2, \lambda_3)| \quad \text{for any } \lambda_i \in S_i.$$

is verified, we obtain the desired estimate (4.16) by applying the nonlinear Loomis–Whitney inequality from Proposition 2.3. Using $|\widehat{\xi}_j| \sim 1$, $|\widehat{\eta}_j| \ll 1$, $|\widetilde{\eta}_j| \lesssim A^{-1}$ and $|\widehat{\xi}_1 \widehat{\eta}_2 - \widehat{\xi}_2 \widehat{\eta}_1| \sim A^{-1}$, we compute

$$\begin{aligned} & |\det N(\widehat{\lambda}_1, \widehat{\lambda}_2, \widehat{\lambda}_3)| \\ & \gtrsim \left| \det \begin{pmatrix} -1 & -1 & -1 \\ 3\widehat{\xi}_1^2 + \widehat{\eta}_1^2 + |\widetilde{\eta}_1|^2 & 3\widehat{\xi}_2^2 + \widehat{\eta}_2^2 + |\widetilde{\eta}_2|^2 & 3\widehat{\xi}^2 + \widehat{\eta}^2 + |\widetilde{\eta}|^2 \\ 2\widehat{\xi}_1 \widehat{\eta}_1 & 2\widehat{\xi}_2 \widehat{\eta}_2 & 2\widehat{\xi} \widehat{\eta} \end{pmatrix} \right| \\ & \gtrsim |(\widehat{\xi}_1 \widehat{\eta}_2 - \widehat{\xi}_2 \widehat{\eta}_1)(3(\widehat{\xi}_1^2 + \widehat{\xi}_1 \widehat{\xi}_2 + \widehat{\xi}_2^2) - (\widehat{\eta}_1^2 + \widehat{\eta}_1 \widehat{\eta}_2 + \widehat{\eta}_2^2)) \\ & \quad + (\widehat{\xi}_1 \widehat{\eta}_2 + \widehat{\xi}_2(\widehat{\eta}_1 + \widehat{\eta}_2))|\widetilde{\eta}_1|^2 - 2(\widehat{\xi}_1 \widehat{\eta}_1 - \widehat{\xi}_2 \widehat{\eta}_2)\widetilde{\eta}_1 \cdot \widetilde{\eta}_2 \\ & \quad - (\widehat{\xi}_1(\widehat{\eta}_1 + \widehat{\eta}_2) + \widehat{\xi}_2 \widehat{\eta}_1)|\widetilde{\eta}_2|^2| \\ & \gtrsim A^{-1}, \end{aligned}$$

which implies (4.20) due to (4.19). In the above computation, we used multi-linearity in the columns to separate the contributions of $\widetilde{\eta}_1$, $\widetilde{\eta}_2$ and $\widetilde{\eta}$ from the main one corresponding to the first line above.

Next, we treat the case $|\xi_1 \boldsymbol{\eta}'_2 - \xi_2 \boldsymbol{\eta}'_1| \sim A^{-1} N_1^2$. Without loss of generality, we assume $|\xi_1 \eta'_2 - \xi_2 \eta'_1| \sim A^{-1} N_1^2$ where η'_1 and η'_2 are the first components of $\boldsymbol{\eta}'_1$ and $\boldsymbol{\eta}'_2$, respectively. By replacing the role of (η_1, η_2) with that of (η'_1, η'_2) in the proof in the previous case, it suffices to show

$$(4.21) \quad \left| \int_{\bar{*}} \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1} |_{S_{j_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2} |_{S_{j_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\bar{\sigma}_1 d\bar{\sigma}_2 \right| \\ \lesssim A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}(\bar{\boldsymbol{\eta}}_1)\|_{L^2_{\tau \xi \eta'}} \|\widehat{v}_{N_2, L_2}(\bar{\boldsymbol{\eta}}_2)\|_{L^2_{\tau \xi \eta'}} \|\widehat{w}_{N_0, L_0}(\bar{\boldsymbol{\eta}})\|_{L^2_{\tau \xi \eta'}},$$

where $\bar{\boldsymbol{\eta}}_j \in \mathbb{R}^{d-2}$ denotes $\boldsymbol{\eta}_j$ excluding η'_j , $d\bar{\sigma}_j = d\tau_j d\xi_j d\eta'_j$ and $\bar{*}$ denotes $(\tau, \xi, \eta') = (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta'_1 + \eta'_2)$. Similarly to the previous case, (4.21) is established by the nonlinear Loomis–Whitney inequality. To avoid redundancy, we here only consider the transversality condition, which is

given by

$$\begin{aligned} & |(\xi_1\eta'_2 - \xi_2\eta'_1)(3(\xi_1^2 + \xi_1\xi_2 + \xi_2^2) - (\eta'_1{}^2 + \eta'_1\eta'_2 + \eta'_2{}^2)) \\ & \quad + (\xi_1\eta'_2 + \xi_2(\eta'_1 + \eta'_2))|\bar{\eta}_1|^2 - 2(\xi_1\eta'_1 - \xi_2\eta'_2)\bar{\eta}_1 \cdot \bar{\eta}_2 \\ & \quad - (\xi_1(\eta'_1 + \eta'_2) + \xi_2\eta'_1)|\bar{\eta}_2|^2| \\ & \gtrsim A^{-1}N_1^4, \end{aligned}$$

where we used $|\xi_1\eta'_2 - \xi_2\eta'_1| \sim A^{-1}N_1^2$, $|\eta'_j| \lesssim A^{-1}N_1$ and $|\bar{\eta}_j| \ll N_1$. \square

PROPOSITION 4.6. — Assume Assumption 4.1. Let $A \gg 1$ be dyadic, $j_1, j_2 \in \Omega_{A,(\text{Ia})}$, $\alpha(j_1, j_2) \sim A^{-1}$. Then we get

$$\begin{aligned} (4.22) \quad & \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Proof. — Proposition 4.5 gives (4.22) if $|\xi_1 + \xi_2| \lesssim A^{-1}N_1$. Thus we assume $|\xi_1 + \xi_2| \gg A^{-1}N_1$. We show that $|\xi_1 + \xi_2| \gg A^{-1}N_1$ provides

$$\begin{aligned} (4.23) \quad & |\Phi(\xi_1, \boldsymbol{\eta}_1, \xi_2, \boldsymbol{\eta}_2)| \\ & := |3\xi_1\xi_2(\xi_1 + \xi_2) + (\xi_1 + \xi_2)|\boldsymbol{\eta}_1 + \boldsymbol{\eta}_2|^2 - \xi_1|\boldsymbol{\eta}_1|^2 - \xi_2|\boldsymbol{\eta}_2|^2| \\ & \gtrsim A^{-1}N_1^3. \end{aligned}$$

Since $L_{012}^{\max} \gtrsim |\Phi(\xi_1, \boldsymbol{\eta}_1, \xi_2, \boldsymbol{\eta}_2)|$, this and Proposition 4.4 verify (4.22). Recall that the assumptions $j_1, j_2 \in \Omega_{A,(\text{Ia})}$ and $|\xi_1 + \xi_2| \gg A^{-1}N_1$ imply $\max(|\boldsymbol{\eta}_1|, |\boldsymbol{\eta}_2|) \ll N_1$, $|\boldsymbol{\eta}_1 + \boldsymbol{\eta}_2| \lesssim |\xi_1 + \xi_2|$. Therefore, we have

$$\begin{aligned} & |(\xi_1 + \xi_2)|\boldsymbol{\eta}_1 + \boldsymbol{\eta}_2|^2 - \xi_1|\boldsymbol{\eta}_1|^2 - \xi_2|\boldsymbol{\eta}_2|^2| \\ & \leqslant |\xi_1 + \xi_2|(|\boldsymbol{\eta}_1 + \boldsymbol{\eta}_2|^2 + |\boldsymbol{\eta}_1|^2) + |\xi_2|(|\boldsymbol{\eta}_1|^2 - |\boldsymbol{\eta}_2|^2)| \\ & \ll N_1^2|\xi_1 + \xi_2|, \end{aligned}$$

which immediately yields (4.23). \square

Proof of (4.1) in the case (Ia). — Assume that $(\xi_j, \boldsymbol{\eta}_j)/|(\xi_j, \boldsymbol{\eta}_j)| \in C_{(\text{Ia})}$. Define

$$I_{j_1, j_2}^A = \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\mathcal{S}_{j_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\mathcal{S}_{j_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right|.$$

We observe

$$\begin{aligned} & \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim \sum_{1 \ll A \leq N_1^{3/2}} \sum_{\alpha(j_1, j_2) \sim A^{-1}} I_{j_1, j_2}^A + \sum_{\alpha(j_1, j_2) \lesssim N_1^{-3/2}} I_{j_1, j_2}^{N_1^{3/2}}. \end{aligned}$$

Note that $\langle |(\xi_1 + \xi_2, \eta_1 + \eta_2)| \rangle \sim N_0 \gtrsim A^{-1} N_1$ if $(\tau_1, \xi_1, \eta_1) \times (\tau_2, \xi_2, \eta_2) \in \mathcal{S}_{j_1}^A \times \mathcal{S}_{j_2}^A$ with $\alpha(j_1, j_2) \sim A^{-1}$. Thus, the former term is estimated by using Proposition 4.6 as

$$\begin{aligned} & \sum_{1 \ll A \leq N_1^{3/2}} \sum_{\alpha(j_1, j_2) \sim A^{-1}} I_{j_1, j_2}^A \\ & \lesssim \sum_{1 \ll A \leq N_1^{3/2}} \sum_{\alpha(j_1, j_2) \sim A^{-1}} N_0^{\frac{d-3}{2}} N_1^{-\frac{3}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|\widehat{u}_{N_1, L_1}\|_{\mathcal{S}_{j_1}^A} \|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{\mathcal{S}_{j_2}^A} \|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2} \\ & \lesssim \sum_{1 \ll A \leq N_1^{3/2}} N_0^{\frac{d-3}{2}} N_1^{-\frac{3}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2} \\ & \lesssim (\log N_1) N_0^{\frac{d-3}{2}} N_1^{-\frac{3}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}. \end{aligned}$$

By using (4.3) in Proposition 4.4 for the latter term, we complete the proof. \square

Next, we treat the case (Ib) $\min(|\eta_1|, |\eta_2|) \ll N_1$, $\max(|\eta_1|, |\eta_2|) \sim N_1$. Without loss of generality, we assume $|\eta_1| \sim N_1$ and $|\eta_2| \ll N_1$. Note that $N_1 \sim N_2$ and $|\eta_2| \ll N_1$ imply $|\xi_2| \sim N_1$. We define

$$\begin{aligned} F(\xi_1, \eta_1, \xi_2, \eta_2) &= (\xi_1 \eta_2 - \xi_2 \eta_1) (3(\xi_1^2 + \xi_1 \xi_2 + \xi_2^2) - (\eta_1^2 + \eta_1 \eta_2 + \eta_2^2)) \\ &+ (\xi_1 \eta_2 + \xi_2 (\eta_1 + \eta_2)) |\eta'_1|^2 - 2(\xi_1 \eta_1 - \xi_2 \eta_2) \eta'_1 \cdot \eta'_2 \\ &\quad - (\xi_1 (\eta_1 + \eta_2) + \xi_2 \eta_1) |\eta'_2|^2. \end{aligned}$$

Recall that this function $F(\xi_1, \eta_1, \xi_2, \eta_2)$ appeared in the proof of Proposition 4.5 and provided a transversality of the three hypersurfaces.

LEMMA 4.7. — Assume Assumption 4.1, $|\eta_1| \sim N_1$ and $|\eta_2| \ll N_1$. Then we have $|F(\xi_1, \eta_1, \xi_2, \eta_2)| \gtrsim N_1^4$.

Proof. — We show $L_{012}^{\max} \gtrsim N_1^3$ if $|\eta_1| \sim N_1$, $|\eta_2| \ll N_1$ and $|F(\xi_1, \eta_1, \xi_2, \eta_2)| \ll N_1^4$. Since $|\eta_1| \sim N_1$, $|\eta_2| \ll N_1$, it is observed that

$$\begin{aligned} (4.24) \quad & |F(\xi_1, \eta_1, \xi_2, \eta_2)| \ll N_1^4 \\ & \implies |\xi_2 \eta_1 (3(\xi_1^2 + \xi_1 \xi_2 + \xi_2^2) - \eta_1^2) - \xi_2 \eta_1 |\eta'_1|^2| \ll N_1^4 \\ & \implies ||\eta_1|^2 - 3(\xi_1^2 + \xi_1 \xi_2 + \xi_2^2)| \ll N_1^2. \end{aligned}$$

We use the function $\Phi(\xi_1, \eta_1, \xi_2, \eta_2)$ which was defined in the proof of Proposition 4.6. It follows from $|\eta_2| \ll N_1$ and (4.24) that there is $0 < c \ll 1$ such that

$$\begin{aligned} |\Phi(\xi_1, \eta_1, \xi_2, \eta_2)| & \geq |3\xi_1 \xi_2 (\xi_1 + \xi_2) + \xi_2 |\eta_1|^2| - |2(\xi_1 + \xi_2) \eta_1 \cdot \eta_2 + \xi_1 |\eta_2|^2| \\ & \geq |\xi_2| |3\xi_1 (\xi_1 + \xi_2) + 3(\xi_1^2 + \xi_1 \xi_2 + \xi_2^2)| - cN_1^3 \\ & = 3|\xi_2| |2\xi_1^2 + 2\xi_1 \xi_2 + \xi_2^2| - cN_1^3 \\ & \gtrsim N_1^3, \end{aligned}$$

which completes the proof. \square

Lemma 4.7 suggests that we can obtain (4.1) by the same argument as in the proof of Proposition 4.5. We omit the details.

Lastly, we consider the case (Ic) $|\eta_1| \sim |\eta_2| \sim N_1$.

In this case, we perform an angular decomposition in the η -space. In the same way as above (see [2]), for $A \in \mathbb{N}$ we choose a maximally separated set $\{\bar{\omega}_A^j\}_{j \in \bar{\Omega}_A}$ of spherical caps of \mathbb{S}^{d-2} of aperture A^{-1} , i.e. the angle $\angle(\theta_1, \theta_2)$ between any two vectors in $\theta_1, \theta_2 \in \bar{\omega}_A^j$ satisfies $|\angle(\theta_1, \theta_2)| \leq A^{-1}$ and the characteristic functions $\{\mathbf{1}_{\bar{\omega}_A^j}\}$ satisfy $1 \leq \sum_{j \in \bar{\Omega}_A} \mathbf{1}_{\bar{\omega}_A^j}(\theta) \leq 2^d$, for all $\theta \in \mathbb{S}^{d-2}$. Further, we define the function

$$\bar{\alpha}(j_1, j_2) = \inf \left\{ |\angle(\pm \theta_1, \theta_2)| : \theta_1 \in \bar{\omega}_A^{j_1}, \theta_2 \in \bar{\omega}_A^{j_2} \right\}.$$

For each $j \in \bar{\Omega}_A$ we define

$$\bar{\mathcal{S}}_j^A = \left\{ (\tau, \xi, \eta) \in \mathbb{R} \times \mathbb{R} \times (\mathbb{R}^{d-1} \setminus \{0\}) : \frac{\eta}{|\eta|} \in \bar{\omega}_A^j \right\}$$

and the corresponding localization operator

$$\mathcal{F}(\bar{R}_j^A u)(\tau, \xi, \eta) = \mathbf{1}_{\omega_j^A} \left(\frac{\eta}{|\eta|} \right) \mathcal{F}u(\tau, \xi, \eta).$$

Let $k = (k_{(1)}, \dots, k_{(d)}) \in \mathbb{Z}^d$. We define regular cubes $\{\mathcal{C}_k^A\}_{k \in \mathbb{Z}^d}$ whose side length is $A^{-1}N_1$ and $\{\tilde{\mathcal{C}}_k^A\}_{k \in \mathbb{Z}^d}$ as

$$\mathcal{C}_k^A = \{x = (x_1, \dots, x_d) \in \mathbb{R}^d \mid x_i \in A^{-1}N_1[k_{(i)}, k_{(i)}+1) \text{ for all } i = 1, \dots, d.\},$$

we set $\tilde{\mathcal{C}}_k^A = \mathbb{R} \times \mathcal{C}_k^A$, and lastly we define $\mathcal{E}_{j,k}^A = \bar{\mathcal{S}}_j^A \cap \tilde{\mathcal{C}}_k^A$.

PROPOSITION 4.8. — Assume Assumption 4.1 and $|\boldsymbol{\eta}_1| \sim |\boldsymbol{\eta}_2| \sim N_1$. Let $\bar{\alpha}(j_1, j_2) \sim A^{-1}$ and $k_1, k_2 \in \mathbb{Z}^d$. Then we get

$$(4.25) \quad \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\mathcal{E}_{j_1, k_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\mathcal{E}_{j_2, k_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

where $d\sigma_j = d\tau_j d\xi_j d\boldsymbol{\eta}_j$ and $*$ denotes $(\tau, \xi, \boldsymbol{\eta}) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2)$.

Proof. — After rotation, we can assume $|\eta_1 \eta'_2 - \eta_2 \eta'_1| \sim A^{-1} N_1^2$ and $|\boldsymbol{\eta}'_j| \lesssim A^{-1} N_1$. Recall that η_j and η'_j are first and second components of $\boldsymbol{\eta}_j$, respectively. For simplicity, we use $\check{\boldsymbol{\eta}}_j \in \mathbb{R}^{d-3}$ which satisfies $\boldsymbol{\eta}_j = (\eta_j, \boldsymbol{\eta}'_j) = (\eta_j, \eta'_j, \check{\boldsymbol{\eta}}_j)$. Similarly to the proof of Proposition 4.5, for fixed $\xi_1, \xi_2, \check{\boldsymbol{\eta}}_1, \check{\boldsymbol{\eta}}_2$, it suffices to show

$$(4.26) \quad \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\mathcal{E}_{j_1, k_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\mathcal{E}_{j_2, k_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\tilde{\sigma}_1 d\tilde{\sigma}_2 \right| \\ \lesssim A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}(\xi_1, \check{\boldsymbol{\eta}}_1)\|_{L^2_{\tau \eta \eta'}} \\ \times \|\widehat{v}_{N_2, L_2}(\xi_2, \check{\boldsymbol{\eta}}_2)\|_{L^2_{\tau \eta \eta'}} \|\widehat{w}_{N_0, L_0}(\xi, \check{\boldsymbol{\eta}})\|_{L^2_{\tau \eta \eta'}},$$

where $d\tilde{\sigma}_j = d\tau_j d\eta_j d\eta'_j$ and $\tilde{*}$ denotes $(\tau, \eta, \eta') = (\tau_1 + \tau_2, \eta_1 + \eta_2, \eta'_1 + \eta'_2)$. We follow the proof of Proposition 4.5. Assume that $\xi_1, \xi_2, \check{\boldsymbol{\eta}}_1, \check{\boldsymbol{\eta}}_2$ are fixed. We use the functions $\tilde{f}_{\xi_1, \check{\boldsymbol{\eta}}_1}, \tilde{g}_{\xi_2, \check{\boldsymbol{\eta}}_2}$ on \mathbb{R}^3 that are defined as

$$\begin{aligned} \tilde{f}_{\xi_1, \check{\boldsymbol{\eta}}_1}(\tau_1, \eta_1, \eta'_1) &= \widehat{u}_{N_1, L_1}|_{\mathcal{E}_{j_1, k_1}^A}(N_1^3 \tau_1, \xi_1, N_1 \eta_1, N_1 \eta'_1, \check{\boldsymbol{\eta}}_1), \\ \tilde{g}_{\xi_2, \check{\boldsymbol{\eta}}_2}(\tau_2, \eta_2, \eta'_2) &= \widehat{v}_{N_2, L_2}|_{\mathcal{E}_{j_2, k_2}^A}(N_1^3 \tau_2, \xi_2, N_1 \eta_2, N_1 \eta'_2, \check{\boldsymbol{\eta}}_2), \end{aligned}$$

and show the following estimate:

$$(4.27) \quad \|\tilde{f}_{\xi_1, \check{\boldsymbol{\eta}}_1}|_{S_1} * \tilde{g}_{\xi_2, \check{\boldsymbol{\eta}}_2}|_{S_2}\|_{L^2(S_3)} \lesssim A^{\frac{1}{2}} \|\tilde{f}_{\xi_1, \check{\boldsymbol{\eta}}_1}\|_{L^2(S_1)} \|\tilde{g}_{\xi_2, \check{\boldsymbol{\eta}}_2}\|_{L^2(S_2)}.$$

Here, $c_0, c_1, c_2 \in \mathbb{R}$, $\tilde{\xi} = N_1^{-1} \xi$, $\tilde{\xi}_j = N_1^{-1} \xi_j$, $\bar{\boldsymbol{\eta}}_j = N_1^{-1} \check{\boldsymbol{\eta}}_j$, $\bar{\boldsymbol{\eta}} = N_1^{-1} \check{\boldsymbol{\eta}}$ and for

$$\phi_{\xi_j, \check{\boldsymbol{\eta}}_j, c_j}(\eta, \eta') = (\xi_j(\xi_j^2 + |\boldsymbol{\eta}|^2) + c_j, \eta, \eta'),$$

the surfaces are given as

$$\begin{aligned} S_1 &= \{\phi_{\tilde{\xi}_1, \bar{\boldsymbol{\eta}}_1, c_1}(\eta_1, \eta'_1) \in \mathbb{R}^3 \mid (\eta_1, \eta'_1) \in \text{supp}(\tilde{f}_{\xi_1, \check{\boldsymbol{\eta}}_1} \circ \phi_{\tilde{\xi}_1, \bar{\boldsymbol{\eta}}_1, c_1})\}, \\ S_2 &= \{\phi_{\tilde{\xi}_2, \bar{\boldsymbol{\eta}}_2, c_2}(\eta_2, \eta'_2) \in \mathbb{R}^3 \mid (\eta_2, \eta'_2) \in \text{supp}(\tilde{g}_{\xi_2, \check{\boldsymbol{\eta}}_2} \circ \phi_{\tilde{\xi}_2, \bar{\boldsymbol{\eta}}_2, c_2})\}, \\ S_3 &= \{(\psi_{\tilde{\xi}, \bar{\boldsymbol{\eta}}}(\eta, \eta'), \eta, \eta') \in \mathbb{R}^3 \mid \psi_{\tilde{\xi}, \bar{\boldsymbol{\eta}}}(\eta, \eta') = \xi(\xi^2 + |\boldsymbol{\eta}|^2) + c_0\}. \end{aligned}$$

Since $\text{diam}(S_1) \lesssim A^{-1}$, $\text{diam}(S_2) \lesssim A^{-1}$, we can assume $\text{diam}(S_3) \lesssim A^{-1}$. We easily confirm that S_1, S_2, S_3 satisfy the necessary regularity and diameter conditions to use the nonlinear Loomis–Whitney inequality. Thus,

here we only confirm that S_1, S_2, S_3 satisfy the suitable transversality condition. We define $\lambda_i \in S_i$ as

$$\lambda_1 = \phi_{\xi_1, \bar{\eta}_1, c_1}(\eta_1, \eta'_1), \quad \lambda_2 = \phi_{\xi_2, \bar{\eta}_2, c_2}(\eta_2, \eta'_2), \quad \lambda_3 = (\psi_{\xi, \bar{\eta}}(\eta, \eta'), \eta, \eta').$$

The unit normals \mathbf{n}_i on λ_i are described explicitly as

$$\mathbf{n}_i(\lambda_i) = \frac{1}{\sqrt{1 + 4\xi_i^2(\eta_i^2 + \eta'^2)}} \left(-1, 2\tilde{\xi}_i \eta_i, 2\tilde{\xi}_i \eta'_i \right),$$

for $i = 1, 2$, and the same for $\mathbf{n}_3(\lambda_3)$. Letting $\widehat{\boldsymbol{\eta}}_1 = (\widehat{\eta}_1, \widehat{\eta}'_1)$, $\widehat{\boldsymbol{\eta}}_2 = (\widehat{\eta}_2, \widehat{\eta}'_2)$, $\widehat{\boldsymbol{\eta}} = (\widehat{\eta}, \widehat{\eta}')$, $\widehat{\boldsymbol{\eta}}_1 + \widehat{\boldsymbol{\eta}}_2 = \widehat{\boldsymbol{\eta}}$ and

$\widehat{\lambda}_1 := \phi_{\xi_1, \bar{\eta}_1, c_1}(\widehat{\boldsymbol{\eta}}_1) \in S_1$, $\widehat{\lambda}_2 := \phi_{\xi_2, \bar{\eta}_2, c_2}(\widehat{\boldsymbol{\eta}}_2) \in S_2$, $\widehat{\lambda}_3 := (\psi_{\xi, \bar{\eta}}(\widehat{\boldsymbol{\eta}}), \widehat{\boldsymbol{\eta}}) \in S_3$, we show

$$A^{-1} \lesssim |\det N(\widehat{\lambda}_1, \widehat{\lambda}_2, \widehat{\lambda}_3)|,$$

which means the transversality of S_1, S_2, S_3 and completes the proof. We observe

$$\begin{aligned} |\det N(\widehat{\lambda}_1, \widehat{\lambda}_2, \widehat{\lambda}_3)| &\gtrsim \left| \det \begin{pmatrix} -1 & -1 & -1 \\ 2\tilde{\xi}_1 \widehat{\eta}_1 & 2\tilde{\xi}_2 \widehat{\eta}_2 & 2\tilde{\xi} \widehat{\eta} \\ 2\tilde{\xi}_1 \widehat{\eta}'_1 & 2\tilde{\xi}_2 \widehat{\eta}'_2 & 2\tilde{\xi} \widehat{\eta}' \end{pmatrix} \right| \\ &\gtrsim |(\widehat{\eta}_1 \widehat{\eta}'_2 - \widehat{\eta}_2 \widehat{\eta}'_1)(\tilde{\xi}_1^2 + \tilde{\xi}_1 \tilde{\xi}_2 + \tilde{\xi}_2^2)| \\ &\gtrsim A^{-1}. \end{aligned}$$

Here we used the assumptions $|\eta_1 \eta'_2 - \eta_2 \eta'_1| \sim A^{-1} N_1^2$ and $\max(|\xi_1|, |\xi_2|) \sim N_1$ which imply $|\widehat{\eta}_1 \widehat{\eta}'_2 - \widehat{\eta}_2 \widehat{\eta}'_1| \sim A^{-1}$ and $\max(|\tilde{\xi}_1|, |\tilde{\xi}_2|) \sim 1$, respectively.

□

PROPOSITION 4.9. — Assume Assumption 4.1 and $|\boldsymbol{\eta}_1| \sim |\boldsymbol{\eta}_2| \sim N_1$. Let $\bar{\alpha}(j_1, j_2) \sim A^{-1}$. Then we get

$$\begin{aligned} &\left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1} |_{\bar{\mathcal{S}}_{j_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2} |_{\bar{\mathcal{S}}_{j_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ &\lesssim N_0^{\frac{d-4}{2}+2\varepsilon} N_1^{-1-\frac{3}{2}\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}, \end{aligned}$$

where $d\sigma_j = d\tau_j d\xi_j d\boldsymbol{\eta}_j$ and $*$ denotes $(\tau, \xi, \boldsymbol{\eta}) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2)$.

Before we state a proof, let us see that Proposition 4.9 establishes (4.1) in the case (Ic).

Proof of (4.1) in the case (Ic). — For convenience, we use

$$\bar{I}_{j_1, j_2}^A := \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1} |_{\bar{\mathcal{S}}_{j_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2} |_{\bar{\mathcal{S}}_{j_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right|.$$

We observe that

$$\begin{aligned} \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim \sum_{2 \leq A \leq N_1^6} \sum_{\bar{\alpha}(j_1, j_2) \sim A^{-1}} \bar{I}_{j_1, j_2}^A + \sum_{\bar{\alpha}(j_1, j_2) \lesssim N_1^{-6}} \bar{I}_{j_1, j_2}^{N_1^6}. \end{aligned}$$

For the former term, by using Proposition 4.9 and the almost orthogonality of j_1, j_2 which satisfy $\bar{\alpha}(j_1, j_2) \sim A^{-1}$, we get

$$\begin{aligned} & \sum_{2 \leq A \leq N_1^6} \sum_{\bar{\alpha}(j_1, j_2) \sim A^{-1}} \bar{I}_{j_1, j_2}^A \\ & \lesssim \sum_{2 \leq A \leq N_1^6} N_0^{\frac{d-4}{2}+2\varepsilon} N_1^{-1-\frac{3}{2}\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2} \\ & \lesssim N_0^{\frac{d-4}{2}+2\varepsilon} N_1^{-1-\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}. \end{aligned}$$

For the latter term, since the size of the set $\{(\xi_1, \eta_1) | (\tau_1, \xi_1, \eta_1) \in \bar{\mathcal{S}}_{j_1}^{N_1^6}\}$ is less than $\sim N_1^{-5(d-2)+2} \leq N_1^{-3}$, we easily obtain

$$\bar{I}_{j_1, j_2}^{N_1^6} \lesssim N_1^{-\frac{3}{2}} L_1^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{\bar{\mathcal{S}}_{j_1}^A} \|\widehat{v}_{N_2, L_2}\|_{\bar{\mathcal{S}}_{j_2}^A} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

which completes the proof of (4.1) in the case (Ic). \square

The next subsection is devoted to the proof of Proposition 4.9. Note that, as in the proof of Proposition 4.5, by rotating η_1, η_2 , we can assume $|\eta_1 \eta'_2 - \eta_2 \eta'_1| \sim A^{-1} N_1^2$ and $|\eta'_j| \lesssim A^{-1} N_1$. Further, by performing the invertible linear transformation $(\xi_j, \eta_j) \rightarrow (\xi_j + \eta_j, \sqrt{3}(\xi_j - \eta_j))$, it is easily observed that Proposition 4.9 is equivalent to Proposition 4.10 below.

4.1. Proof of Proposition 4.9

As justified by the above discussion, in this subsection we assume the following:

Assumption 4.1'.

- (1) $L_{012}^{\max} \ll (N_{012}^{\max})^3$,
- (2) $1 \ll N_0 \lesssim N_1 \sim N_2$,
- (3) $\max(|\xi_1 + \eta_1|, |\xi_2 + \eta_2|) \geq 2^{-5} N_1$.

PROPOSITION 4.10. — *In addition to Assumption 4.1', suppose that $|\xi_j - \eta_j| \sim N_1$, $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$ where $j = 1, 2$ and $|(\xi_1 - \eta_1)\eta'_2 - (\xi_2 - \eta_2)\eta'_1| \sim A^{-1}N_1^2$. Then we get*

$$(4.28) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim N_0^{\frac{d-4}{2} + 2\varepsilon} N_1^{-1 - \frac{3}{2}\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2},$$

where functions f_{N_1, L_1} , g_{N_2, L_2} , h_{N_0, L_0} satisfy

$$(4.29) \quad \text{supp } f_{N_1, L_1} \subset G_{N_1, L_1}, \quad \text{supp } g_{N_2, L_2} \subset G_{N_2, L_2}, \quad \text{supp } h_{N_0, L_0} \subset G_{N_0, L_0},$$

and $G_{N, L}$ is the set

$$\{(\tau, \xi, \boldsymbol{\eta}) \in \mathbb{R}^{d+1} \mid \langle |(\xi, \boldsymbol{\eta})| \rangle \sim N, \langle \tau - (\xi^3 + \eta^3) - (\xi + \eta)|\boldsymbol{\eta}'|^2 \rangle \sim L\}.$$

We consider Proposition 4.10 instead of Proposition 4.9. The advantage in this way is that we can reuse the propositions and lemmas that were established in the paper by the second author [21] which was concerned with the 2D Zakharov–Kuznetsov equation. In [21], the following symmetrized 2D Zakharov–Kuznetsov equation was considered.

$$\partial_t u + (\partial_x^3 + \partial_y^3)u = 4^{-\frac{1}{3}}(\partial_x + \partial_y)(u^2), \quad (t, x, y) \in \mathbb{R} \times \mathbb{R}^2.$$

This equation is equivalent to the original 2D Zakharov–Kuznetsov equation, which can be seen by applying the above linear transformation $(\xi_j, \eta_j) \rightarrow (\xi_j + \eta_j, \sqrt{3}(\xi_j - \eta_j))$ to the original 2D Zakharov–Kuznetsov equation. See [16].

Now we turn to Proposition 4.10. Note that the assumptions in Proposition 4.10 suggest that we can assume $A^{-1}N_1 \lesssim N_0$. We divide the proof into the two cases

$$|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \gtrsim 1 \quad \text{and} \quad |\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \ll 1.$$

First, we consider the case $|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \gtrsim 1$.

DEFINITION 4.11. — *Let $M \gg 1$ be a dyadic number and $\ell = (\ell_{(1)}, \ell_{(2)}) \in \mathbb{Z}^2$. We define square-tiles $\{\mathcal{T}_\ell^M\}_{\ell \in \mathbb{Z}^2}$ whose side length is $M^{-1}N_1$ and $\{\tilde{\mathcal{T}}_\ell^M\}_{k \in \mathbb{Z}^2}$ as follows:*

$$\mathcal{T}_\ell^M := \{(\xi, \eta) \in \mathbb{R}^2 \mid (\xi, \eta) \in M^{-1}N_1([\ell_{(1)}, \ell_{(1)} + 1) \times [\ell_{(2)}, \ell_{(2)} + 1))\} \\ \tilde{\mathcal{T}}_\ell^M := \mathbb{R} \times \mathcal{T}_\ell^M \times \mathbb{R}^{d-2}.$$

DEFINITION 4.12 (Whitney type decomposition). — Let A, M, \widehat{M} be dyadic such that $1 \ll \widehat{M} \leq M \leq A$ and

$$\begin{aligned}\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2) &= \xi_1 \xi_2 (\xi_1 + \xi_2) + \eta_1 \eta_2 (\eta_1 + \eta_2), \\ \bar{F}(\xi_1, \eta_1, \xi_2, \eta_2) &= \xi_1 \eta_2 + \xi_2 \eta_1 + 2(\xi_1 \eta_1 + \xi_2 \eta_2).\end{aligned}$$

We define

$$\begin{aligned}Z_M^1 &= \{(\ell_1, \ell_2) \in (\mathbb{Z}^2)^2 \mid |\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1} N_1^3 \text{ for all } (\xi_j, \eta_j) \in \mathcal{T}_{\ell_j}^M\}, \\ Z_M^2 &= \{(\ell_1, \ell_2) \in (\mathbb{Z}^2)^2 \mid |\bar{F}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1} N_1^2 \text{ for all } (\xi_j, \eta_j) \in \mathcal{T}_{\ell_j}^M\}, \\ Z_M &= Z_M^1 \cup Z_M^2 \subset \mathbb{Z}^2 \times \mathbb{Z}^2, \quad R_M = \bigcup_{(\ell_1, \ell_2) \in Z_M} \mathcal{T}_{\ell_1}^M \times \mathcal{T}_{\ell_2}^M \subset \mathbb{R}^2 \times \mathbb{R}^2.\end{aligned}$$

It is clear that $M_1 \leq M_2 \implies R_{M_1} \subset R_{M_2}$. Further, we define

$$Q_M = \begin{cases} R_M \setminus R_{\frac{M}{2}} & \text{for } M > \widehat{M}, \\ R_{\widehat{M}} & \text{for } M = \widehat{M}. \end{cases}$$

and a set of integer pair $Z'_M \subset Z_M$ as

$$\bigcup_{(\ell_1, \ell_2) \in Z'_M} \mathcal{T}_{\ell_1}^M \times \mathcal{T}_{\ell_2}^M = Q_M.$$

We easily see that Z'_M is uniquely defined and

$$M_1 \neq M_2 \implies Q_{M_1} \cap Q_{M_2} = \emptyset, \quad \bigcup_{\widehat{M} \leq M \leq M_0} Q_M = R_{M_0}$$

where $M_0 \geq \widehat{M}$ is dyadic. Thus, we can decompose $\mathbb{R}^2 \times \mathbb{R}^2$ as

$$\mathbb{R}^2 \times \mathbb{R}^2 = \left(\bigcup_{\widehat{M} \leq M \leq M_0} Q_M \right) \cup (R_{M_0})^c.$$

Lastly, we define

$$\begin{aligned}\mathcal{A} &= \{(\tau_1, \xi_1, \eta_1) \times (\tau_2, \xi_2, \eta_2) \in \mathbb{R}^{d+1} \times \mathbb{R}^{d+1} \mid |\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \gtrsim 1\}, \\ \tilde{Z}_M &= \{(\ell_1, \ell_2) \in Z'_M \mid (\tilde{\mathcal{T}}_{\ell_1}^M \times \tilde{\mathcal{T}}_{\ell_2}^M) \cap (G_{N_1, L_1} \times G_{N_2, L_2}) \cap \mathcal{A} \neq \emptyset\}.\end{aligned}$$

By the same argument as for the 2D case in [21], we can obtain the following estimate.

PROPOSITION 4.13. — Assume $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$ and (4.29). Let A, M be dyadic which satisfy $1 \ll M \leq A$ and $(\ell_1, \ell_2) \in \tilde{Z}_M$. Then we get

$$(4.30) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\tilde{T}_{\ell_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{T}_{\ell_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim A^{-\frac{d-2}{2}} M^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{T}_{\ell_1}^M}\|_{L^2} \|g_{N_2, L_2} |_{\tilde{T}_{\ell_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2},$$

where $d\sigma_j = d\tau_j d\xi_j d\eta_j$ and $*$ denotes $(\tau, \xi, \boldsymbol{\eta}) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2)$.

Proof. — Since $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$, for fixed $\boldsymbol{\eta}'_1, \boldsymbol{\eta}'_2$, it suffices to show

$$(4.31) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\tilde{T}_{\ell_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{T}_{\ell_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\hat{\sigma}_1 d\hat{\sigma}_2 \right| \\ \lesssim M^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{T}_{\ell_1}^M}(\boldsymbol{\eta}'_1)\|_{L^2_{\tau \xi \eta}} \\ \times \|g_{N_2, L_2} |_{\tilde{T}_{\ell_2}^M}(\boldsymbol{\eta}'_2)\|_{L^2_{\tau \xi \eta}} \|h_{N_0, L_0}(\boldsymbol{\eta}')\|_{L^2_{\tau \xi \eta}},$$

where $d\hat{\sigma}_j = d\tau_j d\xi_j d\eta_j$ and $\hat{*}$ denotes $(\tau, \xi, \eta) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2)$. (4.31) is established by the same argument as for Propositions 3.3–3.5 in [21] which considered the Cauchy problem of the 2D Zakharov–Kuznetsov equation. The only difference is that, in [21] it was assumed that $f_{N_1, L_1}, g_{N_2, L_2}, h_{N_0, L_0}$ satisfy

$$\text{supp } f_{N_1, L_1} \subset \bar{G}_{N_1, L_1}, \quad \text{supp } g_{N_2, L_2} \subset \bar{G}_{N_2, L_2}, \quad \text{supp } h_{N_0, L_0} \subset \bar{G}_{N_0, L_0}, \\ \bar{G}_{N, L} := \{(\tau, \xi, \boldsymbol{\eta}) \in \mathbb{R}^{d+1} \mid \langle |\langle \xi, \boldsymbol{\eta} \rangle| \rangle \sim N, \langle \tau - (\xi^3 + \eta^3) \rangle \sim L\},$$

instead of (4.29). We will see that, because of the assumptions $M \leq A$ and $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$, the proofs of Propositions 3.3–3.5 in [21] can be transferred. Firstly, either $|\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1}N_1^3$ or $|\bar{F}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1}N_1^2$ holds under the assumption $(\ell_1, \ell_2) \in \tilde{Z}_M$. We first assume

$$|\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1}N_1^3$$

and show (4.31). For simplicity, we use

$$f_{\boldsymbol{\eta}'_1}(\tau_1, \xi_1, \eta_1) := f_{N_1, L_1} |_{\tilde{T}_{\ell_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1), \\ g_{\boldsymbol{\eta}'_2}(\tau_2, \xi_2, \eta_2) := g_{N_2, L_2} |_{\tilde{T}_{\ell_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2), \\ h_{\boldsymbol{\eta}'}(\tau, \xi, \eta) := h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}).$$

Since

$$\begin{aligned} 3L_{012}^{\max} &\geq \left| \tau_1 + \tau_2 - ((\xi_1 + \xi_2)^3 + (\eta_1 + \eta_2)^3) - (\xi_1 + \xi_2 + \eta_1 + \eta_2)|\boldsymbol{\eta}'_1 + \boldsymbol{\eta}'_2|^2 \right. \\ &\quad \left. - (\tau_1 - (\xi_1^3 + \eta_1^3) - (\xi_1 + \eta_1)|\boldsymbol{\eta}'_1|^2) - (\tau_2 - (\xi_2^3 + \eta_2^3) - (\xi_2 + \eta_2)|\boldsymbol{\eta}'_2|^2) \right| \\ &\gtrsim |\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| + \mathcal{O}(A^{-2}N_1) \gtrsim M^{-1}N_1^3, \end{aligned}$$

the following estimates which correspond to Proposition 3.3 in [21] immediately yields (4.31).

$$\begin{aligned} (4.32) \quad &\left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{\boldsymbol{\eta}'_1}(\tau_1, \xi_1, \eta_1) g_{\boldsymbol{\eta}'_2}(\tau - \tau_1, \xi - \xi_1, \eta - \eta_1) d\hat{\sigma}_1 \right\|_{L_{\tau\xi\eta}^2} \\ &\lesssim (MN_1)^{-\frac{1}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{\boldsymbol{\eta}'_1}\|_{L_{\tau\xi\eta}^2} \|g_{\boldsymbol{\eta}'_2}\|_{L_{\tau\xi\eta}^2}, \end{aligned}$$

and

$$\begin{aligned} (4.33) \quad &\left\| \mathbf{1}_{G_{N_1, L_1} \cap \tilde{\mathcal{T}}_{k_1}^A} \int g_{\boldsymbol{\eta}'_2}(\tau_1, \xi_2, \eta_2) h_{\boldsymbol{\eta}'}(\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2) d\hat{\sigma}_2 \right\|_{L_{\tau\xi\eta}^2} \\ &\lesssim (MN_1)^{-\frac{1}{2}} (L_0 L_2)^{\frac{1}{2}} \|g_{\boldsymbol{\eta}'_2}\|_{L_{\tau\xi\eta}^2} \|h_{\boldsymbol{\eta}'}\|_{L^2}, \end{aligned}$$

as well as

$$\begin{aligned} (4.34) \quad &\left\| \mathbf{1}_{G_{N_2, L_2} \cap \tilde{\mathcal{T}}_{k_2}^A} \int h_{\boldsymbol{\eta}'}(\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2) f_{\boldsymbol{\eta}'_1}(\tau_1, \xi_1, \eta_1) d\hat{\sigma}_1 \right\|_{L_{\tau\xi\eta}^2} \\ &\lesssim (MN_1)^{-\frac{1}{2}} (L_0 L_1)^{\frac{1}{2}} \|h_{\boldsymbol{\eta}'}\|_{L_{\tau\xi\eta}^2} \|f_{\boldsymbol{\eta}'_1}\|_{L_{\tau\xi\eta}^2}. \end{aligned}$$

Here we sketch the proof of (4.32) only. The other estimates (4.33) and (4.34) can be obtained in the same way as for (4.32). We first observe that the assumptions imply

$$(4.35) \quad \max(|(\xi_1^2 - (\xi - \xi_1)^2)|, |(\eta_1^2 - (\eta - \eta_1)^2)|) \gtrsim N_1^2.$$

If (4.35) does not hold, we can assume one of the following.

- (1) $|\xi_1 - (\xi - \xi_1)| \ll N_1$ and $|\eta_1 - (\eta - \eta_1)| \ll N_1$,
- (2) $|\xi_1 - (\xi - \xi_1)| \ll N_1$ and $|\eta_1 + (\eta - \eta_1)| \ll N_1$,
- (3) $|\xi_1 + (\xi - \xi_1)| \ll N_1$ and $|\eta_1 - (\eta - \eta_1)| \ll N_1$,
- (4) $|\xi_1 + (\xi - \xi_1)| \ll N_1$ and $|\eta_1 + (\eta - \eta_1)| \ll N_1$.

(1) and (4) contradict the angular assumption

$$|\sin \angle((\xi_1, \eta_1), (\xi - \xi_1, \eta - \eta_1))| \gtrsim 1.$$

We show (2) contradicts one of the assumptions. Clearly, $\max(|\xi_1|, |\xi - \xi_1|) \gtrsim N_1$ holds under this angular assumption. Without loss of generality,

we can assume $|\xi_1| \gtrsim N_1$. This and the inequality $|\xi_1 - (\xi - \xi_1)| \ll N_1$ in (2) yield $\min(|\xi|, |\xi - \xi_1|) \gtrsim N_1$ which, combined with $|\eta| = |\eta_1 + (\eta - \eta_1)| \ll N_1$ in (2), gives

$$\begin{aligned} & 3L_{012}^{\max} \\ & \geq 3 \max(|\tau - \xi^3 - \eta^3|, |\tau_1 - \xi_1^3 - \eta_1^3|, |\tau - \tau_1 - (\xi - \xi_1)^3 - (\eta - \eta_1)^3|) \\ & \quad + \mathcal{O}(A^{-2}N_1^3) \\ & \geq |\xi\xi_1(\xi - \xi_1) + \eta\eta_1(\eta - \eta_1)| + \mathcal{O}(A^{-2}N_1^3) \\ & \gtrsim N_1^3 + \mathcal{O}(A^{-2}N_1^3) \gtrsim N_1^3 \end{aligned}$$

which contradicts $L_{012}^{\max} \ll N_1^3$. Similarly, we can show that (3) contradicts at least one of the assumptions. Without loss of generality, we assume $|\xi_1^2 - (\xi - \xi_1)^2| \gtrsim N_1^2$.

We turn to show (4.32). By the Cauchy–Schwarz inequality, we get

$$\begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{\boldsymbol{\eta}'_1}(\tau_1, \xi_1, \eta_1) g_{\boldsymbol{\eta}'_2}(\tau - \tau_1, \xi - \xi_1, \eta - \eta_1) d\widehat{\sigma}_1 \right\|_{L_{\tau\xi\eta}^2} \\ & \leq \left\| \mathbf{1}_{G_{N_0, L_0}} \left(|f_{\boldsymbol{\eta}'_1}|^2 * |g_{\boldsymbol{\eta}'_2}|^2 \right)^{1/2} |E(\tau, \xi, \eta)|^{1/2} \right\|_{L_{\tau\xi\eta}^2} \\ & \leq \sup_{(\tau, \xi, \eta) \in G_{N_0, L_0}} |E(\tau, \xi, \eta)|^{1/2} \left\| |f_{\boldsymbol{\eta}'_1}|^2 * |g_{\boldsymbol{\eta}'_2}|^2 \right\|_{L_{\tau\xi\eta}^1}^{1/2} \\ & \leq \sup_{(\tau, \xi, \eta) \in G_{N_0, L_0}} |E(\tau, \xi, \eta)|^{1/2} \|f_{\boldsymbol{\eta}'_1}\|_{L_{\tau\xi\eta}^2} \|g_{\boldsymbol{\eta}'_2}\|_{L_{\tau\xi\eta}^2}, \end{aligned}$$

where $E(\tau, \xi, \eta) \subset \mathbb{R}^3$ is defined by

$$E(\tau, \xi, \eta) := \{(\tau_1, \xi_1, \eta_1) \in \text{supp}(f_{\boldsymbol{\eta}'_1}) \mid (\tau - \tau_1, \xi - \xi_1, \eta - \eta_1) \in \text{supp}(g_{\boldsymbol{\eta}'_2})\}.$$

Thus, it suffices to show

$$(4.36) \quad \sup_{(\tau, \xi, \eta) \in G_{N_0, L_0}} |E(\tau, \xi, \eta)| \lesssim (MN_1)^{-1} L_1 L_2.$$

For fixed (ξ_1, η_1) , we easily have

$$(4.37) \quad \sup_{(\tau, \xi, \eta) \in G_{N_0, L_0}} |\{\tau_1 \mid (\tau_1, \xi_1, \eta_1) \in E(\tau, \xi, \eta)\}| \lesssim \min(L_1, L_2).$$

Let $C(\xi, \boldsymbol{\eta}', \xi_1, \boldsymbol{\eta}'_1) = (\xi_1 + \eta_1)|\boldsymbol{\eta}'_1|^2 + (\xi - \xi_1 + \eta - \eta_1)|\boldsymbol{\eta}' - \boldsymbol{\eta}'_1|^2$. We observe

$$\begin{aligned} & \max(L_1, L_2) \\ & \gtrsim |(\tau_1 - \xi_1^3 - \eta_1^3) + (\tau - \tau_1) - (\xi - \xi_1)^3 - (\eta - \eta_1)^3 - C(\xi, \boldsymbol{\eta}, \xi_1, \boldsymbol{\eta}_1)| \\ & = |(\tau - \xi^3 - \eta^3) + 3(\xi\xi_1(\xi - \xi_1) + \eta\eta_1(\eta - \eta_1)) - C(\xi, \boldsymbol{\eta}, \xi_1, \boldsymbol{\eta}_1)|. \end{aligned}$$

Thus, we deduce from $|\partial_{\xi_1} (\xi \xi_1 (\xi - \xi_1))| = |\xi_1^2 - (\xi - \xi_1)^2| \gtrsim N_1^2$, $|\boldsymbol{\eta}'_1| \lesssim A^{-1} N_1$ and $|\boldsymbol{\eta}' - \boldsymbol{\eta}'_1| \lesssim A^{-1} N_1$ that, for fixed η_1 , it holds that

$$(4.38) \quad \sup_{(\tau, \xi, \eta) \in G_{N_0, L_0}} |\{\xi_1 \mid (\tau_1, \xi_1, \eta_1) \in E(\tau, \xi, \eta)\}| \lesssim N_1^{-2} \max(L_1, L_2).$$

Lastly, since $(\tau_1, \xi_1, \eta_1) \in \text{supp}(f_{\boldsymbol{\eta}'_1})$ implies $(\xi_1, \eta_1) \in \mathcal{T}_{\ell_1}^M$, we have

$$(4.39) \quad \sup_{(\tau, \xi, \eta) \in G_{N_0, L_0}} |\{\eta_1 \mid (\tau_1, \xi_1, \eta_1) \in E(\tau, \xi, \eta)\}| \lesssim M^{-1} N_1.$$

The estimates (4.37)-(4.39) complete the proof of (4.36).

Next we show (4.31) under the assumption $|\bar{F}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1} N_1^2$ by following the proof for Proposition 3.5 in [21]. By Fubini's theorem, (4.31) reduces to

$$(4.40) \quad \left| \int h_{\boldsymbol{\eta}'} (\varphi_{\boldsymbol{\eta}'_1, c_1}(\xi_1, \eta_1) + \varphi_{\boldsymbol{\eta}'_2, c_2}(\xi_2, \eta_2)) f_{\boldsymbol{\eta}'_1} (\varphi_{\boldsymbol{\eta}'_1, c_1}(\xi_1, \eta_1)) \right. \\ \times g(\varphi_{\boldsymbol{\eta}'_2, c_2}(\xi_2, \eta_2)) d\xi_1 d\eta_1 d\xi_2 d\eta_2 \Big| \\ \lesssim M^{\frac{1}{2}} N_1^{-2} \|f_{\boldsymbol{\eta}'_1} \circ \varphi_{\boldsymbol{\eta}'_1, c_1}\|_{L_{\xi\eta}^2} \|g_{\boldsymbol{\eta}'_2} \circ \varphi_{\boldsymbol{\eta}'_2, c_2}\|_{L_{\xi\eta}^2} \|h_{\boldsymbol{\eta}'}\|_{L_{\tau\xi\eta}^2},$$

where $h_{\boldsymbol{\eta}'}(\tau, \xi, \eta)$ is supported in $c_0 \leq \tau - \xi^3 - \eta^3 - (\xi + \eta)|\boldsymbol{\eta}'|^2 \leq c_0 + 1$ and

$$\varphi_{\boldsymbol{\eta}'_j, c_j}(\xi, \eta) = (\xi^3 + \eta^3 + (\xi + \eta)|\boldsymbol{\eta}'_j|^2 + c_j, \xi, \eta) \quad \text{for } j = 1, 2.$$

Note that if $\boldsymbol{\eta}'_1 = \boldsymbol{\eta}'_2 = \boldsymbol{\eta}' = 0$, (4.40) corresponds exactly to the inequality (3.17) in [21]. Similarly to the proof of Proposition 4.5, we define

$$\begin{aligned} \tilde{f}_{\boldsymbol{\eta}'_1}(\tau_1, \xi_1, \eta_1) &= f_{\boldsymbol{\eta}'_1}(N_1^3 \tau_1, N_1 \xi_1, N_1 \eta_1), \\ \tilde{g}_{\boldsymbol{\eta}'_2}(\tau_2, \xi_2, \eta_2) &= g_{\boldsymbol{\eta}'_2}(N_1^3 \tau_2, N_1 \xi_2, N_1 \eta_2), \\ \tilde{h}_{\boldsymbol{\eta}'}(\tau, \xi, \eta) &= h_{\boldsymbol{\eta}'}(N_1^3 \tau, N_1 \xi, N_1 \eta), \end{aligned}$$

and prove

$$(4.41) \quad \|\tilde{f}_{\boldsymbol{\eta}'_1}|_{\bar{S}_1} * \tilde{g}_{\boldsymbol{\eta}'_2}|_{\bar{S}_2}\|_{L^2(\bar{S}_3)} \lesssim M^{\frac{1}{2}} \|\tilde{f}_{\boldsymbol{\eta}'_1}\|_{L^2(\bar{S}_1)} \|\tilde{g}_{\boldsymbol{\eta}'_2}\|_{L^2(\bar{S}_2)},$$

where $\tilde{\boldsymbol{\eta}}_j = N_1^{-1} \boldsymbol{\eta}'_j$, $\tilde{\boldsymbol{\eta}} = N_1^{-1} \boldsymbol{\eta}'$, $\tilde{c}_j = N_1^{-3} c_j$ and

$$\begin{aligned} \bar{S}_1 &= \left\{ \varphi_{\tilde{\boldsymbol{\eta}}_1, \tilde{c}_1}(\xi_1, \eta_1) \in \mathbb{R}^3 \mid (\xi_1, \eta_1) \in \text{supp}(\tilde{f}_{\boldsymbol{\eta}'_1} \circ \varphi_{\tilde{\boldsymbol{\eta}}_1, \tilde{c}_1}) \right\}, \\ \bar{S}_2 &= \left\{ \varphi_{\tilde{\boldsymbol{\eta}}_2, \tilde{c}_2}(\xi_2, \eta_2) \in \mathbb{R}^3 \mid (\xi_2, \eta_2) \in \text{supp}(\tilde{g}_{\boldsymbol{\eta}'_2} \circ \varphi_{\tilde{\boldsymbol{\eta}}_2, \tilde{c}_2}) \right\}, \\ \bar{S}_3 &= \left\{ (\varphi_{\tilde{\boldsymbol{\eta}}}(\xi, \eta), \xi, \eta) \in \mathbb{R}^3 \mid \varphi_{\tilde{\boldsymbol{\eta}}}(\xi, \eta) = \xi^3 + \eta^3 + (\xi + \eta)|\tilde{\boldsymbol{\eta}}|^2 + \frac{c'_0}{N_1^3} \right\}. \end{aligned}$$

Clearly, \bar{S}_1 , \bar{S}_2 , \bar{S}_3 satisfy necessary regularity and diameter conditions to apply the nonlinear Loomis–Whitney inequality. Define $\lambda_i \in \bar{S}_i$ as

$$\lambda_1 = \varphi_{\tilde{\eta}_1, \tilde{c}_1}(\xi_1, \eta_1), \quad \lambda_2 = \varphi_{\tilde{\eta}_2, \tilde{c}_2}(\xi_2, \eta_2), \quad \lambda_3 = (\varphi_{\tilde{\eta}}(\xi, \eta), \xi, \eta),$$

then the unit normals n_i on λ_i can be described explicitly as

$$n_i(\lambda_i) = \frac{1}{\sqrt{1 + (3\xi_i^2 + |\tilde{\eta}_i|^2)^2 + (3\eta_i^2 + |\tilde{\eta}_i|^2)^2}} (-1, 3\xi_i^2 + |\tilde{\eta}_i|^2, 3\eta_i^2 + |\tilde{\eta}_i|^2),$$

for $i = 1, 2$, and the same for $n_3(\lambda_3)$. We define

$$n_i^0(\lambda_i) = \frac{1}{\sqrt{1 + 9\xi_i^4 + 9\eta_i^4}} (-1, 3\xi_i^2, 3\eta_i^2),$$

for $i = 1, 2$, and the same for $n_3^0(\lambda_3)$. Since $|\tilde{\eta}_j| \lesssim A^{-1}$, $|\tilde{\eta}| \lesssim A^{-1}$, we easily get $|n_j(\lambda_j) - n_j^0(\lambda_j)| \ll A^{-1} \leq M^{-1}$. Therefore we only need to show

$$|\det(n_1^0(\lambda_1), n_2^0(\lambda_2), n_3^0(\lambda_3))| \gtrsim M^{-1}$$

with $(\xi_1, \eta_1) + (\xi_2, \eta_2) = (\xi, \eta)$. We calculate

$$\begin{aligned} |\det(n_1^0(\lambda_1), n_2^0(\lambda_2), n_3^0(\lambda_3))| &\gtrsim \left| \det \begin{pmatrix} -1 & -1 & -1 \\ 3\xi_1^2 & 3\xi_2^2 & 3\xi^2 \\ 3\eta_1^2 & 3\eta_2^2 & 3\eta^2 \end{pmatrix} \right| \\ &\gtrsim |\xi_1\eta_2 - \xi_2\eta_1| |\xi_1\eta_2 + \xi_2\eta_1 + 2(\xi_1\eta_1 + \xi_2\eta_2)| \\ &\gtrsim M^{-1}. \end{aligned}$$

Here we used $|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \gtrsim 1$, and $(\xi_1, \eta_1) \in \text{supp}(\tilde{f}_{\eta'_1} \circ \varphi_{\tilde{\eta}_1, \tilde{c}_1})$, $(\xi_2, \eta_2) \in \text{supp}(\tilde{g}_{\eta'_2} \circ \varphi_{\tilde{\eta}_2, \tilde{c}_2})$ which implies $|\bar{F}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1}$. \square

The key ingredient to show (4.28) is the almost orthogonality of ℓ_1 and ℓ_2 which satisfy $(\ell_1, \ell_2) \in \tilde{Z}_M$. However, in [21] it was found that there exist pairs of tiles which do not satisfy the almost orthogonality. Thus we perform the decompositions which was introduced in [21], see [21, Remark 3.3] for the details.

DEFINITION 4.14 ([21, Def. 3]). — Let $\mathcal{K}_0, \mathcal{K}_1, \mathcal{K}_2, \mathcal{K}'_0, \mathcal{K}'_1, \mathcal{K}'_2 \subset \mathbb{R}^2$ and $\tilde{\mathcal{K}}_0, \tilde{\mathcal{K}}_1, \tilde{\mathcal{K}}_2, \tilde{\mathcal{K}}'_0, \tilde{\mathcal{K}}'_1, \tilde{\mathcal{K}}'_2 \subset \mathbb{R}^{d+1}$ be defined as follows:

$$\begin{aligned}\mathcal{K}_0 &= \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid \left| \eta - (\sqrt{2} - 1)^{\frac{4}{3}} \xi \right| \leq 2^{-20} N_1 \right\}, \\ \mathcal{K}_1 &= \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid \left| \eta - (\sqrt{2} + 1)^{\frac{2}{3}} (\sqrt{2} + \sqrt{3}) \xi \right| \leq 2^{-20} N_1 \right\}, \\ \mathcal{K}_2 &= \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid \left| \eta + (\sqrt{2} + 1)^{\frac{2}{3}} (\sqrt{3} - \sqrt{2}) \xi \right| \leq 2^{-20} N_1 \right\}, \\ \mathcal{K}'_0 &= \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid (\eta, \xi) \in \mathcal{K}_0 \right\}, \\ \mathcal{K}'_1 &= \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid (\eta, \xi) \in \mathcal{K}^1 \right\}, \\ \mathcal{K}'_2 &= \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid (\eta, \xi) \in \mathcal{K}^2 \right\}, \\ \tilde{\mathcal{K}}_i &= \mathbb{R} \times \mathcal{K}_i \times \mathbb{R}^{d-2}, \quad \tilde{\mathcal{K}}'_i = \mathbb{R} \times \mathcal{K}'_i \times \mathbb{R}^{d-2} \quad \text{for } i = 0, 1, 2.\end{aligned}$$

We define the subsets of $\mathbb{R}^2 \times \mathbb{R}^2$ and $\mathbb{R}^{d+1} \times \mathbb{R}^{d+1}$ as

$$\begin{aligned}\mathcal{K} &= (\mathcal{K}_0 \times (\mathcal{K}_1 \cup \mathcal{K}_2)) \cup ((\mathcal{K}_1 \cup \mathcal{K}_2) \times \mathcal{K}_0) \subset \mathbb{R}^2 \times \mathbb{R}^2, \\ \tilde{\mathcal{K}} &= (\tilde{\mathcal{K}}_0 \times (\tilde{\mathcal{K}}_1 \cup \tilde{\mathcal{K}}_2)) \cup ((\tilde{\mathcal{K}}_1 \cup \tilde{\mathcal{K}}_2) \times \tilde{\mathcal{K}}_0) \subset \mathbb{R}^{d+1} \times \mathbb{R}^{d+1}, \\ \mathcal{K}' &= (\mathcal{K}'_0 \times (\mathcal{K}'_1 \cup \mathcal{K}'_2)) \cup ((\mathcal{K}'_1 \cup \mathcal{K}'_2) \times \mathcal{K}'_0) \subset \mathbb{R}^2 \times \mathbb{R}^2, \\ \tilde{\mathcal{K}}' &= (\tilde{\mathcal{K}}'_0 \times (\tilde{\mathcal{K}}'_1 \cup \tilde{\mathcal{K}}'_2)) \cup ((\tilde{\mathcal{K}}'_1 \cup \tilde{\mathcal{K}}'_2) \times \tilde{\mathcal{K}}'_0) \subset \mathbb{R}^{d+1} \times \mathbb{R}^{d+1},\end{aligned}$$

and their complements as

$$\begin{aligned}(\mathcal{K})^c &= (\mathbb{R}^2 \times \mathbb{R}^2) \setminus \mathcal{K}, \quad (\tilde{\mathcal{K}})^c = (\mathbb{R}^{d+1} \times \mathbb{R}^{d+1}) \setminus \tilde{\mathcal{K}}, \\ (\mathcal{K}')^c &= (\mathbb{R}^2 \times \mathbb{R}^2) \setminus \mathcal{K}', \quad (\tilde{\mathcal{K}}')^c = (\mathbb{R}^{d+1} \times \mathbb{R}^{d+1}) \setminus \tilde{\mathcal{K}}'.\end{aligned}$$

Lastly, we define

$$\widehat{Z}_M = \left\{ (\ell_1, \ell_2) \in \widetilde{Z}_M \mid (\mathcal{T}_{\ell_1}^M \times \mathcal{T}_{\ell_2}^M) \cap ((\mathcal{K})^c \cap (\mathcal{K}')^c) \neq \emptyset \right\},$$

and \overline{Z}_M as the collection of $(\ell_1, \ell_2) \in \mathbb{Z}^2 \times \mathbb{Z}^2$ which satisfies

$$\begin{aligned}\mathcal{T}_{\ell_1}^M \times \mathcal{T}_{\ell_2}^M &\not\subset \bigcup_{\widehat{M} \leq M'} \bigcup_{(\ell'_1, \ell'_2) \in \widehat{Z}_M} \left(\mathcal{T}_{\ell'_1}^{M'} \times \mathcal{T}_{\ell'_2}^{M'} \right), \\ \left(\widetilde{\mathcal{T}}_{\ell_1}^M \times \widetilde{\mathcal{T}}_{\ell_2}^M \right) \cap (G_{N_1, L_1} \times G_{N_2, L_2}) \cap \mathcal{A} \cap \left((\tilde{\mathcal{K}})^c \cap (\tilde{\mathcal{K}}')^c \right) &\neq \emptyset.\end{aligned}$$

LEMMA 4.15 ([21, Lem. 3.7]). — For fixed $\ell_1 \in \mathbb{Z}^2$, the number of $\ell_2 \in \mathbb{Z}^2$ such that $(\ell_1, \ell_2) \in \widehat{Z}_M$ is finite (uniformly bounded). Furthermore, the same claim holds true if we replace \widehat{Z}_M by \overline{Z}_M .

Now we show (4.28) under the assumption $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in (\mathcal{K})^c \cap (\mathcal{K}')^c$.

PROPOSITION 4.16. — *Assume the same conditions as in Proposition 4.10. Suppose further that $|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \gtrsim 1$ and $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in (\mathcal{K})^c \cap (\mathcal{K}')^c$. Then we have*

$$(4.42) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \eta) f_{N_1, L_1}(\tau_1, \xi_1, \eta_1) g_{N_2, L_2}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2},$$

where functions f_{N_1, L_1} , g_{N_2, L_2} , h_{N_0, L_0} satisfy (4.29).

Proof. — By the definitions of \widehat{Z}_M and \overline{Z}_A , $(G_{N_1, L_1} \times G_{N_2, L_2}) \cap \mathcal{A} \cap (\widetilde{\mathcal{K}})^c \cap (\widetilde{\mathcal{K}}')^c$ are contained in

$$\bigcup_{\widehat{M} \leq M \leq A} \bigcup_{(\ell_1, \ell_2) \in \widehat{Z}_M} \left(\widetilde{\mathcal{T}}_{\ell_1}^M \times \widetilde{\mathcal{T}}_{\ell_2}^M \right) \cup \bigcup_{(\ell_1, \ell_2) \in \overline{Z}_A} \left(\widetilde{\mathcal{T}}_{\ell_1}^A \times \widetilde{\mathcal{T}}_{\ell_2}^A \right).$$

Therefore, we get

(LHS) of (4.42)

$$\leq \sum_{\widehat{M} \leq M \leq A} \sum_{(\ell_1, \ell_2) \in \widehat{Z}_M} \left| \int_* h_{N_0, L_0}(\tau, \xi, \eta) f_{N_1, L_1} |_{\widetilde{\mathcal{T}}_{\ell_1}^M}(\tau_1, \xi_1, \eta_1) \right. \\ \times g_{N_2, L_2} |_{\widetilde{\mathcal{T}}_{\ell_2}^M}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \Big| \\ + \sum_{(\ell_1, \ell_2) \in \overline{Z}_A} \left| \int_* h_{N_0, L_0}(\tau, \xi, \eta) f_{N_1, L_1} |_{\widetilde{\mathcal{T}}_{\ell_1}^A}(\tau_1, \xi_1, \eta_1) \right. \\ \times g_{N_2, L_2} |_{\widetilde{\mathcal{T}}_{\ell_2}^A}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \Big| \\ =: \sum_{\widehat{M} \leq M \leq A} \sum_{(\ell_1, \ell_2) \in \widehat{Z}_M} I_1 + \sum_{(\ell_1, \ell_2) \in \overline{Z}_A} I_2.$$

For the former term, we deduce from Proposition 4.13 and Lemma 4.15 that

$$\sum_{(\ell_1, \ell_2) \in \widehat{Z}_M} I_1 \\ \lesssim \sum_{(\ell_1, \ell_2) \in \widehat{Z}_M} A^{-\frac{d-2}{2}} M^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ \lesssim A^{-\frac{d-2}{2}} M^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}.$$

Consequently, we obtain

$$\begin{aligned} \sum_{\widehat{M} \leq M \leq A} \sum_{(\ell_1, \ell_2) \in \widehat{Z}_M} I_1 \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Next we consider the latter term. The assumption $|\eta'_j| \lesssim A^{-1} N_1$ implies that space variables of $\text{supp}(f_{N_1, L_1}|_{\widetilde{T}_{\ell_1}^A})$ and $\text{supp}(g_{N_2, L_2}|_{\widetilde{T}_{\ell_2}^A})$ are confined to regular cubes which side lengths are comparable to $A^{-1} N_1$, respectively. Since the linear transformation $(\xi_j, \eta_j) \rightarrow (\xi_j + \eta_j, \sqrt{3}(\xi_j - \eta_j))$ is invertible, Proposition 4.8 yields

$$I_2 \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}|_{\widetilde{T}_{\ell_1}^A}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{T}_{\ell_2}^A}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}.$$

Hence, by Lemma 4.15, we get

$$\begin{aligned} \sum_{(\ell_1, \ell_2) \in \overline{Z}_A} I_2 \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \sum_{(\ell_1, \ell_2) \in \overline{Z}_A} \|f_{N_1, L_1}|_{\widetilde{T}_{\ell_1}^A}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{T}_{\ell_2}^A}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

This completes the proof. \square

Next we deal with the case $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in (\mathcal{K} \cup \mathcal{K}')$. The strategy of proof is the same as for the case $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in (\mathcal{K})^c \cap (\mathcal{K}')^c$. By symmetry, it suffices to show the estimate (4.28) for the case $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in (\mathcal{K}_1 \cup \mathcal{K}_2) \times \mathcal{K}_0$.

DEFINITION 4.17 ([21, Def. 4]). — Let $m = (n, z) \in \mathbb{N} \times \mathbb{Z}$. We define the increasing sequence $\{a_{M,n}\}_{n \in \mathbb{N}}$ as

$$a_{M,1} = 0, \quad a_{M,n+1} = a_{M,n} + \frac{N_1}{\sqrt{(n+1)M}}.$$

and sets $\mathcal{R}_{M,m,1}$, $\mathcal{R}_{M,m,2}$ as follows:

$$\begin{aligned} \mathcal{R}_{M,m,1} &= \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid \begin{array}{l} a_{M,n} \leq |\eta - (\sqrt{2} + 1)^{\frac{2}{3}}(\sqrt{2} + \sqrt{3})\xi| < a_{M,n+1}, \\ zM^{-1}N_1 \leq \eta - (\sqrt{2} + 1)^{\frac{2}{3}}\xi < (z+1)M^{-1}N_1 \end{array} \right\}, \\ \mathcal{R}_{M,m,2} &= \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid \begin{array}{l} a_{M,n} \leq |\eta + (\sqrt{2} + 1)^{\frac{2}{3}}(\sqrt{3} - \sqrt{2})\xi| < a_{M,n+1}, \\ zM^{-1}N_1 \leq \eta - (\sqrt{2} + 1)^{\frac{2}{3}}\xi < (z+1)M^{-1}N_1 \end{array} \right\} \\ \widetilde{\mathcal{R}}_{M,m,1} &= \mathbb{R} \times \mathcal{R}_{M,m,1} \times \mathbb{R}^{d-2}, \quad \widetilde{\mathcal{R}}_{M,m,2} = \mathbb{R} \times \mathcal{R}_{M,m,2} \times \mathbb{R}^{d-2}. \end{aligned}$$

We will perform the Whitney type decomposition by using the above sets instead of simple square tiles. We define for $i = 1, 2$ that

$$M_{M,i}^1 = \left\{ (m, \ell) \in (\mathbb{N} \times \mathbb{Z}) \times \mathbb{Z}^2 \mid \begin{array}{l} |\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1} N_1^3 \\ \text{for any } (\xi_1, \eta_1) \in \mathcal{R}_{M,m,i} \\ \text{and } (\xi_2, \eta_2) \in \mathcal{T}_\ell^M \end{array} \right\},$$

$$M_{M,i}^2 = \left\{ (m, \ell) \in (\mathbb{N} \times \mathbb{Z}) \times \mathbb{Z}^2 \mid \begin{array}{l} |\bar{F}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1} N_1^3 \\ \text{for any } (\xi_1, \eta_1) \in \mathcal{R}_{M,m,i} \\ \text{and } (\xi_2, \eta_2) \in \mathcal{T}_\ell^M \end{array} \right\},$$

$$M_{M,i} = M_{M,i}^1 \cup M_{M,i}^2 \subset (\mathbb{N} \times \mathbb{Z}) \times \mathbb{Z}^2,$$

$$R_{M,i} = \bigcup_{(m, \ell) \in M_{M,i}} \mathcal{R}_{M,m,i} \times \mathcal{T}_\ell^M \subset \mathbb{R}^2 \times \mathbb{R}^2.$$

Furthermore, we define $M'_{M,i} \subset M_{M,i}$ as the collection of $(m, \ell) \in (\mathbb{N} \times \mathbb{Z}) \times \mathbb{Z}^2$ such that

$$\mathcal{R}_{M,m,i} \times \mathcal{T}_\ell^M \subset \bigcup_{\widehat{M} \leq M' < M} R_{M',i}.$$

By using $M'_{M,i}$, we define

$$Q_{M,i} = \begin{cases} R_{M,i} \setminus \bigcup_{(m, \ell) \in M'_{M,i}} (\mathcal{R}_{M,m,i} \times \mathcal{T}_\ell^M) & \text{for } M > \widehat{M}, \\ R_{\widehat{M},i} & \text{for } M = \widehat{M}, \end{cases}$$

and $\widetilde{M}_{M,i} = M_{M,i} \setminus M'_{M,i}$. Clearly, the followings hold.

$$\bigcup_{(m, \ell) \in \widetilde{M}_{M,i}} \mathcal{R}_{M,m,i} \times \mathcal{T}_\ell^M = Q_{M,i}, \quad \bigcup_{\widehat{M} \leq M \leq M_0} Q_{M,i} = R_{M_0,i},$$

where $M_0 \geq \widehat{M}$ is dyadic. Lastly, we define

$$\widehat{Z}_{M,i} = \{(m, \ell) \in \widetilde{M}_{M,i} \mid (\widetilde{\mathcal{R}}_{M,m,i} \times \widetilde{\mathcal{T}}_\ell^M) \cap (G_{N_1, L_1} \times G_{N_2, L_2}) \cap (\widetilde{\mathcal{K}}_i \times \widetilde{\mathcal{K}}_0) \neq \emptyset\},$$

$$\overline{Z}_{M,i} = \{(m, \ell) \in M_{M,i}^c \mid (\widetilde{\mathcal{R}}_{M,m,i} \times \widetilde{\mathcal{T}}_\ell^M) \cap (G_{N_1, L_1} \times G_{N_2, L_2}) \cap (\widetilde{\mathcal{K}}_i \times \widetilde{\mathcal{K}}_0) \neq \emptyset\},$$

where $M_{M,i}^c = (\mathbb{N} \times \mathbb{Z}) \times \mathbb{Z}^2 \setminus M_{M,i}$. We easily see that

$$(G_{N_1, L_1} \times G_{N_2, L_2}) \cup (\widetilde{\mathcal{K}}_i \times \widetilde{\mathcal{K}}_0)$$

is contained in the set

$$\bigcup_{(m, \ell) \in \widehat{Z}_{M,i}} (\widetilde{\mathcal{R}}_{M,m,i} \times \widetilde{\mathcal{T}}_\ell^M) \cup \bigcup_{(m, \ell) \in \overline{Z}_{M,i}} (\widetilde{\mathcal{R}}_{M,m,i} \times \widetilde{\mathcal{T}}_\ell^M).$$

LEMMA 4.18 (Lemma 3.9 in [21]). — Let $i = 1, 2$. For fixed $m \in \mathbb{N} \times \mathbb{Z}$, the number of $k \in \mathbb{Z}^2$ such that $(m, k) \in \widehat{Z}_{M,i}$ is finitely many. On the other hand, for fixed $k \in \mathbb{Z}^2$, the number of $m \in \mathbb{N} \times \mathbb{Z}$ such that $(m, k) \in \widehat{Z}_{M,i}$ is finitely many. Furthermore, the claim holds true whether we replace $\widehat{Z}_{M,i}$ by $\overline{Z}_{M,i}$ in the above statements.

PROPOSITION 4.19. — In addition to the hypothesis of Proposition 4.10, assume that $|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \gtrsim 1$ and $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in (\mathcal{K}_1 \cup \mathcal{K}_2) \times \mathcal{K}_0$. Then we have

$$(4.43) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2},$$

where functions f_{N_1, L_1} , g_{N_2, L_2} , h_{N_0, L_0} satisfy (4.29).

Proof. — To avoid redundancy, we only treat the case $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathcal{K}_1 \times \mathcal{K}_0$. The case $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathcal{K}_2 \times \mathcal{K}_0$ can be dealt with in the similar way. Similarly to the proof of Lemma 4.16, by the inclusion of

$$(G_{N_1, L_1} \times G_{N_2, L_2}) \cup (\widetilde{\mathcal{K}}_1 \times \widetilde{\mathcal{K}}_0)$$

in the set

$$\bigcup_{(m, \ell) \in \widehat{Z}_{M,1}} (\widetilde{\mathcal{R}}_{M,m,1} \times \widetilde{\mathcal{T}}_\ell^M) \cup \bigcup_{(m, \ell) \in \overline{Z}_{M,1}} (\widetilde{\mathcal{R}}_{M,m,1} \times \widetilde{\mathcal{T}}_\ell^M),$$

we get

$$\begin{aligned} & (\text{LHS}) \text{ of (4.43)} \\ & \leq \sum_{\widehat{M} \leq M \leq A} \sum_{(m, \ell) \in \widehat{Z}_{M,1}} \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}|_{\widetilde{\mathcal{R}}_{M,m,1}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\ & \quad \times g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_\ell^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \Big| \\ & \quad + \sum_{(m, \ell) \in \overline{Z}_{A,1}} \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}|_{\widetilde{\mathcal{R}}_{A,m,1}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\ & \quad \times g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_\ell^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \Big| \\ & =: \sum_{\widehat{M} \leq M \leq A} \sum_{(m, \ell) \in \widehat{Z}_{M,1}} I_1 + \sum_{(m, \ell) \in \overline{Z}_{A,1}} I_2. \end{aligned}$$

The former term is estimated by Proposition 4.13 and Lemma 4.18 as

$$\begin{aligned} & \sum_{(m,\ell) \in \widehat{Z}_{M,1}} I_1 \\ & \lesssim \sum_{(m,\ell) \in \widehat{Z}_{M,1}} A^{-\frac{d-2}{2}} M^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}|_{\widetilde{\mathcal{R}}_{A,m,1}}\|_{L^2} \| \\ & \quad \times g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_\ell^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ & \lesssim A^{-\frac{d-2}{2}} M^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}, \end{aligned}$$

which yields

$$\begin{aligned} & \sum_{\widehat{M} \leqslant M \leqslant A} \sum_{(m,\ell) \in \widehat{Z}_{M,1}} I_1 \\ & \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

We deal with the latter term in the same manner as that for the proof of Proposition 4.16. The assumption $|\eta'_2| \lesssim A^{-1} N_1$ means that support of $g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_\ell^A}$ is contained in a regular cube which side length is comparable to $A^{-1} N_1$. Thus, by the almost orthogonality and Proposition 4.8, we obtain

$$I_2 \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}|_{\widetilde{\mathcal{R}}_{A,m,1}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_\ell^A}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}.$$

Hence, it follows from Lemma 4.18 that

$$\begin{aligned} & \sum_{(m,\ell) \in \overline{\mathcal{Z}}_{A,1}} I_2 \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \sum_{(m,\ell) \in \overline{\mathcal{Z}}_{A,1}} \|f_{N_1, L_1}|_{\widetilde{\mathcal{R}}_{A,m,1}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_\ell^A}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ & \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

This completes the proof. \square

Next, we consider the case $|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \ll 1$. Similarly to the case $|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \gtrsim 1$, we follow the proof for the 2D Zakharov–Kuznetsov equation.

DEFINITION 4.20. — Let M be dyadic. Define

$$\begin{aligned} \Theta_k^M &= \left[\frac{\pi}{M} (k-2), \frac{\pi}{M} (k+2) \right] \cup \left[-\pi + \frac{\pi}{M} (k-2), -\pi + \frac{\pi}{M} (k+2) \right], \\ \mathfrak{D}_k^M &= \{(r \cos \theta, r \sin \theta) \in \mathbb{R}^2 \mid r \geqslant 0, \theta \in \Theta_k^M\}, \\ \widetilde{\mathfrak{D}}_k^M &= \mathbb{R} \times \mathfrak{D}_k^M \times \mathbb{R}^{d-2}. \end{aligned}$$

Let \mathcal{I} , $(\mathcal{I})^c \subset \mathbb{R}^2 \times \mathbb{R}^2$ be defined as follows:

$$\begin{aligned}\mathcal{I} &= \left(\mathfrak{D}_0^{2^{11}} \times \mathfrak{D}_0^{2^{11}} \right) \cup \left(\mathfrak{D}_{2^{10}}^{2^{11}} \times \mathfrak{D}_{2^{10}}^{2^{11}} \right), \\ \tilde{\mathcal{I}} &= \left(\tilde{\mathfrak{D}}_0^{2^{11}} \times \tilde{\mathfrak{D}}_0^{2^{11}} \right) \cup \left(\tilde{\mathfrak{D}}_{2^{10}}^{2^{11}} \times \tilde{\mathfrak{D}}_{2^{10}}^{2^{11}} \right), \\ (\mathcal{I})^c &= (\mathbb{R}^2 \times \mathbb{R}^2) \setminus \mathcal{I}, \\ (\tilde{\mathcal{I}})^c &= (\mathbb{R}^{d+1} \times \mathbb{R}^{d+1}) \setminus \tilde{\mathcal{I}}.\end{aligned}$$

Note that

$$\begin{aligned}\mathfrak{D}_0^{2^{11}} &= \left\{ (|\langle \xi, \eta \rangle| \cos \theta, |\langle \xi, \eta \rangle| \sin \theta) \in \mathbb{R}^2 \mid \min(|\theta|, |\theta - \pi|) \leq 2^{-10} \pi \right\}, \\ \mathfrak{D}_{2^{10}}^{2^{11}} &= \left\{ (|\langle \xi, \eta \rangle| \cos \theta, |\langle \xi, \eta \rangle| \sin \theta) \in \mathbb{R}^2 \mid \min\left(\left|\theta - \frac{\pi}{2}\right|, \left|\theta + \frac{\pi}{2}\right|\right) \leq 2^{-10} \pi \right\}.\end{aligned}$$

We begin with the case $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in (\mathcal{I})^c$. Note that $\max(|\xi_1 + \eta_1|, |\xi_2 + \eta_2|) \geq 2^{-5} N_1$ in *Assumption 4.1'* allows us to assume

$$(4.44) \quad (\xi_1, \eta_1) \times (\xi_2, \eta_2) \notin \left(\mathfrak{D}_{2^9 \times 3}^{2^{11}} \times \mathfrak{D}_{2^9 \times 3}^{2^{11}} \right).$$

Remark that

$$\mathfrak{D}_{2^9 \times 3}^{2^{11}} = \left\{ (|\langle \xi, \eta \rangle| \cos \theta, |\langle \xi, \eta \rangle| \sin \theta) \in \mathbb{R}^2 \mid \min\left(\left|\theta - \frac{3\pi}{4}\right|, \left|\theta + \frac{\pi}{4}\right|\right) \leq 2^{-10} \pi \right\}.$$

PROPOSITION 4.21. — Assume $|\boldsymbol{\eta}'_j| \lesssim A^{-1} N_1$, (4.44) and (4.29). Let A , M be dyadic which satisfy $1 \ll M \leq A$ and (k_1, k_2) satisfies $\mathfrak{D}_{k_1}^M \times \mathfrak{D}_{k_2}^M \subset (\mathcal{I})^c$, $16 \leq |k_1 - k_2| \leq 32$. Then we get

$$\begin{aligned}(4.45) \quad & \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim A^{-\frac{d-2}{2}} M^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2},\end{aligned}$$

where $d\sigma_j = d\tau_j d\xi_j d\boldsymbol{\eta}_j$ and $*$ denotes $(\tau, \xi, \boldsymbol{\eta}) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2)$.

Proof. — It suffices to show

$$\begin{aligned}(4.46) \quad & \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\hat{\sigma}_1 d\hat{\sigma}_2 \right| \\ & \lesssim M^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\boldsymbol{\eta}'_1)\|_{L^2_{\tau \xi \eta}} \| \\ & \quad \times g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\boldsymbol{\eta}'_2)\|_{L^2_{\tau \xi \eta}} \|h_{N_0, L_0}(\boldsymbol{\eta}')\|_{L^2_{\tau \xi \eta}},\end{aligned}$$

where $d\hat{\sigma}_j = d\tau_j d\xi_j d\eta_j$ and $\hat{*}$ denotes $(\tau, \xi, \eta) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2)$. As we saw in the proof of Proposition 4.13, since $M \leq A$ and $|\boldsymbol{\eta}'_j| \lesssim A^{-1} N_1$, we can show (4.46) by following the proof of Proposition 3.14 in [21]. We omit the proof. \square

PROPOSITION 4.22. — *In addition to the hypothesis of Proposition 4.10, assume that $|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \ll 1$, (4.44) and $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in (\mathcal{I})^c$. Then we have*

$$(4.47) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2},$$

where functions f_{N_1, L_1} , g_{N_2, L_2} , h_{N_0, L_0} satisfy (4.29).

Proof. — We define that

$$J_M^{(\mathcal{I})^c} = \left\{ (k_1, k_2) \middle| \begin{array}{l} 0 \leq k_1, k_2 \leq M-1, \\ (\mathfrak{D}_{k_1}^M \times \mathfrak{D}_{k_2}^M) \subset (\mathcal{I})^c \cap (\mathfrak{D}_{2^9 \times 3}^{2^{11}} \times \mathfrak{D}_{2^9 \times 3}^{2^{11}})^c \end{array} \right\}.$$

We perform the Whitney type decomposition as

$$(\mathcal{I})^c \cap (\mathfrak{D}_{2^9 \times 3}^{2^{11}} \times \mathfrak{D}_{2^9 \times 3}^{2^{11}})^c \\ = \bigcup_{64 \leq M \leq A} \bigcup_{\substack{(k_1, k_2) \in J_M^{(\mathcal{I})^c} \\ 16 \leq |k_1 - k_2| \leq 32}} \mathfrak{D}_{k_1}^M \times \mathfrak{D}_{k_2}^M \cup \bigcup_{\substack{(k_1, k_2) \in J_A^{(\mathcal{I})^c} \\ |k_1 - k_2| \leq 16}} \mathfrak{D}_{k_1}^A \times \mathfrak{D}_{k_2}^A.$$

Note that $|\sin \angle((\xi_1, \eta_1), (\xi_2, \eta_2))| \ll 1$ implies $M \gg 1$. We observe

$$\text{(LHS) of (4.47)} \\ \leq \sum_{1 \ll M \leq A} \sum_{\substack{(k_1, k_2) \in J_M^{(\mathcal{I})^c} \\ 16 \leq |k_1 - k_2| \leq 32}} \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\ \times g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \Big| \\ + \sum_{\substack{(k_1, k_2) \in J_A^{(\mathcal{I})^c} \\ |k_1 - k_2| \leq 16}} \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\ \times g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \Big| \\ := \sum_{1 \ll M \leq A} \sum_{\substack{(k_1, k_2) \in J_M^{(\mathcal{I})^c} \\ 16 \leq |k_1 - k_2| \leq 32}} I_1 + \sum_{\substack{(k_1, k_2) \in J_A^{(\mathcal{I})^c} \\ |k_1 - k_2| \leq 16}} I_2.$$

The former term is dealt with by Proposition 4.21 as follows.

$$\begin{aligned} & \sum_{1 \ll M \leq A} \sum_{\substack{(k_1, k_2) \in J_M^{(\mathcal{I})^c} \\ 16 \leq |k_1 - k_2| \leq 32}} I_1 \\ & \lesssim \sum_{1 \ll M \leq A} A^{-\frac{d-2}{2}} M^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ & \lesssim A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

For the latter term, we only consider the case $|(\xi, \eta)| \gg A^{-1} N_1$. The case $|(\xi, \eta)| \lesssim A^{-1} N_1$ can be treated by Proposition 4.8. By Lemma 3.12 in [21] and $|\eta'_j| \lesssim A^{-1} N_1$, we easily observe that $|(\xi, \eta)| \gg A^{-1} N_1$ gives $|\Phi(\xi_1, \eta_1, \xi_2, \eta_2)| \gtrsim A^{-1} N_1^3$. Thus, it suffices to show the following bilinear estimates.

$$\begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^A} (\tau_1, \xi_1, \eta_1) g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^A} (\tau - \tau_1, \xi - \xi_1, \eta - \eta_1) d\sigma_1 \right\|_{L^2} \\ & \quad \lesssim A^{-\frac{d-2}{2}} N_1^{\frac{d-3}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{\widetilde{\mathfrak{D}}_{k_1}^A} \|g_{N_2, L_2}\|_{\widetilde{\mathfrak{D}}_{k_2}^A} \|_{L^2}, \\ & \left\| \mathbf{1}_{G_{N_1, L_1} \cap \widetilde{\mathfrak{D}}_{k_1}^A} \int g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^A} (\tau_2, \xi_2, \eta_2) h_{N_0, L_0} (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2) d\sigma_2 \right\|_{L^2} \\ & \quad \lesssim A^{-\frac{d-1}{2}} N_1^{\frac{d-3}{2}} (L_0 L_2)^{\frac{1}{2}} \|g_{N_2, L_2}\|_{\widetilde{\mathfrak{D}}_{k_2}^A} \|h_{N_0, L_0}\|_{L^2}, \\ & \left\| \mathbf{1}_{G_{N_2, L_2} \cap \widetilde{\mathfrak{D}}_{k_2}^A} \int h_{N_0, L_0} (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2) f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^A} (\tau_1, \xi_1, \eta_1) d\sigma_1 \right\|_{L^2} \\ & \quad \lesssim A^{-\frac{d-1}{2}} N_1^{\frac{d-3}{2}} (L_0 L_1)^{\frac{1}{2}} \|h_{N_0, L_0}\|_{L^2} \|f_{N_1, L_1}\|_{\widetilde{\mathfrak{D}}_{k_1}^A} \|_{L^2}, \end{aligned}$$

that are verified by showing

$$\begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^A} (\tau_1, \xi_1, \eta_1) g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^A} (\tau - \tau_1, \xi - \xi_1, \eta - \eta_1) d\widehat{\sigma}_1 \right\|_{L^2} \\ & \quad \lesssim N_1^{-\frac{1}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{\widetilde{\mathfrak{D}}_{k_1}^A} (\eta'_1) \|_{L^2} \|g_{N_2, L_2}\|_{\widetilde{\mathfrak{D}}_{k_2}^A} (\eta'_2) \|_{L^2}, \\ & \left\| \mathbf{1}_{G_{N_1, L_1} \cap \widetilde{\mathfrak{D}}_{k_1}^A} \int g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^A} (\tau_2, \xi_2, \eta_2) h_{N_0, L_0} (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2) d\widehat{\sigma}_2 \right\|_{L^2} \\ & \quad \lesssim (AN_1)^{-\frac{1}{2}} (L_0 L_2)^{\frac{1}{2}} \|g_{N_2, L_2}\|_{\widetilde{\mathfrak{D}}_{k_2}^A} (\eta'_2) \|_{L^2} \|h_{N_0, L_0} (\eta')\|_{L^2}, \\ & \left\| \mathbf{1}_{G_{N_2, L_2} \cap \widetilde{\mathfrak{D}}_{k_2}^A} \int h_{N_0, L_0} (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2) f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^A} (\tau_1, \xi_1, \eta_1) d\widehat{\sigma}_1 \right\|_{L^2} \\ & \quad \lesssim (AN_1)^{-\frac{1}{2}} (L_0 L_1)^{\frac{1}{2}} \|h_{N_0, L_0} (\eta')\|_{L^2} \|f_{N_1, L_1}\|_{\widetilde{\mathfrak{D}}_{k_1}^A} (\eta'_1) \|_{L^2}, \end{aligned}$$

respectively. These estimates are established in the same manner as for Proposition 3.13 in [21]. We omit the details. \square

Next we treat the case $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathcal{I}$. By symmetry, we may assume $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathfrak{D}_0^{2^{11}} \times \mathfrak{D}_0^{2^{11}}$ and show the following.

PROPOSITION 4.23. — *Under the hypothesis of Proposition 4.10, we have*

$$(4.48) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_0^{2^{11}}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_0^{2^{11}}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim N_0^{\frac{d-4}{2} + 2\varepsilon} N_1^{-1 - \frac{3}{2}\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_0^{2^{11}}} \|_{L^2} \|g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_0^{2^{11}}} \|_{L^2} \|h_{N_0, L_0} \|_{L^2},$$

where functions f_{N_1, L_1} , g_{N_2, L_2} , h_{N_0, L_0} satisfy (4.29).

We note that the proof is almost the same as that for Proposition 3.18 in [21]. Therefore, we only give a sketch of the proof here.

DEFINITION 4.24. — *Let $M \gg 1$ and K be dyadic which satisfy $2^{10} \leq K \leq 2^{-10}M$. We define that*

$$\mathfrak{K}_M^K = \left\{ k \in \mathbb{N} \mid \frac{M}{K} \leq k \leq 2\frac{M}{K}, \quad M - 2\frac{M}{K} \leq k \leq M - \frac{M}{K} \right\}, \\ \mathfrak{K}_M = \left\{ k \in \mathbb{N} \mid 0 \leq k \leq 2^{10}, \quad M - 2^{10} \leq k \leq M - 1 \right\}.$$

The following proposition corresponds to Proposition 3.19 in [21].

PROPOSITION 4.25. — *Suppose that $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$ and functions f_{N_1, L_1} , g_{N_2, L_2} , h_{N_0, L_0} satisfy (4.29). Let M be dyadic such that $1 \ll M \leq A$, $|k_1 - k_2| \leq 32$ and*

$$(\mathfrak{D}_{k_1}^M \times \mathfrak{D}_{k_2}^M) \subset \mathcal{I}.$$

Then we have

$$(4.49) \quad \left\| \mathbf{1}_{G_{N_1, L_1} \cap \widetilde{\mathfrak{D}}_{k_1}^M} \int g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) h_{N_0, L_0}(\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) d\sigma_2 \right\|_{L^2} \\ \lesssim A^{-\frac{d-2}{2}} M^{-\frac{1}{2}} N_1^{\frac{d-3}{2}} (L_0 L_2)^{\frac{1}{2}} \|g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^M} \|_{L^2} \|h_{N_0, L_0} \|_{L^2},$$

$$(4.50) \quad \left\| \mathbf{1}_{G_{N_2, L_2} \cap \widetilde{\mathfrak{D}}_{k_2}^M} \int h_{N_0, L_0}(\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ \lesssim A^{-\frac{d-2}{2}} M^{-\frac{1}{2}} N_1^{\frac{d-3}{2}} (L_0 L_1)^{\frac{1}{2}} \|h_{N_0, L_0} \|_{L^2} \|f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^M} \|_{L^2}.$$

In addition to the above assumptions,

(1) assume $N_0 \gg M^{-1}N_1$, then we have

$$(4.51) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ & \lesssim A^{-\frac{d-2}{2}} (MN_0)^{-\frac{1}{2}} N_1^{\frac{d-2}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}\|_{L^2}. \end{aligned}$$

(2) assume $k_1 \in \mathfrak{K}_M^K$, then we have

$$(4.52) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ & \lesssim A^{-\frac{d-2}{2}} K^{\frac{1}{4}} N_1^{\frac{d-3}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}\|_{L^2}. \end{aligned}$$

(3) assume $M \ll A$, $k_1 \in \mathfrak{K}_M$ and either $16 \leq |k_1 - k_2| \leq 32$ or $|\xi| \geq M^{-3/2}N_1$, then we have

$$(4.53) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ & \lesssim A^{-\frac{d-2}{2}} M^{\frac{1}{4}} N_1^{\frac{d-3}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}\|_{L^2}. \end{aligned}$$

Proof. — (4.49) and (4.50) are given by

$$\begin{aligned} & \left\| \mathbf{1}_{G_{N_1, L_1} \cap \tilde{\mathfrak{D}}_{k_1}^M} \int g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) h_{N_0, L_0}(\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) d\hat{\sigma}_2 \right\|_{L^2} \\ & \lesssim (MN_1)^{-\frac{1}{2}} (L_0 L_2)^{\frac{1}{2}} \|g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\boldsymbol{\eta}'_2)\|_{L^2} \|h_{N_0, L_0}(\boldsymbol{\eta}')\|_{L^2}, \\ & \left\| \mathbf{1}_{G_{N_2, L_2} \cap \tilde{\mathfrak{D}}_{k_2}^M} \int h_{N_0, L_0}(\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) d\hat{\sigma}_1 \right\|_{L^2} \\ & \lesssim (MN_1)^{-\frac{1}{2}} (L_0 L_1)^{\frac{1}{2}} \|h_{N_0, L_0}(\boldsymbol{\eta}')\|_{L^2} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\boldsymbol{\eta}'_1)\|_{L^2}, \end{aligned}$$

respectively. These estimates are obtained in the same manner as for (4.33) and (4.34) in Proposition 4.13, respectively. We omit the proof.

Next we consider (4.51). We will show

$$(4.54) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\hat{\sigma}_1 \right\|_{L^2} \\ & \lesssim (MN_0)^{-\frac{1}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\boldsymbol{\eta}'_1)\|_{L^2} \|g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\boldsymbol{\eta}'_2)\|_{L^2}. \end{aligned}$$

We write $(\xi_1, \eta_1) = r_1(\cos \theta_1, \sin \theta_1)$, $(\xi - \xi_1, \eta - \eta_1) = r_2(\cos \theta_2, \sin \theta_2)$. Similarly to the proof of (4.32), it suffices to show

$$(4.55) \quad |\partial_{r_1}(\xi \xi_1(\xi - \xi_1) + \eta \eta_1(\eta - \eta_1))| \gtrsim N_0 N_1.$$

We may assume $|(\xi, \eta)| \geq N_0/2 \gg M^{-1} N_1$. By the assumption $|k_1 - k_2| \leq 32$, we easily confirm that $|\eta| \leq 2|\xi| \sim N_0$ which implies (4.55).

Lastly, we consider (4.52) and (4.53). It suffices to show

$$(4.56) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\hat{\sigma}_1 \right\|_{L^2} \\ & \lesssim N_1^{-\frac{1}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\boldsymbol{\eta}'_1)\|_{L^2} \|g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\boldsymbol{\eta}'_2)\|_{L^2}, \end{aligned}$$

for $k_1 \in \mathfrak{K}_M^K$, $|k_1 - k_2| \leq 32$ and

$$(4.57) \quad \begin{aligned} & \left\| \mathbf{1}_{G_{N_0, L_0}} \int f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\hat{\sigma}_1 \right\|_{L^2} \\ & \lesssim M^{\frac{1}{4}} N_1^{-\frac{1}{2}} (L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\boldsymbol{\eta}'_1)\|_{L^2} \|g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\boldsymbol{\eta}'_2)\|_{L^2}, \end{aligned}$$

for $k_1 \in \mathfrak{K}_M$ and either $16 \leq |k_1 - k_2| \leq 32$ or $|\xi| \geq M^{-3/2} N_1$. (4.56) and (4.57) are established in the same way as that for (3.67) and (3.68) in Proposition 3.19 in [21]. Thus, here we only confirm that it holds

$$(4.58) \quad \left\| |\nabla_x|^{\frac{1}{2p}} |\nabla_y|^{\frac{1}{2p}} u_{N,L} \right\|_{L_t^p L_{xy}^q} \lesssim L^{\frac{1}{2}} \|u_{N,L}\|_{L_{x,y,t}^2}, \quad \text{if } \frac{2}{p} + \frac{2}{q} = 1, p > 2,$$

where $\text{supp } \widehat{u}_{N,L} \subset G_{N,L}$. Let $c \in \mathbb{R}$. (4.58) is given by the Strichartz estimates.

$$(4.59) \quad \left\| |\nabla_x|^{\frac{1}{2p}} |\nabla_y|^{\frac{1}{2p}} e^{-t(\partial_x^3 + \partial_y^3 - (\partial_x + \partial_y)c)} \varphi \right\|_{L_t^p L_{xy}^q} \lesssim \|\varphi\|_{L_{x,y}^2},$$

if $2/p + 2/q = 1$, $p > 2$. We can establish (4.59) by applying Theorem 3.1 in [18]. Employing (4.58) with $p = q = 4$, we can show (4.56) and (4.57) by the same argument as that for (3.67) and (3.68) in Proposition 3.19 in [21], respectively. \square

First we consider the case $k_1 \in \mathfrak{K}_M^K$.

PROPOSITION 4.26. — *Let M be dyadic such that $1 \ll M \leq A$. Assume that $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$, $N_0 \gg M^{-1}N_1$, $k_1 \in \mathfrak{K}_M^K$, k_2 satisfy $|k_1 - k_2| \leq 32$. Then we have*

$$(4.60) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim A^{-\frac{d-2}{2}} N_0^{-\frac{1}{2}} N_1^{\frac{d-5}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2},$$

where functions f_{N_1, L_1} , g_{N_2, L_2} , h_{N_0, L_0} satisfy (4.29).

Proof. — We easily confirm that $|\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| \gtrsim N_0 N_1^2$ holds. This and Proposition 4.25 immediately yield (4.60). \square

PROPOSITION 4.27. — *Let M be dyadic such that $1 \ll M \leq A$. Assume that $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$, $N_0 \sim M^{-1}N_1$, $k_1 \in \mathfrak{K}_M^K$, k_2 satisfy $16 \leq |k_1 - k_2| \leq 32$. Then we have*

$$\left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim A^{-\frac{d-2}{2}} K^{\frac{1}{2}} N_0^{-\frac{1}{2}} N_1^{\frac{d-5}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ \times \|f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2},$$

where functions f_{N_1, L_1} , g_{N_2, L_2} , h_{N_0, L_0} satisfy (4.29).

Proof. — It suffices to show

$$(4.61) \quad \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\hat{\sigma}_1 d\hat{\sigma}_2 \right| \\ \lesssim (MK)^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \\ \times \|f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^M}(\boldsymbol{\eta}'_1)\|_{L^2_{\tau \xi \eta}} \|g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^M}(\boldsymbol{\eta}'_2)\|_{L^2_{\tau \xi \eta}} \|h_{N_0, L_0}(\boldsymbol{\eta}')\|_{L^2_{\tau \xi \eta}},$$

for fixed $\boldsymbol{\eta}'_1$, $\boldsymbol{\eta}'_2$. By using Proposition 4.25 and smallness of $|\boldsymbol{\eta}'_1|$ and $|\boldsymbol{\eta}'_2|$, (4.61) can be obtained in the same way as for Proposition 3.20 in [21]. We omit the details. \square

Next we deal with the case $(k_1, k_2) \in \mathfrak{K}_M \times \mathfrak{K}_M$.

DEFINITION 4.28. — *Let M and ν be dyadic such that $1 \ll M \leq A^{2/3}$, $2 \leq \nu \leq AM^{-3/2}$ and $m = (m_{(1)}, m_{(2)}) \in \mathbb{Z}^2$. We define rectangle-tiles $\{\mathcal{T}_m^{M, \nu}\}_{m \in \mathbb{Z}^2}$ whose short side is parallel to ξ -axis and its length is*

$M^{-3/2}\nu^{-1}N_1$, and long side length is $M^{-1}\nu^{-1}N_1$ by

$$\mathcal{T}_m^{M,\nu} := \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid \begin{array}{l} \xi \in M^{-\frac{3}{2}}\nu^{-1}N_1[m_{(1)}, m_{(1)} + 1), \\ \eta \in M^{-1}\nu^{-1}N_1[m_{(2)}, m_{(2)} + 1) \end{array} \right\}$$

and and prisms $\{\tilde{\mathcal{T}}_m^{M,\nu}\}_{m \in \mathbb{Z}^2}$ by $\tilde{\mathcal{T}}_m^{M,\nu} := \mathbb{R} \times \mathcal{T}_m^{M,\nu} \times \mathbb{R}^{d-2}$. Recall that

$$\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2) = \xi_1\xi_2(\xi_1 + \xi_2) + \eta_1\eta_2(\eta_1 + \eta_2),$$

$$\bar{F}(\xi_1, \eta_1, \xi_2, \eta_2) = \xi_1\eta_2 + \xi_2\eta_1 + 2(\xi_1\eta_1 + \xi_2\eta_2).$$

Let $\mathbf{k} := (k_1, k_2) \in \mathfrak{K}_M \times \mathfrak{K}_M$. We define $Z_{M,\nu,\mathbf{k}}^1$ as the set of $(m_1, m_2) \in \mathbb{Z}^2 \times \mathbb{Z}^2$ such that

$$\begin{cases} |\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-\frac{3}{2}}\nu^{-1}N_1^3 & \text{for any } (\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu}, \\ (\mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu}) \cap (\mathfrak{D}_{k_1}^M \times \mathfrak{D}_{k_2}^M) \neq \emptyset, \\ |\xi_1 + \xi_2| \lesssim M^{-3/2}N_1 & \text{for any } (\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu}. \end{cases}$$

Similarly, we define $Z_{M,\nu,\mathbf{k}}^2$ as the set of $(m_1, m_2) \in \mathbb{Z}^2 \times \mathbb{Z}^2$ such that

$$\begin{cases} |\bar{F}(\xi_1, \eta_1, \xi_2, \eta_2)| \geq M^{-1}\nu^{-1}N_1^2 & \text{for any } (\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu}, \\ (\mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu}) \cap (\mathfrak{D}_{k_1}^M \times \mathfrak{D}_{k_2}^M) \neq \emptyset, \\ |\xi_1 + \xi_2| \lesssim M^{-3/2}N_1 & \text{for any } (\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu}, \end{cases}$$

and

$$Z_{M,\nu}^{\mathbf{k}} = Z_{M,\nu,\mathbf{k}}^1 \cup Z_{M,\nu,\mathbf{k}}^2, \quad R_{M,\nu}^{\mathbf{k}} = \bigcup_{(m_1, m_2) \in Z_{M,\nu}^{\mathbf{k}}} \mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu} \subset \mathbb{R}^2 \times \mathbb{R}^2.$$

It is clear that $\nu_1 \leq \nu_2 \implies R_{M,\nu_1}^{\mathbf{k}} \subset R_{M,\nu_2}^{\mathbf{k}}$. Further, we define

$$Q_{M,\nu}^{\mathbf{k}} = \begin{cases} R_{M,\nu}^{\mathbf{k}} \setminus R_{M,\nu/2}^{\mathbf{k}} & \text{for } \nu > 2, \\ R_{M,2}^{\mathbf{k}} & \text{for } \nu = 2. \end{cases}$$

and a set of pairs of integer pair $\hat{Z}_{M,\nu}^{\mathbf{k}} \subset Z_{M,\nu}^{\mathbf{k}}$ as

$$\bigcup_{(m_1, m_2) \in \hat{Z}_{M,\nu}^{\mathbf{k}}} \mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu} = Q_{M,\nu}^{\mathbf{k}}.$$

Clearly, $\hat{Z}_{M,\nu}^{\mathbf{k}}$ is uniquely defined and

$$\nu_1 \neq \nu_2 \implies Q_{M,\nu_1}^{\mathbf{k}} \cap Q_{M,\nu_2}^{\mathbf{k}} = \emptyset, \quad \bigcup_{2 \leq \nu \leq \nu_0} Q_{M,\nu}^{\mathbf{k}} = R_{M,\nu_0}^{\mathbf{k}}$$

where $\nu_0 \geq 2$ is dyadic. Lastly, we define $\bar{Z}_{M,\nu}^{\mathbf{k}}$ as the collection of $(m_1, m_2) \in \mathbb{Z}^2 \times \mathbb{Z}^2$ which satisfies

$$\begin{cases} \mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu} \not\subset \bigcup_{2 \leq \nu' \leq \nu} \bigcup_{(m'_1, m'_2) \in \widehat{Z}_{M,\nu'}^{\mathbf{k}}} \left(\mathcal{T}_{m'_1}^{M,d'} \times \mathcal{T}_{m'_2}^{M,d'} \right), \\ (\mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu}) \cap (\mathfrak{D}_{k_1}^M \times \mathfrak{D}_{k_2}^M) \neq \emptyset, \\ |\xi_1 + \xi_2| \lesssim M^{-3/2} N_1 \text{ for any } (\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathcal{T}_{m_1}^{M,\nu} \times \mathcal{T}_{m_2}^{M,\nu}. \end{cases}$$

PROPOSITION 4.29. — Assume $|\boldsymbol{\eta}'_j| \lesssim A^{-1} N_1$ and (4.29). Let M and ν be dyadic such that $1 \ll M \leq A^{2/3}$, $2 \leq \nu \leq AM^{-3/2}$, $16 \leq |k_1 - k_2| \leq 32$ and $(m_1, m_2) \in \widehat{Z}_{M,\nu}^{\mathbf{k}}$. Then we get

$$\begin{aligned} (4.62) \quad & \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\widetilde{\mathcal{T}}_{m_1}^{M,\nu}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\widetilde{\mathcal{T}}_{m_2}^{M,\nu}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim A^{-\frac{d-2}{2}} M \nu^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\widetilde{\mathcal{T}}_{m_1}^{M,\nu}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_{m_2}^{M,\nu}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}, \end{aligned}$$

where $d\sigma_j = d\tau_j d\xi_j d\boldsymbol{\eta}_j$ and $*$ denotes $(\tau, \xi, \boldsymbol{\eta}) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2)$.

Proof. — It suffices to show the following inequality.

$$\begin{aligned} (4.63) \quad & \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\widetilde{\mathcal{T}}_{m_1}^{M,\nu}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\widetilde{\mathcal{T}}_{m_2}^{M,\nu}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\widehat{\sigma}_1 d\widehat{\sigma}_2 \right| \\ & \lesssim M \nu^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\widetilde{\mathcal{T}}_{m_1}^{M,\nu}}(\boldsymbol{\eta}'_1)\|_{L^2_{\tau \xi \eta}} \|g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_{m_2}^{M,\nu}}(\boldsymbol{\eta}'_2)\|_{L^2_{\tau \xi \eta}} \|h_{N_0, L_0}(\boldsymbol{\eta}')\|_{L^2_{\tau \xi \eta}}. \end{aligned}$$

Similarly to the proof of Proposition 4.13, since $|\boldsymbol{\eta}'_j| \lesssim A^{-1} N_1$, we can reuse the proofs of Propositions 3.22 and 3.23 in [21] with slight modifications to get the estimate (4.63). We omit the proof. \square

LEMMA 4.30. — Let M and d be dyadic such that $1 \ll M \leq A^{2/3}$, $2 \leq \nu \leq AM^{-3/2}$ and $k_1, k_2 \in \mathfrak{K}_M$. For fixed $m_1 \in \mathbb{Z}^2$, the number of $m_2 \in \mathbb{Z}^2$ such that $(m_1, m_2) \in \widehat{Z}_{M,\nu}^{\mathbf{k}}$ is finitely many. Furthermore, the same claim holds true if we replace $\widehat{Z}_{M,\nu}^{\mathbf{k}}$ by $\bar{Z}_{M,\nu}^{\mathbf{k}}$.

PROPOSITION 4.31. — Let $1 \ll M \leq A^{2/3}$. Assume that $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$ and $k_1, k_2 \in \mathfrak{K}_M$ satisfy $16 \leq |k_1 - k_2| \leq 32$. Then we have

$$(4.64) \quad \begin{aligned} & \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim A^{-\frac{d-3}{2}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\tilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2}|_{\tilde{\mathfrak{D}}_{k_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}, \end{aligned}$$

where functions $f_{N_1, L_1}, g_{N_2, L_2}, h_{N_0, L_0}$ satisfy (4.29).

Proof. — First we consider $|\xi_1 + \xi_2| \gg M^{-3/2}N_1$. It is easily observed that $|\xi_1 + \xi_2| \gg M^{-3/2}N_1$ means $|\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| \gg M^{-3/2}N_1^2$. Then, by using (4.53) in Proposition 4.25, we obtain

$$\begin{aligned} & \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \boldsymbol{\eta}_1) g_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim A^{-\frac{d-2}{2}} M^{\frac{1}{4}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\tilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2}|_{\tilde{\mathfrak{D}}_{k_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}, \end{aligned}$$

which completes the proof for the case $|\xi_1 + \xi_2| \gg M^{-3/2}N_1$.

We consider the case $|\xi_1 + \xi_2| \lesssim M^{-3/2}N_1$. For simplicity, we assume $\text{supp } f_{N_1, L_1} \subset \tilde{\mathfrak{D}}_{k_1}^M$ and $\text{supp } g_{N_2, L_2} \subset \tilde{\mathfrak{D}}_{k_2}^M$, and use

$$\begin{aligned} I_{M, \nu}^{m_1, m_2} := & \left| \int_* h_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) f_{N_1, L_1} |_{\tilde{\mathcal{T}}_{m_1}^{M, \nu}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\ & \left. \times g_{N_2, L_2} |_{\tilde{\mathcal{T}}_{m_2}^{M, \nu}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \end{aligned}$$

Let ν_0 denote the maximal dyadic number which satisfies $\nu_0 \leq AM^{-3/2}$. By the definition of $\widehat{Z}_{M, \nu}^{\mathbf{k}}$ and $\overline{Z}_{M, \nu}^{\mathbf{k}}$, we observe that

$$(\text{LHS}) \text{ of (4.64)} \leq \sum_{2 \leq \nu \leq \nu_0} \sum_{(m_1, m_2) \in \widehat{Z}_{M, \nu}^{\mathbf{k}}} I_{M, \nu}^{m_1, m_2} + \sum_{(m_1, m_2) \in \overline{Z}_{M, \nu_0}^{\mathbf{k}}} I_{M, \nu_0}^{m_1, m_2}.$$

It follows from Proposition 4.29 and Lemma 4.30 that

$$\begin{aligned} & \sum_{(m_1, m_2) \in \widehat{Z}_{M, \nu}^k} I_{M, \nu}^{m_1, m_2} \\ & \lesssim A^{-\frac{d-2}{2}} M \nu^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \sum_{(m_1, m_2) \in \widehat{Z}_{M, \nu}^k} \|f_{N_1, L_1}|_{\widetilde{\mathcal{T}}_{m_1}^{M, \nu}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_{m_2}^{M, \nu}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ & \lesssim A^{-\frac{d-2}{2}} M \nu^{\frac{1}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}, \end{aligned}$$

which gives

$$\begin{aligned} & \sum_{2 \leq \nu \leq \nu_0} \sum_{(m_1, m_2) \in \widehat{Z}_{M, \nu}^k} I_{M, \nu}^{m_1, m_2} \\ & \lesssim A^{-\frac{d-3}{2}} M^{\frac{1}{4}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}|_{\widetilde{\mathcal{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathcal{D}}_{k_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Since $|\xi_1 + \xi_2| \lesssim M^{-3/2} N_1$, we can assume $N_0 \sim M^{-1} N_1$. Then this completes the desired estimate for the first term. For the second term, we first note that $\mathcal{T}_m^{M, \nu_0} \subset \mathbb{R}^2$ is a rectangle set whose short-side length is $\sim A^{-1}$ and long-side length is $\sim A^{-1} M^{1/2}$. Then we can decompose \mathcal{T}_m^{M, ν_0} into $\sim M^{1/2}$ number of square tiles whose side length is A^{-1} . Thus, by Proposition 4.8 and the almost orthogonality, we observe

$$\begin{aligned} I_{M, \nu_0}^{m_1, m_2} & \lesssim A^{-\frac{d-3}{2}} M^{\frac{1}{4}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\widetilde{\mathcal{T}}_{m_1}^{M, \nu_0}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathcal{T}}_{m_2}^{M, \nu_0}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Consequently, by Lemma 4.30, we obtain

$$\begin{aligned} & \sum_{(m_1, m_2) \in \widehat{Z}_{M, \nu_0}^k} I_{M, \nu_0}^{m_1, m_2} \\ & \lesssim A^{-\frac{d-3}{2}} M^{\frac{1}{4}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}|_{\widetilde{\mathcal{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathcal{D}}_{k_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}, \end{aligned}$$

which completes the proof. \square

Proof of Proposition 4.23. — We should recall that we can assume $A^{-1} N_1 \lesssim N_0$. Let M be dyadic such that $1 \ll M \leq A^{2/3}$ and M_0 be the maximal dyadic number which satisfies $M_0 \leq A^{2/3}$. We define

$$K_M^{\mathcal{I}} = \{(k_1, k_2) \mid 0 \leq k_1, k_2 \leq M-1, (\mathcal{D}_{k_1}^M \times \mathcal{D}_{k_2}^M) \subset (\mathcal{D}_0^{2^{11}} \times \mathcal{D}_0^{2^{11}})\}.$$

It is observed that

$$\mathfrak{D}_0^{2^{11}} \times \mathfrak{D}_0^{2^{11}} = \bigcup_{1 \ll M \leqslant M_0} \bigcup_{\substack{(k_1, k_2) \in K_M^{\mathcal{I}} \\ 16 \leqslant |k_1 - k_2| \leqslant 32}} \mathfrak{D}_{k_1}^M \times \mathfrak{D}_{k_2}^M \cup \bigcup_{\substack{(k_1, k_2) \in K_{M_0}^{\mathcal{I}} \\ |k_1 - k_2| \leqslant 16}} \mathfrak{D}_{k_1}^{M_0} \times \mathfrak{D}_{k_2}^{M_0}.$$

Let us write

$$I_M^{k_1, k_2} := \left| \int_* h_{N_0, L_0}(\tau, \xi, \eta) f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^M}(\tau_1, \xi_1, \eta_1) g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^M}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right|.$$

We calculate that

$$\begin{aligned} & \left| \int_* h_{N_0, L_0}(\tau, \xi, \eta) f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_0^{2^{11}}}(\tau_1, \xi_1, \eta_1) g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_0^{2^{11}}}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim \sum_{1 \ll M \leqslant M_0} \sum_{\substack{(k_1, k_2) \in K_M^{\mathcal{I}} \\ 16 \leqslant |k_1 - k_2| \leqslant 32}} I_M^{k_1, k_2} + \sum_{\substack{(k_1, k_2) \in K_{M_0}^{\mathcal{I}} \\ |k_1 - k_2| \leqslant 16}} I_{M_0}^{k_1, k_2}. \end{aligned}$$

We consider the former term. Since $M \leqslant A^{2/3}$ and $16 \leqslant |k_1 - k_2|$ we may assume $A^{-2/3} N_1 \lesssim N_0$. This and $2^{10} K \leqslant M$ mean

$$A^{-(d-2)/2} K^{1/2} N_0^{-1/2} N_1^{(d-5)/2} \lesssim A^{-(d-3)/2} N_0^{-1/4} N_1^{(2d-11)/4}.$$

Then by using Propositions 4.26, 4.27 and 4.31, we obtain

$$\begin{aligned} & \sum_{\substack{(k_1, k_2) \in K_M^{\mathcal{I}} \\ 16 \leqslant |k_1 - k_2| \leqslant 32}} I_M^{k_1, k_2} \\ & \lesssim A^{-\frac{d-3}{2}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \sum_{\substack{(k_1, k_2) \in K_M^{\mathcal{I}} \\ 16 \leqslant |k_1 - k_2| \leqslant 32}} \|f_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^M}\|_{L^2} \|g_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^M}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ & \lesssim A^{-\frac{d-3}{2}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} & \sum_{1 \ll M \leqslant M_0} \sum_{\substack{(k_1, k_2) \in K_M^{\mathcal{I}} \\ 16 \leqslant |k_1 - k_2| \leqslant 32}} I_M^{k_1, k_2} \\ & \lesssim (\log A) A^{-\frac{d-3}{2}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

This gives the desired estimate since $A^{-1} N_1 \lesssim N_0$.

For the latter term, letting $2^{10} \leq K \leq 2^{-10}M_0$, we first assume $k_1 \in \mathfrak{K}_{M_0}^K$. Define

$$K_{M,M_0}^{k_1,k_2} = \{(k'_1, k'_2) \mid (\mathfrak{D}_{k'_1}^M \times \mathfrak{D}_{k'_2}^M) \subset (\mathfrak{D}_{k_1}^{M_0} \times \mathfrak{D}_{k_2}^{M_0})\}.$$

Let M' be the maximal dyadic number which satisfies $M' \leq AK^{-1/2}$. If $|k_1 - k_2| \leq 16$, we have

$$\mathfrak{D}_{k_1}^{M_0} \times \mathfrak{D}_{k_2}^{M_0} = \bigcup_{2M_0 \leq M \leq M'} \bigcup_{\substack{(k'_1, k'_2) \in K_{M,M_0}^{k_1,k_2} \\ 16 \leq |k'_1 - k'_2| \leq 32}} \mathfrak{D}_{k'_1}^M \times \mathfrak{D}_{k'_2}^M \cup \bigcup_{\substack{(k'_1, k'_2) \in K_{M',M_0}^{k_1,k_2} \\ |k'_1 - k'_2| \leq 16}} \mathfrak{D}_{k'_1}^{M'} \times \mathfrak{D}_{k'_2}^{M'}.$$

This implies

$$(4.65) \quad I_{M_0}^{k_1,k_2} \lesssim \sum_{2M_0 \leq M \leq M'} \sum_{\substack{(k'_1, k'_2) \in K_{M,M_0}^{k_1,k_2} \\ 16 \leq |k'_1 - k'_2| \leq 32}} I_M^{k'_1,k'_2} + \sum_{\substack{(k'_1, k'_2) \in K_{M',M_0}^{k_1,k_2} \\ |k'_1 - k'_2| \leq 16}} I_{M'}^{k'_1,k'_2}.$$

For the former term, we may assume $N_0 \gtrsim A^{-1}K^{1/2}N_1$. It follows from Propositions 4.26 and 4.27 that

$$\begin{aligned} & \sum_{2M_0 \leq M \leq M'} \sum_{\substack{(k'_1, k'_2) \in K_{M,M_0}^{k_1,k_2} \\ 16 \leq |k_1 - k_2| \leq 32}} I_M^{k'_1,k'_2} \\ & \lesssim (\log A) A^{-\frac{d-2}{2}} K^{\frac{1}{2}} N_0^{-\frac{1}{2}} N_1^{\frac{d-5}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^{M_0}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^{M_0}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ & \lesssim (\log A) A^{-\frac{2d-5}{4}} K^{\frac{3}{8}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^{M_0}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^{M_0}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

We next consider the latter term. If $N_0 \gg N_1/M'$ Proposition 4.26 yields

$$\begin{aligned} I_{M'}^{k_1,k_2} & \lesssim A^{-\frac{d-2}{2}} N_0^{-\frac{1}{2}} N_1^{\frac{d-5}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k'_1}^{M'}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k'_2}^{M'}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ & \lesssim A^{-\frac{2d-5}{4}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|f_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k'_1}^{M'}}\|_{L^2} \|g_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k'_2}^{M'}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Next we assume $N_0 \lesssim N_1/M'$. We divide the proof into the two cases. First we assume

$$|\xi| \gg M'^{-1} K^{-1/2} N_1 \sim A^{-1} N_1$$

which provides $|\bar{\Phi}(\xi_1, \eta_1, \xi_2, \eta_2)| \gtrsim A^{-1}N_1^3$. Thus, by (4.52) in Proposition 4.25 and $N_0 \lesssim N_1/M'$, we obtain

$$\begin{aligned} I_{M'}^{k'_1, k'_2} &\lesssim A^{-\frac{d-3}{2}} K^{\frac{1}{4}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ &\quad \times \|f_{N_1, L_1}|_{\tilde{\mathfrak{D}}_{k'_1}^{M'}}\|_{L^2} \|g_{N_2, L_2}|_{\tilde{\mathfrak{D}}_{k'_2}^{M'}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ &\lesssim A^{-\frac{2d-5}{4}} K^{\frac{3}{8}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ &\quad \times \|f_{N_1, L_1}|_{\tilde{\mathfrak{D}}_{k'_1}^{M'}}\|_{L^2} \|g_{N_2, L_2}|_{\tilde{\mathfrak{D}}_{k'_2}^{M'}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Next we treat the case $|\xi| \lesssim A^{-1}N_1$. Since $N_0 \lesssim N_1/M' \sim A^{-1}K^{1/2}N_1$, $|(\xi, \eta)|$ is confined to a rectangle set whose long-side length is $\sim A^{-1}K^{1/2}N_1$ and short-side length is $\sim A^{-1}N_1$. Therefore, after decomposing $|(\xi, \eta)|$ into $\sim K^{1/2}$ square tiles whose side length is A^{-1} , we utilize Proposition 4.8 and get

$$\begin{aligned} I_{M'}^{k'_1, k'_2} &\lesssim A^{-\frac{d-3}{2}} K^{\frac{1}{4}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ &\quad \times \|f_{N_1, L_1}|_{\tilde{\mathfrak{D}}_{k'_1}^{M'}}\|_{L^2} \|g_{N_2, L_2}|_{\tilde{\mathfrak{D}}_{k'_2}^{M'}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2} \\ &\lesssim A^{-\frac{2d-5}{4}} K^{\frac{3}{8}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ &\quad \times \|f_{N_1, L_1}|_{\tilde{\mathfrak{D}}_{k'_1}^{M'}}\|_{L^2} \|g_{N_2, L_2}|_{\tilde{\mathfrak{D}}_{k'_2}^{M'}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Collecting the above estimates, we obtain

$$(4.66) \quad \begin{aligned} I_{M_0}^{k_1, k_2} &\lesssim (\log A) A^{-\frac{2d-5}{4}} K^{\frac{3}{8}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ &\quad \times \|f_{N_1, L_1}|_{\tilde{\mathfrak{D}}_{k_1}^{M_0}}\|_{L^2} \|g_{N_2, L_2}|_{\tilde{\mathfrak{D}}_{k_2}^{M_0}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Lastly, we assume $k_1 \in \mathfrak{K}_{M_0}$. In the same way as the proof for the latter term of (4.65), we can obtain

$$(4.67) \quad \begin{aligned} I_{M_0}^{k_1, k_2} &\lesssim A^{-\frac{d-3}{2}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ &\quad \times \|f_{N_1, L_1}|_{\tilde{\mathfrak{D}}_{k_1}^{M_0}}\|_{L^2} \|g_{N_2, L_2}|_{\tilde{\mathfrak{D}}_{k_2}^{M_0}}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Consequently, since $K \leqslant A^{2/3}$, (4.66) and (4.67) complete the proof as follows.

$$\begin{aligned} \sum_{\substack{(k_1, k_2) \in K_{M_0}^T \\ |k_1 - k_2| \leqslant 16}} I_{M_0}^{k_1, k_2} &\lesssim \sum_{2^{10} \leqslant K \leqslant 2^{-10} M_0} \sum_{\substack{k_1 \in \mathfrak{K}_{M_0}^K \\ |k_1 - k_2| \leqslant 16}} I_{M_0}^{k_1, k_2} + \sum_{\substack{k_1 \in \mathfrak{K}_{M_0} \\ |k_1 - k_2| \leqslant 16}} I_{M_0}^{k_1, k_2} \\ &\lesssim (\log A) A^{-\frac{d-3}{2}} N_0^{-\frac{1}{4}} N_1^{\frac{2d-11}{4}} (L_0 L_1 L_2)^{\frac{1}{2}} \\ &\quad \times \|f_{N_1, L_1}\|_{L^2} \|g_{N_2, L_2}\|_{L^2} \|h_{N_0, L_0}\|_{L^2}. \quad \square \end{aligned}$$

Proof of Proposition 4.10. — Collecting Propositions 4.16, 4.19, 4.22, 4.23, since $A^{-1}N_1 \lesssim N_0$, we completed the proof of Proposition 4.10. \square

5. Proof of the key bilinear estimate: Case 2

It remains to show (3.3) when the supports of \widehat{u}_{N_1, L_1} and \widehat{v}_{N_2, L_2} are both contained in $\{(\tau, \xi, \eta) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^{d-1} \mid |\xi| \leq 2^{-5}N_{012}^{\max}\}$. Throughout this section, $L_{012}^{\max} \ll (N_{012}^{\max})^3$ and $N_{012}^{\min} \gg 1$ are assumed. Let us start with the case $1 \ll N_0 \lesssim N_1 \sim N_2$.

Assumption 5.1. — Let α be dyadic such that $2^5 \leq \alpha \leq N_1^3$ and we assume that

- (1) $1 \ll N_0 \lesssim N_1 \sim N_2$,
- (2) $\alpha^{-1}N_1 \leq \max(|\xi_1|, |\xi_2|) \leq 2\alpha^{-1}N_1$.

PROPOSITION 5.2. — Assume Assumption 5.1. Then we obtain

$$(5.1) \quad \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim N_0^{\frac{d-2}{2}} N_1^{-1+\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

where $d\sigma_j = d\tau_j d\xi_j d\eta_j$ and $*$ denotes $(\tau, \xi, \eta) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2)$.

We first note that $\max(|\xi_1|, |\xi_2|) \leq 2\alpha^{-1}N_1$ means $|\xi| \leq 4\alpha^{-1}N_1$. Then, if $L_{012}^{\max} \gtrsim \alpha^{-1}N_1^3$, we easily get (5.1) by utilizing the L^4 Strichartz estimate. Hereafter, we assume $L_{012}^{\max} \ll \alpha^{-1}N_1^3$.

DEFINITION 5.3. — Let $k = (k_{(1)}, \dots, k_{(d)}) \in \mathbb{Z}^d$. We define cubes $\{\mathcal{C}_k^{\alpha, A}\}_{k \in \mathbb{Z}^d}$ and $\{\tilde{\mathcal{C}}_k^{\alpha, A}\}_{k \in \mathbb{Z}^d}$ as

$$\mathcal{C}_k^{\alpha, A} = \left\{ x = (x_1, \dots, x_d) \in \mathbb{R}^d \mid \begin{array}{l} x_1 \in A^{-1}\alpha^{-1}N_1[k_{(1)}, k_{(1)}+1), \\ x_i \in A^{-1}N_1[k_{(i)}, k_{(i)}+1) \text{ for } i = 2, \dots, d, \end{array} \right\},$$

and $\tilde{\mathcal{C}}_k^{\alpha, A} := \mathbb{R} \times \mathcal{C}_k^{\alpha, A}$. Lastly we define $\mathcal{E}_{j,k}^{\alpha, A} = \bar{\mathcal{S}}_j^A \cap \tilde{\mathcal{C}}_k^{\alpha, A}$.

PROPOSITION 5.4. — Assume Assumption 5.1. Let $16 \leq |j_1 - j_2| \leq 32$ and $k_1, k_2 \in \mathbb{Z}^d$. Then we get

$$(5.2) \quad \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1}|_{\mathcal{E}_{j_1, k_1}^{\alpha, A}}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2}|_{\mathcal{E}_{j_2, k_2}^{\alpha, A}}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim \alpha A^{-\frac{d-3}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

where $d\sigma_j = d\tau_j d\xi_j d\eta_j$ and $*$ denotes $(\tau, \xi, \eta) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \eta_1 + \eta_2)$.

Proof. — We use the same notations as in the proof of Proposition 4.8. Similarly to the proof of Proposition 4.8, by applying a suitable rotation to $\boldsymbol{\eta}_1, \boldsymbol{\eta}_2$, we can assume $|\eta_1\eta'_2 - \eta_2\eta'_1| \sim A^{-1}N_1^2$, $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$ and, for fixed $\xi_1, \xi_2, \check{\boldsymbol{\eta}}_1, \check{\boldsymbol{\eta}}_2$, we will show

$$(5.3) \quad \begin{aligned} & \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\mathcal{E}_{j_1, k_1}^{\alpha, A}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\mathcal{E}_{j_2, k_2}^{\alpha, A}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\tilde{\sigma}_1 d\tilde{\sigma}_2 \right| \\ & \lesssim \alpha^{\frac{3}{2}} A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}(\xi_1, \check{\boldsymbol{\eta}}_1)\|_{L^2_{\tau \eta \eta'}} \\ & \quad \times \|\widehat{v}_{N_2, L_2}(\xi_2, \check{\boldsymbol{\eta}}_2)\|_{L^2_{\tau \eta \eta'}} \|\widehat{w}_{N_0, L_0}(\xi, \check{\boldsymbol{\eta}})\|_{L^2_{\tau \eta \eta'}}. \end{aligned}$$

Let $\ell = (\ell_{(1)}, \ell_{(2)}) \in \mathbb{Z}^2$ and

$$\begin{aligned} \mathcal{G}_\ell^{\alpha, A} &:= \{(\eta, \eta') \in A^{-1}\alpha^{-\frac{1}{2}}N_1([\ell_{(1)}, \ell_{(1)}+1) \times [\ell_{(2)}, \ell_{(2)}+1))\}, \\ \widetilde{\mathcal{G}}_\ell^{\alpha, A} &:= \mathbb{R}^2 \times \mathcal{G}_\ell^{\alpha, A} \times \mathbb{R}^{d-3}, \quad \mathcal{H}_{j, k, \ell}^{\alpha, A} := \bar{\mathcal{S}}_j^A \cap \bar{\mathcal{C}}_k^{\alpha, A} \cap \widetilde{\mathcal{G}}_\ell^{\alpha, A}. \end{aligned}$$

Define $\mathcal{L}_{k_1}^{\alpha, A} = \{\ell \in \mathbb{Z}^2 \mid \widetilde{\mathcal{C}}_{k_1}^{\alpha, A} \cap \widetilde{\mathcal{G}}_\ell^{\alpha, A} \neq \emptyset\}$. We easily observe that the number of $\ell_1 \in \mathcal{L}_{k_1}^{\alpha, A}$ is comparable to α . Therefore, for fixed $\ell_1 \in \mathcal{L}_{k_1}^{\alpha, A}$, it suffices to show

$$(5.4) \quad \begin{aligned} & \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\mathcal{H}_{j_1, k_1, \ell_1}^{\alpha, A}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\mathcal{E}_{j_2, k_2}^{\alpha, A}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\tilde{\sigma}_1 d\tilde{\sigma}_2 \right| \\ & \lesssim \alpha A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2_{\tau \eta \eta'}} \|\widehat{v}_{N_2, L_2}\|_{L^2_{\tau \eta \eta'}} \|\widehat{w}_{N_0, L_0}\|_{L^2_{\tau \eta \eta'}}. \end{aligned}$$

Indeed, by using this estimate, we have

$$\begin{aligned} (\text{LHS}) \text{ of (5.3)} &\leqslant \sum_{\ell \in \mathcal{L}_{k_1}^{\alpha, A}} (\text{LHS}) \text{ of (5.4)} \\ &\lesssim \alpha A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \sum_{\ell_1 \in \mathcal{L}_{k_1}^{\alpha, A}} \|\widehat{u}_{N_1, L_1}|_{\mathcal{H}_{j_1, k_1, \ell_1}^{\alpha, A}}\|_{L^2_{\tau \eta \eta'}} \\ &\quad \times \|\widehat{v}_{N_2, L_2}\|_{L^2_{\tau \eta \eta'}} \|\widehat{w}_{N_0, L_0}\|_{L^2_{\tau \eta \eta'}} \\ &\lesssim \alpha^{\frac{3}{2}} A^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2_{\tau \eta \eta'}} \\ &\quad \times \|\widehat{v}_{N_2, L_2}\|_{L^2_{\tau \eta \eta'}} \|\widehat{w}_{N_0, L_0}\|_{L^2_{\tau \eta \eta'}}. \end{aligned}$$

We show (5.4). By the almost orthogonality, we can assume that \widehat{v}_{N_2, L_2} is restricted to $\mathcal{H}_{j_2, k_2, \ell_2}^{\alpha, A}$ with fixed $\ell_2 \in \mathcal{L}_{k_2}^{\alpha, A}$. By the same argument as in the proof of Proposition 4.8, we establish

$$(5.5) \quad \|\widetilde{f}_{\xi_1, \check{\boldsymbol{\eta}}_1}|_{S_1} * \widetilde{g}_{\xi_2, \check{\boldsymbol{\eta}}_2}|_{S_2}\|_{L^2(S_3)} \lesssim \alpha A^{\frac{1}{2}} \|\widetilde{f}_{\xi_1, \check{\boldsymbol{\eta}}_1}\|_{L^2(S_1)} \|\widetilde{g}_{\xi_2, \check{\boldsymbol{\eta}}_2}\|_{L^2(S_2)},$$

where we used the similar notations that were defined in the proof of Proposition 4.8: The functions are

$$\begin{aligned}\tilde{f}_{\xi_1, \check{\eta}_1}(\tau_1, \eta_1, \eta'_1) &= \widehat{u}_{N_1, L_1}|_{\mathcal{H}_{j_1, k_1, \ell_1}^{\alpha, A}}(N_1^3 \tau_1, \xi_1, N_1 \eta_1, N_1 \eta'_1, \check{\eta}_1), \\ \tilde{g}_{\xi_2, \check{\eta}_2}(\tau_2, \eta_2, \eta'_2) &= \widehat{v}_{N_2, L_2}|_{\mathcal{H}_{j_2, k_2, \ell_2}^{\alpha, A}}(N_1^3 \tau_2, \xi_2, N_1 \eta_2, N_1 \eta'_2, \check{\eta}_2),\end{aligned}$$

and with $\phi_{\xi_j, \check{\eta}_j, c_j}(\eta, \eta') = (\xi_j(\xi_j^2 + |\eta|^2) + c_j, \eta, \eta')$, we define

$$\begin{aligned}S_j &= \left\{ \phi_{\xi_j, \bar{\eta}_j, c_j}(\eta_j, \eta'_j) \in \mathbb{R}^3 \mid (N_1 \eta_j, N_1 \eta'_j) \in \mathcal{G}_{\ell_j}^{\alpha, A} \right\}, \quad (j = 1, 2), \\ S_3 &= \left\{ (\psi_{\xi, \bar{\eta}}^{\alpha}(\eta, \eta'), \eta, \eta') \in \mathbb{R}^3 \mid \psi_{\xi, \check{\eta}}(\eta, \eta') = \xi(\xi^2 + |\eta|^2) + c_0 \right\},\end{aligned}$$

where $c_0, c_1, c_2 \in \mathbb{R}$, $\xi = N_1^{-1} \xi$, $\check{\xi}_j = N_1^{-1} \xi_j$, $\bar{\eta}_j = N_1^{-1} \check{\eta}_j$, $\bar{\eta} = N_1^{-1} \check{\eta}$. Since $\text{diam}(S_1) \lesssim A^{-1} \alpha^{-1/2}$, $\text{diam}(S_2) \lesssim A^{-1} \alpha^{-1/2}$, we may assume $\text{diam}(S_3) \lesssim A^{-1} \alpha^{-1/2}$. We establish (5.5) by using the nonlinear Loomis–Whitney inequality. However, it is observed that the hypersurfaces S_1, S_2, S_3 do not satisfy the necessary diameter condition. To be specific, the diameters of the three hypersurfaces are all comparable to $A^{-1} \alpha^{-1/2}$ and the transversality is comparable to $A^{-1} \alpha^{-2}$. To overcome this difficulty, we employ new functions.

$$\begin{aligned}\tilde{f}_{\xi_1, \check{\eta}_1}^{\alpha}(\tau_1, \eta_1, \eta'_1) &= \tilde{f}_{\xi_1, \check{\eta}_1}(\tau_1, \alpha^{\frac{1}{2}} \eta_1, \alpha^{\frac{1}{2}} \eta'_1), \\ \tilde{g}_{\xi_2, \check{\eta}_2}^{\alpha}(\tau_2, \eta_2, \eta'_2) &= \tilde{g}_{\xi_2, \check{\eta}_2}(\tau_2, \alpha^{\frac{1}{2}} \eta_2, \alpha^{\frac{1}{2}} \eta'_2).\end{aligned}$$

Then, (5.5) can be rewritten as

$$(5.6) \quad \|\tilde{f}_{\xi_1, \check{\eta}_1}^{\alpha}|_{S_1^{\alpha}} * \tilde{g}_{\xi_2, \check{\eta}_2}^{\alpha}|_{S_2^{\alpha}}\|_{L^2(S_3^{\alpha})} \lesssim (A\alpha)^{\frac{1}{2}} \|\tilde{f}_{\xi_1, \check{\eta}_1}^{\alpha}\|_{L^2(S_1^{\alpha})} \|\tilde{g}_{\xi_2, \check{\eta}_2}^{\alpha}\|_{L^2(S_2^{\alpha})},$$

where $\phi_{\xi_j, \check{\eta}_j, c_j}^{\alpha}(\eta, \eta') = (\xi_j(\xi_j^2 + \alpha(\eta^2 + \eta'^2) + |\check{\eta}|^2) + c_j, \eta, \eta')$ and

$$\begin{aligned}S_j^{\alpha} &= \left\{ \phi_{\xi_j, \bar{\eta}_j, c_j}^{\alpha}(\eta_j, \eta'_j) \in \mathbb{R}^3 \mid (\alpha^{\frac{1}{2}} N_1 \eta_j, \alpha^{\frac{1}{2}} N_1 \eta'_j) \in \mathcal{G}_{\ell_j}^{\alpha, A} \right\}, \quad (j = 1, 2), \\ S_3^{\alpha} &= \left\{ (\psi_{\xi, \bar{\eta}}^{\alpha}(\eta, \eta'), \eta, \eta') \in \mathbb{R}^3 \mid \psi_{\xi, \check{\eta}}^{\alpha}(\eta, \eta') = \xi(\xi^2 + \alpha(\eta^2 + \eta'^2) + |\check{\eta}|^2) + c_0 \right\},\end{aligned}$$

Now we verify that the hypersurfaces $S_1^{\alpha}, S_2^{\alpha}, S_3^{\alpha}$ satisfy the suitable conditions to utilize the nonlinear Loomis–Whitney inequality. Let

$$\begin{aligned}\lambda_1 &= \phi_{\xi_1, \bar{\eta}_1, c_1}^{\alpha}(\eta_1, \eta'_1) \in S_1^{\alpha}, \\ \lambda_2 &= \phi_{\xi_2, \bar{\eta}_2, c_2}^{\alpha}(\eta_2, \eta'_2) \in S_2^{\alpha}, \\ \lambda_3 &= (\psi_{\xi, \bar{\eta}}^{\alpha}(\eta, \eta'), \eta, \eta') \in S_3^{\alpha}.\end{aligned}$$

We can write the unit normals $\mathbf{n}_1(\lambda_1), \mathbf{n}_2(\lambda_2), \mathbf{n}_3(\lambda_3)$ on $\lambda_1, \lambda_2, \lambda_3$ as

$$\mathbf{n}_j(\lambda_j) = \frac{1}{\sqrt{1 + 4\alpha^2 \tilde{\xi}_j^2 |\eta_j|^2}} \left(-1, 2\alpha \tilde{\xi}_j \eta_j, 2\alpha \tilde{\xi}_j \eta'_j \right), \quad (j = 1, 2)$$

and $\mathfrak{n}_3(\lambda_3)$ accordingly. Since $|\tilde{\xi}_1| \leq 2\alpha^{-1}$, $|\tilde{\xi}_2| \leq 2\alpha^{-1}$, we easily observe that the hypersurfaces satisfy the necessary regularity conditions, and the diameters of hypersurfaces are all comparable to $A^{-1}\alpha^{-1}$. Thus, we consider the transversality here. Let $(\hat{\eta}_1, \hat{\eta}_1')$, $(\hat{\eta}_2, \hat{\eta}_2')$, $(\hat{\eta}, \hat{\eta}')$ satisfy $(\hat{\eta}_1, \hat{\eta}_1') + (\hat{\eta}_2, \hat{\eta}_2') = (\hat{\eta}, \hat{\eta}')$ and

$$\begin{aligned}\hat{\lambda}_1 &= \phi_{\xi_1, \bar{\eta}_1, c_1}^\alpha(\hat{\eta}_1, \hat{\eta}_1') \in S_1^\alpha, \\ \hat{\lambda}_2 &= \phi_{\xi_2, \bar{\eta}_2, c_2}^\alpha(\hat{\eta}_2, \hat{\eta}_2') \in S_2^\alpha, \\ \hat{\lambda}_3 &= (\psi_{\xi, \bar{\eta}}^\alpha(\hat{\eta}, \hat{\eta}'), \hat{\eta}, \hat{\eta}') \in S_3^\alpha.\end{aligned}$$

It suffices to show

$$(A\alpha)^{-1} \lesssim |\det N(\hat{\lambda}_1, \hat{\lambda}_2, \hat{\lambda}_3)|.$$

We have

$$\begin{aligned}|\det N(\hat{\lambda}_1, \hat{\lambda}_2, \hat{\lambda}_3)| &\gtrsim \left| \det \begin{pmatrix} -1 & -1 & -1 \\ 2\alpha\tilde{\xi}_1\hat{\eta}_1 & 2\alpha\tilde{\xi}_2\hat{\eta}_2 & 2\alpha\tilde{\xi}\hat{\eta} \\ 2\alpha\tilde{\xi}_1\hat{\eta}_1' & 2\alpha\tilde{\xi}_2\hat{\eta}_2' & 2\alpha\tilde{\xi}\hat{\eta}' \end{pmatrix} \right| \\ &\gtrsim \alpha^2 |(\hat{\eta}_1\hat{\eta}_2' - \hat{\eta}_2\hat{\eta}_1')(\tilde{\xi}_1^2 + \tilde{\xi}_1\tilde{\xi}_2 + \tilde{\xi}_2^2)| \gtrsim A^{-1}\alpha^{-1}.\end{aligned}$$

Here we used the assumptions $\alpha^{-1}N_1 \leq \max(|\xi_1|, |\xi_2|) \leq 2\alpha^{-1}N_1$ which implies $|\tilde{\xi}_1^2 + \tilde{\xi}_1\tilde{\xi}_2 + \tilde{\xi}_2^2| \sim \alpha^{-2}$, and $|(\hat{\eta}_1, \hat{\eta}_1')| \sim |(\hat{\eta}_2, \hat{\eta}_2')| \sim \alpha^{-1/2}$, $|\eta_1\eta'_2 - \eta_2\eta'_1| \sim A^{-1}N_1^2$ which imply $|\hat{\eta}_1\hat{\eta}_2' - \hat{\eta}_2\hat{\eta}_1'| \sim (A\alpha)^{-1}$. \square

PROPOSITION 5.5. — Assume Assumption 5.1. Let $16 \leq |j_1 - j_2| \leq 32$. Then we obtain

$$\begin{aligned}(5.7) \quad &\left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\bar{\mathcal{S}}_{j_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\bar{\mathcal{S}}_{j_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ &\lesssim N_0^{\frac{d-2}{2}} N_1^{-1} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}|_{\bar{\mathcal{S}}_{j_1}^A}\|_{L^2} \|\widehat{v}_{N_2, L_2}|_{\bar{\mathcal{S}}_{j_2}^A}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},\end{aligned}$$

where $d\sigma_j = d\tau_j d\xi_j d\boldsymbol{\eta}_j$ and $*$ denotes $(\tau, \xi, \boldsymbol{\eta}) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2)$.

Proof. — In the case $A \sim 1$, since $|\xi| \lesssim \alpha^{-1}N_1$ and $N_0 \sim N_1 \sim N_2$, Proposition 5.4 immediately gives (5.7). Therefore, we assume $A \gg 1$. Furthermore, without loss of generality, we can assume $|\eta_1\eta'_2 - \eta_2\eta'_1| \sim A^{-1}N_1^2$, $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$. Let M be dyadic such that $2 \leq M \leq A$ and suppose that k_1, k_2 satisfy $16 \leq |k_1 - k_2| \leq 32$. We first show the following

inequality.

$$(5.8) \quad \begin{aligned} & \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau_2, \xi_2, \eta_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim (M^{-\frac{1}{2}} + A^{-\frac{1}{2}} M^{\frac{1}{2}}) N_0^{\frac{d-2}{2}} N_1^{-1} (L_0 L_1 L_2)^{\frac{1}{2}} \\ & \quad \times \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}, \end{aligned}$$

We divide the proof of (5.8) into the two cases $|\eta_1 + \eta_2| \lesssim M^{-1} N_1$ and $|\eta_1 + \eta_2| \gg M^{-1} N_1$.

Case $|\eta_1 + \eta_2| \lesssim M^{-1} N_1$. — Let us fix η'_1, η'_2 . (5.8) is verified by showing the following estimate.

$$(5.9) \quad \begin{aligned} & \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau_2, \xi_2, \eta_2) d\widehat{\sigma}_1 d\widehat{\sigma}_2 \right| \\ & \lesssim M^{-\frac{1}{2}} N_1^{-1} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}(\eta'_1)\|_{L^2_{\tau \xi \eta}} \\ & \quad \times \|\widehat{v}_{N_2, L_2}(\eta'_2)\|_{L^2_{\tau \xi \eta}} \|\widehat{w}_{N_0, L_0}(\eta')\|_{L^2_{\tau \xi \eta}}. \end{aligned}$$

We first observe that $|\eta_1 + \eta_2| \lesssim M^{-1} N_1$ provides $|\xi_1 + \xi_2| \lesssim \alpha^{-1} M^{-1} N_1$. To see this, we assume $M \gg 1$ since $M \sim 1$ is a trivial case. If we write $(\xi_1, \eta_1) = (r_1 \cos \theta_1, r_1 \sin \theta_1)$, $(\xi_2, \eta_2) = (r_2 \cos \theta_2, r_2 \sin \theta_2)$, by the assumptions, we easily check $|r_1 - r_2| \lesssim M^{-1} N_1$, $|\cos \theta_1 + \cos \theta_2| \lesssim \alpha^{-1} M^{-1}$ and $|\cos \theta_1| \lesssim \alpha^{-1}$. Therefore,

$$\begin{aligned} |\xi_1 + \xi_2| &= |r_1 \cos \theta_1 + r_2 \cos \theta_2| \leqslant |(r_1 - r_2) \cos \theta_1| + r_2 |\cos \theta_1 + \cos \theta_2| \\ &\lesssim \alpha^{-1} M^{-1} N_1. \end{aligned}$$

Thus, it suffices to show

$$(5.10) \quad \begin{aligned} & \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \eta) \widehat{u}_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \eta_1) \widehat{v}_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau_2, \xi_2, \eta_2) d\widehat{\sigma}_1 d\widehat{\sigma}_2 \right| \\ & \lesssim \alpha M^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}(\eta'_1)\|_{L^2_{\tau \xi \eta}} \\ & \quad \times \|\widehat{v}_{N_2, L_2}(\eta'_2)\|_{L^2_{\tau \xi \eta}} \|\widehat{w}_{N_0, L_0}(\eta')\|_{L^2_{\tau \xi \eta}}. \end{aligned}$$

To show (5.10), we apply a dyadic decomposition to $|\eta_1 + \eta_2|$. Let $m \in \mathbb{N}_0$ and define

$$\mathbb{S}_\delta^m = \{\eta \in \mathbb{R} \mid m\delta^{-1} N_1 \leqslant |\eta| \leqslant (m+1)\delta^{-1} N_1\}.$$

Since $|\eta_1 + \eta_2| \lesssim M^{-1}N_1$, we can see $\{\eta_1 + \eta_2\} \subset \bigcup_{m \lesssim \alpha} \mathbb{S}_{\alpha M}^m$. Therefore, for fixed $m \in \mathbb{Z}$, we only need to show

$$(5.11) \quad \begin{aligned} & \left| \int_* \mathbf{1}_{\mathbb{S}_{\alpha M}^m}(\eta) \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim (\alpha M)^{\frac{1}{2}} N_1^{-2} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}(\boldsymbol{\eta}'_1)\|_{L^2_{\tau \xi \eta}} \\ & \quad \times \|\widehat{v}_{N_2, L_2}(\boldsymbol{\eta}'_2)\|_{L^2_{\tau \xi \eta}} \|\widehat{w}_{N_0, L_0}(\boldsymbol{\eta}')\|_{L^2_{\tau \xi \eta}}. \end{aligned}$$

By employing the nonlinear Loomis–Whitney inequality, we can establish (5.11) in the same manner as that for Proposition 4.5. We omit the details.

Case $|\eta_1 + \eta_2| \gg M^{-1}N_1$. — Next we consider the case $|\eta_1 + \eta_2| \gg M^{-1}N_1$. It follows from $2N_0 \geq |\eta_1 + \eta_2| \gg M^{-1}N_1$ that

$$(A^{-1}N_1)^{\frac{d-2}{2}} = (A^{-1}M)^{\frac{d-2}{2}} (M^{-1}N_1)^{\frac{d-2}{2}} \lesssim (A^{-1}M)^{\frac{1}{2}} N_0^{\frac{d-2}{2}}.$$

Therefore, since $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$, for fixed $\boldsymbol{\eta}'_1, \boldsymbol{\eta}'_2$, it suffices to prove

$$(5.12) \quad \begin{aligned} & \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim A^{-\frac{d-2}{2}} N_1^{\frac{d-4}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}. \end{aligned}$$

Let a dyadic number \widetilde{M} satisfies $1 \leq \widetilde{M} \ll M$. We apply a dyadic decomposition to $|\eta_1 + \eta_2|$. Suppose that $|\eta_1 + \eta_2|$ satisfies $\widetilde{M}^{-1}N_1 \leq |\eta_1 + \eta_2| \leq 2\widetilde{M}^{-1}N_1$. Then, by the same observation as in the previous case, we get $|\xi_1 + \xi_2| \lesssim \alpha^{-1}\widetilde{M}^{-1}N_1$. Therefore, it suffices to show that for $\widetilde{M}^{-1}N_1 \leq |\eta_1 + \eta_2| \leq 2\widetilde{M}^{-1}N_1$ the following holds true.

$$(5.13) \quad \begin{aligned} & \left| \int_* \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}|_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim \alpha \widetilde{M}^{\frac{1}{2}} A^{-\frac{d-2}{2}} N_1^{\frac{d-6}{2}} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}. \end{aligned}$$

We observe that the condition $\widetilde{M}^{-1}N_1 \leq |\eta_1 + \eta_2| \leq 2\widetilde{M}^{-1}N_1$ yields $|\Phi(\xi_1, \boldsymbol{\eta}_1, \xi_2, \boldsymbol{\eta}_2)| \gtrsim (\alpha\widetilde{M})^{-1}N_1^3$. To see this, we first observe

$$\begin{aligned} & |\xi_1\eta_2(2\eta_1 + \eta_2) + \xi_2\eta_1(\eta_1 + 2\eta_2)| \\ &= \left| \frac{3}{2}(\xi_1\eta_2 + \xi_2\eta_1)(\eta_1 + \eta_2) + \frac{\xi_1\eta_2 - \xi_2\eta_1}{2}(\eta_1 - \eta_2) \right| \\ &\geq |(\xi_1\eta_2 + \xi_2\eta_1)(\eta_1 + \eta_2)| - |(\eta_1 - \eta_2)(\xi_1\eta_2 - \xi_2\eta_1)| \\ &\gtrsim (\alpha\widetilde{M}^{-1})N_1^3. \end{aligned}$$

Here we used $\widetilde{M}^{-1}N_1 \leq |\eta_1 + \eta_2|$ and $|\xi_1\eta_2 - \xi_2\eta_1| \lesssim (\alpha M)^{-1}N_1^2$ which follows from $(\xi_1, \eta_1) \times (\xi_2, \eta_2) \in \mathfrak{D}_{k_1}^{\alpha M} \times \mathfrak{D}_{k_2}^{\alpha M}$ with $|k_1 - k_2| \leq 32$. Hence, since $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$, $|\xi_1 + \xi_2| \ll |\eta_1 + \eta_2|$ and $\max(|\xi_1|, |\xi_2|) \sim \alpha^{-1}N_1$, $|\eta_1| \sim |\eta_2| \sim N_1$, we calculate

$$\begin{aligned} & |\Phi(\xi_1, \boldsymbol{\eta}_1, \xi_2, \boldsymbol{\eta}_2)| \\ &\geq |3\xi_1\xi_2(\xi_1 + \xi_2) + \xi_1\eta_2(2\eta_1 + \eta_2) + \xi_2\eta_1(\eta_1 + 2\eta_2)| + \mathcal{O}(\alpha^{-1}A^{-2}N_1^3) \\ &\geq |\xi_1\eta_2(2\eta_1 + \eta_2) + \xi_2\eta_1(\eta_1 + 2\eta_2)| - 3|\xi_1\xi_2(\xi_1 + \xi_2)| + \mathcal{O}(\alpha^{-1}A^{-2}N_1^3) \\ &\geq (\alpha\widetilde{M})^{-1}N_1^3. \end{aligned}$$

This observation also implies that we can assume $|\eta_1 + \eta_2| \ll N_1$ since $L_{012}^{\max} \ll \alpha^{-1}N_1$. Thus we assume $\widetilde{M} \gg 1$ hereafter. By using the assumptions $\widetilde{M}^{-1}N_1 \leq |\eta_1 + \eta_2| = |\eta| \leq 2\widetilde{M}^{-1}N_1$ and $|\boldsymbol{\eta}'_j| \lesssim A^{-1}N_1$, we show the following bilinear estimates:

$$\begin{aligned} (5.14) \quad & \left\| \mathbf{1}_{G_{N_0, L_0}} \int \widehat{u}_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ &\lesssim A^{-\frac{d-2}{2}} N_1^{\frac{d-3}{2}} (L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}\|_{L^2} \|\widehat{v}_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}\|_{L^2}, \\ & \left\| \mathbf{1}_{G_{N_1, L_1} \cap \widetilde{\mathfrak{D}}_{k_1}^{\alpha M}} \int \widehat{v}_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) \widehat{w}_{N_0, L_0}(\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) d\sigma_2 \right\|_{L^2} \\ &\lesssim A^{-\frac{d-2}{2}} \widetilde{M}^{-\frac{1}{2}} N_1^{\frac{d-3}{2}} (L_0 L_2)^{\frac{1}{2}} \|\widehat{v}_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}, \end{aligned}$$

and

$$\begin{aligned} & \left\| \mathbf{1}_{G_{N_2, L_2} \cap \widetilde{\mathfrak{D}}_{k_2}^{\alpha M}} \int \widehat{w}_{N_0, L_0}(\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2) \widehat{u}_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) d\sigma_1 \right\|_{L^2} \\ &\lesssim A^{-\frac{d-2}{2}} \widetilde{M}^{-\frac{1}{2}} N_1^{\frac{d-3}{2}} (L_0 L_1)^{\frac{1}{2}} \|\widehat{w}_{N_0, L_0}\|_{L^2} \|\widehat{u}_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}\|_{L^2}. \end{aligned}$$

These estimates, combined with $L_{012}^{\max} \gtrsim |\Phi(\xi_1, \boldsymbol{\eta}_1, \xi_2, \boldsymbol{\eta}_2)| \geq (\alpha \tilde{M})^{-1} N_1^3$, imply (5.13). We only consider first estimate (5.14) here. The other estimates can be handled in the similar way since $|\eta_1| \sim |\eta_2| \gg |\eta_1 + \eta_2|$. By the same argument as for the proof of Proposition 4.4, the following estimates establish the claim (5.14).

$$(5.15) \quad \sup_{\tilde{M}^{-1} N_1 \leq |\eta| \leq 2\tilde{M}^{-1} N_1} |E(\tau, \xi, \boldsymbol{\eta})| \lesssim A^{-(d-2)} N_1^{d-3} L_1 L_2,$$

where $E(\tau, \xi, \boldsymbol{\eta})$ is defined as

$$\left\{ (\tau_1, \xi_1, \boldsymbol{\eta}_1) \in G_{N_1, L_1} \cap \tilde{\mathfrak{D}}_{k_1}^{\alpha M} \mid (\tau - \tau_1, \xi - \xi_1, \boldsymbol{\eta} - \boldsymbol{\eta}_1) \in G_{N_2, L_2} \cap \tilde{\mathfrak{D}}_{k_2}^{\alpha M} \right\}.$$

We recall the function $\Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)$ which was defined in the proof of Proposition 4.4 as

$$\begin{aligned} & \max(L_1, L_2) \\ & \gtrsim |(\tau_1 - \xi_1(\xi_1^2 + |\boldsymbol{\eta}_1|^2)) + ((\tau - \tau_1) - (\xi - \xi_1)((\xi - \xi_1)^2 + |\boldsymbol{\eta} - \boldsymbol{\eta}_1|^2))| \\ & = |(\tau - \xi(\xi^2 + |\boldsymbol{\eta}|^2)) + \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)|. \end{aligned}$$

Let $(\tau_1, \xi_1, \boldsymbol{\eta}_1) \in E(\tau, \xi, \boldsymbol{\eta})$. Since $|\xi| \ll |\eta|$ and $|\boldsymbol{\eta}'| \lesssim A^{-1} N_1$, $|\boldsymbol{\eta}'_1| \lesssim A^{-1} N_1$, for fixed $\boldsymbol{\eta}_1$, it is easily observed

$$\begin{aligned} |\partial_{\xi_1} \Phi_{\xi, \boldsymbol{\eta}}(\xi_1, \boldsymbol{\eta}_1)| &= |3\xi(\xi - 2\xi_1) + \eta(\eta - 2\eta_1)| + \mathcal{O}(A^{-2} N_1^2) \\ &\gtrsim |\eta| N_1 \sim \tilde{M}^{-1} N_1^2. \end{aligned}$$

This, $|\eta| \sim \tilde{M}^{-1} N_1$ and $|\boldsymbol{\eta}'_1| \lesssim A^{-1} N_1$ complete the proof of (5.15).

Next we assume $|k_1 - k_2| \leq 16$ and show

$$(5.16) \quad \begin{aligned} & \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1} |_{\tilde{\mathfrak{D}}_{k_1}^{\alpha A}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2} |_{\tilde{\mathfrak{D}}_{k_2}^{\alpha A}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim N_0^{\frac{d-2}{2}} N_1^{-1} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2}, \end{aligned}$$

Similarly to the proof of (5.8), we divide the proof into the two cases.

Case $|\eta_1 + \eta_2| \lesssim A^{-1} N_1$. — As we saw above, we may assume $|\xi_1 + \xi_2| \lesssim (A\alpha)^{-1} N_1$. Thus Proposition 5.4 implies (5.16).

Case $|\eta_1 + \eta_2| \gg A^{-1} N_1$. — We only need to follow the proof of (5.8) in the case $|\eta_1 + \eta_2| \gg M^{-1} N_1$. We omit the details.

We now see that the two estimates (5.8) and (5.16) yield (5.7). For simplicity, we use

$$I_{k_1, k_2}^{\alpha, M} = \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1} |_{\widetilde{\mathfrak{D}}_{k_1}^{\alpha M}}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\ \times \left. \widehat{v}_{N_2, L_2} |_{\widetilde{\mathfrak{D}}_{k_2}^{\alpha M}}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right|.$$

It follows from (5.8) and (5.16) that

(LHS) of (5.7)

$$\leq \sum_{2 \leq M \leq A} \sum_{16 \leq |k_1 - k_2| \leq 32} I_{k_1, k_2}^{\alpha, M} + \sum_{|k_1 - k_2| \leq 16} I_{k_1, k_2}^{\alpha, A} \\ \lesssim \sum_{2 \leq M \leq A} (M^{-\frac{1}{2}} + M^{\frac{1}{2}} A^{-\frac{1}{2}}) N_0^{\frac{d-2}{2}} N_1^{-1} (L_0 L_1 L_2)^{\frac{1}{2}} \\ \times \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2} \\ + N_0^{\frac{d-2}{2}} N_1^{-1} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2} \\ \lesssim N_0^{\frac{d-2}{2}} N_1^{-1} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

which completes the proof of (5.7). \square

In the same manner as in the proof of (4.1) in the case (Ic) (see p. 1225), Proposition 5.5 gives Proposition 5.2. We omit the proof.

6. Proof of the key bilinear estimate: Case 3

Next we deal with $1 \ll N_2 \lesssim N_1 \sim N_0$.

Assumption 6.1. — Let α be dyadic such that $2^5 \leq \alpha \leq N_1^3$ and we assume that

- (1) $1 \ll N_2 \lesssim N_1 \sim N_0$,
- (2) $\alpha^{-1} N_1 \leq \max(|\xi_1|, |\xi_2|) \leq 2\alpha^{-1} N_1$.

PROPOSITION 6.2. — Assume Assumption 6.1. Then we get

$$(6.1) \quad \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{v}_{N_2, L_2}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ \lesssim N_2^{\frac{d-4}{2} + 2\varepsilon} N_1^{-\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},$$

where $d\sigma_j = d\tau_j d\xi_j d\boldsymbol{\eta}_j$ and $*$ denotes $(\tau, \xi, \boldsymbol{\eta}) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2)$.

Since $|\xi| \lesssim \alpha^{-1} N_1$, (6.1) is given by the following estimate.

PROPOSITION 6.3. — Assume Assumption 6.1. Let $16 \leq |j_1 - j_2| \leq 32$. Then we get

$$\begin{aligned} & \left| \int_* \widehat{v}_{N_2, L_2}(\tau, \xi, \boldsymbol{\eta}) \widehat{u}_{N_1, L_1}|_{\bar{\mathcal{S}}_{j_1}^A}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \widehat{w}_{N_0, L_0}|_{\bar{\mathcal{S}}_{j_2}^A}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \right| \\ & \lesssim \alpha N_2^{\frac{d-4}{2} + 2\varepsilon} N_1^{-1-\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}|_{\bar{\mathcal{S}}_{j_1}^A}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}|_{\bar{\mathcal{S}}_{j_2}^A}\|_{L^2}, \end{aligned}$$

where $d\sigma_j = d\tau_j d\xi_j d\boldsymbol{\eta}_j$ and $*$ denotes $(\tau, \xi, \boldsymbol{\eta}) = (\tau_1 + \tau_2, \xi_1 + \xi_2, \boldsymbol{\eta}_1 + \boldsymbol{\eta}_2)$.

By exchanging the roles of \widehat{w}_{N_0, L_0} and \widehat{v}_{N_2, L_2} , we can establish Proposition 6.3 in the same manner as Proposition 5.5. In addition, by following the proof of (4.1) in the case (Ic), Proposition 6.3 yields Proposition 6.2. We omit the details.

Now we show (3.3) with the condition
(6.2)

$$\text{supp } \widehat{u}_{N_1, L_1} \cup \text{supp } \widehat{v}_{N_2, L_2} \subset \{(\tau, \xi, \boldsymbol{\eta}) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^{d-1} \mid |\xi| \leq 2^{-5} N_{012}^{\max}\}.$$

Proof of (3.3) under (6.2). — By symmetry, we can assume $N_2 \leq N_1$. Let us consider $1 \ll N_0 \lesssim N_1 \sim N_2$. We define

$$\begin{aligned} E_\alpha &:= \{(\xi_1, \xi_2) \mid \alpha^{-1} N_1 \leq \max(|\xi_1|, |\xi_2|) \leq 2\alpha^{-1} N_1\}, \\ F &:= \{(\xi_1, \xi_2) \mid \max(|\xi_1|, |\xi_2|) \leq N_1^{-2}\}. \end{aligned}$$

Applying dyadic decomposition to $\max(|\xi_1|, |\xi_2|)$, we see

$$\begin{aligned} & \left| \int (\partial_x w_{N_0, L_0}) u_{N_1, L_1} v_{N_2, L_2} dt dx dy \right| \\ & \lesssim \sum_{2^5 \leq \alpha \leq N_1^3} \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \mathbf{1}_{E_\alpha}(\xi_1, \xi_2) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\ & \quad \times \widehat{v}_{N_2, L_2}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \Big| \\ & \quad + \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \mathbf{1}_F(\xi_1, \xi_2) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\ & \quad \times \widehat{v}_{N_2, L_2}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \Big|. \end{aligned}$$

The first term can be handled by Proposition 5.2 as follows.

$$\begin{aligned}
& \sum_{2^5 \leq \alpha \leq N_1^3} \left| \int_* |\xi| \widehat{w}_{N_0, L_0}(\tau, \xi, \boldsymbol{\eta}) \mathbf{1}_{E_\alpha}(\xi_1, \xi_2) \widehat{u}_{N_1, L_1}(\tau_1, \xi_1, \boldsymbol{\eta}_1) \right. \\
& \quad \times \widehat{v}_{N_2, L_2}(\tau_2, \xi_2, \boldsymbol{\eta}_2) d\sigma_1 d\sigma_2 \Big| \\
& \lesssim \sum_{2^5 \leq \alpha \leq N_1^3} N_0^{\frac{d-2}{2}} N_1^{-1+\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2} \\
& \lesssim N_0^{\frac{d-2}{2}} N_1^{-1+2\varepsilon} (L_0 L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2} \|\widehat{w}_{N_0, L_0}\|_{L^2},
\end{aligned}$$

which implies (3.3) since $s > (d-4)/2$ and $L_{012}^{\max} \ll N_1^3$. Next we consider the second term. The inequality $\max(|\xi_1|, |\xi_2|) \leq N_1^{-2}$ implies $|\xi| \leq 2N_1^{-2}$. Therefore, the L^4 Strichartz estimate is enough to verify the claim.

By using Proposition 6.2, the case $1 \ll N_2 \lesssim N_0 \sim N_1$ can be treated in the similar way. We omit the details. \square

Appendix A. Proof of Proposition 2.2

First we consider (2.3). Set $\widetilde{B}_r(p) = \mathbb{R} \times B_r(p)$ and $\zeta = (\xi, \boldsymbol{\eta})$, $\zeta_j = (\xi_j, \boldsymbol{\eta}_j)$. By Plancherel's theorem, it suffices to show

$$\begin{aligned}
(A.1) \quad & \left\| \int \widehat{u}_{N_1, L_1}|_{\widetilde{B}_r(p)}(\tau_1, \zeta_1) \widehat{v}_{N_2, L_2}(\tau - \tau_1, \zeta - \zeta_1) d\tau_1 d\zeta_1 \right\|_{L^2} \\
& \lesssim r^{\frac{d-1}{2}} K^{-\frac{1}{2}} (L_1 L_2)^{\frac{1}{2}} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2}.
\end{aligned}$$

By performing a harmless decomposition, we may replace r with r' such that $r' \ll d^{-1}r$ in the above. Furthermore, by the almost orthogonality, we can assume that there exists $p' \in \mathbb{R}^d$ such that $\zeta - \zeta_1 \in B_{r'}(p')$. Since φ is a cubic polynomial, we deduce from $N_2 \leq N_1$ that

$$\sup_{1 \leq i, j \leq d} (|\partial_i \partial_j \varphi(\zeta_1)| + |\partial_i \partial_j \varphi(\zeta - \zeta_1)|) \lesssim N_1.$$

Therefore, because $K \gtrsim rN_1$, we easily observe

$$|\nabla \varphi(\zeta) - \nabla \varphi(\zeta')| \ll K \quad \text{if } \zeta, \zeta' \in B_{r'}(p).$$

This implies that there exists $j \in \mathbb{N}$ such that $1 \leq j \leq d$ and

$$(A.2) \quad |\partial_j \varphi(\zeta_1) - \partial_j \varphi(\zeta_2)| \gtrsim K,$$

for all ζ_1, ζ_2 which satisfy that there exist τ_1 and τ_2 such that $(\tau_1, \zeta_1) \in \text{supp } \widehat{u}_{N_1, L_1} \cap \widetilde{B}_{r'}(p)$ and $(\tau_2, \zeta_2) \in \text{supp } \widehat{v}_{N_2, L_2} \cap \widetilde{B}_{r'}(p')$, respectively. Now we turn to show (A.1). By the Cauchy–Schwarz inequality, we get

$$\begin{aligned} & \left\| \int \widehat{u}_{N_1, L_1} |_{\widetilde{B}_{r'}(p)}(\tau_1, \zeta_1) \widehat{v}_{N_2, L_2}(\tau - \tau_1, \zeta - \zeta_1) d\tau_1 d\zeta_1 \right\|_{L^2} \\ & \leq \left\| \left(|\widehat{u}_{N_1, L_1}|^2 * |\widehat{v}_{N_2, L_2}|^2 \right)^{1/2} |E(\tau, \zeta)|^{1/2} \right\|_{L^2} \\ & \leq \sup_{\tau, \zeta} |E(\tau, \zeta)|^{1/2} \left\| |\widehat{u}_{N_1, L_1}|^2 * |\widehat{v}_{N_2, L_2}|^2 \right\|_{L^1}^{1/2} \\ & \leq \sup_{\tau, \zeta} |E(\tau, \zeta)|^{1/2} \|\widehat{u}_{N_1, L_1}\|_{L^2} \|\widehat{v}_{N_2, L_2}\|_{L^2}, \end{aligned}$$

where $E(\tau, \zeta) \subset \mathbb{R}^{d+1}$ is defined by

$$E(\tau, \zeta) := \{(\tau_1, \zeta_1) \in G_{N_1, L_1} \cap \widetilde{B}_{r'}(p) \mid (\tau - \tau_1, \zeta - \zeta_1) \in G_{N_2, L_2}\}.$$

Thus, it suffices to show

$$(A.3) \quad |E(\tau, \zeta)| \lesssim r^{d-1} K^{-1} L_1 L_2.$$

If we fix ζ_1 , it is easily observed that

$$(A.4) \quad |\{\tau_1 \mid (\tau_1, \zeta_1) \in E(\tau, \zeta)\}| \lesssim \min(L_1, L_2).$$

Next, if we fix $(\zeta_{1,1}, \dots, \zeta_{1,j-1}, \zeta_{1,j+1}, \dots, \zeta_{1,d})$, since

$$\max(L_1, L_2) \gtrsim |\varphi(\zeta_1) + \varphi(\zeta - \zeta_1)|,$$

the inequality (A.2) implies that $\zeta_{1,j}$ is confined to an interval whose length is comparable to $\max(L_1, L_2)/K$. This, combined with (A.4) and $\zeta_1 \in B_{r'}(p)$, yields (A.3).

To see (2.4), it suffices to show

$$|\zeta_1| \geq 2|\zeta_2| \implies |\nabla \varphi(\zeta_1) - \nabla \varphi(\zeta_2)| \gtrsim |\zeta_1|^2,$$

which is immediately verified by $|\partial_\xi \varphi(\xi, \eta)| = |3\xi^2 + |\eta|^2| \geq |(\xi, \eta)|^2$ and $|\partial_{\xi_1} \varphi(\xi_1, \eta_1)| \leq 3|(\xi_1, \eta_1)|^2$. \square

BIBLIOGRAPHY

- [1] I. BEJENARU, S. HERR, J. HOLMER & D. TATARU, “On the 2D Zakharov system with L^2 -Schrödinger data”, *Nonlinearity* **22** (2009), no. 5, p. 1063–1089.
- [2] I. BEJENARU & S. HERR, “Convolutions of singular measures and applications to the Zakharov system”, *J. Funct. Anal.* **261** (2011), no. 2, p. 478–506.
- [3] I. BEJENARU, S. HERR & D. TATARU, “A convolution estimate for two-dimensional hypersurfaces”, *Rev. Mat. Iberoam.* **26** (2010), no. 2, p. 707–728.
- [4] P. M. BELLAN, *Fundamentals of Plasma Physics*, Cambridge University Press, Cambridge, 2006.

- [5] J. BENNETT, A. CARBERY & J. WRIGHT, “A non-linear generalisation of the Loomis–Whitney inequality and applications”, *Math. Res. Lett.* **12** (2005), no. 4, p. 443–457.
- [6] D. BHATTACHARYA, L. G. FARAH & S. ROUDENKO, “Global well-posedness for low regularity data in the 2d modified Zakharov–Kuznetsov equation”, <https://arxiv.org/abs/1906.05822>.
- [7] H. A. BIAGIONI & F. LINARES, “Well-posedness results for the modified Zakharov–Kuznetsov equation”, in *Nonlinear equations: methods, models and applications (Bergamo, 2001)*, Progr. Nonlinear Differential Equations Appl., vol. 54, Birkhäuser, Basel, 2003, p. 181–189.
- [8] J. BOURGAIN, “Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. II. The KdV-equation”, *Geom. Funct. Anal.* **3** (1993), no. 3, p. 209–262.
- [9] T. CANDY & S. HERR, “Transference of bilinear restriction estimates to quadratic variation norms and the Dirac-Klein-Gordon system”, *Anal. PDE* **11** (2018), no. 5, p. 1171–1240.
- [10] R. CÔTE, C. MUÑOZ, D. PILOD & G. SIMPSON, “Asymptotic stability of high-dimensional Zakharov–Kuznetsov solitons”, *Arch. Ration. Mech. Anal.* **220** (2016), no. 2, p. 639–710.
- [11] A. V. FAMINSKIĬ, “The Cauchy problem for the Zakharov–Kuznetsov equation”, *Differentsial’ nye Uravneniya* **31** (1995), no. 6, p. 1070–1081, 1103.
- [12] L. G. FARAH, F. LINARES & A. PASTOR, “A note on the 2D generalized Zakharov–Kuznetsov equation: local, global, and scattering results”, *J. Differential Equations* **253** (2012), no. 8, p. 2558–2571.
- [13] J. GINIBRE, Y. TSUTSUMI & G. VELO, “On the Cauchy problem for the Zakharov system”, *J. Funct. Anal.* **151** (1997), no. 2, p. 384–436.
- [14] A. GRÜNROCK, “On the generalized Zakharov–Kuznetsov equation at critical regularity”, <https://arxiv.org/abs/1509.09146>.
- [15] ———, “A remark on the modified Zakharov–Kuznetsov equation in three space dimensions”, *Math. Res. Lett.* **21** (2014), no. 1, p. 127–131.
- [16] A. GRÜNROCK & S. HERR, “The Fourier restriction norm method for the Zakharov–Kuznetsov equation”, *Discrete Contin. Dyn. Syst.* **34** (2014), no. 5, p. 2061–2068.
- [17] A. GRÜNROCK, M. PANTEE & J. DRUMOND SILVA, “On KP-II type equations on cylinders”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **26** (2009), no. 6, p. 2335–2358.
- [18] C. E. KENIG, G. PONCE & L. VEGA, “Oscillatory integrals and regularity of dispersive equations”, *Indiana Univ. Math. J.* **40** (1991), no. 1, p. 33–69.
- [19] ———, “A bilinear estimate with applications to the KdV equation”, *J. Amer. Math. Soc.* **9** (1996), no. 2, p. 573–603.
- [20] S. KINOSHITA, “Well-posedness for the Cauchy problem of the modified Zakharov–Kuznetsov equation”, <https://arxiv.org/abs/1911.13265>.
- [21] ———, “Global well-posedness for the Cauchy problem of the Zakharov–Kuznetsov equation in 2D”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **38** (2021), no. 2, p. 451–505.
- [22] H. KOCH & S. STEINERBERGER, “Convolution estimates for singular measures and some global nonlinear Brascamp–Lieb inequalities”, *Proc. Roy. Soc. Edinburgh Sect. A* **145** (2015), no. 6, p. 1223–1237.
- [23] E. W. LAEDKE & K. H. SPATSCHEK, “Nonlinear ion-acoustic waves in weak magnetic fields”, *Phys. Fluids* **25** (1982), no. 6, p. 985–989.
- [24] D. LANNES, F. LINARES & J.-C. SAUT, “The Cauchy problem for the Euler-Poisson system and derivation of the Zakharov–Kuznetsov equation”, in *Studies in phase space analysis with applications to PDEs*, Progr. Nonlinear Differential Equations Appl., vol. 84, Birkhäuser/Springer, New York, 2013, p. 181–213.

- [25] F. LINARES & A. PASTOR, “Well-posedness for the two-dimensional modified Zakharov-Kuznetsov equation”, *SIAM J. Math. Anal.* **41** (2009), no. 4, p. 1323-1339.
- [26] ———, “Local and global well-posedness for the 2D generalized Zakharov-Kuznetsov equation”, *J. Funct. Anal.* **260** (2011), no. 4, p. 1060-1085.
- [27] F. LINARES & J.-C. SAUT, “The Cauchy problem for the 3D Zakharov-Kuznetsov equation”, *Discrete Contin. Dyn. Syst.* **24** (2009), no. 2, p. 547-565.
- [28] L. H. LOOMIS & H. WHITNEY, “An inequality related to the isoperimetric inequality”, *Bull. Amer. Math. Soc.* **55** (1949), p. 961-962.
- [29] L. MOLINET & D. PILOD, “Bilinear Strichartz estimates for the Zakharov-Kuznetsov equation and applications”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **32** (2015), no. 2, p. 347-371.
- [30] F. RIBAUD & S. VENTO, “A note on the Cauchy problem for the 2D generalized Zakharov-Kuznetsov equations”, *C. R. Math. Acad. Sci. Paris* **350** (2012), no. 9-10, p. 499-503.
- [31] ———, “Well-posedness results for the three-dimensional Zakharov-Kuznetsov equation”, *SIAM J. Math. Anal.* **44** (2012), no. 4, p. 2289-2304.
- [32] C. SULEM & P.-L. SULEM, *The nonlinear Schrödinger equation*, Applied Mathematical Sciences, vol. 139, Springer-Verlag, New York, 1999, Self-focusing and wave collapse, xvi+350 pages.
- [33] V. E. ZAKHAROV & E. A. KUZNETSOV, “Three-dimensional solitons”, *Sov. Phys. JETP* **39** (1974), no. 2, p. 285-286.

Manuscrit reçu le 29 janvier 2020,
révisé le 8 novembre 2021,
accepté le 1^{er} décembre 2021.

Sebastian HERR
Universität Bielefeld
Fakultät für Mathematik
Postfach 10 01 31
33501 Bielefeld (Germany)
herr@math.uni-bielefeld.de
Shinya KINOSHITA
Department of Mathematics
Graduate School of Science and Engineering
Saitama University
Saitama 338-8570 (Japan)
kinoshita@mail.saitama-u.ac.jp