



ANNALES DE L'INSTITUT FOURIER

Maxim ARNOLD, Dmitry FUCHS & Serge TABACHNIKOV

**A family of integrable transformations of centroaffine
polygons: geometrical aspects**

Article à paraître, mis en ligne le 17 mai 2024, 45 p.

Article mis à disposition par ses auteurs selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE



<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les *Annales de l'Institut Fourier* sont membres du
Centre Mersenne pour l'édition scientifique ouverte

www.centre-mersenne.org

e-ISSN : 1777-5310

A FAMILY OF INTEGRABLE TRANSFORMATIONS OF CENTROAFFINE POLYGONS: GEOMETRICAL ASPECTS

by Maxim ARNOLD,
Dmitry FUCHS & Serge TABACHNIKOV (*)

ABSTRACT. — Two polygons, (P_1, \dots, P_n) and (Q_1, \dots, Q_n) in \mathbb{R}^2 are c -related if $\det(P_i, P_{i+1}) = \det(Q_i, Q_{i+1})$ and $\det(P_i, Q_i) = c$ for all i . This relation extends to twisted polygons (polygons with monodromy), and it descends to the moduli space of $\mathrm{SL}(2, \mathbb{R})$ -equivalent polygons. This relation is an equiaffine analog of the discrete bicycle correspondence studied by a number of authors. We study the geometry of this relations, present its integrals, and show that, in an appropriate sense, these relations, considered for different values of the constants c , commute. We relate this topic with the dressing chain of Veselov and Shabat. The case of small-gons is investigated in detail.

RÉSUMÉ. — On appelle deux polygones planaires (P_1, \dots, P_n) et (Q_1, \dots, Q_n) c -correspondants si leurs déterminants $\det(P_i, P_{i+1}) = \det(Q_i, Q_{i+1})$ et $\det(P_i, Q_i) = c$ sont satisfaits pour tous i .

Cette relation est un analogue équiaffine de la correspondance de bicyclette discrète étudiée par un certain nombre d'auteurs. Nous étudions la géométrie de ces relations, présentons ses intégrales, et montrons que – dans un sens approprié – ces relations commutent quand considérées pour différentes valeurs des constantes c . Nous relierons ce sujet à la chaîne de pansement de Veselov et Shabat.

1. Introduction

The motivation for this paper is two-fold.

The first one is the study of the *discrete bicycle correspondence* on polygons in the Euclidean plane [15], a discretization of the *bicycle correspondence* on smooth curves that was studied in [6, 13]. See also [7, 12] for the discrete and [18] for the continuous versions of this correspondence.

Keywords: Liouville integrability, Integrable Systems, Centroaffine geometry, Dressing chain.

2020 *Mathematics Subject Classification:* 37J70, 37J39, 53D30.

(*) S. Tabachnikov is supported by NSF grant DMS-2005444.




Figure 1.1. Left: folding the parallelogram $P_1Q_1Q'_1P_2$ to the trapezoid $P_1Q_1P_2Q_2$. Right: the pentagons \mathbf{P} and \mathbf{Q} are in the discrete bicycle correspondence.

Two n -gons, $\mathbf{P} = (P_1, \dots, P_n)$ and $\mathbf{Q} = (Q_1, \dots, Q_n)$, are in the discrete bicycle correspondence if every quadrilateral $P_iQ_iQ_{i+1}P_{i+1}$ is obtained by folding a parallelogram along a diagonal as shown in Figure 1.1:

$$(1.1) \quad |P_iP_{i+1}| = |Q_iQ_{i+1}|, \quad |P_i, Q_i| = c, \quad i = 1, \dots, n,$$

where c is a fixed parameter. In other words, $P_iQ_iP_{i+1}Q_{i+1}$ is an equilateral trapezoid, perhaps self-intersecting.

The discrete bicycle correspondence is completely integrable. Specifically, in [15] a Lax presentation with a spectral parameter of the discrete bicycle correspondence is described, providing integrals of this correspondence.

It is also shown there that the discrete bicycle correspondence commutes and shares its integrals with the *polygon recutting*, another integrable transformation of polygons, introduced and studied in [1, 2], see Figure 1.2.




Figure 1.2. Polygon recutting on vertex P_1 : point P'_1 is the reflection of P_1 in the perpendicular bisector of the diagonal P_2P_5 . The recutting of the polygon is the result of five such transformations performed cyclically.

The present paper concerns an analog of the discrete bicycle correspondence in the centroaffine geometry, associated with the group $SL(2, \mathbb{R})$ – or $SL(2, \mathbb{C})$, if one works with complex coefficients. That is, we consider two polygons in \mathbb{R}^2 congruent if they are related by a linear transformation with determinant 1. We denote the determinant by bracket.

Let $c \in \mathbb{R}$ be a non-zero number. Two n -gons, \mathbf{P} and \mathbf{Q} , are c -related if

$$(1.2) \quad [P_i, P_{i+1}] = [Q_i, Q_{i+1}], \quad [P_i, Q_i] = c, \quad i = 1, \dots, n,$$

see Figure 1.3. We write $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$.




Figure 1.3. Two c -related pentagons.

Equations (1.2) are centroaffine analogs of equations (1.1): the role of the length is played by the area (i.e., the determinant). The space of centroaffine polygons is foliated by the c -relation invariant subspaces consisting of the polygons whose “side areas” $[P_i, P_{i+1}]$ depend only on i .

Along with closed polygons ($P_{i+n} = P_i$ for all i), we consider *twisted n -gons*. A twisted n -gon \mathbf{P} is an infinite collection of points $P_i \in \mathbb{R}^2$ such that $P_{i+n} = \mathcal{M}_{\mathbf{P}}(P_i)$ for all i ; this map $\mathcal{M}_{\mathbf{P}} \in SL(2, \mathbb{R})$ is called the monodromy of the twisted polygon \mathbf{P} . Twisted polygons \mathbf{P} and \mathbf{Q} are c -related if, in addition to (1.2), they share their monodromies.

The c -relation is a discretization of a relation on centroaffine curves, which is a geometrical realization of the Bäcklund transformation of the KdV equation studied in [5, 14].

Two consecutive pairs of vertices of c -related polygons form a quadrilateral satisfying

$$[P_i, P_{i+1}] = [Q_i, Q_{i+1}], \quad [P_i, Q_i] = [P_{i+1}, Q_{i+1}].$$

We call such quadrilaterals *centroaffine butterflies*, see Figure 1.4 (the term is adopted from [15]). They are centroaffine analogs of the folded parallelograms in Figure 1.1.




Figure 1.4. A centroaffine butterfly: an affine reflection interchanges P_1 with Q_2 and P_2 with Q_1 .

We also consider centroaffine version of polygon recutting. An elementary centroaffine recutting is depicted in Figure 1.4: it is a linear involution that swaps the triangles $P_1Q_1Q_2$ and $P_1P_2Q_2$. The centroaffine recutting of a n -gon is the composition of n elementary recuttings performed cyclically.

Let $\pi : \mathbb{R}^2 \setminus \{O\} \rightarrow \mathbb{RP}^1$ be the natural projection. Abusing notation, we use the same symbol for the projections of centroaffine polygons to polygons in \mathbb{RP}^1 . This projection commutes with the natural actions of $\mathrm{SL}(2, \mathbb{R})$ on \mathbb{R}^2 and \mathbb{RP}^1 .

Let \mathbf{P} and \mathbf{Q} be c -related centroaffine polygons, and \mathbf{p} and \mathbf{q} be their projections to \mathbb{RP}^1 . Then

$$(1.3) \quad [p_i, p_{i+1}, q_i, q_{i+1}] = \frac{[P_i, P_{i+1}][Q_i, Q_{i+1}]}{[P_i, Q_i][P_{i+1}, Q_{i+1}]}$$

where the bracket on the left hand side denotes the cross-ratio (there are six different choices of cross-ratio to make; the right hand side of the formula specifies our choice). If $[P_i, P_{i+1}]$ is the same for all i , then the polygons \mathbf{p} and \mathbf{q} are in the cross-ratio relation: $[p_i, p_{i+1}, q_i, q_{i+1}] = \alpha$ for all i .

The cross-ratio relation on projective polygons was thoroughly studied, starting with [11] and, more recently, in [3] and [4]. This is the second source of our motivation: many results in this paper have analogs in [4].

The cross-ratio relation can be generalized: n -gons \mathbf{p} and \mathbf{q} are related if $[p_i, p_{i+1}, q_i, q_{i+1}] = \alpha_i$, where α_i is an n -periodic sequence (not necessarily constant). Formulas (1.2) and (1.3) imply that the projection π conjugates the c -relation with this generalized cross-ratio relation.

Let us present the main results of the paper.

In Section 2 we introduce coordinates in the moduli space of twisted centroaffine polygons and calculate the monodromy of a twisted polygon: the result is given in terms of continuants (3-diagonal determinants). We

also describe the algebraic relations satisfied by the coordinates of closed polygons.

Let $\mathbf{P} = (P_1, \dots, P_n)$ be a centroaffine n -gon. Choose a test vector Q_1 with $[P_1, Q_1] = c$, and consecutively construct vectors Q_2, \dots, Q_n, Q_{n+1} according to (1.2). We call the map $Q_1 \mapsto Q_{n+1}$ the *Lax transformation* associated with the polygon \mathbf{P} and denote it by $\mathcal{L}_{\mathbf{P},c}$. Then $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$ if and only if Q_1 is a fixed point of $\mathcal{L}_{\mathbf{P},c}$. Similarly one defines the Lax transformation associated with a twisted polygon.

We show that $\mathcal{L}_{\mathbf{P},c}$ is a Möbius map. This makes it possible to consider the c -relation on generic polygons as a 2-2 map.

Theorem 2.11 states that the c -relation satisfies the Bianchi permutability. Informally speaking, it says that the c -relations with different values of the constant c commute (see Section 2.6 for the precise formulation).

Theorem 3.1 states that if \mathbf{P} and \mathbf{Q} are c -related twisted n -gons, then the Lax transformations $\mathcal{L}_{\mathbf{P},\lambda}$ and $\mathcal{L}_{\mathbf{Q},\lambda}$ are conjugated for every value of the spectral parameter λ . This is the source of integrals of the c -relation. The moduli space of twisted centroaffine n -gons with fixed “side areas” has dimension n ; we obtain $\lfloor \frac{n+1}{2} \rfloor$ integrals therein.

In Section 3.2 we study how the c -relation interacts with the centroaffine polygon recutting. We prove that the Lax transformation is preserved by the recutting and that the recutting commutes with the c -relations (Theorem 3.8).

In Section 3.4 we calculate the integrals provided by the conjugacy invariance of the Lax transformations and show that they coincide with the integrals of the dressing chain of Veselov and Shabat [17]. In Section 3.5 we describe two relations between these integrals that hold for closed polygons.

Section 4 concerns the space of closed centroaffine polygons before its factorization by the group $\mathrm{SL}(2, \mathbb{R})$. We construct presymplectic forms on the subspaces of polygons whose “side areas” $[P_i, P_{i+1}]$ depend on i only. These forms are $\mathrm{SL}(2, \mathbb{R})$ -invariant, but they do not descend on the quotient spaces by the group. The forms are invariant under the c -relation and under the polygon recutting.

The reason to introduce these 2-forms is that they provide three additional integrals of the c -relation, quadratic in the coordinates; one polynomial function of these three integrals is invariant under $\mathrm{SL}(2, \mathbb{R})$ and is a pull-back of an integral defined on the moduli space. These integrals are interpreted as the moment map of the Hamiltonian action of $\mathfrak{sl}(2, \mathbb{R})$ on the spaces of polygons with fixed “side areas”.

These integrals define a certain *center* $C(\mathbf{P})$ of a centroaffine polygon, that takes values in quadratic forms on \mathbb{R}^2 . This center is invariant under the c -relation and the recutting, and equivariant with respect to the action of $\mathrm{SL}(2, \mathbb{R})$. It has interesting properties: it is additive with respect to cutting polygons into two, and it coincides with the origin for centroaffine butterflies.

Section 5 is devoted to a study of “small-gons”, triangles, quadrilaterals, and pentagons.

Let us emphasize that we do not prove Liouville integrability of the c -relations here: this would involve the existence of an invariant Poisson structure with respect to which our integrals commute. This, along with the relation with the theory of cluster algebras, will be studied by A. Izosimov in the forthcoming paper [8].

Acknowledgements

We are very grateful to Anton Izosimov for his insights and useful suggestions. It is a pleasure to thank the author of the detailed referee report; the suggestions therein greatly helped to improve the exposition.

2. Spaces and maps

2.1. Spaces and coordinates

In this paper we consider polygons $\mathbf{P} = (\dots P_i P_{i+1} \dots)$ in \mathbb{R}^2 that satisfy $[P_i, P_{i+1}] \neq 0$ for all i (when appropriate, the indices are understood cyclically). Denote by $\tilde{\mathcal{Y}}_n$ and $\tilde{\mathcal{X}}_n$ the spaces of twisted and closed n -gons, and by \mathcal{Y}_n and \mathcal{X}_n their quotient spaces by $\mathrm{SL}(2, \mathbb{R})$.

Let us introduce coordinates in \mathcal{Y}_n :

$$(2.1) \quad s_{2j-1} = [P_{j-1}, P_j], \quad v_{2j} = [P_{j-1}, P_{j+1}].$$

That is, s_{2j-1} are the areas subtended by the sides, and v_{2j} are the areas subtended by the short diagonals of the polygon.

One has a linear recursion

$$(2.2) \quad P_{j+1} = \frac{v_{2j}}{s_{2j-1}} P_j - \frac{s_{2j+1}}{s_{2j-1}} P_{j-1},$$

that is,

$$\begin{pmatrix} P_j \\ P_{j+1} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -\frac{s_{2j+1}}{s_{2j-1}} & \frac{v_{2j}}{s_{2j-1}} \end{pmatrix} \begin{pmatrix} P_{j-1} \\ P_j \end{pmatrix}.$$

It follows that the (conjugacy class of the) monodromy is given by

$$(2.3) \quad \mathcal{M}_{\mathbf{P}} = \begin{pmatrix} 0 & 1 \\ -\frac{s_1}{s_{2n-1}} & \frac{v_{2n}}{s_{2n-1}} \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ -\frac{s_5}{s_3} & \frac{v_4}{s_3} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -\frac{s_3}{s_1} & \frac{v_2}{s_1} \end{pmatrix}.$$

Note that $\det \mathcal{M}_{\mathbf{P}} = 1$.

Let $\mathbf{S} = (s_1, s_3, s_5, \dots, s_{2n-1})$; denote by $\tilde{\mathcal{Y}}_{n, \mathbf{S}}$ the space of twisted n -gons with $[P_i, P_{i+1}] = s_{2i+1}$ for all i , and let $\tilde{\mathcal{X}}_{n, \mathbf{S}}$ be the corresponding space for closed polygons.

Remark 2.1. — The spaces $\tilde{\mathcal{X}}_{n, \mathbf{S}}$ and $\mathcal{X}_{n, \mathbf{S}}$ are centroaffine analogs of the spaces of polygons with fixed side lengths, studied in [9].

When is $\tilde{\mathcal{X}}_{n, \mathbf{S}}$ a smooth n -dimensional manifold? One has a map $\tilde{\mathcal{X}}_n \rightarrow \mathbb{R}^n$ that sends \mathbf{P} to $\mathbf{S} = (\dots, [P_i, P_{i+1}], \dots)$. The next lemma describes the regular values \mathbf{S} of this map.

LEMMA 2.2. — *If n is odd and $s_j \neq 0$ for all j , then $\tilde{\mathcal{X}}_{n, \mathbf{S}}$ is smooth. If n is even and $s_1 s_5 s_9 \cdots \neq \pm s_3 s_7 s_{11} \cdots$, then $\tilde{\mathcal{X}}_{n, \mathbf{S}}$ is smooth.*

Proof. — If $P_i = (x_i, y_i)$ then $s_{2i+1} = x_i y_{i+1} - x_{i+1} y_i$. We need to know when the 1-forms ds_{2i+1} are linearly dependent.

We may assume that the coordinates are chosen so that all x_i and y_i are distinct from zero. Assume that $\sum a_i ds_i = 0$. This expands out as

$$\sum_i a_i (x_i dy_{i+1} + y_{i+1} dx_i - x_{i+1} dy_i - y_i dx_{i+1}) = 0,$$

where not all a_i vanish, hence

$$\sum_i (a_i y_{i+1} - a_{i-1} y_{i-1}) dx_i - (a_i x_{i+1} - a_{i-1} x_{i-1}) dy_i = 0.$$

Therefore

$$\frac{a_{i-1}}{a_i} = \frac{y_{i+1}}{y_{i-1}} = \frac{x_{i+1}}{x_{i-1}}, \quad i = 1, \dots, n,$$

(and hence all a_i are different from zero).

This implies that P_{i-1} and P_{i+1} are collinear for all i . If n is odd, then all vertices of \mathbf{P} are collinear, and $s_{2i+1} = 0$ for all i .

If n is even, then the odd vertices of \mathbf{P} are collinear, and so are the even vertices. Without loss of generality, assume that

$$P_1 = (x_1, 0), P_2 = (0, y_2), P_3 = (x_3, 0), P_4 = (0, y_4), \dots$$

Then

$$s_3 = x_1 y_2, s_5 = -y_2 x_3, s_7 = x_3 y_4, s_9 = -y_4 x_5 \dots$$

and $s_3 s_7 \cdots = \pm s_1 s_5 \cdots$. This completes the proof. \square

2.2. Monodromy of a twisted polygon

Set

$$a_j = \frac{v_{2j}}{s_{2j-1}}, \quad b_j = \frac{s_{2j+1}}{s_{2j-1}}.$$

Then

$$\mathcal{M}_n = \begin{pmatrix} 0 & 1 \\ -b_{n-1} & a_{n-1} \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ -b_1 & a_1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -b_0 & a_0 \end{pmatrix},$$

as in (2.3).

For $0 \leq i \leq j$, consider the continuants

$$(2.4) \quad D_{i,j+1} = \det \begin{pmatrix} a_i & 1 & 0 & \cdots & 0 \\ b_{i+1} & a_{i+1} & 1 & 0 & \cdots \\ 0 & b_{i+2} & a_{i+2} & 1 & \cdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \cdots & 0 & b_{j-1} & a_{j-1} & 1 \\ 0 & \cdots & 0 & b_j & a_j \end{pmatrix}.$$

We describe the monodromy of a twisted polygon in terms of these continuants.

Expanding by the last two rows, we see that

$$(2.5) \quad D_{i,j+1} = a_j D_{i,j} - b_j D_{i,j-1},$$

the same recursion as (2.2). Let us add the boundary conditions $D_{i,i} = 1$, $D_{i,i-1} = 0$.

PROPOSITION 2.3. — *One has*

$$\mathcal{M}_n = \begin{pmatrix} -b_0 D_{1,n-1} & D_{0,n-1} \\ -b_0 D_{1,n} & D_{0,n} \end{pmatrix}.$$

Proof. — Induction on n . For $n = 1$, the claim holds due to the boundary conditions $D_{1,0} = 0$, $D_{0,0} = D_{1,1} = 1$, $D_{0,1} = a_0$.

Next,

$$\mathcal{M}_{n+1} = \begin{pmatrix} 0 & 1 \\ -b_n & a_n \end{pmatrix} \mathcal{M}_n,$$

and the result follows from the recurrence (2.5). \square

If \mathbf{P} is a twisted n -gon, then the sequences a_i and b_i are n -periodic and $\prod_i b_i = 1$. Proposition 2.3 implies the following statement.

COROLLARY 2.4. — *For a closed n -gon, the coordinates (s_{2i-1}, v_{2i}) satisfy $D_{i,n+i-1} = 0$ for all i .*

In fact, any three of these identities imply the rest (the codimension of the space of closed polygons is three).

Example 2.5. — Consider the case $n = 3$. Corollary 2.4 implies

$$v_0v_2 = s_3s_5, \quad v_2v_4 = s_5s_1, \quad v_4v_0 = s_1s_3,$$

hence $v_0^2 = s_3^2$. We have two solutions;

$$v_0 = -s_3, \quad v_2 = -s_5, \quad v_4 = -s_1, \quad \text{and} \quad v_0 = s_3, \quad v_2 = s_5, \quad v_4 = s_1.$$

The first one corresponds to a closed triangle (the monodromy is Id), and the second one to a centrally symmetric hexagon (the monodromy is $-\text{Id}$).

Next, consider the case $n = 4$. Corollary 2.4 implies

$$v_0v_2v_4 = s_3s_7v_4 + s_1s_5v_0$$

and its three cyclic permutations. Rewrite it as

$$\frac{s_3s_7}{v_0v_2} + \frac{s_1s_5}{v_2v_4} = 1$$

and its cyclic permutations. This is a system of four linear equations on the variables

$$\frac{1}{v_0v_2}, \quad \frac{1}{v_2v_4}, \quad \frac{1}{v_4v_6}, \quad \frac{1}{v_6v_0}$$

with coefficients s_3s_7 and s_1s_5 . This system implies

$$\frac{1}{v_0v_2} = \frac{1}{v_4v_6}, \quad \frac{1}{v_2v_4} = \frac{1}{v_6v_0},$$

and hence $v_4^2 = v_0^2$. As before, one has two choices of signs, one corresponding to the monodromy Id, and another to $-\text{Id}$. In the former case of closed quadrilaterals, one has

$$v_4 = -v_0, \quad v_6 = -v_2 \quad \text{and} \quad v_0v_2 = s_7s_3 - s_1s_5,$$

the latter being the Ptolemy–Plücker relation.

2.3. Lax transformation is fractional-linear

Given a non-zero vector P_i , the i th vertex of a polygon \mathbf{P} , the vectors Q with $[P_i, Q] = c$ comprise a line L_{P_i} parallel to P_i . Identify L_{P_i} with \mathbb{R} by parameterizing it as

$$\frac{cP_{i+1}}{[P_i, P_{i+1}]} + tP_i, \quad t \in \mathbb{R}.$$

Let $\mathbf{P} = (P_1, \dots, P_n)$ be a closed n -gon. As described in Introduction, choose a test vector Q_1 with $[P_1, Q_1] = c$, and consecutively construct vectors Q_2, \dots, Q_n, Q_{n+1} according to (1.2), which we recall here:

$$[P_i, P_{i+1}] = [Q_i, Q_{i+1}], \quad [P_i, Q_i] = c, \quad i = 1, \dots, n.$$

For this to work, that is, for each next point Q_{i+1} to be defined, we need the lines given by the equations $[Q_i, Q_{i+1}] = [P_i, P_{i+1}]$ and $[P_{i+1}, Q_{i+1}] = c$ to intersect. This is the case if the vectors P_{i+1} and Q_i are not parallel. This is a general position condition that we assume to hold for all i .

We use the notation

$$\mathcal{L}_{P_1 P_2, c}(Q_1) = Q_2, \mathcal{L}_{P_2 P_3, c}(Q_2) = Q_3, \dots$$

(we omit c from the notation when it does not lead to confusion). The Lax transformation $\mathcal{L}_{\mathbf{P}, c} : L_{P_1} \rightarrow L_{P_1}$ is the composition of these maps. The polygonal line \mathbf{Q} closes up, that is, $Q_{n+1} = Q_1$, if and only if Q_1 is a fixed point of $\mathcal{L}_{\mathbf{P}, c} : \mathbb{R} \rightarrow \mathbb{R}$. Then we have $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$.

Likewise, if \mathbf{P} is a twisted n -gon with monodromy $\mathcal{M}_{\mathbf{P}}$, then $\mathcal{M}_{\mathbf{P}}$ sends L_{P_1} to $L_{P_{n+1}}$ and, as a map $\mathbb{R} \rightarrow \mathbb{R}$, it is the identity. Hence it is still true that the fixed points of $\mathcal{L}_{\mathbf{P}, c}$ give rise to twisted polygons \mathbf{Q} such that $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$.

LEMMA 2.6. — *The Lax transformations are fractional-linear.*

Proof. — Denote by R_{Q_1, P_2} the centroaffine reflection that interchanges Q_1 and P_2 . This map is given by the formula

$$(2.6) \quad R_{Q_1, P_2}(X) = \frac{[Q_1, X]Q_1 + [X, P_2]P_2}{[Q_1, P_2]},$$

and one has $Q_2 = R_{Q_1, P_2}(P_1)$, that is, $\mathcal{L}_{P_1 P_2}(Q_1) = R_{Q_1, P_2}(P_1)$.

Using the identifications of L_{P_1} and L_{P_2} with \mathbb{R} , the map $\mathcal{L}_{P_1 P_2} : Q_1 \mapsto Q_2$ becomes

$$t \mapsto -\frac{c[P_1, P_3]}{[P_1, P_2][P_2, P_3]} + \frac{[P_1, P_2]^2 - c^2}{t[P_1, P_2]^2} = -\frac{cv_4}{s_3 s_5} + \frac{s_3^2 - c^2}{t s_3^2},$$

a fractional-linear transformation. Hence $\mathcal{L}_{\mathbf{P}, c}$ is fractional-linear as well. \square

The above fractional-linear transformation is represented by the matrix

$$(2.7) \quad \begin{pmatrix} -\frac{cv_4}{s_3 s_5} & 1 - \frac{c^2}{s_3^2} \\ 1 & 0 \end{pmatrix},$$

its determinant equals $\frac{c^2}{s_3^2} - 1$.

If the ground field is \mathbb{C} , a fractional-linear transformation has two fixed points, perhaps, coinciding (unless it is the identity). Over the reals, this number is 0, 1, 2, or ∞ .

Over \mathbb{C} , Lemma 2.6 makes it possible to consider c -relation as a map (defined in a Zariski open set of polygons): start with \mathbf{P} and choose one

of the two c -related polygons, say \mathbf{Q}_1 . This polygon also has two c -related ones, one of which is $-\mathbf{P}$, the polygon that is centrally symmetric to \mathbf{P} . Choose the other polygon c -related to \mathbf{Q}_1 and continue in a similar way. If at the beginning we choose \mathbf{Q}_2 instead of \mathbf{Q}_1 then $-\mathbf{Q}_2 \stackrel{c}{\sim} \mathbf{P}$, that is, up to central symmetry, we obtain the inverse of the same map.

Over \mathbb{R} , a polygon may have no c -related ones. To have a map, one needs to assume that a c -related polygon exists. We shall see in Section 3.1 that the c -related polygons have conjugated Lax transformations. This makes it possible to continue in the same way as in the complex case.

2.4. Centroaffine butterflies

We show that centroaffine butterflies have trivial Lax transformations for all c and classify all such quadrilaterals. Here and in what follows we say that two vertices are not collinear if they do not lie on the same line through the origin.

LEMMA 2.7. — *The Lax transformation for a quadrilateral is the identity for every c if and only if the quadrilateral is a centroaffine butterfly, or it is obtained from a centroaffine butterfly by reflecting one of the vertices in the origin (an “anti-butterfly”), or its two opposite vertices are symmetric with respect to the origin.*

Proof. — Assume first that a pair of opposite vertices of a quadrilateral \mathbf{P} is not collinear. Applying a linear transformation, assume that

$$P_1 = (1, 0), P_2 = (a, b), P_3 = (0, 1), P_4 = (u, v), Q_1 = (x, y),$$

so $y = c$. Applying equation (2.6) twice, we find

$$Q_2 = \frac{1}{bx - ay}(ab - xy, b^2 - y^2), Q_3 = \left(-y, \frac{-y(a^2 + b^2) + abx + y^3}{ab - xy} \right).$$

Going in the opposite direction, that is, replacing (a, b) with (u, v) , we obtain point

$$\left(-y, \frac{-y(u^2 + v^2) + uvx + y^3}{uv - xy} \right).$$

This point coincides with Q_3 for all x , which implies

$$ab = uv, a^2 + b^2 = u^2 + v^2.$$

Hence either $u = b, v = a$, a butterfly, or $u = -b, v = -a$, an anti-butterfly, or $u = -a, v = -b$, symmetric opposite vertices.

It remains to consider the case when both pairs of opposite vertices of \mathbf{P} are collinear. Then one may assume that

$$P_1 = (1, 0), P_2 = (0, 1), P_3 = (a, 0), P_4 = (0, b), Q_1 = (x, y).$$

Using equation (2.6) consecutively, one calculates

$$Q_5 = \left(\frac{x(a^2 - y^2)(b^2 - y^2)}{(a^2b^2 - y^2)(1 - y^2)}, y \right).$$

Equating it to Q_1 yields

$$(a^2 - y^2)(b^2 - y^2) = (a^2b^2 - y^2)(1 - y^2),$$

hence $(a^2 - 1)(b^2 - 1) = 0$. If $a = 1$ or $b = 1$, then $P_1 = P_3$ or $P_2 = P_4$, and if $a = -1$ or $b = -1$, we have the already considered symmetric case. \square

Thus if \mathbf{P} is a centroaffine butterfly, then for every choice of the seed vertex Q_1 , the quadrilateral \mathbf{Q} closes up. In fact, it is a centroaffine butterfly as well.

LEMMA 2.8. — *If \mathbf{P} is a centroaffine butterfly and \mathbf{Q} is c -related to it, then \mathbf{Q} is also a butterfly, see Figure 2.1.*




Figure 2.1. Two centroaffine butterflies are c -related.

Proof. — We need to show that

$$[Q_1, Q_2] = [Q_4, Q_3], [Q_1, Q_4] = [Q_2, Q_3].$$

By definition of c -relation,

$$[Q_1, Q_2] = [P_1, P_2], [Q_4, Q_3] = [P_4, P_3],$$

and $[P_1, P_2] = [P_4, P_3]$ because \mathbf{P} is a butterfly. This implies the first of the claimed equalities. The second follows in the same way. \square

2.5. Constructing c -related polygons

In this section we describe a construction that yields a pair of c -related n -gons. This is a centroaffine analog of a construction that yields pairs of polygons in the discrete bicycle correspondence that is described in [15].

Assume first that n is odd. Start with an n -gon (pentagon \mathbf{A} in Figure 2.2). Connect the midpoints of its sides with the origin. Consider the affine reflections in these lines that interchange the vertices of the respective sides of \mathbf{A} . Let R be the composition of these reflections taken around the polygon.




Figure 2.2. Constructing a c -related pair of n -gons, odd n .

LEMMA 2.9. — *The map R is an affine reflection.*

Proof. — Since n is odd, R is orientation reversing and its determinant equals -1 . Hence it has two eigendirections. Also R has a fixed point, a vertex of \mathbf{A} . Therefore the eigenvalues of R are 1 and -1 , and it is an affine reflection. \square

It follows that $R^2 = \text{Id}$. The construction of a c -related pair follows.

Start with an arbitrary point P_1 and apply consecutive affine reflections around the polygon \mathbf{A} twice. This produces $2n$ points

$$P_1 \mapsto Q_2 \mapsto P_3 \mapsto Q_4 \mapsto \cdots \mapsto Q_n,$$

see Figure 2.2. Each quadrilateral $P_i P_{i+1} Q_i Q_{i+1}$ is a centroaffine butterfly, therefore $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$.

One has $c = [P_1, Q_1] = [P_1, R(P_1)]$. The locus of points P_1 for which $[P_1, R(P_1)]$ is fixed is a hyperbola (indeed, R is conjugated to the map $(x, y) \mapsto (-x, y)$, in which case this claim is obvious).

Now assume that n is even. We repeat the above construction, but this time the transformation R has determinant 1. It still has an eigendirection with eigenvalue 1 (a vertex of \mathbf{A} is a fixed point), but it is not necessarily the identity. We need to assume that $R = \text{Id}$, see below.




Figure 2.3. Constructing a c -related pair of n -gons, even n .

With this assumption, we choose two starting points, P_1 and Q_1 and apply consecutive affine reflections to obtain polygons $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$ with $c = [P_1, Q_1]$ as before, see Figure 2.3.

We describe when the transformation R is the identity. Recall that we consider the case of an even-gon.

LEMMA 2.10. — One has $R = \text{Id}$ if and only if

$$\sum_{i=1}^n (-1)^i \frac{v_{2i}}{s_{2i-1} s_{2i+1}} = 0,$$

where the coordinates v_{2i}, s_{2i+1} are associated with the polygon \mathbf{A} .

Proof. — Since R has a fixed point, a vertex of polygon \mathbf{A} , the transformation R is the identity if and only if it has another fixed point not

collinear with the first one. Thus we need to learn when a polygon \mathbf{B} exists, not homothetic to \mathbf{A} , whose sides are parallel to those of \mathbf{A} and the midpoints of whose sides are collinear to those of \mathbf{A} .

These conditions are written as

$$[A_{i+1} + A_i, B_{i+1} + B_i] = 0 \quad \text{and} \quad [A_{i+1} - A_i, B_{i+1} - B_i] = 0,$$

or

$$(2.8) \quad [A_i, B_i] + [A_{i+1}, B_{i+1}] = 0, \quad [A_i, B_{i+1}] + [A_{i+1}, B_i] = 0.$$

We will look for points B_i written as $B_i = a_i A_{i-1} + b_i A_i$. Then $B_{i+1} = a_{i+1} A_i + b_{i+1} A_{i+1}$. Taking cross-products with A_i and A_{i+1} , we also find that $B_i = b_{i+1} A_i + a_{i+1} A_{i+1}$. Since

$$[A_{i-1}, A_i] = s_{2i-1}, \quad [A_i, A_{i+1}] = s_{2i+1}, \quad [A_{i-1}, A_{i+1}] = v_{2i},$$

we also have

$$a_{i+1} s_{2i+1} = -a_i s_{2i-1}, \quad (b_{i+1} - b_i) s_{2i+1} = a_i v_{2i}.$$

It follows that $a_i = (-1)^i t / s_{2i-1}$ for some non-zero constant t (non-zero since \mathbf{B} is not homothetic to \mathbf{A}), and

$$b_{i+1} - b_i = t \frac{(-1)^i v_{2i}}{s_{2i-1} s_{2i+1}}.$$

This system of linear equations on b_i has a solution if and only if the sum of the right hand sides vanishes, as needed. \square

We add that the above described construction is a general method of constructing c -related pairs. Indeed, a pair of c -related n -gons with odd n gives rise to an affine reflection R , as described above and illustrated in Figure 2.2. Then one can chose a fixed point of R as a vertex of n -gon \mathbf{A} and obtain the other vertices by consecutive affine reflections. A similar remark applies to the case of even n as well.

2.6. Bianchi permutability

In the following theorem, the polygons are either closed or twisted.

THEOREM 2.11. — *Assume that $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$ and $\mathbf{P} \stackrel{d}{\sim} \mathbf{R}$. Then there exists a polygon \mathbf{S} such that $\mathbf{Q} \stackrel{d}{\sim} \mathbf{S}$ and $\mathbf{R} \stackrel{c}{\sim} \mathbf{S}$.*




Figure 2.4. *Bianchi permutability: one can see four butterflies in a coordinated “flight”.*

Proof. — Recall that \mathcal{L} is the Lax transformation. The idea of the proof is to define polygon \mathbf{S} such that $P_i Q_i S_i R_i$ is a centroaffine butterfly for all i and to show that this polygon is a Lax c -orbit with respect to \mathbf{R} and a Lax d -orbit with respect to \mathbf{Q} .

Thus we define point S_1 by requiring $P_1 Q_1 S_1 R_1$ to be a centroaffine butterfly. Then $[R_1, S_1] = c, [Q_1, S_1] = d$. Let $\mathcal{L}_{Q_1 Q_2, d}(S_1) = S_2$ and $\mathcal{L}_{R_1 R_2, c}(S_1) = S'_2$. We claim that $S_2 = S'_2$ and that $P_2 Q_2 S_2 R_2$ is a centroaffine butterfly.

Indeed, one has

$$\mathcal{L}_{S_1 R_1}(S'_2) = R_2, \mathcal{L}_{R_1 P_1}(R_2) = P_2, \mathcal{L}_{P_1 Q_1}(P_2) = Q_2, \mathcal{L}_{Q_1 S_1}(Q_2) = S_2.$$

Since $P_1 Q_1 S_1 R_1$ is a centroaffine butterfly, Lemma 2.7 implies that the quadrilateral $S'_2 R_2 P_2 Q_2 S_2$ closes up and that it is a centroaffine butterfly.

This shows that one can define the polygon \mathbf{S} so that $P_i Q_i S_i R_i$ is a centroaffine butterfly for all i . \square

We leave it to the reader to make sure that Theorem 2.11 can be interpreted as a configuration theorem depicted in Figure 2.4.

3. Integrals

3.1. Invariance of the Lax transformations

The next result provide a Lax presentation of the c -relation.

THEOREM 3.1. — *If \mathbf{P} and \mathbf{Q} are c -related twisted n -gons, then the Lax transformations $\mathcal{L}_{\mathbf{P},\lambda}$ and $\mathcal{L}_{\mathbf{Q},\lambda}$ are conjugated for every value of the parameter λ .*

Proof. — We know that $P_i P_{i+1} Q_{i+1} Q_i$ is a centroaffine butterfly. Lemma 2.7 shows that the product of the Lax transformations along its sides is trivial. This can be rearranged to show that $\mathcal{L}_{P_i P_{i+1}}$ and $\mathcal{L}_{Q_i Q_{i+1}}$ are conjugate:

$$\mathcal{L}_{Q_i Q_{i+1},\lambda} = \mathcal{L}_{P_{i+1} Q_{i+1},\lambda} \mathcal{L}_{P_i P_{i+1},\lambda} \mathcal{L}_{P_i Q_i,\lambda}^{-1}.$$

Taking composition over $i = 1, \dots, n$ yields the result. □

Since $\mathcal{L}_{\mathbf{P},\lambda}$ is a fractional-linear transformation, one can realize it as a 2×2 matrix that, abusing notation, we also call \mathcal{L} . Then $(\text{Tr } \mathcal{L})^2 / \det \mathcal{L}$ is a well-defined conjugacy invariant function. It depends on λ , and expanding in a Taylor series in λ , one obtains integrals of the c -relation.

Note that the determinant of the Lax transformation, $\det \mathcal{L}$, depends only on the variables s_{2i-1} (and, of course, λ), see formula 2.7. This determinant is itself an integral, therefore we may – and will – use $\text{Tr } \mathcal{L}$, rather than $(\text{Tr } \mathcal{L})^2 / \det \mathcal{L}$, to obtain integrals of the c -relation.

Example 3.2. — Consider the case of twisted triangles. Let v_2, v_4, v_6 and s_1, s_3, s_5 be the respective coordinates in the moduli space. Consider the space $\mathcal{Y}_{3,\mathbf{s}}$.

Multiply the three matrices (2.7) (replacing c by λ) to find the Lax matrix \mathcal{L} depending on this spectral parameter λ . One has

$$\text{Tr } \mathcal{L} = \lambda^3 \left(\frac{v_2 v_4 v_6}{s_1 s_3 s_5} - \frac{v_2}{s_5} - \frac{v_4}{s_1} - \frac{v_6}{s_3} \right) + \lambda(v_2 s_5 + v_4 s_1 + v_6 s_3),$$

and

$$\det \mathcal{L} = (\lambda^2 - s_1^2)(\lambda^2 - s_3^2)(\lambda^2 - s_5^2).$$

The determinant does not depend on the v -variables, and since the s -variables are preserved by the c -relation, the trace is invariant. We obtain two integrals:

$$I = \frac{v_2 v_4 v_6}{s_1 s_3 s_5} - \left(\frac{v_2}{s_5} + \frac{v_4}{s_1} + \frac{v_6}{s_3} \right)$$

and

$$J = \frac{v_2}{s_1 s_3} + \frac{v_4}{s_3 s_5} + \frac{v_6}{s_5 s_1}.$$

Thus $\mathcal{Y}_{3,\mathbf{s}}$ is foliated by the common level curves of the functions I and J .

For comparison, let \mathcal{M} be the monodromy of the twisted triangle. Using formula (2.3), we find that $\text{Tr } \mathcal{M} = I$.

Example 3.3. — Similarly, in the case of twisted quadrilaterals, we have variables v_2, v_4, v_6, v_8 and parameters s_1, s_3, s_5, s_7 . The trace of the Lax matrix is a biquadratic polynomial in λ , and its three coefficients are integrals. The free term is $2s_1s_3s_5s_7$, a constant on $\mathcal{Y}_{4,\mathbf{s}}$, and the other two terms give two integrals

$$I = \frac{v_2v_4v_6v_8}{s_1s_3s_5s_7} - \left(\frac{v_2v_4}{s_3s_7} + \frac{v_4v_6}{s_5s_1} + \frac{v_6v_8}{s_7s_3} + \frac{v_8v_2}{s_1s_5} \right)$$

and

$$J = \left(\frac{v_2v_4}{s_1s_3^2s_5} + \frac{v_4v_6}{s_3s_5^2s_7} + \frac{v_6v_8}{s_5s_7^2s_1} + \frac{v_8v_2}{s_7s_1^2s_3} \right)$$

(omitting the terms that do not depend on the v -variables). Thus $\mathcal{Y}_{4,\mathbf{s}}$ is foliated by the common level surfaces of the functions I and J . Note that again one has $\text{Tr } \mathcal{M} = I$.

If the quadrilateral is closed, according to Example 2.5, we have

$$v_6 = -v_2, v_8 = -v_4, v_2v_4 = s_1s_5 - s_3s_7,$$

and both integrals, I and J , become functions of the parameters s_j only.

Example 3.4. — Consider the case of twisted pentagons. As before, we calculate $\text{Tr } \mathcal{L}$ and decompose it in homogeneous components in the spectral parameter λ . There are three terms, of degrees 1, 3, and 5. This gives three integrals:

$$I = \frac{v_2v_4v_6v_8v_{10}}{s_1s_3s_5s_7s_9} - \left(\frac{v_2v_4v_6}{s_3s_5s_9} + \frac{v_4v_6v_8}{s_5s_7s_1} + \frac{v_6v_8v_{10}}{s_7s_9s_3} + \frac{v_8v_{10}v_2}{s_9s_1s_5} + \frac{v_{10}v_2v_4}{s_1s_3s_7} \right) \\ + \left(\frac{v_2s_7}{s_5s_9} + \frac{v_4s_9}{s_7s_1} + \frac{v_6s_1}{s_9s_3} + \frac{v_8s_3}{s_1s_5} + \frac{v_{10}s_5}{s_3s_7} \right)$$

(this also comes from the trace of the monodromy),

$$J = \frac{v_2v_4v_6}{s_1s_3^2s_5^2s_7} + \frac{v_4v_6v_8}{s_3s_5^2s_7^2s_9} + \frac{v_6v_8v_{10}}{s_5s_7^2s_9^2s_1} + \frac{v_8v_{10}v_2}{s_7s_9^2s_1^2s_3} + \frac{v_{10}v_2v_4}{s_9s_1^2s_3^2s_5} \\ - \frac{v_2}{s_1s_3} \left(\frac{1}{s_5^2} + \frac{1}{s_9^2} \right) - \frac{v_4}{s_3s_5} \left(\frac{1}{s_7^2} + \frac{1}{s_1^2} \right) - \frac{v_6}{s_5s_7} \left(\frac{1}{s_9^2} + \frac{1}{s_3^2} \right) \\ - \frac{v_8}{s_7s_9} \left(\frac{1}{s_1^2} + \frac{1}{s_5^2} \right) - \frac{v_{10}}{s_9s_1} \left(\frac{1}{s_3^2} + \frac{1}{s_7^2} \right).$$

and

$$K = \frac{v_2}{s_1s_3} + \frac{v_4}{s_3s_5} + \frac{v_6}{s_5s_7} + \frac{v_8}{s_7s_9} + \frac{v_{10}}{s_9s_1}.$$

The common level surfaces of these integrals foliate the 5-dimensional space $\mathcal{Y}_{5,\mathbf{s}}$.

In the case of closed pentagons, we have the Ptolemy–Plücker relations

$$v_2 v_4 = s_1 s_5 - s_3 v_8$$

and its cyclic permutations. This makes the integrals functionally dependent, and leaves a single integral K on the 2-dimensional space $\mathcal{X}_{5,\mathbf{s}}$.

The formulas of Examples 3.2, 3.3, and 3.4 are extended to arbitrary values of n in Sections 3.4 and 3.5 below.

3.2. Centroaffine polygon recutting

As mentioned in Introduction, by the elementary recutting of a closed polygon \mathbf{P} we mean the linear transformation that changes only one vertex: $P'_j = R_{P_{j-1}P_{j+1}}(P_j)$. Denote this transformation by R_j . Elementary recuttings are involutions. The recutting R of the polygon \mathbf{P} is the composition $R_n \circ \dots \circ R_1$.

The next lemma is a centroaffine analog of [1, Corollary 2].

LEMMA 3.5. — *Let R_j be the elementary recutting at the vertex j . Then $(R_j R_{j+1})^3 = \text{Id}$, $R_j^2 = \text{Id}$ and $R_j R_k = R_k R_j$ for $|k - j| \geq 2$.*

Proof. — The only non-trivial fact is the $(R_j R_{j+1})^3 = \text{Id}$. We have two approaches to the proof. The first approach consists in normalizing two vertices involved to be the standard basis in \mathbb{R}^2 , to denote the other vertices using variables, such as $(a, b), (c, d)$, etc., and to work out the necessary calculations in Mathematica (or another CAS). The calculations are not too complicated.

The alternative, hand calculation based, approach is presented below.

Consider Figure 3.1 that depicts the configuration theorem described by this identity. One fixes the frame made of the vectors P_{j-1} and P_{j+2} .




Figure 3.1. To Lemma 3.5.

Then we have six points $P_j, P'_j, P''_j, P_{j+1}, P'_{j+1}, P''_{j+1}$ that should satisfy 12 relations (cf. (1.2)):

$$\begin{aligned} [P_{j-1}, P_j] &= [P'_j, P_{j+1}], & [P_{j-1}, P'_j] &= [P_j, P_{j+1}], \\ [P_{j-1}, P'_j] &= [P''_j, P'_{j+1}], & [P_{j-1}, P''_j] &= [P'_j, P'_{j+1}], \\ [P_{j-1}, P''_j] &= [P_j, P''_{j+1}], & [P_{j-1}, P_j] &= [P''_j, P''_{j+1}], \\ [P''_{j+1}, P_{j+2}] &= [P_j, P_{j+1}], & [P_{j+1}, P_{j+2}] &= [P_j, P''_{j+1}], \\ [P_{j+1}, P_{j+2}] &= [P'_j, P'_{j+1}], & [P'_{j+1}, P_{j+2}] &= [P'_j, P_{j+1}], \\ [P'_{j+1}, P_{j+2}] &= [P''_j, P''_{j+1}], & [P''_{j+1}, P_{j+2}] &= [P''_j, P'_{j+1}]. \end{aligned}$$

This system is equivalent to the system of 9 relations

$$(3.1) \quad \begin{aligned} [P_{j-1}, P_j] &= [P'_j, P_{j+1}] = [P''_j, P''_{j+1}] = [P'_{j+1}, P_{j+2}] := x, \\ [P_{j-1}, P'_j] &= [P''_j, P'_{j+1}] = [P_j, P_{j+1}] = [P''_{j+1}, P_{j+2}] := y, \\ [P_{j-1}, P''_j] &= [P_j, P''_{j+1}] = [P'_j, P'_{j+1}] = [P_{j+1}, P_{j+2}] := z. \end{aligned}$$

Assume, without loss of generality, that $[P_{j-1}, P_{j+2}] = 1$ and express the remaining points as linear combinations of P_{j-1} and P_{j+2} :

$$\begin{aligned} P_j &= \lambda P_{j-1} + x P_{j+2}, & P'_j &= \lambda_1 P_{j-1} + y P_{j+2}, \\ P''_j &= \lambda_2 P_{j-1} + z P_{j+2}, & P_{j+1} &= \mu P_{j+2} + z P_{j-1}, \\ P'_{j+1} &= \mu_1 P_{j+2} + x P_{j-1}, & P''_{j+1} &= \mu_2 P_{j+2} + y P_{j-1}. \end{aligned}$$

Then the rest of the equations (3.1) become the following six equations on the nine variables $x, y, z, \lambda_i, \mu_i, i = 1, 2, 3$:

$$\lambda_1 \mu = x + yz = \lambda_2 \mu_2, \lambda_2 \mu_1 = y + xz = \lambda \mu, \lambda \mu_2 = z + xy = \lambda_1 \mu_1.$$

These equations are not independent: the product of the three left hand sides equals the product of the three right hand sides.

Overall, one has nine variables satisfying five relations. The resulting four degrees of freedom make it possible to choose point P_j and P_{j+1} arbitrarily, proving the existence of the configuration of the lemma. \square

Remark 3.6. — The relations of Lemma 3.5 show that one has a representation of the group of permutations S_{n+1} on the space of centroaffine n -gons, similar to the situation with the Euclidean polygon recutting, see [1].

The next lemma states that elementary recuttings commute with the c -relations (an analog of one of the statements of Theorem 4 in [15]).

LEMMA 3.7. — *If $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$ then $R_j(\mathbf{P}) \stackrel{c}{\sim} R_j(\mathbf{Q})$ for all j .*

Proof. — The situation with the proof of this lemma is similar to that of the preceding lemma: one can delegate the calculation to CAS, or can do it by hand. We did both, and the latter approach is presented below.

One only has to check that $[P'_j, Q'_j] = c$. From (2.2) it follows that

$$P_j = \frac{s_{2j+1}(\mathbf{P})}{v_{2j}(\mathbf{P})} P_{j-1} + \frac{s_{2j-1}(\mathbf{P})}{v_{2j}(\mathbf{P})} P_{j+1}.$$

Since $[P'_j, P_{j+1}] = [P_{j-1}, P_j]$ and $[P_{j-1}, P'_j] = [P_j, P_{j+1}]$, one has

$$(3.2) \quad P'_j = \frac{s_{2j-1}(\mathbf{P})}{v_{2j}(\mathbf{P})} P_{j-1} + \frac{s_{2j+1}(\mathbf{P})}{v_{2j}(\mathbf{P})} P_{j+1}.$$

Likewise, for \mathbf{Q} one has

$$Q_j = \frac{s_{2j+1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} Q_{j-1} + \frac{s_{2j-1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} Q_{j+1},$$

and

$$Q'_j = \frac{s_{2j-1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} Q_{j-1} + \frac{s_{2j+1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} Q_{j+1}.$$

Since $\mathbf{Q} \stackrel{c}{\sim} \mathbf{P}$ we have

$$\begin{aligned} c = [P_j, Q_j] &= \left(\frac{s_{2j+1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j+1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} + \frac{s_{2j-1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j-1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} \right) c \\ &\quad + \frac{s_{2j+1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j-1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} [P_{j-1}, Q_{j+1}] \\ &\quad + \frac{s_{2j-1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j+1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} [P_{j+1}, Q_{j-1}]. \end{aligned}$$

For $[P'_j, Q'_j]$, we have

$$\begin{aligned} [P'_j, Q'_j] &= \left(\frac{s_{2j+1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j+1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} + \frac{s_{2j-1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j-1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} \right) c \\ &\quad + \frac{s_{2j-1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j+1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} [P_{j-1}, Q_{j+1}] \\ &\quad + \frac{s_{2j+1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j-1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} [P_{j+1}, Q_{j-1}]. \end{aligned}$$

But

$$\begin{aligned} \frac{s_{2j-1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j+1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})} &= \frac{[P_{j-1}, P_j][Q_j, Q_{j+1}]}{[P_{j-1}, P_{j+1}][Q_{j-1}, Q_{j+1}]} \\ &= \frac{[Q_{j-1}, Q_j][P_j, P_{j+1}]}{[P_{j-1}, P_{j+1}][Q_{j-1}, Q_{j+1}]} = \frac{s_{2j+1}(\mathbf{P})}{v_{2j}(\mathbf{P})} \frac{s_{2j-1}(\mathbf{Q})}{v_{2j}(\mathbf{Q})}. \end{aligned}$$

Hence the result. \square

Lemma 3.7 also can be interpreted as a configuration theorem, see Figure 3.2. As before, we leave it to the reader to make sure of it.




Figure 3.2. Elementary recutting commutes with c -relation.

THEOREM 3.8.

- (1) The Lax transformation is preserved by recutting;
- (2) Recutting commutes with c -relations.

Proof. — The first statement follows from Lemma 2.7:

$$\mathcal{L}_{P_i P_{i+1}} \mathcal{L}_{P_{i-1} P_i} = \mathcal{L}_{P'_i P'_{i+1}} \mathcal{L}_{P_{i-1} P'_i}$$

for all i . The second statement follows from Lemma 3.7: if $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$ then $\mathbf{R}_1(\mathbf{P}) \stackrel{c}{\sim} \mathbf{R}_1(\mathbf{Q})$, therefore $\mathbf{R}_2 \mathbf{R}_1(\mathbf{P}) \stackrel{c}{\sim} \mathbf{R}_2 \mathbf{R}_1(\mathbf{Q})$, etc. \square

3.3. Odd-gons, infinitesimal map

If the constant c is infinitesimal, we obtain an $\mathrm{SL}(2, \mathbb{R})$ -invariant vector field on the space of twisted polygons, that is, a vector field on the moduli space \mathcal{Y}_n . In this section we calculate this field.

Let \mathbf{P} be a twisted n -gon with odd n . Let the field be given by vectors ξ_i with foot points P_i , $i = 1, \dots, n$. The conditions $[P_i, Q_i] = c$, $[P_i, P_{i+1}] = [Q_i, Q_{i+1}]$ become

$$(3.3) \quad [P_i, \xi_i] = 1, \quad [P_i, \xi_{i+1}] + [\xi_i, P_{i+1}] = 0$$

(we normalize the field so the constant in the first equation is 1).

THEOREM 3.9. — *In terms of the (v, s) -coordinates, the field ξ is given by*

$$\dot{v}_{2i} = v_{2i} \left(\sum_{k=1}^{n-1} (-1)^{k-1} \frac{v_{2i+2k}}{s_{2i+2k-1} s_{2i+2k+1}} \right) + \frac{s_{2i+1}}{s_{2i-1}} - \frac{s_{2i-1}}{s_{2i+1}}.$$

Proof. — Let $\xi_i = a_i P_i + b_i P_{i+1}$. Then (3.3) implies

$$b_i s_{2i+1} = 1, (a_i + a_{i+1}) s_{2i+1} + b_{i+1} v_{2i+2} = 0.$$

Since n is odd, there is a unique solution

$$(3.4) \quad a_i = \frac{1}{2} \sum_{k=1}^n (-1)^k \frac{v_{2i+2k}}{s_{2i+2k-1} s_{2i+2k+1}}, \quad b_i = \frac{1}{s_{2i+1}}.$$

Since $v_{2i} = [P_{i-1}, P_{i+1}]$, we have

$$(3.5) \quad \begin{aligned} \dot{v}_{2i} &= [P_{i-1}, \xi_{i+1}] + [\xi_{i-1}, P_{i+1}] \\ &= [P_{i-1}, a_{i+1} P_{i+1} + b_{i+1} P_{i+2}] + [a_{i-1} P_{i-1} + b_{i-1} P_i, P_{i+1}] \\ &= (a_{i-1} + a_{i+1}) v_{2i} + b_{i+1} [P_{i-1}, P_{i+2}] + b_{i-1} s_{2i+1}. \end{aligned}$$

By the Prolomy–Plücker relation,

$$[P_{i-1}, P_{i+2}] = \frac{v_{2i} v_{2i+2} - s_{2i-1} s_{2i+3}}{s_{2i+1}},$$

and the values of a_i and b_i are found in (3.4). Substitute in (3.5) to obtain the result. \square

It follows from Theorems 2.11 and 3.8 that the flow of the field ξ commutes with the c -relations and with the polygon recutting.

Remark 3.10. — If n is even, then a necessary condition for a_i to exist is

$$\sum_{i=1}^n (-1)^n \frac{v_{2i}}{s_{2i-1} s_{2i+1}} = 0,$$

the equation that appeared in Lemma 2.10. If this necessary condition holds, the vector field is defined modulo a 1-dimensional space, the kernel of the matrix that gives the linear equations on the variables a_i .

Set

$$g_i = \frac{v_{2i}}{s_{2i-1} s_{2i+1}}, \quad \beta_i = -\frac{1}{s_{2i-1}^2}.$$

Then, after scaling, the vector field ξ is given by

$$\dot{g}_i = -g_i (g_{i+1} - g_{i+2} + g_{i+3} - \cdots - g_{i+n-1}) + \beta_i - \beta_{i+1}.$$

This is the dressing chain of Veselov–Shabat, formula (12) in [17].

3.4. Integrals for twisted polygons

In view of Theorem 3.1, we calculate the trace of the Lax matrix \mathcal{L} of an n -gon, the product of n matrices (2.7). Notice that

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -b & a \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}^{-1} = \begin{pmatrix} a & -b \\ 1 & 0 \end{pmatrix}.$$

That is, the Lax matrices are conjugated to the monodromy matrices, and we can use the calculations of the latter from Section 2.2.

Set

$$(3.6) \quad a_i = \frac{\lambda v_{2i}}{s_{2i-1} s_{2i+1}}, \quad b_i = \frac{\lambda^2}{s_{2i-1}^2} - 1, \quad i = 0, \dots, n-1,$$

and consider the continuants (2.4). Then Proposition 2.3 implies that

$$\mathrm{Tr} \mathcal{L} = D_{0,n} - b_0 D_{1,n-1}.$$

The homogeneous in λ components are integrals of the c -relations for all values of c .

A combinatorial rule for calculation of the general continuants

$$\begin{vmatrix} a_1 & c_1 & 0 & \dots & 0 & 0 \\ b_1 & a_2 & c_2 & \dots & 0 & 0 \\ 0 & b_2 & a_3 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & a_{n-1} & c_{n-1} \\ 0 & 0 & 0 & \dots & b_{n-1} & a_n \end{vmatrix}$$

is as follows: one term of the continuant is $a_1 a_2 \dots a_n$, and the other terms are obtained from it by replacing any number of disjoint pairs $(a_i a_{i+1})$ by $(-b_i c_i)$, see [10]. That is, this continuant can be written as

$$(3.7) \quad \prod_{i=1}^{n-1} \left(E - b_i c_i \frac{\partial^2}{\partial a_i \partial a_{i+1}} \right) (a_1 a_2 \dots a_n),$$

where E is the identity operator. Note that the differential operators involved in this formula commute with each other.

As in [4], a subset of the set $\{0, 1, \dots, n-1\}$ is called *cyclically sparse* if it contains no pairs of consecutive indices, and the indices are understood cyclically mod n (the empty set is also sparse).

The above rule implies

LEMMA 3.11. — *One has*

$$\begin{aligned}
 \text{Tr } \mathcal{L} &= \sum_{I \text{ sparse}} (-1)^{|I|} \lambda^{n-2|I|} \prod_{j,j+1 \notin I} \left(\frac{v_{2j}}{s_{2j-1}s_{2j+1}} \right) \prod_{i \in I} \left(\frac{\lambda^2}{s_{2i-1}^2} - 1 \right) \\
 (3.8) \quad &= \lambda^n \sum_{I \text{ sparse}} \prod_{j,j+1 \notin I} \left(\frac{v_{2j}}{s_{2j-1}s_{2j+1}} \right) \prod_{i \in I} \left(\frac{1}{\lambda^2} - \frac{1}{s_{2i-1}^2} \right).
 \end{aligned}$$

Thus, for $t = \lambda^{-2}$,

$$(3.9) \quad \sum_{I \text{ sparse}} \prod_{j,j+1 \notin I} \left(\frac{v_{2j}}{s_{2j-1}s_{2j+1}} \right) \prod_{i \in I} \left(\frac{1}{\lambda^2} - \frac{1}{s_{2i-1}^2} \right) =: \sum_k t^k F_k,$$

is a generating function of the integrals F_k of the c -relation on twisted n -gons.

Note that, as a polynomial in the variables v_{2i} , one has $\deg F_k = n - 2k$. In particular, if $n = 2q + 1$, then

$$F_q = \sum_i \frac{v_{2i}}{s_{2i-1}s_{2i+1}},$$

and one has $q + 1$ integrals in this case. If $n = 2q$, then

$$F_{q-1} = \sum_i \frac{v_{2i}v_{2i+2}}{s_{2i-1}s_{2i+1}^2s_{2i+3}},$$

and one has q integrals in this case. Compare with the examples in Section 3.1.

To summarize, we obtain $\lfloor \frac{n+1}{2} \rfloor$ integrals on the moduli space of twisted n -gons. We do not dwell on the independence and completeness of this set of integrals here; see [8] and the remark below.

Remark 3.12. — If one has $s_{2i-1} = 1$ for all i , corresponding – as explained in Introduction – to the case studied in [4], then formula (3.8) simplifies. The integrals in [4] are given by the formulas

$$F_k = \sum_{I \text{ sparse}, |I|=k} \prod_{i \in I} c_i,$$

where, in our notation, $c_i = 1/(v_{2i}v_{2i+2})$. Multiplying by $(\prod c_i)^{-1/2}$ (which is also an integral), this becomes

$$\sum_{I \text{ sparse}, |I|=k} \prod_{j,j+1 \notin I} v_{2j},$$

precisely what (3.8) yields when $s_{2i-1} = 1$ for all i .

As before, we write

$$g_i = \frac{v_{2i}}{s_{2i-1}s_{2i+1}}, \quad \beta_i = -\frac{1}{s_{2i-1}^2}.$$

Modifying formula (3.7) to account for the cyclic symmetry, we write the generating function of the integrals F_k as

$$\prod_{i=0}^{n-1} \left[E + (t + \beta_{i+1}) \frac{\partial^2}{\partial g_i \partial g_{i+1}} \right] \left(\prod_{j=0}^{n-1} g_j \right).$$

Comparing this formula with formula (22) in [17], we conclude that our integrals coincide with the integrals of the dressing chain obtained in [17]. This is not unexpected, given the appearance of the dressing chain as the infinitesimal version of the c -relation in the preceding Section 3.3. Note that the independence of the integrals is asserted in [17].

3.5. Integrals for closed polygons

When restricted to the moduli space of closed polygons, the integrals from Section 3.4 become dependent. This follows from the next general observation.

Let X be a manifold, $M_t : X \rightarrow \mathrm{SL}(2, \mathbb{R})$ be a 1-parameter family of smooth maps, analytically depending on parameter t . Assume that the identity matrix $E \in \mathrm{SL}(2, \mathbb{R})$ is a regular value of M_0 , and let $Y = M_0^{-1}(E)$. Consider a 1-parameter family of smooth functions $G_t = \mathrm{Tr} M_t : X \rightarrow \mathbb{R}$. Let prime denote d/dt .

LEMMA 3.13. — *One has:*

$$G_0|_Y = 2, \quad G'_0|_Y = 0, \quad dG_0|_Y = 0.$$

Proof. — Since M_0 sends Y to the unit matrix, G_0 equals 2 on Y .

Fix $y \in Y$ and consider the curve $M_t(y)$ in $\mathrm{SL}(2, \mathbb{R})$. The tangent vector $M'_t(y)$ at $t = 0$ lies in $\mathfrak{sl}(2, \mathbb{R})$, and this matrix has zero trace. This proves the second equality.

For the third equality, let the eigenvalues of $M_t(x)$ be $e^{\pm\mu(x,t)}$. Then $G_t(x) = e^{\mu(x,t)} + e^{-\mu(x,t)}$, hence $dG_t = (e^{\mu(x,t)} - e^{-\mu(x,t)}) d\mu(x,t)$. Since $\mu(x,0) = 0$ for $x \in Y$, one has $dG_0|_Y = 0$, as claimed. \square

We shall apply this lemma to a modified Lax matrix of an n -gon. Recall that $t = \lambda^{-2}$. Let

$$(3.10) \quad \bar{\mathcal{L}} = \frac{\mathcal{L}}{\sqrt{\prod \left(\frac{\lambda^2}{s_{2i-1}^2} - 1 \right)}} = \frac{\left(\prod_{i=1}^n s_{2i-1} \right) \mathcal{L}}{\lambda^n \sqrt{\prod (1 - t s_{2i-1}^2)}}.$$

Then, according to (2.7), $\bar{\mathcal{L}} \in \text{SL}(2, \mathbb{R})$. We write $\bar{\mathcal{L}}_t$ to emphasize the dependence on t . In what follows we consider the limit $t \rightarrow 0$, that is, $\lambda \rightarrow \infty$.

LEMMA 3.14. — *One has $\bar{\mathcal{L}}_0 = E$ on closed polygons.*

Proof. — We will show that the Lax matrix $\bar{\mathcal{L}}_t$ is a deformation of the monodromy \mathcal{M} .

The monodromy was calculated in Proposition (2.3), where the variables in the continuants were

$$a_i = \frac{v_{2i}}{s_{2ji1}}, \quad b_i = \frac{s_{2i+1}}{s_{2i-1}}.$$

One can rewrite these continuants as follows:

$$D_{i,j+1} = (s_{2i+1}s_{2i+3} \cdots s_{2j+1})\bar{D}_{i,j+1},$$

where

$$\bar{D}_{i,j+1} = \det \begin{pmatrix} \frac{v_{2i}}{s_{2i-1}s_{2i+1}} & 1 & 0 & \cdots & 0 \\ \frac{1}{s_{2i+1}^2} & \frac{v_{2i+2}}{s_{2i+1}s_{2i+3}} & 1 & 0 & \cdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \cdots & 0 & \frac{1}{s_{2j-3}^2} & \frac{v_{2j-2}}{s_{2j-3}s_{2j-1}} & 1 \\ 0 & \cdots & 0 & \frac{1}{s_{2j-1}^2} & \frac{v_{2j}}{s_{2j-1}s_{2j+1}} \end{pmatrix}.$$

Since $\mathcal{M} = E$ for closed n -gons, we have

$$(3.11) \quad - \left(\frac{\prod_{i=1}^n s_{2i-1}}{s_{2n-1}^2} \right) \bar{D}_{1,n-1} = \left(\prod_{i=1}^n s_{2i-1} \right) \bar{D}_{0,n} = 1, \quad \bar{D}_{1,n} = \bar{D}_{0,n-1} = 0.$$

On the other hand, \mathcal{L}_0 is also given by Proposition (2.3), where the variables in the continuants are as in (3.6). This matrix equals

$$\lambda^n \begin{pmatrix} - \left(\frac{1}{s_{2n-1}^2} - t \right) \bar{D}_{1,n-1}(t), & \frac{1}{\lambda} \bar{D}_{0,n-1}(t) \\ -\lambda \left(\frac{1}{s_{2n-1}^2} - t \right) \bar{D}_{1,n}(t), & \bar{D}_{0,n}(t) \end{pmatrix},$$

where

$$\bar{D}_{i,j+1}(t) = \det \begin{pmatrix} \frac{v_{2i}}{s_{2i-1}s_{2i+1}} & 1 & 0 & \cdots & 0 \\ \frac{1}{s_{2i+1}^2} - t & \frac{v_{2i+2}}{s_{2i+1}s_{2i+3}} & 1 & 0 & \cdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \cdots & 0 & \frac{1}{s_{2j-3}^2} - t & \frac{v_{2j-2}}{s_{2j-3}s_{2j-1}} & 1 \\ 0 & \cdots & 0 & \frac{1}{s_{2j-1}^2} - t & \frac{v_{2j}}{s_{2j-1}s_{2j+1}} \end{pmatrix}.$$

Since $\bar{D}_{1,n} = 0$, one has $\bar{D}_{1,n}(t) = O(t)$, and hence $\lim_{t \rightarrow 0} \lambda \bar{D}_{1,n}(t) = 0$. This, along with equations (3.11), implies that $\bar{\mathcal{L}}_0 = E$, as claimed. \square

We apply Lemma 3.13 to $\bar{\mathcal{L}}_t$. Recall that $\text{Tr} \mathcal{L}_t = \lambda^n (F_0 + tF_1 + \dots)$ where F_i are integrals of the c -relation on twisted n -gons.

PROPOSITION 3.15. — *Restricted to the space \mathcal{X}_n of closed n -gons, one has the following relations:*

$$F_0|_{\mathcal{X}} = 2 \left(\prod_{i=1}^n s_{2i-1} \right)^{-1/2}, \quad F_1|_{\mathcal{X}} = - \left(\frac{1}{2} \sum_{i=1}^n s_{2i-1} \right) F_0|_{\mathcal{X}}, \quad dF_0|_{\mathcal{X}} = 0.$$

Proof. — One has

$$\prod_{i=1}^n (1 - ts_{2i-1})^{-1/2} = 1 + \left(\frac{1}{2} \sum_{i=1}^n s_{2i-1} \right) t + O(t^2),$$

and then, according to (3.10),

$$\begin{aligned} \text{Tr} \bar{\mathcal{L}}_t &= \left(\prod_{i=1}^n s_{2i-1} \right) \left[1 + \left(\frac{1}{2} \sum_{i=1}^n s_{2i-1} \right) t + O(t^2) \right] [F_0 + tF_1 + O(t^2)] \\ &= \left(\prod_{i=1}^n s_{2i-1} \right) \left[F_0 + \left(F_1 + \left(\frac{1}{2} \sum_{i=1}^n s_{2i-1} \right) F_0 \right) t + O(t^2) \right]. \end{aligned}$$

Now Lemma 3.13 implies the result. \square

As before, we do not dwell here on the question whether the relations from Proposition 3.15 are the only ones satisfied by the integrals when restricted to the moduli space of closed polygons (similarly to [4], we do expect this to be the case).

4. Closed centroaffine polygons, before factorizing by $\text{SL}(2, \mathbb{R})$

4.1. Presymplectic forms

Recall that $\tilde{\mathcal{X}}_{n,S}$ is the space of closed n -gons with fixed “side areas” s_{2i-1} , $i = 1, \dots, n$.

Choose a coordinate system in \mathbb{R}^2 , and let $P_i = (x_i, y_i)$ be the vertices of an n -gon. Consider the differential 2-form

$$(4.1) \quad \omega = \sum_{i=1}^n s_{2i+1} (dx_i \wedge dy_{i+1} + dx_{i+1} \wedge dy_i)$$

in $\tilde{\mathcal{X}}_n$. The restriction of ω to $\tilde{\mathcal{X}}_{n,\mathbf{s}}$, which we denote by the same letter, is closed, that is, a presymplectic form in $\tilde{\mathcal{X}}_{n,\mathbf{s}}$.

The generators of the Lie algebra $sl(2, \mathbb{R})$ are the vector fields

$$x\partial/\partial y, \quad x\partial/\partial x - y\partial/\partial y, \quad y\partial/\partial x,$$

and the generators of its diagonal action on polygons are the vector fields

$$e = \sum_i x_i \partial/\partial y_i, \quad h = \sum_i (x_i \partial/\partial x_i - y_i \partial/\partial y_i), \quad f = \sum_i y_i \partial/\partial x_i.$$

These vector fields are tangent to the submanifolds $\tilde{\mathcal{X}}_{n,\mathbf{s}}$.

Let

$$I = \sum_i s_{2i+1} x_i x_{i+1}, \quad J = \sum_i s_{2i+1} (x_i y_{i+1} + x_{i+1} y_i), \quad K = \sum_i s_{2i+1} y_i y_{i+1}.$$

The restrictions of these functions on $\tilde{\mathcal{X}}_{n,\mathbf{s}}$ are the Hamiltonian functions of the above vector fields:

$$(4.2) \quad i_e \omega = -dI, \quad i_h \omega = dJ, \quad i_f \omega = dK,$$

as shown by a straightforward calculation.

THEOREM 4.1.

- (1) *The restriction of ω to $\tilde{\mathcal{X}}_{n,\mathbf{s}}$ is $SL(2, \mathbb{R})$ -invariant, but it is not basic: it does not descend on the moduli space $\mathcal{X}_{n,\mathbf{s}}$.*
- (2) *The form ω is invariant under the c -relations for all values of c and under the polygon recutting.*

Proof. — Equations (4.2) and the Cartan formula imply that $L_e(\omega) = L_h(\omega) = L_f(\omega) = 0$, where L denotes the Lie derivative. Therefore ω is $SL(2, \mathbb{R})$ -invariant, but it is not basic, that is, it is not a pull-back of a differential form defined on the quotient space, since it is not annihilated by $sl(2, \mathbb{R})$.

To prove the invariance of ω under the c -relations, let \mathbf{Q} be an n -gon such that $\mathbf{Q} \stackrel{c}{\sim} \mathbf{P}$, and let $Q_i = (u_i, v_i)$. In the calculations below we tacitly consider \mathbf{Q} to be a function of \mathbf{P} , so when we write a differential form, such as du_i or dv_i , we mean their pull back to \mathbf{P} , that is, differential forms that are linear combinations of dx_j and dy_j , with $j = 1, \dots, n$. Now, to the calculations.

One has

$$s_{2i+1} Q_i = cP_{i+1} + [Q_i, P_{i+1}]P_i, \quad s_{2i+1} P_{i+1} = cQ_i + [Q_i, P_{i+1}]Q_{i+1}.$$

Take the bracket (i.e., the determinant) of the first equality with dQ_{i+1} , bracket of the second equality with dP_i , subtract the second from the first, and sum up over i :

$$(4.3) \quad \begin{aligned} \sum s_{2i+1}([Q_i, dQ_{i+1}] - [P_{i+1}, dP_i]) \\ = c \sum ([P_{i+1}, dQ_{i+1}] - [Q_i, dP_i]) \\ + \sum ([Q_i, P_{i+1}]([P_i, dQ_{i+1}] - [Q_{i+1}, dP_i])). \end{aligned}$$

The differential of the left hand side of (4.3) is the difference of ω evaluated at \mathbf{Q} and \mathbf{P} . Therefore it suffices to show that the right hand side is an exact 1-form on $\tilde{\mathcal{X}}_{n,\mathbf{S}}$.

Indeed,

$$\begin{aligned} \sum ([P_{i+1}, dQ_{i+1}] - [Q_i, dP_i]) &= \sum ([P_i, dQ_i] - [Q_i, dP_i]) \\ &= \sum d(x_i v_i + y_i u_i). \end{aligned}$$

Next, $[P_i, dQ_{i+1}] - [Q_{i+1}, dP_i] = d[P_i, Q_{i+1}]$. Equation (2.6) implies

$$[P_i, Q_{i+1}] = \frac{s_{2i+1}^2 - c^2}{[Q_i, P_{i+1}]}.$$

It follows that, up to a constant,

$$\begin{aligned} [Q_i, P_{i+1}]([P_i, dQ_{i+1}] - [Q_{i+1}, dP_i]) &= [Q_i, P_{i+1}]d([Q_i, P_{i+1}]^{-1}) \\ &= -d \ln([Q_i, P_{i+1}]), \end{aligned}$$

therefore the second sum on the right hand side of (4.3) is also an exact 1-form.

To show that the form ω is invariant under the polygon recutting, consider the difference of the form evaluated at \mathbf{P} and at $\mathbf{R}_1(\mathbf{P})$. Let $P'_1 = (x, y) = \mathbf{R}_1(\mathbf{P})_1$. Then

$$x = \frac{x_n s_1 + x_2 s_3}{v_2}, \quad y = \frac{y_n s_1 + y_2 s_3}{v_2}.$$

We get

$$\begin{aligned}
 \omega(\mathbf{P}) - \omega(\mathbf{R}_1(\mathbf{P})) &= s_3(dx_1 \wedge dy_2 + dx_2 \wedge dy_1 - dx_n \wedge dy - dx \wedge dy_n) \\
 &\quad + s_1(dx_n \wedge dy_1 + dx_1 \wedge dy_n - dx \wedge dy_2 - dx_2 \wedge dy) \\
 &= (s_3dx_1 - s_1dx) \wedge dy_2 + dx_2 \wedge (s_3dy_1 - s_1dy) \\
 &\quad + (s_1dx_1 - s_3dx) \wedge dy_n + dx_n \wedge (s_1dy_1 - s_3dy) \\
 &= \frac{s_3^2}{v_2}(dx_n \wedge dy_2 + dx_2 \wedge dy_n) - \frac{s_3^2}{v_2}(dx_n \wedge dy_2 + dx_2 \wedge dy_n) \\
 &= 0,
 \end{aligned}$$

as needed. □

Remark 4.2. — It was pointed out by A. Izosimov that the 2-form ω is also well defined on the space $\tilde{\mathcal{X}}_{n,\mathbf{s},\mathcal{M}}$ of twisted n -gons with monodromy \mathcal{M} .

4.2. Additional integrals

The next theorem, providing two additional integrals of the c -relation, is a discrete version of Proposition 3.4 in [14], concerning a continuous version of the c -relation on centroaffine curves. It is also an analog of Theorem 16 in [4].

THEOREM 4.3.

- (1) *The functions I, J, K are integrals of the c -relations for all values of c and of the polygon recutting.*
- (2) *The integral $4IK - J^2$ descends to the moduli space $\mathcal{X}_{n,\mathbf{s}}$.*

Proof. — We use the notations from the proof of Theorem 4.1. One has

$$x_i v_i - y_i u_i = c, \quad x_i y_{i+1} - y_i x_{i+1} = s_{2i+1} = u_i v_{i+1} - v_i u_{i+1}$$

for all i . Hence $v_i = (c + y_i u_i)/x_i$ and

$$\frac{u_i(c + y_{i+1} u_{i+1})}{x_{i+1}} - \frac{u_{i+1}(c + y_i u_i)}{x_i} = s_{2i+1}.$$

It follows that

$$c(x_i u_i - x_{i+1} u_{i+1}) + u_i u_{i+1}(x_i y_{i+1} - y_i x_{i+1}) = s_{2i+1} x_i x_{i+1}$$

or

$$c[x_i u_i - x_{i+1} u_{i+1}] + s_{2i+1} u_i u_{i+1} = s_{2i+1} x_i x_{i+1}.$$

Taking sum over i eliminates the first summand on the left hand side and shows that I is an integral.

Next we show that I is also an integral for the polygon recutting. Indeed,

$$\begin{aligned} I(\mathbf{P}) - I(\mathbf{R}_1(\mathbf{P})) &= (x_1 x_2 s_3 + x_n x_1 s_1) - (x x_2 s_1 + x_n x s_3) \\ &= x_2 (x_1 s_3 - x s_1) + x_n (x_1 s_1 - x s_3) = 0. \end{aligned}$$

Therefore $I(\mathbf{P}) = I(\mathbf{R}_1(\mathbf{P})) = I(\mathbf{R}_2 \mathbf{R}_1(\mathbf{P})) = \dots = \mathbf{R}(\mathbf{P})$.

One also has

$$(4.4) \quad \begin{aligned} e(I) = 0, e(J) = 2I, e(K) = J, \quad h(I) = 2I, h(J) = 0, h(K) = -2K, \\ f(I) = J, f(J) = 2K, f(K) = 0. \end{aligned}$$

Since the c -relations and the recutting commute with $\mathrm{SL}(2, \mathbb{R})$, it follows that J and K are also integrals.

Finally, (4.4) imply that $4IK - J^2$ is an $\mathfrak{sl}(2, \mathbb{R})$ -invariant function. This proves the last claim. \square

The space spanned by I, J, K is the irreducible 3-dimensional representation of the Lie algebra $\mathfrak{sl}(2, \mathbb{R})$, the symmetric square of its standard 2-dimensional representation, or the coadjoint representation. The map $\mathcal{X}_{n, \mathbf{S}} \rightarrow \mathbb{R}^3$, whose components are the functions I, J, K , is the moment map of the Hamiltonian action of $\mathfrak{sl}(2, \mathbb{R})$ on $\mathcal{X}_{n, \mathbf{S}}$.

Remark 4.4. — One has

$$4IK - J^2 = - \sum_{k,l} s_{2k+1} s_{2l+1} ([P_k, P_l][P_{k+1} P_{l+1}] - [P_k, P_{l+1}][P_{k+1}, P_l]).$$

4.3. Center of a polygon

Define the *center* of a polygon \mathbf{P} as the quadratic form on \mathbb{R}^2

$$C(\mathbf{P}) = I(\mathbf{P})x^2 - J(\mathbf{P})xy + K(\mathbf{P})y^2.$$

The center is invariant under the c -relation and the polygon recutting, and it conjugates the diagonal action of $\mathrm{SL}(2, \mathbb{R})$ on polygons and on its action on quadratic forms. In more detail, the group $\mathrm{SL}(2, \mathbb{R})$ naturally acts on quadratic forms, that is, on $S^2(\mathbb{R}^2)$, and it acts diagonally on centroaffine polygons in \mathbb{R}^2 . The map C that assigns the center to a polygon conjugates these two actions.

In this section we present some properties of the center, somewhat analogous to those of the circumcenter of mass, see [16].

The center is additive in the following sense.

LEMMA 4.5. — *Let $\mathbf{P} = (P_1, \dots, P_k, \dots, P_n)$. Cut \mathbf{P} into two polygons $\mathbf{P}_1 = (P_1, \dots, P_k)$ and $\mathbf{P}_2 = (P_1, P_k, \dots, P_n)$. Then $C(\mathbf{P}) = C(\mathbf{P}_1) + C(\mathbf{P}_2)$.*

Proof. — Each of the three components of the sum $C(\mathbf{P}_1) + C(\mathbf{P}_2)$ contains all the terms of the respective component of $C(\mathbf{P})$ plus the additional terms (two or four), that appear due to the cut P_1P_k . Since the “side area” $[P_1P_k]$ changes the sign when the orientation of the side is reversed, these additional terms cancel pairwise. \square

According to the preceding lemma, the calculation of the center of a polygon reduces to that of a triangle. The next result gives a geometrical interpretation to the center of a triangle.

LEMMA 4.6. — *A triangle $\mathbf{P} = (P_1P_2P_3)$ admits a unique circumscribed central conic given by the equation $ax^2 - bxy + cy^2 = 1$, that is, the conic whose center is at the origin and that passes through the vertices of the triangle. The center of the triangle \mathbf{P} is $2A(ax^2 - bxy + cy^2)$, where A is the oriented area of \mathbf{P} .*

Proof. — Let $P_i = (x_i, y_i)$, $i = 1, 2, 3$. We recall that the sides do not pass through the origin.

To find the circumscribed central conic one needs to solve the linear system $M(a, -b, c)^T = (1, 1, 1)^T$ where

$$M = \begin{pmatrix} x_1^2 & x_1y_1 & y_1^2 \\ x_2^2 & x_2y_2 & y_2^2 \\ x_3^2 & x_3y_3 & y_3^2 \end{pmatrix}.$$

One has $\det M = s_1s_3s_5$. Denote by N the cofactor matrix of M . Then

$$M^{-1} = \frac{1}{s_1s_3s_5}N,$$

and hence

$$(a, -b, c)^T = \frac{1}{s_1s_3s_5}N(1, 1, 1)^T.$$

Note that $s_1s_3s_5$ is twice the oriented area of the triangle.

On the other hand, one notices that $N(1, 1, 1)^T = (K, -J, I)^T$. This implies the result. \square

LEMMA 4.7. — *The center of a centroaffine butterfly is the origin.*




Figure 4.1. Triangles $P_1P_2Q_2$ and $P_1Q_1Q_2$ share the circumscribed central conic.

Proof. — An affine reflection in a line through the origin interchanges points P_1 with Q_2 , and P_2 with Q_1 , see Figure 4.1. This reflection preserves the central conic circumscribed about triangle $P_1P_2Q_2$, therefore triangles $P_1P_2Q_2$ and $P_1Q_1Q_2$ share the circumscribed central conic. Now the result follows from Lemmas 4.5 and 4.6. \square

Lemmas 4.5 and 4.7 provide an alternative proof that the center is invariant under polygon recutting.

5. Small-gons

5.1. Triangles

In this section we investigate closed triangles.

THEOREM 5.1.

(1) A triangle admits a c -related triangle if and only if

$$c^2(s_1 + s_3 + s_5)(s_1 + s_3 - s_5)(s_3 + s_5 - s_1)(s_5 + s_1 - s_3) \leq 4(s_1s_3s_5)^2.$$

No triangles have infinitely many c -related ones for any $c \neq 0$.

(2) Two c -related triangles are $\mathrm{SL}(2, \mathbb{R})$ -equivalent.

(3) The linear transformation M that relates them and that defines the dynamics is elliptic if and only if

$$(5.1) \quad (s_1 + s_3 + s_5)(s_1 + s_3 - s_5)(s_3 + s_5 - s_1)(s_5 + s_1 - s_3) > 0.$$

M is parabolic if and only if the origin is located on the lines that bisect two sides of the triangle, see Figure 5.1.

- (4) Let triangle $A'B'C'$ be the recutting of triangle ABC done in the order $A \rightarrow B \rightarrow C$, and let $A''B''C''$ be the second iteration of this recutting. Then there exists a transformation $M \in \text{SL}(2, \mathbb{R})$ that takes ABC to $A''B''C''$.




Figure 5.1. The shaded regions are where the origin should be located for the linear map M to be elliptic.

We cannot help noticing that the left hand side of (5.1) reminds one of Heron’s formula that expresses the area of a triangle in terms of its side lengths. We do not know whether this similarity has a meaning or it is a pure coincidence.

Proof. — A triangle is uniquely determined, modulo $\text{SL}(2, \mathbb{R})$, by the areas s_1, s_3, s_5 . These numbers are preserved by the c -relation, proving the second claim.

Let $\mathbf{Q} = \mathbf{M}\mathbf{P}$ where $M = \begin{pmatrix} m & n \\ k & l \end{pmatrix} \in \text{SL}(2, \mathbb{R})$, and let $P_j = (x_j, y_j)$. Then the relation the $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$ implies

$$\begin{cases} (m-l)x_1y_1 - kx_1^2 + ny_1^2 = c \\ (m-l)x_2y_2 - kx_2^2 + ny_2^2 = c \\ (m-l)x_3y_3 - kx_3^2 + ny_3^2 = c. \end{cases}$$

Here is the solution:

$$\begin{pmatrix} m-l \\ k \\ n \end{pmatrix} = c \begin{pmatrix} x_1y_1 & -x_1^2 & y_1^2 \\ x_2y_2 & -x_2^2 & y_2^2 \\ x_3y_3 & -x_3^2 & y_3^2 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$

For M to exist, one needs the relation $ml - kn = 1$ to hold. Since $m-l, k, n$ are already determined, this reduces to a quadratic equation on m that has real roots if and only if

$$(m-l)^2 + 4kn + 4 \geq 0.$$

One has a remarkable identity:

$$(m-l)^2 + 4kn = -c^2 \frac{(s_1 + s_3 + s_5)(s_1 + s_3 - s_5)(s_3 + s_5 - s_1)(s_5 + s_1 - s_3)}{(s_1 s_3 s_5)^2},$$

that we verified using Mathematica. This implies the first claim of the theorem. Furthermore, M is elliptic if and only if

$$\text{Tr}(M)^2 - 4 \det(M) = (m+l)^2 - 4 < 0$$

or, which is equivalent, $(m-l)^2 + 4kn < 0$. This implies the third claim.

The right hand side vanishes when $s_1 = s_3 + s_5$ or its cyclic permutation, that is, the origin lies on one of the three middle lines of the triangle. These lines separate the elliptic and hyperbolic regions.

For the last claim, one has $[A, B] = [B', C']$ and $[B, C] = [A', B']$, see Figure 5.2. Since the total area is preserved by recutting, one also has $[C, A] = [C', A']$.




Figure 5.2. Triangle recutting.

Repeating this argument going from $A'B'C'$ to $A''B''C''$, we see that

$$[A, B] = [A'', B''], [B, C] = [B'', C''], [C, A] = [C'', A''].$$

Therefore the triangles ABC and $A''B''C''$ are $\text{SL}(2, \mathbb{R})$ -equivalent. \square

Remark 5.2. — The first claim in Theorem 5.1 has the following geometric interpretation. Given a triangle, there exists a central conic through its vertices. Assume that this conic is an ellipse and apply a transformation from $\text{SL}(2, \mathbb{R})$ to make this ellipse into a circle of radius R . Then a c -related triangle is also inscribed in this circle, and one has $c \leq R^2$.

Thus one expects the following identity to hold:

$$(5.2) \quad R^4 (s_1 + s_3 + s_5)(s_1 + s_3 - s_5)(s_3 + s_5 - s_1)(s_5 + s_1 - s_3) = 4(s_1 s_3 s_5)^2.$$

Let α, β, γ be the (signed) angles under which the sides of the triangle are seen from the origin. Then $\alpha + \beta + \gamma = 2\pi$,

$$s_1 = R^2 \sin \alpha, \quad s_3 = R^2 \sin \beta, \quad s_5 = R^2 \sin \gamma,$$

and (5.2) becomes a true trigonometric identity

$$\begin{aligned} &(\sin \alpha + \sin \beta + \sin \gamma)(\sin \alpha + \sin \beta - \sin \gamma)(\sin \alpha - \sin \beta + \sin \gamma) \\ &\quad \times (-\sin \alpha + \sin \beta + \sin \gamma) = 4 \sin^2 \alpha \sin^2 \beta \sin^2 \gamma. \end{aligned}$$

5.2. Quadrilaterals

Let us consider the dynamics of the c -relation on closed quadrilaterals. Let \mathbf{P} be a quadrilateral, and assume that $\mathcal{L}_{\mathbf{P},c} \neq \text{Id}$.

PROPOSITION 5.3.

- (1) Let $\mathbf{P} \stackrel{c}{\sim} \mathbf{Q}$. Then there exist homothetic central conics \mathcal{C}_1 and \mathcal{C}_2 such that $P_1, Q_2, P_3, Q_4 \in \mathcal{C}_1$ and $Q_1, P_2, Q_3, P_4 \in \mathcal{C}_2$, see Figure 5.3.
- (2) The conics in questions are ellipses if and only if

$$(s_1 + s_3 + s_5 + s_7)(s_1 + s_3 - s_5 - s_7)(s_3 + s_5 - s_7 - s_1)(s_5 + s_7 - s_1 - s_3) < 0.$$




Figure 5.3. Homothetic concentric conics containing the vertices of c -related quadrilaterals.

Proof. — As in Section 2.5, one has four affine reflections R_i whose composition is the identity map. Consider R_i as an orientation reversing isometry of the hyperbolic plane, a reflection in a line ℓ_i . Then $R_2 \circ R_1$ is an orientation preserving isometry. One has four cases: this isometry is elliptic, hyperbolic, parabolic, or the identity.

In the elliptic case, the isometry is a rotation about a point in \mathbb{H}^2 . Hence $R_4 \circ R_3$ is a rotation about the same point, and therefore the four lines ℓ_i are concurrent at this point. It follows that R_i belong to group $G \subset \text{GL}(2, \mathbb{R})$ that is conjugated to $O(2)$, that is, the group generated by the rotations

$$\begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix} \text{ and by } \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix},$$

and preserving the quadratic form $x^2 + y^2$. Thus G preserves a positive-definite quadratic form whose circles are the homothetic ellipses preserved by the reflections R_i . Since R_i swaps P_i with Q_{i+1} and Q_i with P_{i+1} , we are done in this case.

In the hyperbolic case, the argument is similar. The isometries R_i of \mathbb{H}^2 are reflections in the lines ℓ_i that share a common perpendicular (and the lines are concurrent at a point of the projective plane outside of the absolute). In this case the argument is similar with the group G being conjugated to $O(1, 1)$ and generated by

$$\begin{pmatrix} \cosh t & \sinh t \\ \sinh t & \cosh t \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \text{ and by } \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix},$$

preserving the quadratic form $x^2 - y^2$. Thus G preserves a non-degenerate sign-indefinite quadratic form whose level curves are the desired homothetic hyperbolas (note that one of the level curves is singular: it is a pair of lines intersecting at the origin).




Figure 5.4. The parabolic case.

In the parabolic case, the conics degenerate to pairs of origin-symmetric parallel lines, see Figure 5.4. Another degenerate case is when two opposite vertices of \mathbf{P} are collinear, this happens in the hyperbolic case when the zero level curve of the sign-indefinite quadratic form is a pair of intersecting lines.

Finally, in the case of the identity, one has $P_1 = (R_2 \circ R_1)(P_1) = P_3$, contradicting the non-degeneracy of the quadrilateral.

We write the conics in the format $\langle P, MP \rangle = \text{const}$, where $P = (x, y)$ and $M = \begin{pmatrix} m & n \\ n & k \end{pmatrix}$. The homogeneous equation for the matrix elements

$$\begin{cases} m(x_1^2 - x_3^2) + 2n(x_1y_1 - x_3y_3) + k(y_1^2 - y_3^2) = 0 \\ m(x_2^2 - x_4^2) + 2n(x_2y_2 - x_4y_4) + k(y_2^2 - y_4^2) = 0 \end{cases}$$

has a solution

$$\left[\begin{matrix} m \\ n \\ k \end{matrix} \right] = \left[\begin{matrix} y_1y_2s_3 + y_2y_3s_5 + y_3y_4s_7 + y_4y_1s_9 \\ -(1/2)((x_1^2 - x_3^2)(y_2^2 - y_4^2) - (x_2^2 - x_4^2)(y_1^2 - y_3^2)) \\ x_1x_2s_3 + x_2x_3s_5 + x_3x_4s_7 + x_4x_1s_9 \end{matrix} \right],$$

where the square bracket denotes the projective equivalence class. A calculation shows that

$$\begin{aligned} n^2 - mk &= (s_1 + s_3 + s_5 + s_7)(s_1 + s_3 - s_5 - s_7)(s_3 + s_5 - s_7 - s_1)(s_5 + s_7 - s_1 - s_3), \end{aligned}$$

which implies the second result. □

Arguing as in the preceding section, Proposition 5.3 has the following corollary.

COROLLARY 5.4. — *Let \mathbf{Q} be the recutting of the quadrilateral \mathbf{P} . Then the odd vertices of \mathbf{Q} lie on the same central conic as the odd vertices of \mathbf{P} , and the even vertices of \mathbf{Q} lie on the same homothetic central conic as the even vertices of \mathbf{P} .*

Let \mathbf{P} be a quadrilateral with coordinates (s_1, s_3, s_5, s_7) and (v_2, v_4) .

THEOREM 5.5.

(1) *A quadrilateral admits a c -related quadrilateral if and only if*

$$(5.3) \quad c^2(s_1 + s_3 - s_5 - s_7)(s_1 - s_3 + s_5 - s_7)(s_1 - s_3 - s_5 + s_7) \times (s_1 + s_3 + s_5 + s_7) \geq 4(s_3s_5 - s_1s_7)(s_3s_7 - s_1s_5)(s_1s_3 - s_5s_7).$$

This condition is symmetric in $\{s_1, s_3, s_5, s_7\}$, and it has solution in c for every S .

- (2) The second iteration of the c -transformation of \mathbf{P} is $\mathrm{SL}(2, \mathbb{R})$ -equivalent to \mathbf{P} .
- (3) The third iteration of the recutting of \mathbf{P} is $\mathrm{SL}(2, \mathbb{R})$ -equivalent to \mathbf{P} .

Proof. — To prove the first statement, note that \mathbf{P} is $\mathrm{SL}(2, \mathbb{R})$ -equivalent to the following one:

$$P_0 = (1, 0), P_1 = (0, s_1), P_2 = \left(-\frac{s_3}{s_1}, v_2\right), P_3 = \left(-\frac{v_4}{s_1}, -s_7\right),$$

where $v_2 v_4 = s_1 s_5 - s_3 s_7$ (the Ptolemy–Plücker relation).

Let $Q_0 = (b, c)$. A calculation using (2.6) shows that $Q_4 = (b', c)$, and we need $b' = b$. This is a quadratic equation $b^2 + ub + v = 0$ on b whose coefficients are given by the formulas

$$u = \frac{c}{v_2} \left[\frac{s_1^2 - s_3^2 - s_5^2 - s_7^2}{s_5 s_7 - s_1 s_3} + \frac{2s_3 s_5 s_7}{s_1 (s_5 s_7 - s_1 s_3)} \right],$$

$$v = \frac{(c^2 - s_1^2)(s_3 s_5 - s_1 s_7)(s_3 s_7 - s_1 s_5)}{v_2^2 s_1^2 (s_5 s_7 - s_1 s_3)}.$$

One calculates the discriminant $D = u^2 - 4v$, and, after some cancellation, this results in (5.3) (we used Mathematica to clean-up the formulas).

One also has $D = c^2 P + 4s_1^2 Q$, where

$$Q(s_1, s_3, s_5, s_7) = (s_5 s_7 - s_1 s_3)(s_3 s_7 - s_1 s_5)(s_3 s_5 - s_1 s_7),$$

$$P(s_1, s_3, s_5, s_7) = s_1^2 (s_1^2 - s_3^2 - s_5^2 - s_7^2)^2 - Q(s_1, s_3, s_5, s_7).$$

Therefore one cannot have $P < 0$ and $Q < 0$ simultaneously, and this implies that (5.3) always has a solution.

The group of permutations of the "four elements of the set \mathbf{S} " is generated by the involutions (13), (35), (57) that leave inequality (5.3) intact.

The second statement of the theorem follows from Proposition 5.3: if $\mathbf{R} \stackrel{\sim}{\sim} \mathbf{Q}$, then the vertices of the quadrilateral \mathbf{R} lie, alternating, on the same homothetic conics as those of \mathbf{P} , and the respective "side areas" of these quadrilaterals are equal. This implies that \mathbf{R} and \mathbf{P} are $\mathrm{SL}(2, \mathbb{R})$ -equivalent.

For the third statement, using the Ptolemy–Plücker relation, one calculates that after the first recutting the coordinates become

$$(s_1, s_5, s_7, s_3), \left(v_2 \frac{s_1 s_3 - s_5 s_7}{s_1 s_5 - s_3 s_7}, v_4 \frac{s_1 s_7 - s_3 s_5}{s_1 s_3 - s_5 s_7} \right).$$

Then after the second recutting we have

$$(s_1, s_7, s_3, s_5), \left(v_2 \frac{s_1 s_3 - s_5 s_7}{s_1 s_7 - s_3 s_5}, v_4 \frac{s_1 s_7 - s_3 s_5}{s_1 s_5 - s_3 s_7} \right),$$

so after the third recutting one obtains $(s_1, s_3, s_5, s_7), (v_2, v_4)$, as claimed. \square

5.3. Pentagons: invariant area form

In this section we consider the moduli space $\mathcal{X}_{5,\mathbf{s}}$. This material is parallel to the one in [4, Section 7.1.3].

The space $\mathcal{X}_{5,\mathbf{s}}$ is 2-dimensional: the variables v_{2i} satisfy the Ptolemy–Plücker relation $v_2 v_4 + v_8 s_3 = s_1 s_5$ and its cyclic permutations. We break the cyclic symmetry by setting $v_2 = x, v_8 = y$. Then

$$(5.4) \quad v_4 = \frac{s_1 s_5 - s_3 y}{x}, \quad v_6 = \frac{s_5 s_9 - s_7 x}{y}, \quad v_0 = \frac{s_1 s_7 x + s_3 s_9 y - s_1 s_5 s_9}{xy}.$$

Recall (Example 3.4) that

$$K := \frac{v_2}{s_1 s_3} + \frac{v_4}{s_3 s_5} + \frac{v_6}{s_5 s_7} + \frac{v_8}{s_7 s_9} + \frac{v_{10}}{s_9 s_1},$$

the only integral of the c -relation on the moduli space of closed pentagons. In terms of (x, y) -coordinates, one has

$$K = \frac{x}{s_1 s_3} + \frac{y}{s_7 s_9} + \frac{(s_1^2 + s_3^2)}{s_1 s_3 x} + \frac{(s_7^2 + s_9^2)}{s_7 s_9 y} - \frac{x}{s_5 y} - \frac{y}{s_5 x} - \frac{s_5}{xy}.$$

Let

$$\omega = \frac{dx \wedge dy}{xy}.$$

The origin of the area form ω on $\mathcal{X}_{5,\mathbf{s}}$ is in the theory of cluster algebras, and we do not dwell on it here.

THEOREM 5.6. — *The c -relation preserves the form ω .*

Proof. — Recall the vector field ξ from Section 3.3. Using the formula from Theorem 3.9, one has

$$(5.5) \quad \begin{aligned} \dot{x} &= x \left(\frac{v_4}{s_3 s_5} - \frac{v_6}{s_5 s_7} + \frac{v_8}{s_7 s_9} - \frac{v_0}{s_9 s_1} \right) + \frac{s_3}{s_1} - \frac{s_1}{s_3}, \\ \dot{y} &= y \left(\frac{v_0}{s_9 s_1} - \frac{v_2}{s_1 s_3} + \frac{v_4}{s_3 s_5} - \frac{v_6}{s_5 s_7} \right) + \frac{s_3}{s_1} - \frac{s_1}{s_3}. \end{aligned}$$

We claim that $i_\xi \omega = dK$, that is, the integral K is the Hamiltonian function of the vector field ξ . This claim is verified by a direct calculation, after substitution of the formulas (5.4) into (5.5).

Consider the c -relation as a transformation T . By the Bianchi permutability, Φ_t , the t -flow of the field ξ , commutes with T , and these maps preserve the level curves of the function K . Since Φ_t is symplectic, we have

$$\Phi_t^* T^*(\omega) = T^* \Phi_t^*(\omega) = T^*(\omega),$$

that is, $T^*(\omega)$ is also invariant under the flow of ξ . Hence $T^*(\omega) = H\omega$, where the function H is an integral of Φ_t and of T that has the same level curves as K . We want to show that $H \equiv 1$.

Assume that a level curve $K = c$ is closed. Consider an infinitesimally close level curve $K = c + \varepsilon$. Both curves are preserved by T , hence the area between them remains the same. On the other hand, this area is multiplied by the value of the function H on the curve $K = c$. Hence this value equals 1, as needed.




Figure 5.5. Three level curves $K = -10, -9, -8.2$ on $\mathcal{X}_{5,\mathbf{S}}$ with $\mathbf{S} = (1, 1, 1, 1, 1)$.

Thus ω is invariant under T near local maxima or minima of I , see Figure 5.5. The c -relation is an algebraic relation, and we can use analytic continuation to conclude that c -relation preserves ω everywhere. \square

This theorem implies a Poncelet-style porism: if a level curve of the integral K contains a periodic point of the c -relation, then every point of this curve is periodic with the same period.

5.4. Pentagons: when the c -relation is not defined

Let $\mathbf{P} = (P_0, P_1, \dots, P_4)$ be a pentagon. Using an appropriate $\mathrm{SL}(2, \mathbb{R})$ -transformation, we can make $P_0 = (1, 0)$ and $P_1 = (0, s_1)$. Then

$$P_2 = \frac{(-s_3, s_1 v_2)}{s_1}, \quad P_3 = \frac{(s_3 s_7 - s_5 v_4, s_1 s_5 s_9 - s_1 s_7 v_4)}{v_2 v_4 - s_3 s_9}, \quad P_4 = \frac{(v_4, -s_1 s_9)}{s_1}.$$

To find a pentagon $\mathbf{Q} = (Q_0, Q_1, \dots, Q_4)$ c -related to \mathbf{P} , we first put $Q_0 = (b, c)$ (with some unknown b) and compute the remaining vertices using the formulas from Section 2.3.

Our goal is to find a value of b such that $Q_5 = Q_0$; but for an arbitrary b we will have $Q_5 = (b', c)$ with some b' which may be different from b (because $[Q_5, P_5] = [Q_5, P_0]$ must be equal to c). This b' will depend on b, c , and all s, v , and it is not hard to see that it will be, actually, a *quadratic* polynomial in b with coefficients depending on c, s , and v . Then the equation $b' = b$ is quadratic with respect to b . (For $n = 4$, a similar equation was explicitly calculated in the proof of Theorem 5.5.)

We were able to make these calculations, but the result looks depressing, and we do not present it here. Actually, we are more interested in the *discriminant* of this quadratic equation. This discriminant D depends on $c, s_1, s_3, s_5, s_7, s_9$, and v_2, v_4 , but in reality its dependence on v_2, v_4 may be reduced to the dependence on

$$K = \frac{v_2}{s_1 s_3} + \frac{v_4}{s_3 s_5} + \frac{v_6}{s_5 s_7} + \frac{v_8}{s_7 s_9} + \frac{v_{10}}{s_9 s_1}.$$

Moreover, D turns out to be also a *quadratic* function of K with coefficients depending on c , and s_1, s_3, s_5, s_7, s_9 . An explicit expression for this function $D = D(K)$ is less awkward, it can be derived from Propositions 5.7 and 5.8 below.

What we really need is the discriminant of $D(K)$, for which we will use a weird notation $\mathcal{D}(D)$. Indeed, *if, for some $c, s_1, s_3, s_5, s_7, s_9$, $\mathcal{D}(D) \leq 0$, then c -related pentagons exist for all pentagons with these s_1, s_3, s_5, s_7, s_9 ; if $\mathcal{D}(D) > 0$, then the equation $D(K) = 0$ has two different real roots K_1 and K_2 and no c -related pentagons exist for pentagons with K between K_1 and K_2 .*

Notice that both D and $\mathcal{D}(D)$ are defined up to positive factors, which we ignore in the formulas below.

PROPOSITION 5.7. — *One has*

$$\mathcal{D}(D) = \prod_{j=1}^5 (c^2 - s_{2j-1}^2).$$

Thus, if $0 < s_1 < \dots < s_9$ (this condition is not really restrictive, since all our results are not sensitive to permutations and sign changes of s_1, \dots, s_9) then the relation may be undefined only if $s_1 < |c| < s_3$, or $s_5 < |c| < s_7$, or $|c| > s_9$. See Figure 5.6.




Figure 5.6. The zones where the c -relation is not defined; here $\mathbf{S} = \{1, 3, 5, 7, 9\}$. The horizontal and vertical axes are c and K .

PROPOSITION 5.8. — The solutions of the equation $D(K) = 0$ are

$$K = \frac{((\sum_j s_{2j-1}^2) - 2c^2)c^2}{\prod_j s_{2j-1}} \pm \frac{2\sqrt{\mathcal{D}(D)}}{c \prod_j s_{2j-1}}.$$

The proofs consist in tedious but explicit calculations.

By the way, to check a reliable formula, after it has been obtained, we need, as a rule, to prove the equality between two polynomials of the same degree, and for this it is sufficient to verify the equality for a certain number of integral variables, which is an easy task for a computer program.

BIBLIOGRAPHY

- [1] V. E. ADLER, “Recuttings of polygons”, *Funct. Anal. Appl.* **27** (1993), no. 2, p. 141-145.
- [2] ———, “Integrable deformations of a polygon”, *Physica D* **87** (1995), no. 1-4, p. 52-57.
- [3] N. AFFOLTER, T. GEORGE & S. RAMASSAMY, “Cross-ratio dynamics and the dimer cluster integrable system”, <https://arxiv.org/abs/2108.12692>, 2021.

- [4] M. ARNOLD, D. FUCHS, I. IZMESTIEV & S. TABACHNIKOV, “Cross-ratio dynamics on ideal polygons”, *Int. Math. Res. Not.* (2022), no. 9, p. 6770-6853.
- [5] M. BIALY, G. BOR & S. TABACHNIKOV, “Self-Bäcklund curves in centroaffine geometry and Lamé’s equation”, *Commun. Am. Math. Soc.* **2** (2022), p. 232-282.
- [6] G. BOR, M. LEVI, R. PERLINE & S. TABACHNIKOV, “Tire tracks and integrable curve evolution”, *Int. Math. Res. Not.* (2020), no. 9, p. 2698-2768.
- [7] T. HOFFMANN, “Discrete Hashimoto surfaces and a doubly discrete smoke-ring flow”, in *Discrete differential geometry*, Oberwolfach Seminars, vol. 38, Birkhäuser, 2008, p. 95-115.
- [8] A. IZOSIMOV, “In preparation”.
- [9] M. KAPOVICH & J. MILLSON, “On the moduli space of polygons in the Euclidean plane”, *J. Differ. Geom.* **42** (1995), no. 1, p. 133-164.
- [10] T. MUIR, *A treatise on the theory of determinants*, Dover Publications, 1960, revised and enlarged by William H. Metzler, vii+766 pages.
- [11] F. NIJHOFF & H. CAPEL, “The discrete Korteweg-de Vries equation”, *Acta Appl. Math.* **39** (1995), no. 1-3, p. 133-158.
- [12] U. PINKALL, B. SPRINGBORN & S. WEISSMANN, “A new doubly discrete analogue of smoke ring flow and the real time simulation of fluid flow”, *J. Phys. A. Math. Gen.* **40** (2007), no. 42, p. 12563-12576.
- [13] S. TABACHNIKOV, “On the bicycle transformation and the filament equation: results and conjectures”, *J. Geom. Phys.* **115** (2017), p. 116-123.
- [14] ———, “On centro-affine curves and Bäcklund transformations of the KdV equation”, *Arnold Math. J.* **4** (2018), no. 3-4, p. 445-458.
- [15] S. TABACHNIKOV & E. TSUKERMAN, “On the discrete bicycle transformation”, *Publ. Mat. Urug.* **14** (2013), p. 201-219.
- [16] ———, “Circumcenter of mass and generalized Euler line”, *Discrete Comput. Geom.* **51** (2014), no. 4, p. 815-836.
- [17] A. P. VESELOV & A. B. SHABAT, “Dressing chains and the spectral theory of the Schrödinger operator”, *Funkts. Anal. Prilozh.* **27** (1993), no. 2, p. 81-96.
- [18] F. J. WEGNER, “Floating Bodies of Equilibrium in 2D, the Tire Track Problem and Electrons in a Parabolic Magnetic Field”, 2007, <https://arxiv.org/abs/physics/0701241>.

Manuscrit reçu le 17 décembre 2021,
révisé le 12 juin 2022,
accepté le 23 septembre 2022.

Maxim ARNOLD
Department of Mathematical Sciences, University of
Texas at Dallas, 800 West Campbell Road,
Richardson, TX 75080 (USA)
maxim.arnold@utdallas.edu

Dmitry FUCHS
Department of Mathematics, University of
California, Davis, CA 95616 (USA)
fuchs@math.ucdavis.edu

Serge TABACHNIKOV
Department of Mathematics, Pennsylvania State
University, University Park, PA 16802 (USA)
tabachni@math.psu.edu