



ANNALES DE L'INSTITUT FOURIER

Olivier DUDAS & Emily NORTON

Decomposition numbers for the principal Φ_{2n} -block of $\mathrm{Sp}_{4n}(q)$ and $\mathrm{SO}_{4n+1}(q)$

Tome 75, n° 2 (2025), p. 579-622.

<https://doi.org/10.5802/aif.3659>

Article mis à disposition par ses auteurs selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE



<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les *Annales de l'Institut Fourier* sont membres du
Centre Mersenne pour l'édition scientifique ouverte

www.centre-mersenne.org

e-ISSN : 1777-5310

DECOMPOSITION NUMBERS FOR THE PRINCIPAL Φ_{2n} -BLOCK OF $\mathrm{Sp}_{4n}(q)$ AND $\mathrm{SO}_{4n+1}(q)$

by Olivier DUDAS & Emily NORTON (*)

ABSTRACT. — We compute the decomposition numbers of the unipotent characters lying in the principal ℓ -block of a finite group of Lie type $B_{2n}(q)$ or $C_{2n}(q)$ when q is an odd prime power and ℓ is an odd prime number such that the order of $q \bmod \ell$ is $2n$. Along the way, we extend to these finite groups the results of [12] on the branching graph for Harish-Chandra induction and restriction.

RÉSUMÉ. — Nous calculons les nombres de décomposition des caractères unipotents du ℓ -bloc unipotent d'un groupe fini de type de Lie $B_{2n}(q)$ ou $C_{2n}(q)$ lorsque q est une puissance d'un nombre premier impair et ℓ est un nombre premier impair avec q d'ordre $2n$ dans \mathbb{F}_ℓ . En cours de route, nous étendons à ces groupes finis les résultats de [12] sur le graphe de branchement pour l'induction et la restriction de Harish-Chandra.

Introduction

The representation theory of a finite group of Lie type $G := \mathbf{G}(\mathbb{F}_q)$ over a field of positive characteristic ℓ coprime to q has a close relationship to the representation theory of the Hecke algebra of its Weyl group. The decomposition matrix of the Hecke algebra always embeds as a submatrix of the decomposition matrix of G . When $G = \mathrm{GL}_n(q)$ is the finite general linear group, the square unitriangular submatrix of the decomposition matrix of the unipotent blocks is the same as the decomposition matrix of the q -Schur algebra, a quasihereditary cover of the Hecke algebra of the symmetric group S_n . This is related to the fact that in characteristic 0, $\mathrm{GL}_n(q)$ has exactly one cuspidal irreducible unipotent representation as n

Keywords: Finite groups of Lie type, Decomposition numbers.

2020 Mathematics Subject Classification: 20C20, 20C33.

(*) O. Dudas gratefully acknowledges financial support by the ANR, Project No ANR-16-CE40-0010-01E and by the grant SFB-TRR 195. E. Norton was supported by the grant SFB-TRR 195.

ranges over \mathbb{N} , namely, the trivial representation of $\mathrm{GL}_1(q)$. When G is not of type A , less is understood about the decomposition matrix of the unipotent blocks of G . There are more cuspidal unipotent representations in characteristic 0 which give rise to multiple Hecke and quasihereditary algebras, all of which play a role in the unipotent blocks of G . However, in the general case, the knowledge of the decomposition numbers for these algebras is not enough to determine those of G . In [7] the first author initiated the use of Deligne–Lusztig characters to find the missing numbers. This proved successful in determining decomposition matrices for finite groups of Lie type in small rank, see [8, 9, 10, 26].

In this paper, we are concerned with groups of type B_m and C_m such as $G = \mathrm{SO}_{2m+1}(q)$ and $G = \mathrm{Sp}_{2m}(q)$ for odd q . If n is the order of q^2 in \mathbb{F}_ℓ^\times , the complexity of the decomposition matrix grows with m/n . For that reason we will consider the case where $m = 2n$, which is somehow the simplest case outside of the cyclic defect case. Two situations arise:

- (*linear prime case*) q has order n in \mathbb{F}_ℓ^\times , in which case n is necessarily odd, and the representation theory of G behaves as a type A phenomenon and can be deduced from the representation theory of q -Schur algebras of symmetric groups [25];
- (*unitary prime case*) q has order $2n$ in \mathbb{F}_ℓ^\times . Explicit decomposition matrices in that case were obtained by Okuyama–Waki for $m = 2$ [30] and by Malle and the first author for $m = 4$ [9] and $m = 6$ [10].

Our main result provides a generalisation of the unitary prime case to any even m . To our knowledge it is the first general result for defect 2 blocks of finite groups of Lie type outside of type A phenomena.

Let $\Phi_d(q)$ be the d -th cyclotomic polynomial evaluated at q .

MAIN THEOREM (Theorems 2.1, 2.2, 2.3, 2.6, 2.7, and 2.9). — *Let G be a finite group of Lie type over \mathbb{F}_q of type B_{2n} or C_{2n} for q an odd prime power. Let ℓ be an odd prime number such that the order of q in \mathbb{F}_ℓ^\times is $2n$. Then all but two decomposition numbers of the unipotent characters in the principal ℓ -block of G are known. If $\Phi_{2n}(q)_\ell > 4n$ then both these numbers are 2 and the decomposition numbers are completely known.*

A consequence of our theorem is that this decomposition matrix only depends on the order of $q \bmod \ell$ outside of the case that the prime ℓ is small. This is expected to hold in general, see for example [31, Section 6.2.2].

Under our assumptions on ℓ and q , studying the unipotent characters in the principal ℓ -block is a reasonable restriction. First, any other unipotent

ℓ -block has defect 0 or 1, and its decomposition matrix is known by [19]. Second, the decomposition numbers of the non-unipotent characters in the principal ℓ -block of G may be recovered from those of the unipotent characters and from partial knowledge of the character table of G by [20, 21].

The methods used to obtain the decomposition matrices are two-fold:

- (1) First, we use Harish-Chandra induction and restriction to produce projective indecomposable modules (PIMs);
- (2) Second, we compute the missing PIMs (corresponding to cuspidal simple modules) using some partial information on the decomposition of Deligne–Lusztig characters on PIMs.

For both of these steps, we use a truncated version of the Harish-Chandra induction and restriction coming from the categorical $\widehat{\mathfrak{sl}}_{2n}$ -action on unipotent representations defined in [11]. The recent unitriangularity result in [4] allows us to compute the branching graph for this truncated induction, which provides the missing information for step (2) to be successful.

Acknowledgments

We thank Gunter Malle and Raphaël Rouquier for helpful conversations, and Gunter Malle for perspicacious comments on a draft of this paper. E. Norton also thanks the workshop “Categorification in quantum topology and beyond” at the Erwin Schrödinger Institut, Vienna, January 2019.

1. Representation theory for types B and C

1.1. Combinatorics

1.1.1. Partitions and symbols

Let m be a non-negative integer. A partition λ of m is a non-increasing sequence of non-negative integers $\lambda_1 \geq \lambda_2 \geq \dots \geq 0$ which add up to m . We call m the size of λ and we denote it by $|\lambda|$. Let $s \in \mathbb{Z}$ and λ be a partition of m . The charged β -set of λ is the set

$$\beta_s(\lambda) := \{\lambda_1 + s, \lambda_2 + s - 1, \dots, \lambda_i + s - i + 1, \dots\}.$$

It is a subset of \mathbb{Z} which contains all $z \in \mathbb{Z}$ such that $z \leq s - \#\{\text{non-zero parts of } \lambda\}$.

A bipartition $\lambda = \lambda^1.\lambda^2$ of m consists of a pair (λ^1, λ^2) of partitions such that $|\lambda| := |\lambda^1| + |\lambda^2|$ equals m . If λ^2 (resp. λ^1) is the empty partition we will write $\lambda = \lambda^1$. (resp. $\lambda = .\lambda^2$). Given $\mathbf{s} = (s_1, s_2)$ in \mathbb{Z}^2 , a *charged symbol* with charge \mathbf{s} is a pair $\Lambda = (X, Y)$ where $X = \beta_{s_1}(\lambda^1)$ and $Y = \beta_{s_2}(\lambda^2)$ for some bipartition $\lambda = \lambda^1.\lambda^2$. The *defect* of the charged symbol Λ is $D = s_1 - s_2$. We set $\Lambda^\dagger = (Y, X)$. It is a symbol with charge $\mathbf{s}^* := (s_2, s_1)$ and it is associated to the bipartition $\lambda^2.\lambda^1$. Note that the defect of a symbol should not be confused with the other use of the word “defect” arising in representation theory of finite groups, namely the defect of a block.

Throughout this paper we shall only be working with charged symbols with odd defect and with a specific charge. Given $t \in \mathbb{Z}$, let us define

$$(1.1) \quad \sigma_t := \begin{cases} (t, -1 - t) & \text{if } t \text{ is even,} \\ (-1 - t, t) & \text{if } t \text{ is odd.} \end{cases}$$

Note that $\sigma_{-1-t} = \sigma_t$. A *symbol* $\Lambda = (X, Y)$ is a charged symbol with charge σ_t for some $t \in \mathbb{Z}$. If $X = \{x_1, x_2, x_3, \dots\}$ and $Y = \{y_1, y_2, y_3, \dots\}$, we will represent Λ by

$$\Lambda = \begin{pmatrix} x_1 & x_2 & x_3 & \dots \\ y_1 & y_2 & y_3 & \dots \end{pmatrix}.$$

This convention differs from the usual convention, for example the one in Chevie [29] since we allow both positive and negative defect, and since a symbol in Chevie is necessarily truncated on the right whereas our symbols are infinite to the right. Nevertheless this will be needed to have a consistent action of the i -induction operators on all symbols from the various Harish-Chandra series, and to allow n to grow arbitrarily large, see Section 1.3 and especially Remark 1.4.

Remark 1.1. — The 2-row convention for representing the charged symbol Λ of a bipartition λ with charge \mathbf{s} is the 180°-rotation of the 2-abacus of λ with charge \mathbf{s} as in [23].

We will sometimes find it useful to drop the notation of symbols and work with Young diagrams. The Young diagram of the bipartition λ is the set of triples

$$Y(\lambda) := \left\{ (x, y, j) \in \mathbb{N} \times \mathbb{N} \times \{1, 2\} \mid \begin{array}{l} 1 \leq x \leq \#\{\text{nonzero parts of } \lambda^j\}, \\ 1 \leq y \leq \lambda_x^j \end{array} \right\}.$$

An element $b = (x, y, j) \in Y(\lambda)$ is called a *box* of the Young diagram. We will draw the Young diagram of a bipartition λ by putting the diagrams of λ^1 and λ^2 side by side. For a box (x, y, j) in $Y(\lambda)$, x represents the row

and y the column, with the convention that rows are decreasing in length from top to bottom, as illustrated below for the example $\lambda = 543.21$:

$$Y(543.21) = \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline \end{array} \cdot \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array}$$

1.1.2. Cores and co-cores

Let d be a positive integer and let $\Lambda = (X, Y)$ be a symbol. A d -hook in the top row (resp. in the bottom row) of Λ is a pair $(x, x + d)$ such that $x + d \in X$ and $x \notin X$ (resp. $x + d \in Y$ and $x \notin Y$). Removing a d -hook in the top row amounts to changing Λ to $((X \setminus \{x + d\}) \cup \{x\}, Y)$, and similarly for the bottom row. The d -core is the symbol obtained by recursively removing all possible d -hooks. Removing or adding d -hooks does not change the defect of the symbol.

A d -co-hook of Λ is a pair $(x, x + d)$ such that $x + d \in X$ and $x \notin Y$ or $x + d \in Y$ and $x \notin X$. The co-hook is removed from Λ by removing $x + d$ from X and adding x to Y , or removing $x + d$ from Y and adding x to X , and then exchanging X and Y . Recursively removing all d -co-hooks yields the d -co-core of Λ .

1.1.3. Families

Let $\Lambda = (X, Y)$ be a symbol. The *composition* ϖ_Λ attached to the symbol is the non-increasing sequence $\varpi_\Lambda := (\varpi_1 \geq \varpi_2 \geq \varpi_3 \geq \dots)$ obtained by considering the union of X and Y as a multiset. Since X and Y are β -sets of some partitions, any term in the composition ϖ_Λ occurs at most twice (and all but finitely many terms appearing do).

The dominance order on compositions defines a relation on symbols. We say that two symbols Λ and Λ' lie in the same *family* and we write $\Lambda \equiv \Lambda'$ if $\varpi_\Lambda = \varpi_{\Lambda'}$. In other words, two symbols are in the same family if their multisets of entries are the same. We write $\Lambda \triangleleft \Lambda'$ and we say that Λ' dominates Λ if $\varpi_\Lambda \triangleleft \varpi_{\Lambda'}$, by which we mean $\varpi_\Lambda \neq \varpi_{\Lambda'}$ and $\sum_{i=1}^j \varpi_i \leq \sum_{i=1}^j \varpi'_i$ for all $j \geq 1$. This defines a strict partial order on the set of symbols. We will write $\Lambda \trianglelefteq \Lambda'$ if $\Lambda = \Lambda'$ or $\Lambda \triangleleft \Lambda'$.

Example 1.2. — The following four symbols

$$\begin{pmatrix} 1 & 0 & & -2 & \dots \\ & & -1 & -2 & \dots \end{pmatrix}, \begin{pmatrix} 1 & & -1 & -2 & \dots \\ & 0 & & -2 & \dots \end{pmatrix},$$

$$\begin{pmatrix} & 0 & -1 & -2 & \dots \\ 1 & & & -2 & \dots \end{pmatrix}, \begin{pmatrix} & & & -2 & \dots \\ 1 & 0 & -1 & -2 & \dots \end{pmatrix}$$

form a family attached to the composition $(1, 0, -1, -2, -2, \dots)$. The first three symbols have charge $(0, -1)$ and correspond to the bipartitions 1^2 ., 1.1 and $.2$, whereas the fourth symbol has charge $(-2, 1)$ and corresponds to the empty bipartition. Note that we have only included symbols that have charge σ_t for some $t \in \mathbb{Z}$.

1.2. Unipotent representations of finite reductive groups of type B and C

1.2.1. Representations of finite groups


Let G be any finite group and Λ a commutative ring with unit. We denote by $\Lambda G\text{-mod}$ the abelian category of finitely generated left ΛG -modules. The set of isomorphism classes of irreducible (or simple) objects will be denoted by $\text{Irr}_{\Lambda}G$. We will write $K_0(\Lambda G\text{-mod})$ for the Grothendieck group of the category $\Lambda G\text{-mod}$.

Let ℓ be a prime number. We shall work with representations over fields of characteristic zero and ℓ . For that purpose we fix an ℓ -modular system $(\mathbb{K}, \mathbb{O}, \mathbb{k})$ where \mathbb{K} is an extension of \mathbb{Q}_{ℓ} , the ring of integers \mathbb{O} of \mathbb{K} over \mathbb{Z}_{ℓ} is a complete d.v.r and its residue field \mathbb{k} has characteristic ℓ . Throughout this paper we will assume that this modular system is sufficiently large, so that the algebras $\mathbb{K}G$ and $\mathbb{k}G$ split for any finite group G considered, that is, so that all irreducible representations of G over \mathbb{K} (resp. \mathbb{k}) remain irreducible over any field extension of \mathbb{K} (resp. \mathbb{k}). For example, we may take $\mathbb{k} = \overline{\mathbb{k}}$ and \mathbb{O} to be the Witt ring of \mathbb{k} . We will usually identify $K_0(\mathbb{K}G\text{-mod})$ with the space of virtual characters of G , and its basis $\text{Irr}_{\mathbb{K}}G$ by the set of (ordinary) irreducible characters. We will denote by $\langle -; - \rangle_G$ the usual inner product on $K_0(\mathbb{K}G\text{-mod})$.

1.2.2. Finite reductive groups and Deligne–Lusztig characters

We fix a non-negative integer m . Let \mathbf{G} be a connected reductive group, quasi-simple of type B_m or C_m , defined over the finite field \mathbb{F}_q . Let $F : \mathbf{G} \rightarrow \mathbf{G}$ be the corresponding Frobenius endomorphism. The finite group $G := \mathbf{G}^F$ is a *finite reductive group*. If \mathbf{H} is any closed subgroup of \mathbf{G} we will denote by $H := \mathbf{H}^F$ the corresponding finite group. We fix an F -stable maximal torus \mathbf{T} of \mathbf{G} contained in an F -stable Borel subgroup \mathbf{B} of \mathbf{G} . We denote by $W_m := N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$ the corresponding Weyl group,

which is of type B_m . The choice of \mathbf{B} defines a subset of simple reflections $S = \{s_1, s_2, \dots, s_m\}$ on which F acts trivially. They are labeled according to the following Coxeter diagram:



The G -conjugacy classes of F -stable maximal tori are parametrized by the conjugacy classes of W_m . Given $w \in W_m$ we will denote by \mathbf{T}_w a maximal torus of type w . Given θ a \mathbb{K} -linear character of T_w , Deligne–Lusztig defined in [6] a virtual character $R_{T_w}^G(\theta)$ of G over \mathbb{K} . We will write $R_w := R_{T_w}^G(1_{T_w})$ for the Deligne–Lusztig character associated to the trivial character of T_w . The irreducible constituents of the various R_w ’s are the *unipotent characters* of G .

1.2.3. Harish-Chandra induction and restriction

Given $I \subseteq S$, we write W_I for the subgroup of W generated by I and $\mathbf{P}_I = \mathbf{B}W_I\mathbf{B}$ for the corresponding standard parabolic subgroup of \mathbf{G} . It has a Levi decomposition $\mathbf{P}_I = \mathbf{L}_I \ltimes \mathbf{U}_I$ where \mathbf{L}_I is the unique Levi complement of \mathbf{P}_I containing \mathbf{T} . The Harish-Chandra induction and restriction functors are defined by

$$R_{L_I}^G := \mathbb{A}G/U_I \otimes_{\mathbb{A}L_I} - \quad \text{and} \quad {}^*R_{L_I}^G := \mathrm{Hom}_G(\mathbb{A}G/U_I, -)$$

where \mathbb{A} is any of the rings $\mathbb{K}, \mathbb{O}, \mathbb{k}$. A $\mathbb{A}G$ -module V is said to be *cuspidal* if ${}^*R_{L_I}^G(V) = 0$ for all $I \subsetneq S$. When $\ell \nmid q$, the functors $({}^*R_{L_I}^G, R_{L_I}^G)$ form a biadjoint pair of exact functors between $\mathbb{A}L_I\text{-mod}$ and $\mathbb{A}G\text{-mod}$.

When $\mathbb{A} = \mathbb{K}$ these functors yield linear maps on characters of G which we will still denote by $R_{L_I}^G$ and ${}^*R_{L_I}^G$. When $I = \emptyset$ we have that $\mathbf{L}_I = \mathbf{T}$ is a split torus. In that case the Harish-Chandra induction and the Deligne–Lusztig map $R_{\mathbf{T}}^G$ defined above coincide, which justifies our notation.

1.2.4. Unipotent characters

We recall here Lusztig’s parametrization of unipotent characters of G (see for example [28, Section 4]). Recall that \mathbf{G} is a quasi-simple group of type B_m or C_m . The finite group G admits a cuspidal unipotent character if and only if $m = t^2 + t$ for some $t \geq 0$. We label such a character by

the charged symbol of the empty bipartition with the charge σ_t , see (1.1), which is given by

$$\Lambda = \begin{cases} \begin{pmatrix} t & t-1 & \dots & \dots & \dots \\ & & -1-t & -2-t & \dots \end{pmatrix} & \text{if } t \text{ is even,} \\ \begin{pmatrix} & & -1-t & -2-t & \dots \\ t & t-1 & \dots & \dots & \dots \end{pmatrix} & \text{if } t \text{ is odd.} \end{cases}$$

More generally, if $t^2+t \leq m$, one can consider the standard Levi subgroup L of G of type B_{t^2+t} or C_{t^2+t} . Then the unipotent characters of G lying in the Harish–Chandra series of the cuspidal unipotent character of L correspond to bipartitions $\lambda = \lambda^1.\lambda^2$ of size $r = m - t^2 - t$. In that case will we write $[\lambda^1.\lambda^2]_{B_{t^2+t}}$ for the corresponding unipotent character, or $[\Lambda]$ if Λ is the symbol of charge σ_t attached to λ . With our convention, the character with smallest degree in the series is $[r.]_{B_{t^2+t}}$ when t is even but $[.r]_{B_{t^2+t}}$ when t is odd. It agrees with the convention in Chevie [29] when t is even but when t is odd, the components of the bipartition must be swapped.

The decomposition of the Deligne–Lusztig characters R_w in terms of symbols was determined by Lusztig. Given an irreducible character χ of W_m over \mathbb{K} we can form the *almost character*

$$R_\chi := \frac{1}{|W_m|} \sum_{w \in W_m} \chi(w)R_w.$$

Then using [28, Theorem 4.23] one can compute the multiplicity $\langle R_\chi; [\Lambda] \rangle_G$ of the unipotent character $[\Lambda]$ for any symbol Λ . Two examples of computations of Deligne–Lusztig characters are given in the appendix.

1.2.5. Unipotent ℓ -blocks

By an ℓ -block B of G we mean a minimal 2-sided ideal of the group algebra $\mathbb{O}G$. We have $B = \mathbb{O}Gb$ for a unique primitive central idempotent b of $\mathbb{O}G$. We will write $\text{Irr}_{\mathbb{K}}B$ for the ordinary irreducible characters lying in B , that is, those irreducible characters $\chi \in \text{Irr}_{\mathbb{K}}G$ such that $\chi(b) \neq 0$. An ℓ -block B is *unipotent* if it contains at least one unipotent character. In particular, the principal ℓ -block, which is the block containing the trivial character, is unipotent. We will denote by $\mathbb{O}G\text{-umod}$ the category of representations over the sum of the unipotent ℓ -blocks of G .

Assume now that ℓ and q are odd. The ℓ -blocks of G were classified by Fong–Srinivasan in [18]. There are two situations, depending on whether ℓ is “linear” or “unitary”. Let d be the multiplicative order of q in \mathbb{F}_ℓ^\times .

- If d is odd, ℓ is said to be a *linear prime* for G . In that case two unipotent characters $[\Lambda]$ and $[\Lambda']$ lie in the same block if and only if the symbols Λ and Λ' have the same d -core. The number of d -hooks that must be removed to reach the d -core is called the weight of the block.
- If d is even, ℓ is said to be a *unitary prime* for G . Set $e := d/2$, the order of q^2 in \mathbb{F}_ℓ^\times . Then two unipotent characters $[\Lambda]$ and $[\Lambda']$ lie in the same block if and only if the symbols Λ and Λ' have the same e -co-core. The number of e -co-hooks that must be removed to reach the e -co-core is called the weight of the block.

We will often refer to an ℓ -block B as a Φ_d -block, where Φ_d stands for the d -th cyclotomic polynomial. This is justified by the fact that many of the properties of B depend only on d rather than on ℓ , see [3, Theorem 5.24]. For example, if $\ell > m$ then any defect group of B is isomorphic to $(\mathbb{Z}/\Phi_d(q)_\ell)^r$ where r is the weight of the block.

1.2.6. Decomposition matrix

Recall that $(\mathbb{K}, \mathbb{O}, \mathbb{k})$ is an ℓ -modular system which is sufficiently large for G . Then every $\mathbb{K}G$ -module admits an integral form over \mathbb{O} , which can then be reduced modulo ℓ to a $\mathbb{k}G$ -module. The image of that module in the Grothendieck group does not depend on the choice of the integral form and we obtain a linear map

$$\mathrm{dec} : K_0(\mathbb{K}G\text{-mod}) \longrightarrow K_0(\mathbb{k}G\text{-mod})$$

called the decomposition map. The decomposition matrix is the matrix of this map in the bases $\mathrm{Irr}_{\mathbb{K}}G$ and $\mathrm{Irr}_{\mathbb{k}}G$. It respects the block decomposition so that we can talk about the decomposition matrix of an ℓ -block. Dually, every projective $\mathbb{k}G$ -module P lifts to a unique projective $\mathbb{O}G$ -module \tilde{P} , up to isomorphism. By the character of P we mean the character of the $\mathbb{K}G$ -module $\mathbb{K}\tilde{P}$. Brauer reciprocity states that the decomposition matrix of G is also the matrix whose columns are the characters of the PIMs (the projective indecomposable $\mathbb{k}G$ -modules) in the basis $\mathrm{Irr}_{\mathbb{K}}G$.

In this paper we shall only be interested in the decomposition matrix of unipotent ℓ -blocks. It is a reasonable restriction since any block is conjecturally Morita equivalent to some unipotent block [2]. This was proved for a large class of non-unipotent blocks in [1]. We say that a PIM is unipotent if it belongs to a unipotent ℓ -block. When ℓ is odd, the unipotent characters form a basic set of the unipotent ℓ -blocks [20, 21]. If in addition q is odd, this basic set is unitriangular with respect to the order on families [4]

(see [32, Section 4] for the description of the order in terms of symbols). This means that there is a labeling of the unipotent PIMs by symbols such that

$$\langle \mathbb{K}\tilde{P}_\Lambda; [\Lambda'] \rangle_G = \begin{cases} 1 & \text{if } \Lambda = \Lambda', \\ 0 & \text{if } \Lambda' \not\leq \Lambda. \end{cases}$$

To avoid cumbersome notation we will denote by $\Psi_{[\Lambda]}$ the unipotent part of the character of the PIM corresponding to the unipotent character $[\Lambda]$ by unitriangularity.

Similarly, we say that a simple $\mathbb{K}G$ -module is unipotent if it belongs to a unipotent ℓ -block. The unitriangularity of the decomposition matrix gives a natural labeling of the unipotent simple $\mathbb{K}G$ -modules by unipotent characters. If $[\Lambda]$ is a unipotent character, we will denote by $S_{[\Lambda]}$ the corresponding simple $\mathbb{K}G$ -module.

1.3. Branching rules

We recall and complete in this section the main result in [11, Section 6] on the branching rules for Harish-Chandra induction and restriction for unipotent representations of groups of type B and C . Throughout this section we will assume that ℓ and q are odd, and that d , the multiplicative order of q in \mathbb{F}_ℓ^\times , is even. In particular we have $d > 1$.

To avoid cumbersome notation we will work with symplectic groups but under our assumption that ℓ is odd, the main result of this section, Theorem 1.7, remains valid for any group of type B or C .

1.3.1. Level 2 Fock spaces

Let $\lambda = \lambda^1.\lambda^2$ be a bipartition, and $\mathbf{s} \in \mathbb{Z}^2$. The *charged content* of a box $b = (x, y, j)$ in the Young diagram $Y(\lambda)$ is

$$\text{co}^{\mathbf{s}}(b) = y - x + s_j.$$

Given another bipartition μ and $c \in \mathbb{Z}$, we write $\mu \setminus \lambda = c$ if there exists a box b of μ with charged content $\text{co}^{\mathbf{s}}(b) = c$ such that the Young diagram of λ is obtained from the Young diagram of μ by removing the box b .

Let $\{e_i, f_i\}_{i \in 0, \dots, d-1}$ be the Chevalley generators of the affine Lie algebra $\widehat{\mathfrak{sl}}_d$. The Fock space $F(\mathbf{s})$ with charge \mathbf{s} is the $\widehat{\mathfrak{sl}}_d$ -module equipped with a

\mathbb{C} -basis $|\lambda, \mathbf{s}\rangle$ labeled by bipartitions on which the Chevalley generators act by

$$e_i|\lambda, \mathbf{s}\rangle = \sum_{\substack{j \equiv i \pmod d \\ \lambda \setminus \mu = j}} |\mu, \mathbf{s}\rangle \quad \text{and} \quad f_i|\lambda, \mathbf{s}\rangle = \sum_{\substack{j \equiv i \pmod d \\ \mu \setminus \lambda = j}} |\mu, \mathbf{s}\rangle.$$

Note that the action of $\widehat{\mathfrak{sl}}_d$ depends only on the class of s_1 and s_2 in \mathbb{Z}/d .

1.3.2. Order on bipartitions

We consider here an order on bipartitions defined by Dunkl–Griffeth in [13, Section 4]. This order depends both on d and a pair of integers $\mathbf{s} = (s_1, s_2) \in \mathbb{Z}^2$. We will omit d from the notation since d is fixed. Let λ be a bipartition and let $Y(\lambda)$ be its Young diagram. If $b = (x, y, j)$ is a box in $Y(\lambda)$ we write $j(b) := j$. If μ is another bipartition and $\mathbf{s} \in \mathbb{Z}^2$ we write $\lambda \preceq_{\mathbf{s}} \mu$ if for all $\alpha \in \mathbb{R}$ and $j = 1, 2$

$$\begin{aligned} & \# \left\{ b \in Y(\lambda) \mid \mathrm{co}^{\mathbf{s}}(b) - j(b)\frac{d}{2} > \alpha \text{ or } \mathrm{co}^{\mathbf{s}}(b) - j(b)\frac{d}{2} = \alpha \text{ and } j(b) \leq j \right\} \\ & \leq \# \left\{ b \in Y(\mu) \mid \mathrm{co}^{\mathbf{s}}(b) - j(b)\frac{d}{2} > \alpha \text{ or } \mathrm{co}^{\mathbf{s}}(b) - j(b)\frac{d}{2} = \alpha \text{ and } j(b) \leq j \right\}. \end{aligned}$$

This is exactly the order \leq_c defined in [13] with $r = 2$, $c_0 = d^{-1}$ and $d_0 = -d_1 = (s_1 - s_2)/d + 1/2$. Note that changing \mathbf{s} to $\mathbf{s} + (s, s)$ for any $s \in \mathbb{Z}$ does not change the order. We will need the following lemma which relates the Dunkl–Griffeth order on bipartitions to the dominance order on charged symbols defined in Section 1.1.3.

LEMMA 1.3. — *Let $\sigma \in \mathbb{Z}^2$ and set $\mathbf{s} := \sigma + \frac{1}{2}(0, d)$. Let λ, μ be two bipartitions, and let Λ, Λ' be the corresponding symbols of charge σ . Then*

$$\lambda \preceq_{\mathbf{s}} \mu \implies \Lambda \triangleleft \Lambda' \text{ or } \Lambda \equiv \Lambda'.$$

Proof. — First observe that given a box b in the Young diagram of λ we have

$$\mathrm{co}^{\mathbf{s}}(b) - j(b)\frac{d}{2} = \mathrm{co}^{\sigma}(b) - \frac{d}{2}.$$

We deduce that $\lambda \preceq_{\mathbf{s}} \mu$ if and only if for all $\alpha \in \mathbb{R}$ and $j = 1, 2$ we have

$$\begin{aligned} (1.2) \quad & \# \{ b \in Y(\lambda) \mid \mathrm{co}^{\sigma}(b) > \alpha \text{ or } \mathrm{co}^{\sigma}(b) = \alpha \text{ and } j(b) \leq j \} \\ & \leq \# \{ b \in Y(\mu) \mid \mathrm{co}^{\sigma}(b) > \alpha \text{ or } \mathrm{co}^{\sigma}(b) = \alpha \text{ and } j(b) \leq j \}. \end{aligned}$$

Let us consider the extended Young diagram $\widetilde{Y}(\lambda)$, defined as the set of boxes $b = (x, y, j)$ with $j = 1, 2$, $x \geq 1$ and $y \leq \lambda_x^j$. Unlike the usual Young diagram we do not assume $y \geq 1$ for the boxes, which means that each row

is infinite on the left. The set of boxes in $\tilde{Y}(\boldsymbol{\lambda}) \setminus Y(\boldsymbol{\lambda})$ does not depend on the bipartition $\boldsymbol{\lambda}$, and the number of boxes with a given content is finite, therefore one can replace $Y(\boldsymbol{\lambda})$ and $Y(\boldsymbol{\mu})$ by $\tilde{Y}(\boldsymbol{\lambda})$ and $\tilde{Y}(\boldsymbol{\mu})$ in (1.2).

Working with extended Young diagrams makes the computations easier in (1.2). Indeed, if $\varpi_\Lambda := (\varpi_1 \geq \varpi_2 \geq \varpi_3 \geq \dots)$ is the composition attached to Λ , that is, the multiset given by the union of $\beta_{\sigma_1}(\lambda^1)$ and $\beta_{\sigma_2}(\lambda^2)$ (see Section 1.1.3), then we claim that

$$\#\{b \in \tilde{Y}(\boldsymbol{\lambda}) \mid \text{co}^\sigma(b) \geq \alpha\} = \sum_{\varpi_k \geq \alpha} (\varpi_k - \lceil \alpha \rceil).$$

To show the claim we can assume without loss of generality that $\alpha \in \mathbb{Z}$ since the contents are integers. Each row in $\tilde{Y}(\boldsymbol{\lambda})$ corresponds to an element ϖ_k , and the highest content in that row equals $\varpi_k - 1$. Consequently, this row contains a box of content β (and only one) if and only if $\varpi_k - 1 \geq \beta$. Therefore

$$\begin{aligned} \#\{b \in \tilde{Y}(\boldsymbol{\lambda}) \mid \text{co}^\sigma(b) \geq \alpha\} &= \sum_{\beta \geq \alpha} \#\{b \in \tilde{Y}(\boldsymbol{\lambda}) \mid \text{co}^\sigma(b) = \beta\} \\ &= \sum_{\beta \geq \alpha} \#\{k \geq 1 \mid \varpi_k - 1 \geq \beta\} \\ &= \sum_{\substack{k, \beta \\ \varpi_k - 1 \geq \beta \geq \alpha}} 1 = \sum_k \sum_{\substack{\beta \\ \varpi_k - 1 \geq \beta \geq \alpha}} 1 \\ &= \sum_{\substack{k \\ \varpi_k - 1 \geq \alpha}} (\varpi_k - \alpha). \end{aligned}$$

Let $\varpi_{\Lambda'} = (\varpi'_1 \geq \varpi'_2 \geq \varpi'_3 \geq \dots)$ be the composition attached to the symbol Λ' . We deduce from (1.2) with $j = 2$ that for all $\alpha \in \mathbb{R}$

$$(1.3) \quad \sum_{\varpi_k \geq \alpha} (\varpi_k - \lceil \alpha \rceil) \leq \sum_{\varpi'_k \geq \alpha} (\varpi'_k - \lceil \alpha \rceil).$$

Now, if $i \geq 1$ we have

$$\sum_{k=1}^i (\varpi_k - \varpi'_i) \leq \sum_{\varpi_k \geq \varpi'_i} (\varpi_k - \varpi'_i) \leq \sum_{\varpi'_k \geq \varpi'_i} (\varpi'_k - \varpi'_i) = \sum_{k=1}^i (\varpi'_k - \varpi'_i)$$

where we used (1.3) with $\alpha = \varpi'_i$ for the second inequality. This shows that $\sum_{k=1}^i \varpi_k \leq \sum_{k=1}^i \varpi'_k$ for all $i \geq 1$ and completes the proof. \square

1.3.3. Categorification of unipotent representations

We recall here the categorification result of [11, Section 6]. For $m \geq 0$ we set $G_m := \text{Sp}_{2m}(q)$. Using the (unique) standard Levi subgroup $G_{m-1} \times$

$\mathrm{GL}_1(q)$ of G_m we can form the chain of subgroups $\{1\} = G_0 \subset G_1 \subset \dots \subset G_{m-1} \subset G_m \subset \dots$. Since d , the multiplicative order of q in \mathbb{F}_ℓ^\times is even, the group $\mathrm{GL}_1(q)$ is an ℓ' -group and the Harish-Chandra induction and restriction induce exact functors between $\mathbb{k}G_{m-1}\text{-mod}$ and $\mathbb{k}G_m\text{-mod}$ for all m . We can form the abelian category

$$\mathbb{k}G_\bullet\text{-mod} := \bigoplus_{m \geq 0} \mathbb{k}G_m\text{-mod}$$

of the modules over all the unipotent ℓ -blocks of the various groups G_m . We will denote by F and E the endofunctors of this category induced by Harish-Chandra induction and restriction respectively.

Since ℓ is odd, the unipotent characters form a basic set for the unipotent blocks [20, 21]. In particular $K_0(\mathbb{k}G_\bullet\text{-mod})$ has a \mathbb{Z} -basis given by the image of the unipotent characters under the decomposition map, see Section 1.2.6. Recall from Section 1.2.4 that the unipotent characters in a B_{t^2+t} -series are labeled by bipartitions, or equivalently by symbols of charge σ_t . Therefore they are in bijection with the standard basis of any Fock space $F(\mathbf{s})$. For our purpose we will consider the charges \mathbf{s}_t defined by

$$(1.4) \quad \mathbf{s}_t := \sigma_t + \frac{1}{2}(0, d) = \begin{cases} (t, -1 - t + \frac{d}{2}) & \text{if } t \text{ is even,} \\ (-1 - t, t + \frac{d}{2}) & \text{if } t \text{ is odd,} \end{cases}$$

for all $t \geq 0$. The previous discussion shows that there is an isomorphism of \mathbb{C} -vector spaces

$$(1.5) \quad \bigoplus_{t \geq 0} F(\mathbf{s}_t) \xrightarrow{\sim} \mathbb{C} \otimes_{\mathbb{Z}} K_0(\mathbb{k}G_\bullet\text{-mod})$$

$$|\boldsymbol{\lambda}, \mathbf{s}_t\rangle \mapsto \mathrm{dec}[\boldsymbol{\lambda}]_{B_{t^2+t}}$$

which sends an element of the standard basis to the image under the decomposition map of the corresponding unipotent character. One of the main results in [11, Section 6] is a categorification of (1.5). It gives, for every $i = 0, \dots, d - 1$ a construction of a biadjoint pair of exact endofunctors (F_i, E_i) of $\mathbb{k}G_\bullet\text{-mod}$ such that

$$F = \bigoplus_{i=0}^{d-1} F_i \quad \text{and} \quad E = \bigoplus_{i=0}^{d-1} E_i$$

called *i*-induction and *i*-restriction functors, which induce an action of $\widehat{\mathfrak{sl}}_d$ on $K_0(\mathbb{k}G_\bullet\text{-mod})$ making (1.5) an isomorphism of $\widehat{\mathfrak{sl}}_d$ -modules.

Remark 1.4. — In [11, (6.3)] the authors used the charge \mathbf{s}_t^* instead of \mathbf{s}_t when t is odd. This does not affect the categorification result since the

Fock spaces $F(\mathbf{s}_i^*)$ and $F(\mathbf{s}_i)$ are clearly isomorphic, but it explains the discrepancy in our notation for unipotent characters with the one in Chevie. With our convention, under the isomorphism (1.5), the action of f_i on a symbol

$$\Lambda = \begin{pmatrix} x_1 & x_2 & x_3 & \dots \\ y_1 & y_2 & y_3 & \dots \end{pmatrix}$$

is given by increasing by 1 any x_j equal to i modulo d or any y_j equal to $i + d/2$ modulo d , when possible.

1.3.4. Crystal graph and branching rules

Fix a charge $\mathbf{s} = (s_1, s_2) \in \mathbb{Z}^2$, let $d \in \mathbb{Z}_{\geq 2}$ (here we do not require d to be even), and consider the Fock space $F(\mathbf{s})$. Let $|\lambda, \mathbf{s}\rangle$ be a charged bipartition in $F(\mathbf{s})$. A box b of λ is *removable* if $\mu := \lambda \setminus b$ is a bipartition. Then b is called an *addable* box of the bipartition μ . For each $i \in \mathbb{Z}/d\mathbb{Z}$, define the *i-word* of $|\lambda, \mathbf{s}\rangle$ as follows: list all the addable and removable boxes b of $|\lambda, \mathbf{s}\rangle$ such that $\text{co}^{\mathbf{s}}(b) \equiv i \pmod d$ in increasing order from left to right according to their value in \mathbb{Z} , with the convention that if $\text{co}^{\mathbf{s}}(b) = \text{co}^{\mathbf{s}}(b')$ and $b \in \lambda^1, b' \in \lambda^2$, then b' is smaller than b . Now replace each addable box in the list by the symbol $+$ and each removable box in the list by the symbol $-$. The resulting string of pluses and minuses is called the *i-word* of $|\lambda, \mathbf{s}\rangle$. The *reduced i-word* of $|\lambda, \mathbf{s}\rangle$ is then found from the *i-word* by recursively canceling all adjacent pairs $(-+)$. The reduced *i-word* is of the form $(+)^a(-)^b$ for some $a, b \in \mathbb{Z}_{\geq 0}$. The *Kashiwara operator* \tilde{f}_i adds the addable *i*-box corresponding to the rightmost $+$ in the reduced *i-word* of $|\lambda, \mathbf{s}\rangle$, or if there is no $+$ in the reduced *i-word* then it acts by 0. Likewise, the Kashiwara operator \tilde{e}_i removes the removable *i*-box corresponding to the leftmost $-$ in the reduced *i-word* of $|\lambda, \mathbf{s}\rangle$, or if there is no $-$ in the reduced *i-word* then it acts by 0. The directed graph with vertices all bipartitions and $(\mathbb{Z}/d\mathbb{Z}$ -colored) edges $\lambda \xrightarrow{i} \mu$ if and only if $\mu = \tilde{f}_i(\lambda)$, $i \in \mathbb{Z}/d\mathbb{Z}$, is called the $\widehat{\mathfrak{sl}}_d$ -*crystal* on $F(\mathbf{s})$ [27, Section 3], [16, Theorem 2.8].

Example 1.5. — Let $\lambda = 421^4.4^232^21^3$, $\mathbf{s} = (1, 0)$, and $d = 4$.

$$|\lambda, \mathbf{s}\rangle = \begin{array}{cccc} \boxed{1} & \boxed{2} & \boxed{3} & \boxed{4} \\ \boxed{0} & \boxed{1} & & \\ \boxed{-1} & & & \\ \boxed{-2} & & & \\ \boxed{-3} & & & \\ \boxed{-4} & & & \end{array} \cdot \begin{array}{cccc} \boxed{0} & \boxed{1} & \boxed{2} & \boxed{3} \\ \boxed{-1} & \boxed{0} & \boxed{1} & \boxed{2} \\ \boxed{-2} & \boxed{-1} & \boxed{0} & \\ \boxed{-3} & \boxed{-2} & & \\ \boxed{-4} & \boxed{-3} & & \\ \boxed{-5} & & & \\ \boxed{-6} & & & \\ \boxed{-7} & & & \end{array}$$

Let us find the 0-word of $|\lambda, \mathbf{s}\rangle$. The addable 0-boxes (x, y, j) , where x is the row, y is the column, and j is the component, are:

$$(9, 1, 2), (6, 2, 2), (3, 2, 1), (1, 5, 2).$$

The removable 0-boxes are:

$$(6, 1, 1), (3, 3, 2), (1, 4, 1).$$

Ordering them we obtain the 0-word:

$$\begin{array}{ccccccc} (9, 1, 2) & (6, 2, 2) & (6, 1, 1) & (3, 3, 2) & (3, 2, 1) & (1, 5, 2) & (1, 4, 1) \\ + & + & - & - & + & + & - \end{array}$$

Iterating cancellations of all adjacent $(-+)$, we obtain the reduced 0-word:

$$\begin{array}{ccc} (9, 1, 2) & (6, 2, 2) & (1, 4, 1) \\ + & + & - \end{array}$$

Thus \tilde{f}_0 adds the box $(6, 2, 2)$ to $|\lambda, \mathbf{s}\rangle$, and \tilde{e}_0 removes the box $(1, 4, 1)$ from $|\lambda, \mathbf{s}\rangle$. That is, $\tilde{f}_0(\lambda) = 421^4.4^232^31^2$ and $\tilde{e}_0(\lambda) = 321^4.4^232^21^3$.

We will often work with symbols instead of Young diagrams, so it is useful to describe how the operators \tilde{f}_i and \tilde{e}_i act on symbols. Let $[\lambda]_{B_{t^2+t}}$ be a bipartition in the B_{t^2+t} series, and let $\Lambda = (X, Y)$ be its symbol with charge σ_t . Recall that this does not depend on d . Instead of changing the symbol to depend on d , we define the action so that it depends on d .

Let $d \in 2\mathbb{N}$ with $d \geq 2$. An addable i -box of λ^1 is $x \in X$ such that $x + 1 \notin X$ and $x \equiv i \pmod{d}$. A removable i -box of λ^1 is $x \in X$ such that $x + 1 \in X$ and $x \equiv i \pmod{d}$. The next condition records the dependence on d . An addable i -box of λ^2 is $y \in Y$ such that $y + 1 \notin Y$ and $y + \frac{d}{2} \equiv i \pmod{d}$. A removable i -box of λ^2 is $y \in Y$ such that $y + 1 \in Y$ and $y + \frac{d}{2} \equiv i \pmod{d}$. We then order all addable and removable i -boxes of λ by the following rule:

- x_1 is less than x_2 if and only if $x_1 < x_2$ in \mathbb{Z} ,
- y_1 is less than y_2 if and only if $y_1 < y_2$ in \mathbb{Z} ,
- x is less than y if and only if $y + \frac{d}{2} > x$ in \mathbb{Z} ,
- y is less than x if and only if $x \geq y + \frac{d}{2}$ in \mathbb{Z} ,

for all addable and removable boxes x, x_1, x_2 of λ^1 and y, y_1, y_2 of λ^2 . We then form the i -word of λ by listing its addable and removable i -boxes in increasing order as just defined. The cancellation rule is the same as before, and we obtain the good addable i -box and the good removable i -box of λ (if they exist) as described before.

Example 1.6. — Take $\lambda = 31^2.2$ and consider it as a bipartition in the principal series, so that $t = 0$. We have $\sigma_0 = (0, -1)$ and thus the symbol associated to $[\lambda]$ is

$$\Lambda = \begin{pmatrix} X \\ Y \end{pmatrix} = \begin{pmatrix} 3 & & 0 & -1 & & -3 & -4 & \dots \\ & 1 & & & -2 & -3 & -4 & \dots \end{pmatrix}$$

Now suppose that $d = 6$. Let us calculate the action of \tilde{f}_3 and \tilde{e}_3 on Λ . The addable 3-boxes of λ are given by $3, -3 \in X$, and the removable 3-boxes of λ are given by $0 \notin Y$. Ordering them according to our rule, the 3-word of λ is:

$$\begin{array}{ccccc} + & & - & & + \\ -3 \in X & & 0 \notin Y & & 3 \in X \end{array}$$

We then cancel the occurrence of $(-+)$, yielding $+$ as the reduced 3-word and $-3 \in X$ as the good addable 3-box. There is no good removable 3-box, so $\tilde{e}_3\Lambda = 0$. Adding the good addable 3-box to λ yields:

$$\tilde{f}_3\Lambda = \begin{pmatrix} 3 & & 0 & -1 & -2 & & -4 & \dots \\ & 1 & & & -2 & -3 & -4 & \dots \end{pmatrix}.$$

Now we recall how the combinatorics of crystals relates to the representation theory of unipotent blocks. We can define the *colored branching graph* whose vertices are labeled by the unipotent simple $\mathbb{k}G_m$ -modules for all $m \geq 0$ and arrows are given by $T \xrightarrow{i} T'$ if T' appears in the head of $F_i(T)$, or equivalently if T appears in the socle of $E_i(T')$.

THEOREM 1.7. — *The map*

$$|\lambda, \mathbf{s}_t\rangle \mapsto S_{[\lambda]_{B_t 2+t}}$$

induces an isomorphism between the union of the crystal graphs of $F(\mathbf{s}_t)$ for $t \geq 0$ and the colored branching graph of Harish-Chandra induction and restriction.

Proof. — The proof follows the arguments given in the proof of [12, Theorem 4.37]. Let $t \geq 0$. There is a perfect basis of the Fock space $F(\mathbf{s}_t)$ coming from an upper global basis of a quantum deformation of $F(\mathbf{s}_t)$ defined by Uglov in [34]. We denote this basis by $b^\vee(|\lambda, \mathbf{s}_t\rangle)$. It is unitriangular in the standard basis, with respect to the order $\preceq_{\mathbf{s}_t}$ defined in Section 1.3. In other words we have

$$(1.6) \quad b^\vee(|\lambda, \mathbf{s}_t\rangle) \in |\lambda, \mathbf{s}_t\rangle + \sum_{\lambda \prec_{\mathbf{s}_t} \mu} \mathbb{C}|\mu, \mathbf{s}_t\rangle.$$

On the other hand, since the decomposition matrix is unitriangular, we have, for every unipotent character $[\Lambda]$ attached to a symbol Λ

$$(1.7) \quad \mathrm{dec}([\Lambda]) \in S_{[\Lambda]} + \sum_{\Lambda \triangleleft \Lambda'} \mathbb{Z}S_{[\Lambda']}$$

in $K_0(\mathbb{k}G_{\bullet}\text{-umod})$. Now let ψ be the isomorphism in (1.5) sending $|\boldsymbol{\lambda}, \mathbf{s}_t\rangle$ to $\mathrm{dec}([\Lambda])$ where Λ is the symbol of charge $\boldsymbol{\sigma}_t$ attached to $\boldsymbol{\lambda}$. Lemma 1.3 tells us that for all $\boldsymbol{\mu}$ such that $\boldsymbol{\lambda} \prec_{\mathbf{s}_t} \boldsymbol{\mu}$, it holds that $\Lambda \triangleleft \Lambda'$ or $\Lambda \equiv \Lambda'$, where Λ' is the symbol of charge $\boldsymbol{\sigma}_t$ attached to $\boldsymbol{\mu}$. Since $\psi(|\boldsymbol{\mu}, \mathbf{s}_t\rangle) \in S_{[\Lambda']} + \sum_{\Lambda' \triangleleft \Lambda''} \mathbb{C}S_{[\Lambda']}$ by (1.7) we deduce that

$$\psi(|\boldsymbol{\mu}, \mathbf{s}_t\rangle) \in S_{[\Lambda']} + \sum_{\Lambda \triangleleft \Lambda''} \mathbb{C}S_{[\Lambda']}$$

This together with (1.6) gives

$$(1.8) \quad \psi(b^\vee(|\boldsymbol{\lambda}, \mathbf{s}_t\rangle)) \in S_{[\Lambda]} + \sum_{\Lambda \prec \Lambda'} \mathbb{C}S_{[\Lambda']}$$

where $\Lambda \prec \Lambda'$ is the transitive closure of the relation

$$\Lambda \triangleleft \Lambda', \text{ or } \Lambda' \leftrightarrow \boldsymbol{\mu} \text{ with } \boldsymbol{\lambda} \prec_{\mathbf{s}_t} \boldsymbol{\mu}.$$

If we define φ to be the bijection that sends $S_{[\Lambda]}$ to $\psi(b^\vee(|\boldsymbol{\lambda}, \mathbf{s}_t\rangle))$, two perfect bases of $\bigoplus_{t \geq 0} \mathbb{F}(\mathbf{s}_t)$, then (1.8) shows that φ satisfies the assumptions of [12, Proposition 1.14] with respect to the order \prec and therefore induces a crystal isomorphism. \square

2. The decomposition matrix of the principal Φ_{2n} -block of $\mathrm{Sp}_{4n}(q)$ and $\mathrm{SO}_{4n+1}(q)(\mathbf{q})$

We fix an integer $n \geq 0$. Throughout this section $G = G_{2n}$ will denote one of the finite groups $\mathrm{SO}_{4n+1}(q)$ or $\mathrm{Sp}_{4n}(q)$ for q a power of an odd prime. We are interested in the decomposition matrix of the principal ℓ -block of G when the order of q in \mathbb{F}_ℓ^\times equals $2n$. We will often refer to this block as the principal Φ_{2n} -block of G (see Section 1.2.5).

2.1. Unipotent characters in the principal Φ_{2n} -block

Let B be the principal Φ_{2n} -block of G and b be the corresponding block idempotent. By Section 1.2.5 the block has weight 2 and the unipotent

characters lying in B are labeled by symbols of rank $2n$ and n -co-core equal to the symbol

$$(2.1) \quad \begin{pmatrix} 0 & -1 & -2 & \dots \\ & -1 & -2 & \dots \end{pmatrix}.$$

Consequently one obtains such symbols by adding two n -co-hooks to the symbol (2.1). Depending on which rows the co-hooks are inserted in, one gets symbols of defect 1, -3 or 5 which label unipotent characters in the principal series, the B_2 -series, or the B_6 -series of G respectively, see Section 1.2.4.

(1) *Principal series.* There are three families of symbols of defect 1 obtained by adding two n -co-hooks to the symbol (2.1).

- For $0 \leq j \leq n$ and $0 < i < n$, the symbol

$$\begin{pmatrix} n-i & 0 & -1 & \dots & \widehat{-j} & \dots \\ n-j & & -1 & \dots & \widehat{-i} & \dots \end{pmatrix}$$

corresponding to the principal series character

$$[(n-i)1^j.(n-j+1)1^{i-1}].$$

There are $n^2 - 1$ such symbols.

- For $0 \leq j < 2n$, the symbol

$$\begin{pmatrix} 2n-j & 0 & -1 & \dots & \widehat{-j} & \dots \\ & & -1 & \dots & \dots & \dots \end{pmatrix}$$

corresponding to the principal series character

$$[(2n-j)1^j].$$

There are $2n$ such symbols.

- For $0 < i \leq 2n$, the symbol

$$\begin{pmatrix} & 0 & -1 & \dots & \dots & \dots \\ 2n-i & & -1 & \dots & \widehat{-i} & \dots \end{pmatrix}$$

corresponding to the principal series character

$$[(2n-i+1)1^{i-1}].$$


Again, there are $2n$ such symbols.

(2) *B_2 -series.* The symbols of defect -3 are given by

$$\begin{pmatrix} & 0 & -1 & \dots & \widehat{-i} & \dots & \widehat{-j} & \dots \\ n-i & n-j & -1 & \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

for $0 \leq i < j \leq n$. There are $n(n + 1)/2$ such symbols. They correspond to the unipotent characters $[2^i 1^{j-i-1} \cdot (n-i-1)(n-j)]_{B_2}$, all of which lie in the B_2 -series.

In terms of bipartitions and their Young diagrams, the unipotent characters $[\lambda]_{B_2}$ in the principal Φ_{2n} -block are labeled by λ such that λ^1 fits inside the rectangle 2^{n-1} , and λ^2 is the reflection across a diagonal line of slope 1 of the skew shape $2^{n-1} \setminus \lambda^1$ (in particular, λ^2 fits in the rectangle $(n - 1)^2$). This is best illustrated with a picture (here, for $n = 7$, we have shaded $\lambda = 2^3 1^2 \cdot 31$):



(3) B_6 -series. The symbols of defect 5 are given by


$$\left(\begin{matrix} n-i & n-j & 0 & -1 & \dots & \dots & \dots & \dots & \dots \\ & & & -1 & \dots & \widehat{-i} & \dots & \widehat{-j} & \dots \end{matrix} \right)$$

for $0 < i < j < n$. There are $(n - 1)(n - 2)/2$ such symbols. They correspond to the unipotent characters

$$[(n-i-2)(n-j-1) \cdot 2^{i-1} 1^{j-i-1}]_{B_6},$$

all lying in the B_6 -series.

In terms of bipartitions and their Young diagrams, the unipotent characters $[\lambda]_{B_6}$ in the principal Φ_{2n} -block are labeled by λ such that λ^2 fits inside the rectangle 2^{n-3} , and λ^1 is the reflection across a diagonal line of slope 1 of the skew shape $2^{n-3} \setminus \lambda^2$ (in particular, λ^1 fits in the rectangle $(n - 3)^2$). Again, this is best illustrated with a picture (here, for $n = 7$, we have shaded $\lambda = 41 \cdot 1^3$):



Consequently there are $2n^2 + 3n$ unipotent characters in the principal Φ_{2n} -block. By [3], the number of unipotent characters in the principal Φ_{2n} -block may also be counted by the number of complex irreducible representations of the complex reflection group $G(2n, 1, 2)$.

2.2. Induced columns

In this section, we describe the columns of the decomposition matrix of the principal Φ_{2n} -block obtained by Harish-Chandra induction from proper Levi subgroups. These account for all but four columns of the decomposition matrix. In all cases, we find that each non-cuspidal column in the principal Φ_{2n} -block is the character of an induced PIM, cut to the principal block.

Since a group of type A_r has no cuspidal unipotent module (over \mathbb{k}) unless $r = 0$ or $r \geq 2n - 1$, a cuspidal support of a simple unipotent $\mathbb{k}G$ -module corresponds to either a Levi subgroup of type A_{2n-1} or a Levi subgroup of type BC_r for some $r \leq 2n$. In particular, if Λ is a symbol of defect $D \neq 1$ then $S_{[\Lambda]}$ is a cuspidal $\mathbb{k}G$ -module whenever $\tilde{e}_i \Lambda = 0$ for all i .

2.2.1. The B_2 -series submatrix

THEOREM 2.1. — *Let λ be a bipartition such that $[\lambda]_{B_2}$ belongs to the principal Φ_{2n} -block. If $\lambda = 2^{n-1}$. then $S_{[\lambda]_{B_2}}$ is cuspidal and so $\Psi_{[\lambda]_{B_2}}$ cannot be obtained by Harish-Chandra induction. Otherwise, using truncated induction we obtain the unipotent parts of projective indecomposable characters in the B_2 -series as follows:*

- If $\lambda = 2^k 1^{n-1-k} . (n-1-k)$ for some $0 \leq k < n-1$, then

$$\Psi_{[\lambda]_{B_2}} = [\lambda]_{B_2} + [2^{k+1} 1^{n-2-k} . (n-2-k)]_{B_2}.$$

- If $\lambda = 2^{n-2} . 1^2$ then

$$\Psi_{[\lambda]_{B_2}} = [\lambda]_{B_2} + [2^{n-2} 1 \cdot 1]_{B_2}.$$

- If $\lambda = 2^{n-1-k} . k^2$ for some $2 \leq k \leq n-1$ then

$$\begin{aligned} \Psi_{[\lambda]_{B_2}} = [\lambda]_{B_2} + [2^{n-1-k} 1 . k(k-1)]_{B_2} + [2^{n-k} 1 . (k-1)(k-2)]_{B_2} \\ + [2^{n+1-k} . (k-2)^2]_{B_2}. \end{aligned}$$

- If $\lambda = 2^i 1^{j-i-1} . (n-i-1)(n-j)$ with $0 < i+1 < j < n$ then

$$\begin{aligned} \Psi_{[\lambda]_{B_2}} = [\lambda]_{B_2} + [2^{i+1} 1^{j-i-2} . (n-i-2)(n-j)]_{B_2} \\ + [2^i 1^{j-i} . (n-i-1)(n-j-1)]_{B_2} + [2^{i+1} 1^{j-i-1} . (n-i-2)(n-j-1)]_{B_2}. \end{aligned}$$

Proof. — If $\lambda = 2^{n-1}$, then $\tilde{e}_i \lambda = 0$ for all $i \in \mathbb{Z}/2n\mathbb{Z}$. Since $S_{[\lambda]_{B_2}}$ does not have cuspidal support on a type A parabolic, it follows that $S_{[\lambda]_{B_2}}$ is cuspidal. For the unipotent part of the characters of the projective covers of non-cuspidal simple representations in the B_2 -series, we will find $\Psi_{[\lambda]_{B_2}}$ by taking the i -induction of $\Psi_{[\mu]_{B_2}}$ for some $i \in \mathbb{Z}/2n\mathbb{Z}$ such that $\tilde{f}_i \mu = \lambda$. We find that in this case, $f_i \Psi_{[\mu]_{B_2}}$ is indecomposable and equals $\Psi_{[\lambda]_{B_2}}$. Note that the ℓ -blocks of G_{2n-1} have either trivial or cyclic defect groups, so that the PIMs can be easily obtained from [19].

Let $\lambda = 2^k 1^{n-1-k} \cdot (n-1-k)$ for some $0 \leq k < n-1$. Set

$$\mu = 2^k 1^{n-1-k} \cdot (n-2-k).$$

Then $\lambda = \tilde{f}_{2n-k-1} \mu$. We have $\Psi_{[\mu]_{B_2}} = [\mu]_{B_2}$ by [19]. (Note that this implies the cuspidal support of $S_{[\mu]_{B_2}}$, and thus also of $S_{[\lambda]_{B_2}}$, is different from $[B_2]$, and is labeled by some non-empty bipartition in the B_2 -series, see [19].) Then $\Psi_{[\lambda]_{B_2}} = \Psi_{[\tilde{f}_{2n-k-1} \mu]_{B_2}}$ is a summand of

$$f_{2n-k-1} \Psi_{[\mu]_{B_2}} = f_{2n-k-1} [\mu]_{B_2} = [\lambda]_{B_2} + [2^{k+1} 1^{n-2-k} \cdot (n-2-k)]_{B_2}.$$

Thus either $f_{2n-k-1} \Psi_{[\mu]_{B_2}}$ is the unipotent part of an indecomposable projective character and equals $\Psi_{[\lambda]_{B_2}}$, or it is the sum of the unipotent parts of two indecomposable projective characters, each of which would have to contain a single unipotent character. If $k = n-2$ then $\Psi_{[2^{k+1} 1^{n-2-k} \cdot (n-2-k)]_{B_2}} = \Psi_{[2^{n-1}]_{B_2}}$ does not appear as a summand of any induced projective character by cuspidality. Therefore $\Psi_{[\lambda]_{B_2}} = [\lambda]_{B_2} + [2^{k+1} 1^{n-2-k} \cdot (n-2-k)]_{B_2}$ when $k = n-2$. Then by downwards induction on k , $f_{2n-1-k} \Psi_{[\mu]_{B_2}} = \Psi_{[\lambda]_{B_2}}$ for all $0 \leq k < n-2$ as well, and in particular, the desired character formula holds.

Next, let $\lambda = 2^{n-2} \cdot 1^2$. Set $\mu = 2^{n-2} \cdot 1$. Then $\lambda = \tilde{f}_n \mu$. We have $\Psi_{[\mu]_{B_2}} = [\mu]_{B_2}$ by [19]. Therefore $\Psi_{[\lambda]_{B_2}} = \Psi_{[\tilde{f}_n \mu]_{B_2}}$ is a summand of

$$f_n \Psi_{[\mu]_{B_2}} = f_n [\mu]_{B_2} = [\lambda]_{B_2} + [2^{n-2} \cdot 1]_{B_2},$$

thus either $\Psi_{[\lambda]_{B_2}} = [\lambda]_{B_2}$ and $[2^{n-2} \cdot 1]_{B_2} = \Psi_{[2^{n-2} \cdot 1]_{B_2}}$, or $\Psi_{[\lambda]_{B_2}} = f_n \Psi_{[\mu]_{B_2}}$. We showed in the previous paragraph that

$$\Psi_{[2^{n-2} \cdot 1]_{B_2}} = [2^{n-2} \cdot 1]_{B_2} + [2^{n-1}]_{B_2}.$$

Therefore $\Psi_{[\lambda]_{B_2}} = f_n \Psi_{[\mu]_{B_2}}$ and the desired character formula holds.

In the case that $\lambda = 2^{n-1-k} \cdot k^2$ for $2 \leq k \leq n-1$, we have $\lambda = \tilde{f}_{n+k-1} 2^{n-1-k} \cdot k(k-1)$. In the case that $\lambda = 2^i 1^{j-i-1} \cdot (n-i-1)(n-j)$ for $1 < j-i < n$, we have $\lambda = \tilde{f}_{2n-i-1} 2^i 1^{j-i-1} \cdot (n-i-2)(n-j)$. By [19] the

following characters are the unipotent part of PIMs of $G_{2n-1}(q)$:

$$\begin{aligned} \Psi_{[2^{n-1-k}.k(k-1)]_{B_2}} &= [2^{n-k-1}.k(k-1)]_{B_2} + [2^{n-k}1.(k-2)^2]_{B_2}, \\ \Psi_{[2^i 1^{j-i-1}.(n-i-2)(n-j)]_{B_2}} &= [2^i 1^{j-i-1}.(n-i-2)(n-j)]_{B_2} \\ &\quad + [2^i 1^{j-i}.(n-i-2)(n-j-1)]_{B_2}. \end{aligned}$$

In either case, the cuspidal support of $S_{[\lambda]_{B_2}}$ is then equal to $[B_2]$, see [19] (and the source vertex of λ in the $\widehat{\mathfrak{sl}}_{2n}$ -crystal is the empty bipartition). This means that if ν is one of the bipartitions of the first two types listed in the theorem or $\nu = 2^{n-1}.$, then $\Psi_{[\nu]_{B_2}}$ cannot appear as a summand of $f_{n+k-1}\Psi_{[2^{n-1-k}.k(k-1)]_{B_2}}$ or $f_{2n-i-1}\Psi_{[2^i 1^{j-i-1}.(n-i-2)(n-j)]_{B_2}}$.

We compute that

$$\begin{aligned} f_{n+k-1}\Psi_{[2^{n-1-k}.k(k-1)]_{B_2}} &= [2^{n-1-k}.k^2]_{B_2} + [2^{n-1-k}1.(k)(k-1)]_{B_2} \\ &\quad + [2^{n-k}1.(k-1)(k-2)]_{B_2} + [2^{n+1-k}.(k-2)^2]_{B_2}, \\ f_{2n-i-1}\Psi_{[2^i 1^{j-i-1}.(n-i-2)(n-j)]_{B_2}} &= [2^i 1^{j-i-1}.(n-i-1)(n-j)]_{B_2} + [2^{i+1}1^{j-i-2}.(n-i-2)(n-j)]_{B_2} \\ &\quad + [2^i 1^{j-i}.(n-i-1)(n-j-1)]_{B_2} + [2^{i+1}1^{j-i-1}.(n-i-2)(n-j-1)]_{B_2}. \end{aligned}$$

By induction on the number of boxes in λ^2 , together with the fact that no $\Psi_{[\nu]_{B_2}}$ as above can be a summand of these induced projective characters, it follows that for none of the bipartitions ρ appearing on the right-hand side of these formulas can $\Psi_{[\rho]_{B_2}}$ be a summand except for $\rho = \lambda$. That is,

$$f_{n+k-1}\Psi_{[2^{n-1-k}.k(k-1)]_{B_2}} = \Psi_{[2^{n-1-k}.k^2]_{B_2}}$$

and

$$f_{2n-i-1}\Psi_{[2^i 1^{j-i-1}.(n-i-2)(n-j)]_{B_2}} = \Psi_{[2^i 1^{j-i-1}.(n-i-1)(n-j)]_{B_2}}.$$

We thus get the desired character formulas for $\Psi_{[\lambda]_{B_2}}$ in the final two cases listed in the theorem. This concludes the proof. \square

2.2.2. The B_6 -series submatrix

The proof of Theorem 2.2 is identical to the proof of Theorem 2.1, except that the roles of λ^1 and λ^2 are switched.

THEOREM 2.2. — *Let λ be a bipartition such that $[\lambda]_{B_6}$ belongs to the principal Φ_{2n} -block. If $\lambda = .2^{n-3}$ then $S_{[\lambda]_{B_6}}$ is cuspidal and so $\Psi_{[\lambda]_{B_6}}$*

cannot be obtained by Harish-Chandra induction. Otherwise, using truncated induction we obtain the unipotent parts of projective indecomposable characters in the B_6 -series as follows:

- If $\lambda = (n-3-k).2^k 1^{n-3-k}$ for some $0 \leq k < n-3$ then

$$\Psi_{[\lambda]_{B_6}} = [\lambda]_{B_6} + [(n-4-k).2^{k+1} 1^{n-4-k}]_{B_6}.$$

- If $\lambda = 1^2.2^{n-4}$ then

$$\Psi_{[\lambda]_{B_6}} = [\lambda]_{B_6} + [1.2^{n-4} 1]_{B_6}.$$

- If $\lambda = k^2.2^{n-3-k}$ for some $2 \leq k \leq n-3$ then

$$\begin{aligned} \Psi_{[\lambda]_{B_6}} = [\lambda]_{B_6} + [(k)(k-1).2^{n-3-k} 1]_{B_6} + [(k-1)(k-2).2^{n-2-k} 1]_{B_6} \\ + [(k-2)^2.2^{n-1-k}]_{B_6}. \end{aligned}$$

- If $\lambda = (n-i-2)(n-j-1).2^{i-1} 1^{j-i-1}$ for some $0 < i < j-1 < n-2$ then

$$\begin{aligned} \Psi_{[\lambda]_{B_6}} = [\lambda]_{B_6} + [(n-i-3)(n-j-1).2^i 1^{j-i-2}]_{B_6} \\ + [(n-i-2)(n-j-2).2^{i-1} 1^{j-i}]_{B_6} \\ + [(n-i-3)(n-j-2).2^i 1^{j-i-1}]_{B_6}. \end{aligned}$$

2.2.3. The principal series submatrix

We will give explicit formulas in the theorem below, but first, for the sake of intuition we sketch a visual and conceptual way to describe the unipotent constituents of the projective indecomposable characters in the principal Φ_{2n} -block that belong to the principal series, inspired by [5]. One can make a simple graph with vertices the bipartitions λ of $2n$ belonging to the principal Φ_{2n} -block, and an edge between λ and μ if and only if μ is obtained from λ by moving either a single row or a single column of boxes preserving their charged contents mod d . In this case, we place μ below λ if $\mu \prec_{s_0} \lambda$. We refer the reader to [24, Section 5.4–5.5] for a picture of these posets (where they are drawn as simple directed graphs). The resulting graph is planar.

One can then make a CW-complex where 2-cells fill the diamond-and-triangle-shaped regions of the graph, each edge is a 1-cell, and each vertex a 0-cell. Each vertex λ in the graph is then the top vertex of the closure of a cell of maximal possible dimension, or the top vertex of a union of closures of cells sharing that top vertex. These shapes are either a diamond, a triangle, a single edge, a pair of edges (this happens only for $\lambda = 1^{n+1}.1^{n-1}$), or

simply the vertex itself. There turn out to be two such shapes that are points, and these are the two cuspidals. For the λ that are top vertices of 1- and 2-dimensional shapes, taking all vertices in the boundary of the shape and reading off their labels μ yields the unipotent constituents $[\mu]$ of $\Psi_{[\lambda]}$. Moreover, those λ at the top of 2-dimensional shapes label the irreducible representations of the Hecke algebra, while the λ that are only at the top of 1-dimensional shapes label the simple modules with cuspidal support in some non-trivial, proper parabolic. All of this is just an interpretation of the formulas in the following theorem, which are the same formulas as for the principal block of Category \mathcal{O} of the rational Cherednik algebra of B_{2n} at parameters $(\frac{1}{2n}, \frac{1}{2n})$ [24].

THEOREM 2.3. — *Let λ be a bipartition of $2n$ such that $[\lambda]$ belongs to the principal Φ_{2n} -block. If $\lambda = 1^{2n}$. or $.1^{2n}$ then $S_{[\lambda]}$ is cuspidal and so $\Psi_{[\lambda]}$ cannot be obtained by Harish-Chandra induction. Otherwise, the column of the decomposition matrix labeled by λ is given by the appropriate choice of formula below:*

- If $\lambda = (n+k)1^{n-k}$. for some $1 \leq k \leq n$, then

$$\Psi_{[(n+k)1^{n-k}.]} = [(n+k)1^{n-k}.] + [(n+k-1)1^{n-k+1}.] \\ + [(n-1)1^{n-k}.k+1] + [(n-1)1^{n-k+1}.k].$$

- If $\lambda = n1^n$. then

$$\Psi_{[n1^n.]} = [n1^n.] + [(n-1)1^n.1] + [(n-1)1^{n+1}.].$$

- If $\lambda = .(2n-k)1^k$ for some $0 \leq k \leq n-3$ then

$$\Psi_{[.(2n-k)1^k]} = [.(2n-k)1^k] + [n-1-k.(n+1)1^k] \\ + [n-k-2.(n+1)1^{k+1}] + [.(2n-k-1)1^{k+1}].$$

- If $\lambda = .(n+2)1^{n-2}$ then

$$\Psi_{[.(n+2)1^{n-2}]} = [.(n+2)1^{n-2}] + [1.(n+1)1^{n-2}] + [.(n+1)1^{n-1}].$$

- If $\lambda = 1^{k+1}.(n+1-k)1^{n-2}$ for some $0 \leq k \leq n-1$ then

$$\Psi_{[1^{k+1}.(n+1-k)1^{n-2}]} = [1^{k+1}.(n+1-k)1^{n-2}] \\ + [1^{k+2}.(n-k)1^{n-2}] + [.(n+1-k)1^{n-1+k}] + [.(n-k)1^{n+k}].$$

- If $\lambda = (n-k)1^n.1^k$ for some $1 \leq k \leq n-2$ then

$$\Psi_{[(n-k)1^n.1^k]} = [(n-k)1^n.1^k] + [(n-k-1)1^n.1^{k+1}] \\ + [(n-k)1^{n+k}.] + [(n-k-1)1^{n+k+1}.].$$

- If $\lambda = (n - i)1^j.(n - j + 1)1^{i-1}$ for some $1 \leq i \leq n - 2$ and some $0 \leq j \leq n - 1$ then

$$\Psi_{[\lambda]} = [(n-i)1^j.(n-j+1)1^{i-1}] + [(n-i)1^{j+1}.(n-j)1^{i-1}] \\ + [(n-i-1)1^j.(n-j+1)1^i] + [(n-i-1)1^{j+1}.(n-j)1^i].$$

- If $\lambda = 1^{n+1}.1^{n-1}$ then

$$\Psi_{[1^{n+1}.1^{n-1}]} = [1^{n+1}.1^{n-1}] + [1^{2n}.] + [1^{2n}].$$

- If $\lambda = k1^{2n-k}$. for some $2 \leq k \leq n - 1$ then

$$\Psi_{[(k)1^{2n-k}.]} = [k1^{2n-k}.] + [(k-1)1^{2n-k+1}.].$$

- If $\lambda = .k1^{2n-k}$ for some $2 \leq k \leq n + 1$ then

$$\Psi_{[.k1^{2n-k}]} = [.k1^{2n-k}] + [(k-1)1^{2n-k+1}].$$

Proof. — The first seven cases deal with λ such that λ labels an irreducible representation of the Hecke algebra. The formulas are then given by [14]. They may easily be checked to coincide with $F_i\Psi_{[\mu]}$ for μ and i such that $\tilde{f}_i\mu = \lambda$.

Let L be the standard Levi subgroup of G of type A_{2n-1} . If $\lambda = 1^{n+1}.1^{n-1}$, we find that $\Psi_{[\lambda]}$ occurs as a summand of $R_L^G\Psi_{1^{2n}}$ by maximality of the family of λ among the families of the three unipotent characters occurring in $[\lambda] + [1^{2n}.] + [1^{2n}]$, which equals $R_L^G\Psi_{1^{2n}}$ cut to the principal Φ_{2n} -block. Write

$$\Psi_{[1^{n+1}.1^{n-1}]} = [1^{n+1}.1^{n-1}] + \alpha[1^{2n}.] + \beta[1^{2n}]$$

for some $\alpha, \beta \in \{0, 1\}$. We calculate that

$$e_n\Psi_{[1^{n+1}.1^{n-1}]} = [1^n.1^{n-1}] + \beta[1^{2n-1}]$$

which is only a projective character if $\beta \geq 1$. Then we calculate that

$$e_1\Psi_{[1^{n+1}.1^{n-1}]} = [1^{n+1}.1^{n-2}] + \alpha[1^{2n-1}.]$$

which is only a projective character if $\alpha \geq 1$. We conclude that $\alpha = \beta = 1$.

Now we are ready to show that $S_{[1^{2n}.]}$ and $S_{[1^{2n}]}$ are cuspidal. Since their projective covers are not summands of $R_L^G\Psi_{1^{2n}}$, these two simple modules are cuspidal if their Harish-Chandra restriction to a standard Levi subgroup of type BC_{2n-1} is 0. This is confirmed by checking that $\tilde{e}_i(1^{2n}.) = 0 = \tilde{e}_i(1^{2n})$ for all $i \in \mathbb{Z}/2n\mathbb{Z}$. Indeed, for either bipartition, there is only one removable box, and it is canceled by the addable box of the same residue in the first row of the same component.

It remains to verify the formulas in the last two cases listed in the theorem. Consider first the case $\lambda = k1^{2n-k}$. for some $2 \leq k \leq n - 1$.

We find that $\tilde{e}_{k-1}\lambda = (k - 1)1^{2n-k}$, which belongs to a defect 1 block. By [19], $\Psi_{[(k-1)1^{2n-k}]} = [(k - 1)1^{2n-k}]$. Therefore $\Psi_{[\lambda]}$ is a summand of $f_{k-1}\Psi_{[(k-1)1^{2n-k}]} = f_{k-1}[(k - 1)1^{2n-k}] = [k1^{2n-k}] + [(k-1)1^{2n-k+1}]$. Now we argue by induction on k that this is a projective indecomposable character. The base case is $k = 2$: since $S_{[1^{2n}]}$ is cuspidal, $\Psi_{[1^{2n}]}$ is not a summand of the projective character $f_0\Psi_{[1^{2n-1}]} = [21^{2n-2}] + [1^{2n}]$. The induction step says that $\Psi_{[k1^{2n-k}]}$ has the desired formula, and therefore is not a summand of the projective character $f_k\Psi_{[k1^{2n-k-1}]} = [(k + 1)1^{2n-k-1}] + [k1^{2n-k}]$. Finally, the argument for the case $\lambda = .k1^{2n-k}$ for some $2 \leq k \leq n + 1$ is similar. \square

2.3. Cuspidal columns

In the previous section we have accounted for all the columns of the decomposition matrix of the principal Φ_{2n} -block except the ones corresponding by unitriangularity to the unipotent characters

$$[.1^{2n}], \quad [1^{2n}.], \quad [2^{n-1}.]_{B_2}, \quad [.2^{n-3}]_{B_6}.$$

The corresponding projective indecomposable modules (PIMs) are the projective covers of cuspidal simple modules. The purpose of this section is to determine those remaining columns explicitly. As before, we will denote by

$$\Psi_{[.1^{2n}]}, \quad \Psi_{[1^{2n}.]}, \quad \Psi_{[2^{n-1}.]_{B_2}}, \quad \Psi_{[.2^{n-3}]_{B_6}}$$

the unipotent part of the characters of the corresponding PIMs. By unitriangularity of the decomposition matrix and maximality of $.1^{2n}$ in the partial order (by a -value, or by the reverse dominance order on families), it holds that $\Psi_{[.1^{2n}]} = [.1^{2n}]$, the Steinberg character. For the remaining three columns we determine the decomposition numbers using arguments from Deligne–Lusztig theory and from Kac–Moody categorification (truncated induction). The latter technique allows us to study the image of the corresponding PIMs under certain sequences of i -induction functors, providing us with an argument that all or all but one of the decomposition numbers (excepting the one on the diagonal) are zero. Then, two different Deligne–Lusztig characters give an upper and lower bound for the unique non-zero decomposition number, which turn out to coincide. Before treating each PIM in turn, we start with the following lemma.

LEMMA 2.4. — Let Λ be a symbol corresponding to a unipotent character $[\Lambda]$ of G . Let \mathcal{S} be a set of symbols $\Lambda' < \Lambda$ such that

$$\Psi_{[\Lambda]} \in [\Lambda] + \sum_{\Lambda' \in \mathcal{S}} \mathbb{N}[\Lambda'].$$

Let $\mathbf{i} = (i_1, i_2, \dots, i_r)$ be a tuple of elements of $\mathbb{Z}/2n$ and let $\mathbf{i}^* = (i_r, i_{r-1}, \dots, i_1)$ be the reverse tuple. Assume that

- (i) $f_{\mathbf{i}}\Lambda = \widetilde{f}_{\mathbf{i}}\Lambda = \Theta$ for some symbol Θ ;
- (ii) if $\Lambda' \in \mathcal{S}$ and Θ' occurs in $f_{\mathbf{i}}\Lambda'$ with $\Theta' < \Theta$ then $e_{\mathbf{i}^*}\Theta' = \Lambda'$.

Then $\Psi_{[\Lambda]}$ contains only $[\Lambda]$ and the characters $[\Lambda']$ for symbols $\Lambda' \in \mathcal{S}$ such that there exists Θ' occurring in $f_{\mathbf{i}}\Lambda'$ with $\Theta' < \Theta$.

Proof. — Recall that $P_{[\Lambda]}$ denotes the PIM corresponding to the unipotent character $[\Lambda]$ by unitriangularity, and that $\Psi_{[\Lambda]}$ is the unipotent part of its character. Since i -induction functors are exact, the module $f_{\mathbf{i}}P_{[\Lambda]}$ is projective, and by (i) it contains $P_{[\Theta]}$ as a direct summand. Therefore any unipotent character $[\Theta'] \neq [\Theta]$ occurring in $\Psi_{[\Theta]}$ corresponds to a symbol Θ' which satisfies the following two conditions:

- there exists $\Lambda' \in \mathcal{S}$ such that Θ' occurs in $f_{\mathbf{i}}\Lambda'$;
- $\Theta' < \Theta$ (by unitriangularity of the decomposition matrix).

Now, by (ii), such a symbol satisfies $e_{\mathbf{i}^*}\Theta' = \Lambda'$. The result follows from the fact that $P_{[\Lambda]}$ is a direct summand of $e_{\mathbf{i}^*}P_{[\Theta]}$ by (i). □

Remark 2.5. — If Λ has a unique addable i -box (i.e. $f_i\Lambda$ is a symbol) and no removable i -box (i.e. $e_i\Lambda = 0$) then $\widetilde{f}_i\Lambda = f_i\Lambda$ by Section 1.3.4. This is often helpful for checking condition (i) of Lemma 2.4.

2.3.1. The column corresponding to $[1^{2n}]$

The symbol corresponding to this unipotent character is

$$\begin{pmatrix} 1 & 0 & -1 & \dots & \widehat{-2n+1} & \dots \\ & & -1 & \dots & \dots & \dots \end{pmatrix}$$

so that the composition of the associated family is $(1, 0, -1, -1, -2, -2, \dots, \widehat{-2n+1}, \dots)$. It dominates $(0, 0, -1, -1, -2, -2, \dots, \widehat{-2n}, \dots)$, which is the family of the Steinberg character $[.1^{2n}]$, and does not dominate any other family of unipotent characters of G . Therefore $\Psi_{[1^{2n}]} = [1^{2n}] + \alpha[.1^{2n}]$ for some $\alpha \geq 0$, by unitriangularity. We shall prove that $\alpha = 0$.

PROPOSITION 2.6. — We have

$$\Psi_{[1^{2n}]} = [1^{2n}].$$

Proof. — We apply Lemma 2.4 to the symbol Λ associated to $[1^{2n}]$ and $\mathbf{i} = (0)$. We have

$$\begin{aligned} \Theta := f_0\Lambda &= f_0 \begin{pmatrix} 1 & 0 & -1 & \dots & \widehat{-2n+1} & \dots \\ & & -1 & \dots & \dots & \dots \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & -1 & \dots & \widehat{-2n} & \dots \\ & & -1 & \dots & \dots & \dots \end{pmatrix}. \end{aligned}$$

By Remark 2.5 it is also equal to $\widetilde{f}_0\Lambda$, therefore condition (i) of Lemma 2.4 is satisfied. On the other hand, the symbol Λ' associated to $[.1^{2n}]$ is the unique symbol smaller than Λ and it satisfies

$$f_0 \begin{pmatrix} 0 & -1 & \dots & \dots & \dots \\ 0 & -1 & \dots & \widehat{-2n} & \dots \end{pmatrix} = \begin{pmatrix} 1 & -1 & \dots & \dots & \dots \\ & 0 & -1 & \dots & \widehat{-2n} & \dots \end{pmatrix}.$$

Since $f_0\Lambda$ and $f_0\Lambda'$ lie in the same family, condition (ii) of Lemma 2.4 is empty hence automatically satisfied. As a consequence $[.1^{2n}]$ is not a constituent of $\Psi_{[1^{2n}]}$ and the proposition is proved. \square

2.3.2. The column corresponding to $[2^{n-1}.]_{B_2}$

The unipotent character $[2^{n-1}.]_{B_2}$ of the B_2 -series corresponds to the following symbol of defect -3

$$\Lambda = \begin{pmatrix} & 0 & -1 & \dots & \widehat{-n+1} & \widehat{-n} & \dots \\ 1 & 0 & -1 & \dots & \dots & \dots & \dots \end{pmatrix}.$$

The associated composition is $(1, 0, 0, -1, -1, \dots, -n+1, \widehat{-n+1}, \widehat{-n}, -n, \dots)$. From the description in Section 2.1 one checks that the unipotent characters in the principal Φ_{2n} -block lying in smaller families are $[1^{2n}]$, $[.21^{2n-2}]$ and $[.1^{2n}]$. Consequently there exist non-negative integers α, β , and γ such that

$$\Psi_{[2^{n-1}.]_{B_2}} = [2^{n-1}.]_{B_2} + \alpha[1^{2n}] + \beta[.21^{2n-2}] + \gamma[.1^{2n}].$$

The following theorem explicitly determines these integers whenever the ℓ -part of $\Phi_{2n}(q)$ is not too small.

THEOREM 2.7. — *There exists $\gamma \leq 2$ such that*

$$\Psi_{[2^{n-1}.]_{B_2}} = [2^{n-1}.]_{B_2} + \gamma[.1^{2n}].$$

Furthermore, if $\Phi_{2n}(q)_\ell > 4n$, then $\gamma = 2$.

Note that by a result of Feit, the condition $\Phi_{2n}(q)_\ell > 4n$ is satisfied for at least one prime number ℓ except for finitely many pairs (q, n) [15].

Proof.

Step 1. — The first step of the proof establishes that $[.1^{2n}]$ is the only unipotent character different from $[2^{n-1}.]_{B_2}$ that can occur in $\Psi_{[2^{n-1}.]_{B_2}}$ with a nonzero coefficient. For that purpose we use Lemma 2.4 with Λ being the symbol attached to $[2^{n-1}.]_{B_2}$ and $\mathbf{i} = (1, 2, \dots, n - 1)$. We have

$$f_{\mathbf{i}}\Lambda = f_1 f_2 \cdots f_{n-1} \Lambda = \begin{pmatrix} 0 & \cdots & \widehat{-n+1} & \cdots & \widehat{-2n+1} & \cdots \\ 1 & 0 & \cdots & \cdots & \cdots & \cdots \end{pmatrix}$$

which also equals $\tilde{f}_1 \tilde{f}_2 \cdots \tilde{f}_{n-1} \Lambda$ by Remark 2.5 so that condition (i) of Lemma 2.4 holds. We list below the symbols obtained by inducing the ones associated to the unipotent characters $[1^{2n}.]$, $[.21^{2n-2}]$, and $[.1^{2n}]$.

$[\mu]$	Λ'	$f_1 f_2 \cdots f_{n-1} \Lambda'$
$[1^{2n}.]$	$\begin{pmatrix} 1 & 0 & \cdots & \widehat{-2n+1} & \cdots \\ & & -1 & \cdots & \cdots \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & \cdots & \cdots & \widehat{-2n+1} & \cdots \\ 0 & \cdots & \widehat{-n+1} & \cdots & \cdots & \cdots \end{pmatrix} + \begin{pmatrix} 2 & 0 & \cdots & \cdots & \widehat{-2n+1} & \cdots \\ 0 & \cdots & \widehat{-n+2} & \cdots & \cdots & \cdots \end{pmatrix}$
$[.21^{2n-2}]$	$\begin{pmatrix} 0 & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & 0 & \cdots & \widehat{-2n+1} & \cdots & \cdots \end{pmatrix}$	$\begin{pmatrix} 0 & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & 0 & \cdots & \widehat{-n+1} & \cdots & \widehat{-2n+1} & \cdots \end{pmatrix}$
$[.1^{2n}]$	$\begin{pmatrix} 0 & \cdots & \cdots & \cdots \\ 0 & \cdots & \widehat{-2n} & \cdots \end{pmatrix}$	$\begin{pmatrix} 0 & \cdots & \cdots & \cdots \\ 0 & \cdots & \widehat{-3n+1} & \cdots \end{pmatrix}$

The only symbol in the last column which is smaller than $f_{\mathbf{i}}\Lambda$ corresponds to the induction of the symbol Λ' of $[.1^{2n}]$ (the last row in the table). One checks that $e_{\mathbf{i}^*} f_{\mathbf{i}} \Lambda' = \Lambda'$ for that symbol, so that assumption (ii) of Lemma 2.4 is satisfied. We deduce that $[.1^{2n}]$ is indeed the only unipotent constituent of $\Psi_{[2^{n-1}.]_{B_2}}$ apart from $[2^{n-1}.]_{B_2}$.

Step 2. — We now use the method in [7] to show that $\gamma \leq 2$. This requires to know how to decompose certain Deligne–Lusztig characters on the basis of PIMs, or at least to know the coefficient of the PIMs corresponding to cuspidal modules. The following lemma will be useful to deal with PIMs that can be obtained by induction from proper Levi subgroups of G .

LEMMA 2.8. — *Let L (resp. M) be a 1-split Levi of G of type BC_{2n-1} (resp. A_{2n-1}). We have*

$$\begin{aligned}
 U_1 &:= bR_L^G \left(\sum_{i=1}^{2n-1} (-1)^{i-1} [i1^{2n-i-1}.] \right) \\
 &= [1^{2n}.] + [2n.] + (-1)^n [(n-1)1^n.1], \\
 U_2 &:= bR_L^G \left(\sum_{j=1}^{n-1} (-1)^{j+n} [(n-j)1^n.1^{j-1}] \right) \\
 &= (-1)^n [(n-1)1^n.1] - [1^{n+1}.1^{n-1}], \\
 U_3 &:= bR_L^G \left(\sum_{k=0}^{n-2} (-1)^{k+n} [2^k 1^{n-k-1}.n-k-2]_{B_2} \right) \\
 &= (-1)^n [1^{n-1}.n-1]_{B_2} + [2^{n-1}.]_{B_2}, \\
 U_4 &:= bR_M^G ([1^{2n}]) = [1^{2n}.] + [.1^{2n}] + [1^{n+1}.1^{n-1}].
 \end{aligned}$$

In particular the character $[2n.] - [.1^{2n}] = U_1 - U_2 - U_4$ is a combination of characters induced from proper Levi subgroups of G .

Proof. — The Harish-Chandra induction from L to G of a unipotent character associated to a bipartition $\lambda = \lambda^1.\lambda^2$ is described by adding one box to λ^1 or λ^2 in all possible ways so that the result is a bipartition. The formulas for U_1 , U_2 and U_3 are easily deduced from that rule and the description of the principal Φ_{2n} -block given in Section 2.1. By [22, Proposition 6.1.4] we have $R_M^G ([1^{2n}]) = \sum_i [1^i.1^{2n-i}]$ from which we deduce the value of U_4 from Section 2.1. □

With the notation in Section 1.2.2 we consider the Coxeter element $c = s_1 s_2 \cdots s_{2n}$ of W_{2n} . The decomposition of the Deligne–Lusztig character of G attached to c can be deduced from [17, (3.2)]. Most of its constituents do not belong to the principal Φ_{2n} -block and we have

$$bR_c = [2n.] + [.1^{2n}] + (-1)^{n-1} [1^{n-1}.n-1]_{B_2}.$$

Using the combination of unipotent characters defined in Lemma 2.8, it can be rewritten as

$$\begin{aligned}
 bR_c &= U_1 - U_2 - U_3 - U_4 + [2^{n-1}.]_{B_2} + 2[.1^{2n}] \\
 &= U_1 - U_2 - U_3 - U_4 + \Psi_{[2^{n-1}.]_{B_2}} + (2 - \gamma)\Psi_{[.1^{2n}]}.
 \end{aligned}$$

Since the U_i 's are obtained by induction from proper Levi subgroups of G , they cannot involve $\Psi_{[1^{2n}]}$ since it corresponds to a cuspidal module. Therefore the coefficient of $\Psi_{[1^{2n}]}$ in R_c is exactly $2 - \gamma$. By [7, Proposition 1.5], we must have $2 - \gamma \geq 0$ therefore $\gamma \leq 2$.

Step 3. — In this final step of the proof, we show that $\gamma \geq 2$ under the assumption that $\Phi_{2n}(q)_\ell > 4n$. We consider the element

$$w := (s_1 s_2 \cdots s_{2n})^2 = c^2.$$

By [33, Section 5.2], it is a $2n$ -regular element. Therefore any torus \mathbf{T}_w of type w is the centraliser of a Φ_{2n} -Sylow subgroup of \mathbf{G} and the constituents of the Deligne–Lusztig character R_w are exactly the unipotent characters in the principal Φ_{2n} -block, see [3, Theorem 5.24]. The exact decomposition of R_w can be deduced for example from [17, (3.2)]. We find in particular

$$\langle R_w; [2^{n-1} \cdot]_{B_2} \rangle_G = -2 \quad \text{and} \quad \langle R_w; [1^{2n}] \rangle_G = 1.$$

On the other hand, if $\Phi_{2n}(q)_\ell > 4n$ then there exists by [10, Example 1.17] an ℓ -character of T_w in general position. By [10, Lemma 1.13] this forces $\langle R_w; \Psi_{[2^{n-1} \cdot]_{B_2}} \rangle_G \geq 0$, which gives

$$0 \leq \langle R_w; \Psi_{[2^{n-1} \cdot]_{B_2}} \rangle_G = \langle R_w; [2^{n-1} \cdot]_{B_2} \rangle_G + \gamma \langle R_w; [1^{2n}] \rangle_G = -2 + \gamma$$

and therefore proves that $\gamma \geq 2$. Note that it also shows that $\Psi_{[1^{2n}]}$ does not occur in R_c . □

2.3.3. The column corresponding to $[2^{n-3}]_{B_6}$

We now focus on the last column, corresponding to the unipotent character $[2^{n-3}]_{B_6}$ of the B_6 -series. It corresponds to the following symbol of defect 5

$$\Lambda = \left(\begin{array}{cccccccc} 2 & 1 & 0 & -1 & \cdots & \cdots & \cdots & \cdots \\ & & & -1 & \cdots & \widehat{-n+2} & \widehat{-n+1} & \cdots \end{array} \right)$$

whose associated composition is $(2, 1, 0, -1, -1, \dots, \widehat{-n+2}, \widehat{-n+1}, \dots)$. From the description in Section 2.1 one checks that it is minimal among the unipotent characters in the B_6 -series lying in the principal Φ_{2n} -block. There are 3 unipotent characters in the principal Φ_{2n} -block belonging to the B_2 -series which are smaller for the order on families. We list these characters with their associated symbol:

$$\begin{aligned} [2^{n-1}.]_{B_2} &\longleftrightarrow \begin{pmatrix} 0 & \dots & \widehat{-n+1} & \widehat{-n} & \dots \\ 1 & 0 & \dots & \dots & \dots \end{pmatrix}, \\ [2^{n-2}.1.1]_{B_2} &\longleftrightarrow \begin{pmatrix} 0 & \dots & \widehat{-n+2} & -n+1 & \widehat{-n} & \dots \\ 2 & 0 & \dots & \dots & \dots & \dots \end{pmatrix}, \\ [2^{n-2}.1^2]_{B_2} &\longleftrightarrow \begin{pmatrix} 0 & \dots & \widehat{-n+2} & \widehat{-n+1} & \dots \\ 2 & 1 & -1 & \dots & \dots \end{pmatrix}. \end{aligned}$$

Finally, there are 8 unipotent characters in the principal series belonging to the principal Φ_{2n} -block which are smaller than $[.2^{n-3}]_{B_6}$:

$$\begin{aligned} [21^{2n-2}.] &\longleftrightarrow \begin{pmatrix} 2 & 0 & \dots & \widehat{-2n+2} \\ & -1 & \dots & \dots \end{pmatrix}, \\ [.31^{2n-3}] &\longleftrightarrow \begin{pmatrix} 0 & \dots & \dots & \dots \\ 2 & -1 & \dots & \widehat{-2n+2} \end{pmatrix}, \\ [.21^{2n-2}] &\longleftrightarrow \begin{pmatrix} 0 & \dots & \dots & \dots \\ 1 & -1 & \dots & \widehat{-2n+1} \end{pmatrix}, \\ [1^{2n}] &\longleftrightarrow \begin{pmatrix} 0 & \dots & \dots \\ 0 & \dots & \widehat{-2n} \end{pmatrix}, \\ [1^{2n}.] &\longleftrightarrow \begin{pmatrix} 1 & \dots & \widehat{-2n+1} \\ & -1 & \dots \end{pmatrix}, \\ [21^n.1^{n-2}] &\longleftrightarrow \begin{pmatrix} 2 & 0 & \dots & \dots & \widehat{-n} \\ & 0 & \dots & \widehat{-n+2} & \dots \end{pmatrix}, \\ [1^n.21^{n-2}] &\longleftrightarrow \begin{pmatrix} 1 & \dots & \dots & \widehat{-n+1} \\ 1 & -1 & \dots & \widehat{-n+1} \end{pmatrix}, \\ [1^{n+1}.1^{n-1}] &\longleftrightarrow \begin{pmatrix} 1 & \dots & \dots & \dots & \widehat{-n} \\ & 0 & \dots & \widehat{-n+1} & \dots \end{pmatrix}. \end{aligned}$$

We will show that none of these characters contribute to $\Psi_{[.2^{n-3}]_{B_6}}$ with the exception of $[1^{2n}.]$

THEOREM 2.9. — *There exists $\beta \geq 0$ such that*

$$\Psi_{[.2^{n-3}]_{B_6}} = [.2^{n-3}]_{B_6} + \beta[1^{2n}.].$$

Furthermore, if $\Phi_{2n}(q)_\ell > 4n$ then $\beta = 2$.

Proof. — The proof follows that of Theorem 2.7, but more computations are needed since more characters are involved.

Step 1. — We start by computing the image of the various symbols involved under the operator $f_{\mathbf{i}}$ for $\mathbf{i} = (1, 0)$. For the symbol Λ corresponding to the unipotent character in the B_6 -series we have

$$(2.2) \quad f_{\mathbf{i}}\Lambda = \begin{pmatrix} 2 & 1 & 0 & -1 & \dots & \dots & \dots & \dots \\ & & & -1 & \dots & \widehat{-n+1} & \widehat{-n} & \dots \end{pmatrix}$$

which also equals $\widetilde{f}_{\mathbf{i}}\Lambda$ by Remark 2.5. For the symbols corresponding to characters in the B_2 -series we obtain

$[\Lambda']$	$f_{\mathbf{i}}\Lambda'$
$[2^{n-1}.]_{B_2}$	$\begin{pmatrix} 2 & & -1 & \dots & \widehat{-n+1} & \widehat{-n} & \dots \\ & 1 & 0 & -1 & \dots & \dots & \dots \end{pmatrix}$
$[2^{n-2}1.1]_{B_2}$	$\begin{pmatrix} 2 & & -1 & \dots & \widehat{-n+2} & -n+1 & \widehat{-n} & \dots \\ 2 & 0 & -1 & \dots & \dots & \dots & \dots & \dots \end{pmatrix}$
$[2^{n-2}.1^2]_{B_2}$	$\begin{pmatrix} 2 & & -1 & \dots & \widehat{-n+2} & \widehat{-n+1} & \dots \\ 2 & 1 & -1 & \dots & \dots & \dots & \dots \end{pmatrix}$

None of these symbol is smaller than $f_{\mathbf{i}}\Lambda$. The case of the principal series characters is given in the following table:

$[\Lambda']$	$f_{\mathbf{i}}\Lambda'$
$[21^{2n-2}.]$	$\begin{pmatrix} 2 & 1 & -1 & \dots & \widehat{-2n+1} \\ & & -1 & \dots & \dots \end{pmatrix}$
$[\cdot 31^{2n-3}]$	$\begin{pmatrix} 2 & -1 & \dots & \dots \\ 2 & -1 & \dots & \widehat{-2n+2} \end{pmatrix}$
$[\cdot 21^{2n-2}]$	$\begin{pmatrix} 2 & -1 & \dots & \dots \\ & 1 & -1 & \dots & \widehat{-2n+1} \end{pmatrix}$
$[\cdot 1^{2n}]$	$\begin{pmatrix} 2 & -1 & \dots & \dots \\ & 0 & \dots & \dots & \widehat{-2n} \end{pmatrix}$
$[1^{2n}.]$	$\begin{pmatrix} 2 & 0 & \dots & \widehat{-2n} & \dots \\ & -1 & \dots & \dots & \dots \end{pmatrix}$
$[21^n \cdot 1^{n-2}]$	$\begin{pmatrix} 2 & 1 & -1 & \dots & \dots & \widehat{-n} \\ & & 0 & \dots & \dots & \widehat{-n+1} & \dots \end{pmatrix}$
$[1^n \cdot 21^{n-2}]$	$\begin{pmatrix} 2 & 0 & \dots & \dots & \widehat{-n+1} \\ & 1 & -1 & \dots & \widehat{-n} \end{pmatrix}$
$[1^{n+1} \cdot 1^{n-1}]$	$\begin{pmatrix} 2 & \dots & \dots & \dots & \widehat{-n} \\ & 0 & \dots & \widehat{-n} & \dots \end{pmatrix}$

The induction of the symbols corresponding to the characters $[1^n \cdot 21^{n-2}]$, $[\cdot 31^{2n-3}]$, $[21^n \cdot 1^{n-2}]$ and $[1^{n+1} \cdot 1^{n-1}]$ are not strictly dominated by the symbol $f_{\mathbf{i}}\Lambda$ computed in (2.2). The other symbols Λ' satisfy $e_{\mathbf{i}^*} f_{\mathbf{i}}\Lambda' = \Lambda'$, therefore Lemma 2.4 shows that there exist non-negative integers $\beta_1, \beta_2, \beta_3$ and β_4 such that

$$\Psi_{[\cdot 2^{n-3}]_{B_6}} = [\cdot 2^{n-3}]_{B_6} + \beta_1 [21^{2n-2}.] + \beta_2 [\cdot 21^{2n-2}] + \beta_3 [1^{2n}.] + \beta_4 [\cdot 1^{2n}].$$

We now consider the induction with respect to the sequence $\mathbf{i} = (-1, -2, 0, -1, 1, 0)$. We find

$$(2.3) \quad \tilde{f}_{\mathbf{i}}\Lambda = f_{\mathbf{i}}\Lambda = \begin{pmatrix} 2 & 1 & 0 & -1 & \dots & \dots & \dots \\ & & & -1 & \dots & \widehat{-n-1} & \widehat{-n-2} & \dots \end{pmatrix}$$

using Remark 2.5. For the four remaining principal series characters we obtain the following:

$[\Lambda']$	Λ'	$f_{\mathbf{i}}\Lambda'$
$[21^{2n-2}.]$	$\begin{pmatrix} 2 & 0 & \dots & \widehat{-2n+2} \\ & -1 & \dots & \dots \end{pmatrix}$	$\begin{pmatrix} 2 & 1 & 0 & -1 & -3 & \dots & \widehat{-2n-1} \\ & -1 & \dots & \dots & \dots & \dots & \dots \end{pmatrix}$
$[.21^{2n-2}]$	$\begin{pmatrix} 0 & \dots & \dots & \dots \\ 1 & -1 & \dots & \widehat{-2n+1} \end{pmatrix}$	$\begin{pmatrix} 2 & 1 & 0 & -3 & \dots \\ & 1 & -1 & \dots & \widehat{-2n+1} \end{pmatrix}$
$[1^{2n}.]$	$\begin{pmatrix} 1 & \dots & \widehat{-2n+1} \\ & -1 & \dots \end{pmatrix}$	$\begin{pmatrix} 2 & 1 & 0 & -2 & \dots & \widehat{-2n-2} \\ & -1 & \dots & \dots & \dots & \dots \end{pmatrix}$
$[.1^{2n}]$	$\begin{pmatrix} 0 & \dots & \dots \\ 0 & \dots & \widehat{-2n} \end{pmatrix}$	$\begin{pmatrix} 2 & 1 & 0 & -3 & \dots & \dots \\ & 0 & \dots & \dots & \dots & \widehat{-2n} \end{pmatrix}$

Note that for the computations we can use that f_i and f_j commute whenever $i \neq j \pm 1$ modulo $2n$ so that $f_{\mathbf{i}} = (f_{-1}f_0f_1)(f_{-2}f_{-1}f_0)$. The induction of the symbols corresponding to the unipotent characters $[.21^{2n-2}]$ and $[.1^{2n}]$ are not strictly dominated by the symbol $f_{\mathbf{i}}\Lambda$ computed in (2.3). The other symbols satisfy the conditions of Lemma 2.4 which shows that $\beta_2 = \beta_4 = 0$. It remains to show that $\beta_1 = 0$. For that purpose we use again Lemma 2.4 for Λ and $\mathbf{i} = (-n + 2, \dots, -2, -1, 0)$. The induced symbols are given in the following table.

$[\Lambda']$	Λ'	$f_{\mathbf{i}}\Lambda'$
$[.2^{n-3}]_{B_6}$	$\begin{pmatrix} 2 & \dots & \dots & \dots & \dots \\ & -1 & \dots & \widehat{-n+2} & \widehat{-n+1} \end{pmatrix}$	$\begin{pmatrix} 2 & \dots & \dots & \dots & \dots & \dots \\ & -1 & \dots & \widehat{-n+2} & \dots & \widehat{-2n+2} \end{pmatrix}$
$[21^{2n-2}.]$	$\begin{pmatrix} 2 & 0 & \dots & \widehat{-2n+2} \\ & -1 & \dots & \dots \end{pmatrix}$	$\begin{pmatrix} 2 & \dots & \widehat{-n+2} & \dots & \widehat{-2n+2} \\ & -1 & \dots & \dots & \dots \end{pmatrix}$
$[1^{2n}.]$	$\begin{pmatrix} 1 & \dots & \widehat{-2n+1} \\ & -1 & \dots \end{pmatrix}$	$\begin{pmatrix} 1 & \dots & \widehat{-3n+2} \\ & -1 & \dots \end{pmatrix}$

The induced symbols corresponding to the characters $[.2^{n-3}]_{B_6}$ and $[21^{2n-2}.]$ lie in the same family. In addition, if Λ' is the symbol attached to $[1^{2n}.]$ then $e_{\mathbf{i}^*}f_{\mathbf{i}}\Lambda' = \Lambda'$. Finally, one can check that $\tilde{f}_{\mathbf{i}}\Lambda = f_{\mathbf{i}}\Lambda$ using Section 1.3.4 and invoke Lemma 2.4 to conclude that $\beta_1 = 0$ (note that Remark 2.5 does not apply to the last step of the computation of $\tilde{f}_{\mathbf{i}}\Lambda$). For the remainder of the proof we will write $\beta := \beta_3$.

Step 2. — Assume now that $\Phi_{2n}(q)_\ell > 4n$. Then $\Psi_{[2^{n-1}.]_{B_2}} = [2^{n-1}.]_{B_2} + 2[.1^{2n}]$ by Theorem 2.7. In addition, it was observed in the proof of that theorem that the Deligne–Lusztig character associated to a Coxeter element c decomposes as

$$R_c = U_1 - U_2 - U_3 - U_4 + \Psi_{[2^{n-1}.]_{B_2}}.$$

This shows that apart from $\Psi_{[2^{n-1}.]_{B_2}}$, none of the PIMs corresponding to cuspidal modules appear in R_c . In order to decompose the next Deligne–Lusztig characters we shall use the following identities, whose proofs are identical to those for Lemma 2.8.

LEMMA 2.10. — *Let L be a 1-split Levi of G of type B_{2n-1} . We have*

$$\begin{aligned}
 U_5 &:= bR_L^G\left([2n-1.]\right) = [2n.] + [(2n-1)1.], \\
 U_6 &:= bR_L^G\left([1^{n-1}.n-2]_{B_2} + [1^{n-2}.n-1]_{B_2}\right) \\
 &= 2[1^{n-1}.n-1]_{B_2} + [21^{n-2}.n-2]_{B_2} + [1^{n-2}.(n-1)1]_{B_2}, \\
 U_7 &:= bR_L^G\left([(2n-2)1.]\right) = [(2n-1)1.] + [(2n-2)1^2.], \\
 U_8 &:= bR_L^G\left(\sum_{j=1}^n (-1)^{j+n} [1^j.(n+1-j)1^{n-2}]\right) \\
 &= (-1)^{n+1} [1.(n+1)1^{n-2}.] + [1^{n+1}.1^{n-1}], \\
 U_9 &:= bR_L^G\left([1^{n-3}.(n-1)1]_{B_2} + [1^{n-2}.(n-2)1]_{B_2}\right. \\
 &\quad \left.+ [21^{n-2}.n-3]_{B_2} - 2[1^{n-2}.n-1]_{B_2} - [1^{n-1}.n-2]_{B_2}\right), \\
 &= [1^{n-3}.(n-1)2]_{B_2} + [21^{n-3}.(n-2)1]_{B_2} \\
 &\quad + [2^2 1^{n-3}.n-3]_{B_2} - 3[1^{n-1}.n-1]_{B_2}, \\
 U_{10} &:= bR_L^G\left(\sum_{r=0}^{n-4} (-1)^{r+n} [n-r-4.2^r 1^{n-r-3}]_{B_6}\right) \\
 &= (-1)^n [n-3.1^{n-3}]_{B_6} + [.2^{n-3}]_{B_6}.
 \end{aligned}$$

Since we want to use the result in [7] to get information on the PIMs with cuspidal head, it is enough to consider the Deligne–Lusztig characters R_w for elements $w \in W_{2n}$ whose conjugacy class does not meet any proper parabolic subgroup. Such classes are called cuspidal (or elliptic) and are described in [22, Proposition 3.4.6]. With the notation in [22, Section 3.4.2], minimal length elements of the first 3 cuspidal classes, ordered by length, are given by $c = w_{(2n)}^-$, $v_n = s_1 s_2 c = w_{(1,2n-1)}^-$, and $w_n := s_1 s_2 s_3 s_2 c = w_{(2,2n-2)}$. Let us first consider the Deligne–Lusztig character associated to v_n . From the decomposition of R_{v_n} given in Lemma A.11 and the characters

defined in Lemma 2.8 and 2.10 we have

$$\begin{aligned} bR_{w_n} &= 2(U_1 - U_2 - U_3 - U_4) - U_5 - D_G(U_5) + (-1)^n U_6 \\ &\quad + 2[2^{n-1} \cdot]_{B_2} + 4[1^{2n}] \\ &= 2(U_1 - U_2 - U_3 - U_4) - U_5 - D_G(U_5) + (-1)^n U_6 + 2\Psi_{[2^{n-1} \cdot]_{B_2}}. \end{aligned}$$

Again, apart from $\Psi_{[2^{n-1} \cdot]_{B_2}}$, none of the PIMs corresponding to cuspidal modules appear in R_{w_n} . We move on to the element $w_n := s_1 s_2 s_3 s_2 c$. Using Lemma A.12 and Lemma 2.8 we can decompose R_{w_n} as

$$\begin{aligned} bR_{w_n} &= 3U_1 - 3U_2 - U_3 - 4U_4 - 2U_5 + U_7 + U_8 + D_G(U_1) \\ &\quad - 2D_G(U_5) + D_G(U_7) + (-1)^{n-1} U_9 - U_{10} \\ &\quad + [2^{n-3}]_{B_6} + 3[2^{n-1} \cdot]_{B_2} + 2[1^{2n} \cdot] + 3[1^{2n}] \\ &= 3U_1 - 3U_2 - U_3 - 4U_4 - 2U_5 + U_7 + U_8 + D_G(U_1) \\ &\quad - 2D_G(U_5) + D_G(U_7) + (-1)^{n-1} U_9 - U_{10} \\ &\quad + \Psi_{[2^{n-3}]_{B_6}} + 3\Psi_{[2^{n-1} \cdot]_{B_2}} + (2 - \beta)\Psi_{[1^{2n} \cdot]}. \end{aligned}$$

Here we have also used Proposition 2.6 and Theorem 2.7 which give the decomposition of $\Psi_{[2^{n-1} \cdot]_{B_2}}$ and $\Psi_{[1^{2n} \cdot]}$. It follows from [7, Proposition 1.5] that $2 - \beta \geq 0$ hence $\beta \leq 2$.

Step 3. — The argument is entirely similar to that given in the proof of Theorem 2.7 with the exception that one uses the multiplicities

$$\langle R_w; [2^{n-3}]_{B_6} \rangle_G = -2 \quad \text{and} \quad \langle R_w; [1^{2n} \cdot] \rangle_G = 1.$$

This shows that $\beta \geq 2$, hence $\beta = 2$. □

Appendix: Computation of Deligne–Lusztig characters

We fix an integer n . As in Section 1.2.2 we denote by W_{2n} the Weyl group of type B_{2n} with generators s_1, \dots, s_{2n} . We explain here how to compute the Deligne–Lusztig characters associated with the elements

$$\begin{aligned} v_n &:= (s_1 s_2)(s_1 s_2 s_3 \cdots s_{2n-1} s_{2n}) \\ \text{and} \quad w_n &:= (s_1 s_2 s_3 s_2)(s_1 s_2 s_3 \cdots s_{2n-1} s_{2n}) \end{aligned}$$

of respective lengths $2n + 2$ and $2n + 4$.

A.1. The Deligne–Lusztig character associated to v_n

Using the notation in [22, Section 3.4.2], the element decomposes as $v_n = w_{(1,2n-1)}^- = s_1 b_{1,2n-1}^- = b_{0,1}^- b_{1,2n-1}^-$ and one can compute the values of irreducible characters of W_{2n} at the element v_n using the Murnaghan–Nakayama rule. More precisely, if λ is a bipartition of $2n$ and χ_λ is the corresponding irreducible character of W_{2n} then by [22, Theorem 10.3.1] we have

$$(A.1) \quad \chi_\lambda(v_n) = \sum_{\gamma} (-1)^{h(\gamma)} \chi_{\lambda \setminus \gamma}(s_1)$$

where γ runs over all the $(2n-1)$ -hooks of λ , and $h(\gamma)$ equals the leg length of γ plus 1 if γ is in the second component. From the description of the principal Φ_{2n} -block B given in Section 2.1 one can determine the bipartitions λ which have a $(2n-1)$ -hook and whose family has a non-trivial intersection with $\text{Irr } B$. They are listed in Table A.1, together with the value of $h(\gamma)$, the $(2n-1)$ -core $\lambda \setminus \gamma$, the value of $\chi_{\lambda \setminus \gamma}(s_1 s_2)$ and that of $\chi_\lambda(v_n)$ using (A.1). Note that the second half of the table can be obtained from the first using the relation $\chi_{\lambda^*}(v_n) = (-1)^{l(v_n)} \chi_\lambda(v_n) = \chi_\lambda(v_n)$, where $l(w)$ is the Coxeter length of $w \in W_{2n}$.

Table A.1. Bipartitions λ with a $(2n-1)$ -hook whose family intersects the principal Φ_{2n} -block

λ	$h(\gamma)$	$\lambda \setminus \gamma$	$\chi_{\lambda \setminus \gamma}(s_1)$	$\chi_\lambda(v_n)$
$2n.$	0			1
$n21^{n-2}.$	$n - 1$			$(-1)^{n-1}$
$1^{2n}.$	$2n - 2$	1.	1	1
$1.(n+1)1^{n-2}$	$n - 2 + 1$			$(-1)^{n-1}$
1.1^{2n-1}	$2n - 2 + 1$			-1
$.1^{2n}$	$2n - 2 + 1$			1
$.n21^{n-2}$	$n - 1 + 1$			$(-1)^{n-1}$
$.2n$	$0 + 1$.1	-1	1
$(n-1)1^n.1$	n			$(-1)^{n-1}$
$2n-1.1$	0			-1

The families corresponding to the bipartitions $2n.$ and $.1^{2n}$ contain only one unipotent character each, the trivial and the Steinberg character respectively. The other families have 4 elements and are listed in Table A.2.

By convention the special character is the first one in each list. The corresponding almost characters can be computed from [28, Section 4]. They are given in Table A.3.

Table A.2. Families \mathfrak{F} with 4 elements occurring in bR_{v_n}

λ	\mathfrak{F}
$n21^{n-2}$	$[n.21^{n-2}], [n21^{n-2}.], [1.(n+1)1^{n-2}], [1^{n-2}.(n-1)1]_{B_2}$
1^{2n}	$[1.1^{2n-1}], [1^{2n}.], [.21^{2n-2}], [1^{2n-2}]_{B_2}$
$1.(n+1)1^{n-2}$	$[n.21^{n-2}], [n21^{n-2}.], [1.(n+1)1^{n-2}], [1^{n-2}.(n-1)1]_{B_2}$
1.1^{2n-1}	$[1.1^{2n-1}], [1^{2n}.], [.21^{2n-2}], [1^{2n-2}]_{B_2}$
$.n21^{n-2}$	$[(n-1)1.1^n], [.n21^{n-2}], [(n-1)1^n.1], [21^{n-2}.n-2]_{B_2}$
$.2n$	$[2n-1.1], [.2n], [(2n-1)1.], [2n-2.]_{B_2}$
$(n-1)1^n.1$	$[(n-1)1.1^n], [.n21^{n-2}], [(n-1)1^n.1], [21^{n-2}.n-2]_{B_2}$
$2n-1.1$	$[2n-1.1], [.2n], [(2n-1)1.], [2n-2.]_{B_2}$

Table A.3. Some almost characters occurring in bR_{v_n}

λ	bR_{χ_λ}
$n21^{n-2}$	$-\frac{1}{2} \left([1.(n+1)1^{n-2}] + [1^{n-2}.(n-1)1]_{B_2} \right)$
1^{2n}	$\frac{1}{2} \left([1^{2n}.] - [.21^{2n-2}] \right)$
$1.(n+1)1^{n-2}$	$\frac{1}{2} \left([1.(n+1)1^{n-2}] - [1^{n-2}.(n-1)1]_{B_2} \right)$
1.1^{2n-1}	$\frac{1}{2} \left([1^{2n}.] + [.21^{2n-2}] \right)$
$.n21^{n-2}$	$-\frac{1}{2} \left([(n-1)1^n.1] + [21^{n-2}.n-2]_{B_2} \right)$
$.2n$	$\frac{1}{2} \left([.2n] - [(2n-1)1.] \right)$
$(n-1)1^n.1$	$\frac{1}{2} \left([(n-1)1^n.1] - [21^{n-2}.n-2]_{B_2} \right)$
$2n-1.1$	$\frac{1}{2} \left([(2n-1)1.] - [.2n] \right)$

Using the data in these tables, together with the formula $R_w = \sum_{\lambda} \chi_{\lambda}(w)R_{\chi_{\lambda}}$, we obtain the explicit decomposition of R_{v_n} on the block.

LEMMA A.11. — *The decomposition of the Deligne–Lusztig character R_{v_n} of G_{2n} associated to $v_n = s_1 s_2 s_1 s_2 \cdots s_{2n}$ on the principal Φ_{2n} -block is given by*

$$bR_{v_n} = [2n.] + [.1^{2n}] - [(2n-1)1.] - [.21^{2n-2}] + (-1)^n \left([1^{n-2}.(n-1)1]_{B_2} + [21^{n-2}.n-2]_{B_2} \right).$$

A.2. The Deligne–Lusztig character associated to w_n

Using again the notation in [22, Section 3.4.2], we can write w_n as $w_n = w_{(2,2n-2)}^- = b_{0,2}^- b_{2,2n-2}^- = s_1 s_2 b_{2,2n-2}^-$. Let λ be a bipartition of $2n$. By [22, Theorem 10.3.1] we have

$$(A.2) \quad \chi_\lambda(w_n) = \sum_{\gamma} (-1)^{h(\gamma)} \chi_{\lambda \setminus \gamma}(s_1 s_2)$$

where γ runs over all the $(2n-2)$ -hooks of λ , and $h(\gamma)$ equals the leg length of γ plus 1 if γ is in the second component. We list in Table A.4 those bipartitions λ which have a $(2n-2)$ -hook γ , the value of $h(\gamma)$, the $(2n-2)$ -core $\lambda \setminus \gamma$, the value of $\chi_{\lambda \setminus \gamma}(s_1 s_2)$ and that of $\chi_\lambda(w_n)$ using (A.2). We do not list the ones with $(2n-2)$ -core 1.1 since the corresponding character of B_2 vanishes on the element $s_1 s_2$. In addition, since $\chi_{\lambda^*}(w_n) = (-1)^{\ell(w_n)} \chi_\lambda(w_n) = \chi_\lambda(w_n)$ it is enough to deal with the cores 2. and 1^2 .

Table A.4. Bipartitions λ with $(2n-2)$ -core equal to 2. or 1^2 .

λ	$h(\gamma)$	$\lambda \setminus \gamma$	$\chi_{\lambda \setminus \gamma}(s_1 s_2)$	$\chi_\lambda(w_n)$
$2n.$	0			1
$i31^{2n-i-3}.$	$2n - i - 2$	2.	1	$(-1)^i$
$21^{2n-2}.$	$2n - 3$			-1
$2.j1^{2n-2-j}$	$2n - 2 - j + 1$			$(-1)^{j-1}$
$1^{2n}.$	$2n - 3$	$1^2.$	-1	1
$k2^2 1^{2n-k-4}.$	$2n - i - 2$			$(-1)^{i-1}$
$(2n-1)1.$	0			-1
$1^2.l1^{2n-2-l}$	$2n - 2 - j + 1$			$(-1)^j$

For each bipartition λ in Table A.4 one can easily compute the family associated to each of these bipartitions and check using Section 2.1 that

only those with $i = l = n$, $j = 1$ or $n + 1$ and $k = n - 1$ correspond to families which have a non-trivial intersection with the principal Φ_{2n} -block B . The family associated to the bipartition $2n$. has size one and contains only the trivial character. We give in Table A.5 the families with 4 elements, starting with the special character in the family.

Table A.5. Families \mathfrak{F} with 4 elements occurring in bR_{w_n} (up to Alvis-Curtis duality)

λ	\mathfrak{F}
$n31^{n-3}$.	$[n.31^{n-3}], [n31^{n-3}.], [2.(n+1)1^{n-3}], [1^{n-3}.(n-1)2]_{B_2}$
21^{2n-2} .	$[2.1^{2n-2}], [.31^{2n-3}], [21^{2n-2}.], [1^{2n-3}.1]_{B_2}$
$2.(n+1)1^{n-3}$	$[n.31^{n-3}], [n31^{n-3}.], [2.(n+1)1^{n-3}], [1^{n-3}.(n-1)2]_{B_2}$
2.1^{2n-2}	$[2.1^{2n-2}], [.31^{2n-3}], [21^{2n-2}.], [1^{2n-3}.1]_{B_2}$
1^{2n} .	$[1.1^{2n-1}], [1^{2n}.], [.21^{2n-2}], [1^{2n-2}.]_{B_2}$
$(2n-1)1$.	$[2n-1.1], [.2n], [(2n-1)1.], [.2n-2]_{B_2}$

The two remaining bipartitions $(n-1)2^21^{n-3}$. and $1^2.n1^{n-2}$ correspond to unipotent characters lying in a family with 16 elements. Using [28, Section 4] one can deduce from Table A.5 the almost character corresponding to each bipartition. These are listed in Table A.6. For the family with 16 elements we get

$$\begin{aligned}
 bR_{\chi_{(n-1)2^21^{n-3}}} &= -\frac{1}{4} \left([1^2.n1^{n-2}] + [(n-1)1^{n-1}.2] \right. \\
 &\quad \left. + [21^{n-3}.(n-2)1]_{B_2} + [n-3.1^{n-3}]_{B_6} \right), \\
 bR_{\chi_{1^2.n1^{n-2}}} &= \frac{1}{4} \left([1^2.n1^{n-2}] + [(n-1)1^{n-1}.2] \right. \\
 &\quad \left. - [21^{n-3}.(n-2)1]_{B_2} - [n-3.1^{n-3}]_{B_6} \right).
 \end{aligned}$$

Putting this all together we obtain the decomposition of R_{w_n} on the block.

LEMMA A.12. — *The decomposition of the Deligne-Lusztig character R_{w_n} of G_{2n} associated to*

Table A.6. Some almost characters occurring in bR_{w_n}

λ	bR_{χ_λ}
$n31^{n-3}$.	$-\frac{1}{2} \left([2.(n+1)1^{n-3}] + [1^{n-3}.(n-1)2]_{B_2} \right)$
21^{2n-2} .	$\frac{1}{2} \left([21^{2n-2}.] - [.31^{2n-3}] \right)$
$2.(n+1)1^{n-3}$	$\frac{1}{2} \left([2.(n+1)1^{n-3}] - [1^{n-3}.(n-1)2]_{B_2} \right)$
2.1^{2n-2}	$\frac{1}{2} \left([21^{2n-2}.] + [.31^{2n-3}] \right)$
1^{2n} .	$\frac{1}{2} \left([1^{2n}.] - [.21^{2n-2}] \right)$
$(2n-1)1$.	$\frac{1}{2} \left([(2n-1)1.] - [.2n] \right)$

$w_n = s_1 s_2 s_3 s_2 s_1 s_2 \cdots s_{2n}$ on the principal Φ_{2n} -block is given by

$$\begin{aligned}
 bR_{w_n} = & [2n.] - [(2n-1)1.] + [(2n-2)1^2.] + [1^{2n}.] + [.1^{2n}] - [.21^{2n-2}] \\
 & + [.31^{2n-3}] + [.2n] + (-1)^{n-1} \left([1^{n-3}.(n-1)2]_{B_2} + [2^2 1^{n-3}.n-3]_{B_2} \right. \\
 & \left. + [21^{n-3}.(n-2)1]_{B_2} + [n-3.1^{n-3}]_{B_6} \right).
 \end{aligned}$$

BIBLIOGRAPHY

- [1] C. BONNAFÉ, J.-F. DAT & R. ROQUIER, “Derived categories and Deligne–Lusztig varieties II”, *Ann. Math.* **185** (2017), no. 2, p. 609-670.
- [2] M. BROUÉ, “Isométries de caractères et équivalences de Morita ou dérivées”, *Publ. Math., Inst. Hautes Étud. Sci.* (1990), no. 71, p. 45-63.
- [3] M. BROUÉ, G. MALLE & J. MICHEL, “Generic blocks of finite reductive groups”, in *Représentations unipotentes génériques et blocs des groupes réductifs finis*, Astérisque, no. 212, Société Mathématique de France, 1993, p. 7-92.
- [4] O. BRUNAT, O. DUDAS & J. TAYLOR, “Unitriangular shape of decomposition matrices of unipotent blocks”, *Ann. Math.* **192** (2020), no. 2, p. 583-663.
- [5] J. CHUANG & W. TURNER, “Cubist algebras”, *Adv. Math.* **217** (2008), no. 4, p. 1614-1670.
- [6] P. DELIGNE & G. LUSZTIG, “Representations of reductive groups over finite fields”, *Ann. Math.* **103** (1976), no. 1, p. 103-161.
- [7] O. DUDAS, “A note on decomposition numbers for groups of Lie type of small rank”, *J. Algebra* **388** (2013), p. 364-373.
- [8] O. DUDAS & G. MALLE, “Decomposition matrices for low-rank unitary groups”, *Proc. Lond. Math. Soc.* **110** (2015), no. 6, p. 1517-1557.
- [9] ———, “Decomposition matrices for exceptional groups at $d = 4$ ”, *J. Pure Appl. Algebra* **220** (2016), no. 3, p. 1096-1121.

- [10] ———, “Decomposition matrices for groups of Lie type in non-defining characteristic”, <https://arxiv.org/abs/2001.06395>, 2020.
- [11] O. DUDAS, M. VARAGNOLO & E. VASSEROT, “Categorical actions on unipotent representations of finite classical groups”, in *Categorification and higher representation theory*, Contemporary Mathematics, vol. 683, American Mathematical Society, 2017, p. 41-104.
- [12] ———, “Categorical actions on unipotent representations of finite unitary groups”, *Publ. Math., Inst. Hautes Étud. Sci.* **129** (2019), p. 129-197.
- [13] C. DUNKL & S. GRIFFETH, “Generalized Jack polynomials and the representation theory of rational Cherednik algebras”, *Sel. Math., New Ser.* **16** (2010), no. 4, p. 791-818.
- [14] M. FAYERS, “Weight two blocks of Iwahori–Hecke algebras of type B”, *J. Algebra* **303** (2006), no. 1, p. 154-201.
- [15] W. FEIT, “On large Zsigmondy primes”, *Proc. Am. Math. Soc.* **102** (1988), no. 1, p. 29-36.
- [16] O. FODA, B. LECLERC, M. OKADO, J.-Y. THIBON & T. A. WELSH, “Branching functions of $A_{n-1}^{(1)}$ and Jantzen–Seitz problem for Ariki–Koike algebras”, *Adv. Math.* **141** (1999), no. 2, p. 322-365.
- [17] P. FONG & B. SRINIVASAN, “Generalized Harish-Chandra theory for unipotent characters of finite classical groups”, *J. Algebra* **104** (1986), no. 2, p. 301-309.
- [18] ———, “The blocks of finite classical groups”, *J. Reine Angew. Math.* **396** (1989), p. 122-191.
- [19] ———, “Brauer trees in classical groups”, *J. Algebra* **131** (1990), no. 1, p. 179-225.
- [20] M. GECK, “Basic sets of Brauer characters of finite groups of Lie type. II”, *J. Lond. Math. Soc.* **47** (1993), no. 2, p. 255-268.
- [21] M. GECK & G. HISS, “Basic sets of Brauer characters of finite groups of Lie type”, *J. Reine Angew. Math.* **418** (1991), p. 173-188.
- [22] M. GECK & G. PFEIFFER, *Characters of finite Coxeter groups and Iwahori–Hecke algebras*, London Mathematical Society Monographs. New Series, vol. 21, Clarendon Press, 2000, xvi+446 pages.
- [23] T. GERBER, “Triple crystal action in Fock spaces”, *Adv. Math.* **329** (2018), p. 916-954.
- [24] S. GRIFFETH & E. NORTON, “Character formulas and Bernstein–Gelfand–Gelfand resolutions for Cherednik algebra modules”, *Proc. Lond. Math. Soc.* **113** (2016), no. 6, p. 868-906.
- [25] J. GRUBER & G. HISS, “Decomposition numbers of finite classical groups for linear primes”, *J. Reine Angew. Math.* **485** (1997), p. 55-91.
- [26] F. HIMSTEDT & F. NOESKE, “Decomposition numbers of $\mathrm{SO}_7(q)$ and $\mathrm{Sp}_6(q)$ ”, *J. Algebra* **413** (2014), p. 15-40.
- [27] M. JIMBO, K. C. MISRA, T. MIWA & M. OKADO, “Combinatorics of representations of $U_q(\widehat{\mathfrak{sl}}(n))$ at $q = 0$ ”, *Commun. Math. Phys.* **136** (1991), no. 3, p. 543-566.
- [28] G. LUSZTIG, *Characters of reductive groups over a finite field*, Annals of Mathematics Studies, vol. 107, Princeton University Press, 1984, xxi+384 pages.
- [29] J. MICHEL, “The development version of the CHEVIE package of GAP3”, *J. Algebra* **435** (2015), p. 308-336.
- [30] T. OKUYAMA & K. WAKI, “Decomposition numbers of $\mathrm{Sp}(4, q)$ ”, *J. Algebra* **199** (1998), no. 2, p. 544-555.
- [31] R. ROUQUIER, “Modular representations of finite groups and Lie theory”, <https://arxiv.org/abs/2202.08451>, 2022.

- [32] N. SPALTENSTEIN, “A property of special representations of Weyl groups”, *J. Reine Angew. Math.* **343** (1983), p. 212-220.
- [33] T. A. SPRINGER, “Regular elements of finite reflection groups”, *Invent. Math.* **25** (1974), p. 159-198.
- [34] D. UGLOV, “Canonical bases of higher-level q -deformed Fock spaces and Kazhdan–Lusztig polynomials”, in *Physical combinatorics (Kyoto, 1999)*, Progress in Mathematics, vol. 191, Birkhäuser, 2000, p. 249-299.

Manuscrit reçu le 19 février 2021,
révisé le 1^{er} juillet 2022,
accepté le 23 septembre 2022.

Olivier DUDAS
Aix-Marseille Université, CNRS,
Institut de Mathématiques de Marseille – I2M
Campus de Luminy,
Avenue de Luminy,
Case 930,
13288 Marseille Cedex 9 (France)
olivier.dudas@univ-amu.fr

Emily NORTON
School of Mathematics, Statistics
and Actuarial Science,
University of Kent,
Canterbury, Kent, CT2 7NZ (United Kingdom)
e.norton@kent.ac.uk