

ANNALES DE L'institut fourier

Nuria Corral Jacobian curve of singular foliations Tome 75, nº 1 (2025), p. 229-289.

https://doi.org/10.5802/aif.3665

Article mis à disposition par son auteur selon les termes de la licence CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE [CC] BY-ND http://creativecommons.org/licenses/by-nd/3.0/fr/



Les Annales de l'Institut Fourier sont membres du Centre Mersenne pour l'édition scientifique ouverte www.centre-mersenne.org e-ISSN : 1777-5310

JACOBIAN CURVE OF SINGULAR FOLIATIONS

by Nuria CORRAL (*)

Dedicated to Felipe Cano, with admiration and gratitude

ABSTRACT. — Topological properties of the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ of two foliations \mathcal{F} and \mathcal{G} are described in terms of invariants associated to the foliations. The main result gives a decomposition of the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ which depends on how similar are the foliations \mathcal{F} and \mathcal{G} . The similarity between foliations is codified in terms of the Camacho–Sad indices of the foliations with the notion of collinear point or divisor. Our approach allows to recover the results concerning the factorization of the jacobian curve of two plane curves and of the polar curve of a curve or a foliation.

RÉSUMÉ. — Nous décrivons des propriétés topologiques de la courbe jacobienne $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ de deux feuilletages \mathcal{F} et \mathcal{G} en termes des invariants associés aux feuilletages. Le resultat principal donne une décomposition de la courbe jacobienne $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ qui dépend de la similitude des feuilletages \mathcal{F} et \mathcal{G} . Cette similitude entre les feuilletages est codifiée en termes des indices de Camacho–Sad des feuilletages avec la notion de point ou diviseur colinéaire. Notre approche permet de récupérer les résultats concernant la factorisation de la courbe jacobienne de deux courbes planes et de la courbe polaire d'une courbe ou d'un feuilletage.

1. Introduction

Given two germs of holomorphic functions $f, g \in \mathbb{C}\{x, y\}$, the Jacobian determinant

$$J(f,g) = f_x g_y - f_y g_x$$

defines a curve called the *jacobian curve* of f and g (see [8, 23] for instance). The analytic type of the jacobian curve is an invariant of the analytic type of the pair of curves f = 0 and g = 0 but its topological type is not

Keywords: Jacobian curve, singular foliation, polar curve, Camacho–Sad index, equisingularity data.

²⁰²⁰ Mathematics Subject Classification: 32S65, 32S50, 14H20.

^(*) The author is supported by the Spanish research project PID2019-105621GB-I00/AEI/10.13039/501100011033 funded by the Agencia Estatal de Investigación – Ministerio de Ciencia e Innovación.

a topological invariant of the pair of curves (see [24]). Properties of the jacobian curve have been studied by several authors in terms of properties of the curves f = 0 and g = 0 (see for instance [8, 20], and [14] when g is a characteristic approximated root of f).

This notion can be studied in the more general context given by the theory of singular foliations: given two germs of foliations \mathcal{F} and \mathcal{G} in $(\mathbb{C}^2, 0)$, defined by the 1-forms $\omega = 0$ and $\eta = 0$, the *jacobian curve* $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ of \mathcal{F} and \mathcal{G} is the curve given by

$$\omega \wedge \eta = 0.$$

Note that this is the curve of tangency between both foliations. It is easy to show that the branches of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ are not separatrices of \mathcal{F} or \mathcal{G} provided that the foliations \mathcal{F} and \mathcal{G} do not have common separatrices.

If the foliation \mathcal{G} is non-singular, the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ coincides with the polar curve of the foliation \mathcal{F} . Properties of the equisingularity type of polar curves of foliations have been studied in [9, 10, 12, 30]. Moreover, if the foliation \mathcal{F} is given by df = 0 with $f \in \mathbb{C}\{x, y\}$, we recover the notion of polar curve of a plane curve. The local study of these curves has also been widely treated by many authors (see for instance [7, 13, 21, 25] or the recent works [2, 18]).

Moreover, the use of polar curves of foliations allowed to describe properties of foliations. In [6], the study of intersection properties of polar curves of foliations permitted to characterize generalized curve foliations as well as second type foliations; an expression of the GSV-index can also be given in terms of these invariants (see also [15] for the dicritical case). There are also some recent works that show the interest of jacobian curves or polar curves of foliations in the study of analytic invariants of curves (see for instance [16]) or singular foliations (see [26]).

The aim of this paper is to describe properties of the equisingularity type of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ in terms of invariants associated to the foliations \mathcal{F} and \mathcal{G} . Note that, in general, the locus $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ cannot be described from the data of \mathcal{F} and \mathcal{G} . It is enough to consider the non-singular foliations \mathcal{F} given by dx = 0and \mathcal{G} defined by dx + h(x, y) dy = 0, hence the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ is defined by h(x, y) = 0.

To illustrate the kind of conditions we are going to ask to the foliations and the type of results that we can obtain, let us explain the relationship between the multiplicity at the origin of the jacobian curve $\nu_0(\mathcal{J}_{\mathcal{F},\mathcal{G}})$ and the multiplicities of the foliations \mathcal{F} and \mathcal{G} . If the 1-forms ω and η defining \mathcal{F} and \mathcal{G} are given by $\omega = A(x, y) dx + B(x, y) dy$ and $\eta = P(x, y) dx + Q(x, y) dy$ respectively, the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ is defined by J(x, y) = 0

230

where

(1.1)
$$J(x,y) = \begin{vmatrix} A(x,y) & B(x,y) \\ P(x,y) & Q(x,y) \end{vmatrix}.$$

Thus the multiplicity at the origin $\nu_0(\mathcal{J}_{\mathcal{F},\mathcal{G}})$ of the jacobian curve satisfies

(1.2)
$$\nu_0(\mathcal{J}_{\mathcal{F},\mathcal{G}}) \ge \nu_0(\mathcal{F}) + \nu_0(\mathcal{G})$$

where $\nu_0(\mathcal{F})$, $\nu_0(\mathcal{G})$ denote the multiplicity at the origin of the foliations \mathcal{F} and \mathcal{G} respectively. One of the first results describing the properties of the jacobian curve shows that equality in (1.2) holds, that is,

$$\nu_0(\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \nu_0(\mathcal{F}) + \nu_0(\mathcal{G})$$

provided that the foliations \mathcal{F} and \mathcal{G} have different Camacho–Sad index at any singular point in the exceptional divisor E^1 obtained after one blow-up (see Lemma 2.3).

The main result in this paper, Theorem 6.4, gives a factorization of the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ of two generalized curve foliations \mathcal{F} and \mathcal{G} in terms of invariants given by the dual graph of the common minimal reduction of singularities of \mathcal{F} and \mathcal{G} . This result gives a decomposition of the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ in two classes of components: one for which we can control some properties of the topology from the data of \mathcal{F} and \mathcal{G} and another one for which such a control is impossible. The properties of the components in that decomposition depend on how "similar" are the foliations \mathcal{F} and \mathcal{G} in terms of its singularities and Camacho–Sad indices at the common singularities. We introduce the notion of collinear point and collinear divisor to measure this similarity between the foliations (see Section 4 where properties of collinear and non-collinear divisors are given).

The strategy used to prove the decomposition result is to study first the case when the foliations \mathcal{F} and \mathcal{G} have separatrices with non-singular irreducible components (Section 5). In this case, thanks to the hypothesis over the separatrices, we can compute "by hand" the infinitely near points of the jacobian curve under certain hypothesis over the foliations related with the notion of collinearity (a key point is Lemma 4.13 relating the weighted initial part of the 1-forms defining the foliations \mathcal{F} and \mathcal{G} and the one of the equation of the jacobian curve). In these computations, we use the existence of logarithmic models for generalized curve foliations (proved in [9]) and the properties shared by a foliation and its logarithmic model. Moreover, we describe the relationship between the jacobian curve of two foliations and the one of its logarithmic models (see Lemmas 4.14 and 4.16). These results will allow us to do some computations for the jacobian curve of two logarithmic foliations (see Theorem 5.2) and thus we get it for the jacobian curve of any non-dicritical generalized curve foliations.

Then we use a ramification $\rho : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ to reduce the general case to the previous one (Section 6). This strategy works since we can prove that the curves $\rho^{-1}\mathcal{J}_{\mathcal{F},\mathcal{G}}$ and $\mathcal{J}_{\rho^*\mathcal{F},\rho^*\mathcal{G}}$ "share" the same infinitely near points in the common reduction of singularities of $\rho^*\mathcal{F}$ and $\rho^*\mathcal{G}$ (see Lemma 6.1). Thus the results obtained in Section 5 can be used to describe properties of $\rho^{-1}\mathcal{J}_{\mathcal{F},\mathcal{G}}$ and hence, recover properties of the curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ since the equisingularity data of a curve can be recovered from the one obtained after ramification (see [10]). We include an appendix (Appendix A) devoted to explain all the details concerning the ramification process.

Section 2 is devoted to introduce notations and local invariants of curves and foliations which will be used throughout the paper. In Section 3 we recall the notion of logarithmic model (introduced in [9]) and some properties of logarithmic foliations.

In the last part of the article (Section 7) we show the role that the Camacho–Sad indices play to explain some behaviours of jacobian curves of plane curves. In particular, we show how our results imply the results of T.-C. Kuo and A. Parusiński concerning jacobian curves of plane curves [20], the results of E. García Barroso and J. Gwoździewicz about the jacobian curve of a plane curve and its approximate roots [14] and also previous results about polar curves of foliations (given in [9, 30]). All these results can be consider as particular cases of the results in this paper.

The article finishes with two appendices. The first one contains results concerning ramification. The second one is devoted to prove some formulas which describe the multiplicity of intersection of the jacobian curve with the separatrices of the foliations \mathcal{F} and \mathcal{G} in terms of the local invariants associated to \mathcal{F} and \mathcal{G} . These formulas generalize some properties of polar curves of a foliation given in [6, 9] which were key in the proof of the characterization of generalized curve foliations and second type foliations given in [6].

Acknowledgements

The author thanks the referee for their carefully reading of the paper and for their contribution to a good presentation of it.

2. Local invariants

2.1. Foliations

Let \mathbb{F} be the space of singular foliations in $(\mathbb{C}^2, 0)$. An element $\mathcal{F} \in \mathbb{F}$ is defined by a 1-form $\omega = 0$, with $\omega = A(x, y) dx + B(x, y) dy$, or by the vector field $\mathbf{v} = -B(x, y)\partial/\partial x + A(x, y)\partial/\partial y$ where $A, B \in \mathbb{C}\{x, y\}$ are relatively prime. The origin is a singular point if A(0) = B(0) = 0. The multiplicity $\nu_0(\mathcal{F})$ of \mathcal{F} at the origin is the minimum of the orders $\nu_0(A)$, $\nu_0(B)$ at the origin. Thus, the origin is a singular point of \mathcal{F} if $\nu_0(\mathcal{F}) \ge 1$.

Consider a germ of irreducible analytic curve S at $(\mathbb{C}^2, 0)$. We say that S is a separatrix of \mathcal{F} at the origin if S is an invariant curve of the foliation \mathcal{F} . Therefore, if f = 0 is a reduced equation of S, we have that f divides $\omega \wedge \mathrm{d} f$.

Let us now recall the desingularization process of a foliation. We say that the origin is a simple singularity of \mathcal{F} if there are local coordinates (x, y)in $(\mathbb{C}^2, 0)$ such that \mathcal{F} is given by a 1-form of the type

$$\lambda y \, \mathrm{d}x - \mu x \, \mathrm{d}y + \text{h.o.t}$$

with $\mu \neq 0$ and $\lambda/\mu \notin \mathbb{Q}_{>0}$. If $\lambda = 0$, the singularity is called a saddlenode. There are two formal invariant curves Γ_x and Γ_y which are tangent to x = 0 and y = 0 respectively, and such that they are both convergent in the case that $\lambda \mu \neq 0$. In the saddle-node situation with $\lambda = 0$ and $\mu \neq 0$, we say that the saddle-node is well oriented with respect to the curve Γ_y .

Let $\pi_1 : X_1 \to (\mathbb{C}^2, 0)$ be the blow-up of the origin with $E_1 = \pi_1^{-1}(0)$ the exceptional divisor. We say that the blow-up π_1 (or the exceptional divisor E_1) is non-dicritical if E_1 is invariant by the strict transform $\pi_1^* \mathcal{F}$ of \mathcal{F} ; otherwise, the exceptional divisor E_1 is generically transversal to $\pi_1^* \mathcal{F}$ and we say that the blow-up π_1 (or the divisor E_1) is dicritical.

A reduction of singularities of \mathcal{F} is a morphism $\pi : X \to (\mathbb{C}^2, 0)$, composition of a finite number of punctual blow-ups, such that the strict transform $\pi^* \mathcal{F}$ of \mathcal{F} verifies that

- each irreducible component of the exceptional divisor $\pi^{-1}(0)$ is either invariant by $\pi^* \mathcal{F}$ or transversal to $\pi^* \mathcal{F}$;
- all the singular points of $\pi^* \mathcal{F}$ are simple and do not belong to a discritical component of the exceptional divisor.

There exists a reduction of singularities as a consequence of Seidenberg's Desingularization Theorem [32]. Moreover, there is a minimal morphism π such that any other reduction of singularities of \mathcal{F} factorizes through the minimal one. The centers of the blow-ups of a reduction of singularities of

 \mathcal{F} are called *infinitely near points* of \mathcal{F} . If all the irreducible components of the exceptional divisor are invariant by $\pi^* \mathcal{F}$, we say that the foliation \mathcal{F} is non-dicritical; otherwise, \mathcal{F} is called a *dicritical foliation*.

A non-dicritical foliation \mathcal{F} is called a generalized curve foliation if there are not saddle-node singularities in the reduction of singularities (see [3]). We will denote \mathbb{G} the space of non-dicritical generalized curve foliations in $(\mathbb{C}^2, 0)$. The foliation \mathcal{F} is of second type if all saddle-nodes of $\pi^* \mathcal{F}$ are well oriented with respect to the exceptional divisor $E = \pi^{-1}(0)$ (see [22]).

In order to describe properties of generalized curve foliations and second type foliations, let us recall some local invariants used in the local study of foliations in dimension two (see for instance [5]). The *Milnor number* $\mu_0(\mathcal{F})$ is given by

$$\mu_0(\mathcal{F}) = \dim_{\mathbb{C}} \frac{\mathbb{C}\{x, y\}}{(A, B)} = (A, B)_0,$$

where $(A, B)_0$ stands for the intersection multiplicity. Note that, if the foliation is defined by df = 0, the Milnor number of the foliation coincides with the one of the curve given by f = 0. Given an irreducible curve S and a primitive parametrization $\gamma : (\mathbb{C}, 0) \to (\mathbb{C}^2, 0)$ of S with $\gamma(t) = (x(t), y(t))$, we have that S is a separatrix of \mathcal{F} if and only if $\gamma^*\omega = 0$. In this case, the Milnor number $\mu_0(\mathcal{F}, S)$ of \mathcal{F} along S is given by

(2.1)
$$\mu_0(\mathcal{F}, S) = \begin{cases} \operatorname{ord}_t(B(\gamma(t))) - \operatorname{ord}_t(x(t)) + 1 & \text{if } x(t) \neq 0, \\ \operatorname{ord}_t(A(\gamma(t))) - \operatorname{ord}_t(y(t)) + 1 & \text{if } y(t) \neq 0, \end{cases}$$

(this number is also called *multiplicity of* **v** along S, see [3, p. 152-153]). If S is not a separatrix, we define the tangency order $\tau_0(\mathcal{F}, S)$ by

(2.2)
$$\tau_0(\mathcal{F}, S) = \operatorname{ord}_t(\alpha(t))$$

where $\gamma^* \omega = \alpha(t) dt$. If S = (y = 0) is a non-singular invariant curve of the foliation \mathcal{F} , the Camacho–Sad index of \mathcal{F} relative to S at the origin is given by

(2.3)
$$\mathcal{I}_0(\mathcal{F}, S) = -\operatorname{Res}_0 \frac{a(x, 0)}{b(x, 0)}$$

where the 1-form defining \mathcal{F} is written as ya(x, y) dx + b(x, y) dy (see [4]).

Next result summarizes some of the properties of second type and generalized curve foliations that we will use throughout the text:

THEOREM 2.1 ([3, 6, 22]). — Let \mathcal{F} be a non-dicritical foliation and consider \mathcal{G}_f the foliation defined by df = 0 where f is a reduced equation of the curve $S_{\mathcal{F}}$ of separatrices of \mathcal{F} . Let $\pi : X \to (\mathbb{C}^2, 0)$ be the minimal reduction of singularities of \mathcal{F} .

- (i) π is a reduction of singularities of S_F. Moreover, π is the minimal reduction of singularities of the curve S_F if and only if F is of second type;
- (ii) ν₀(F) ≥ ν₀(G_f) and the equality holds if and only if F is of second type;
- (iii) $\mu_0(\mathcal{F}) \ge \mu_0(\mathcal{G}_f)$ and the equality holds if and only if \mathcal{F} is a generalized curve foliation;
- (iv) if S is an irreducible curve which is not a separatrix of \mathcal{F} , then $\tau_0(\mathcal{F}, S) \ge \tau_0(\mathcal{G}_f, S)$ and the equality holds if and only if \mathcal{F} is of second type.

Recall that for the hamiltonian foliation \mathcal{G}_f we have that $\nu_0(\mathcal{G}_f) = \nu_0(S_F) - 1$, $\mu_0(\mathcal{G}_f) = \mu_0(S_F)$ and $\tau_0(\mathcal{G}_f, S) = (S_F, S)_0 - 1$ where $\nu_0(S_F)$ is the multiplicity of the curve S_F at the origin, $\mu_0(S_F)$ is the Milnor number of the curve S_F and $(S_F, S)_0$ denotes the intersection multiplicity of the curves S_F and S at the origin.

Notation. — Given a plane curve C in $(\mathbb{C}^2, 0)$, we denote by \mathbb{F}_C the subspace of \mathbb{F} composed by the foliations having C as curve of separatrices and \mathbb{G}_C the foliations of \mathbb{F}_C which are generalized curve foliations.

Moreover, for generalized curve foliations we have that

LEMMA 2.2 ([11]). — Assume that \mathcal{F} is a non-dicritical generalized curve foliation. Let $\pi : (X, P) \to (\mathbb{C}^2, 0)$ be a morphism composition of a finite number of punctual blow-ups and take an irreducible component Eof the exceptional divisor $\pi^{-1}(0)$ with $P \in E$. Then, the strict transforms $\pi^* \mathcal{F}$ and $\pi^* \mathcal{G}_f$ satisfy that

(i)
$$\nu_P(\pi^*\mathcal{F}) = \nu_P(\pi^*\mathcal{G}_f);$$

(ii)
$$\mu_P(\pi^*\mathcal{F}, E) = \mu_P(\pi^*\mathcal{G}_f, E).$$

where f = 0 is a reduced equation of the curve $S_{\mathcal{F}}$ of separatrices of \mathcal{F} .

2.2. Weighted initial forms and Jacobian curves.

Fix coordinates (x, y) in $(\mathbb{C}^2, 0)$. Given a 1-form ω , we can write $\omega = \sum_{i,j} \omega_{ij}$ where $\omega_{ij} = A_{ij} x^{i-1} y^j \, dx + B_{ij} x^i y^{j-1} \, dy$. We denote $\Delta(\omega) = \Delta(\omega; x, y) = \{(i, j) : \omega_{ij} \neq 0\}$ and the Newton polygon $\mathcal{N}(\mathcal{F}; x, y) = \mathcal{N}(\mathcal{F}) = \mathcal{N}(\omega)$ is given by the convex envelop of $\Delta(\omega) + (\mathbb{R}_{\geq 0})^2$.

Given a rational number $\alpha \in \mathbb{Q}$, we define the initial form of ω with weight α

$$\operatorname{In}_{\alpha}(\omega; x, y) = \sum_{i+\alpha j=k} \omega_{ij}$$

TOME 75 (2025), FASCICULE 1

where $i + \alpha j = k$ is the equation of the first line of slope $-1/\alpha$ which intersects the Newton polygon $\mathcal{N}(\mathcal{F})$ in the coordinates (x, y). Note that $k = \nu_{(1,\alpha)}(\omega)$ where $\nu_{(1,\alpha)}(\omega) = \min\{i + \alpha j : \omega_{ij} \neq 0\}$ is the $(1, \alpha)$ -degree of ω . Hence, we have that the multiplicity $\nu_0(\mathcal{F}) = \nu_{(1,1)}(\omega)$.

In a similar way, given any function $f = \sum_{ij} f_{ij}x^i y^j \in \mathbb{C}\{x, y\}$, we denote $\Delta(f) = \Delta(f; x, y) = \{(i, j) : f_{ij} \neq 0\}$ and the Newton polygon $\mathcal{N}(C; x, y) = \mathcal{N}(C)$ of the curve C = (f = 0) is the convex envelop of $\Delta(f) + (\mathbb{R}_{\geq 0})^2$. Note that $\mathcal{N}(C; x, y) = \mathcal{N}(df; x, y)$ and $\mathcal{N}(\mathcal{F}) = \mathcal{N}(C)$ if $\mathcal{F} \in \mathbb{G}_C$. Thus, we can define the *initial form* $\operatorname{In}_{\alpha}(f; x, y) = \sum_{(i,j) \in L} f_{ij}x^i y^j$ where L is the first line of slope $-1/\alpha$ which intersects $\mathcal{N}(C)$. Note that, if f = 0 is an equation of the curve C, then $\operatorname{In}_1(f; x, y)$ gives an equation of the tangent cone of C, and hence $\operatorname{In}_1(f; x, y) = \sum_{i+j=\nu_0(C)} f_{ij}x^i y^j$.

With these notations we can state the first result which illustrates the type of conditions we are going to ask the foliations \mathcal{F} and \mathcal{G} in order to be able to describe properties of the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$.

LEMMA 2.3. — Let \mathcal{F} and \mathcal{G} be two foliations in $(\mathbb{C}^2, 0)$ and consider $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ its jacobian curve. Let $\pi_1 : X_1 \to (\mathbb{C}^2, 0)$ be the blow-up of the origin and $E^1 = \pi_1^{-1}(0)$ be the exceptional divisor. If there is a point $R \in E^1$ such that the Camacho–Sad indices verify that $\mathcal{I}_R(\pi_1^*\mathcal{F}, E^1) \neq \mathcal{I}_R(\pi_1^*\mathcal{G}, E^1)$, then

$$\nu_0(\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \nu_0(\mathcal{F}) + \nu_0(\mathcal{G}).$$

Proof. — Take (x, y) coordinates such that x = 0 is not tangent to the foliations \mathcal{F} and \mathcal{G} and let (x_1, y_1) be coordinates in the first chart of the blow-up such that $\pi_1(x_1, y_1) = (x_1, x_1y_1)$ and $E^1 = (x_1 = 0)$. Assume that $\nu_0(\mathcal{J}_{\mathcal{F},\mathcal{G}}) > \nu_0(\mathcal{F}) + \nu_0(\mathcal{G})$, then $\operatorname{In}_1(\omega) \wedge \operatorname{In}_1(\eta) \equiv 0$. Thus, if write

$$In_1(\omega) = A_{\nu_0(\mathcal{F})}(x, y) \, \mathrm{d}x + B_{\nu_0(\mathcal{F})}(x, y) \, \mathrm{d}y$$

$$In_1(\eta) = P_{\nu_0(\mathcal{G})}(x, y) \, \mathrm{d}x + Q_{\nu_0(\mathcal{G})}(x, y) \, \mathrm{d}y$$

then we have that

(2.4)
$$A_{\nu_0(\mathcal{F})}(x,y)Q_{\nu_0(\mathcal{G})}(x,y) - B_{\nu_0(\mathcal{F})}(x,y)P_{\nu_0(\mathcal{G})}(x,y) \equiv 0.$$

The computation of the Camacho–Sad index at a point R given by (0, c) in the coordinates (x_1, y_1) gives

$$\mathcal{I}_{R}(\pi_{1}^{*}\mathcal{F}, E^{1}) = -\operatorname{Res}_{y=c} \frac{B_{\nu_{0}(\mathcal{F})}(1, y)}{A_{\nu_{0}(\mathcal{F})}(1, y) + yB_{\nu_{0}(\mathcal{F})}(1, y)}$$
$$\mathcal{I}_{R}(\pi_{1}^{*}\mathcal{G}, E^{1}) = -\operatorname{Res}_{y=c} \frac{Q_{\nu_{0}(\mathcal{G})}(1, y)}{P_{\nu_{0}(\mathcal{G})}(1, y) + yQ_{\nu_{0}(\mathcal{G})}(1, y)}$$

The equality in (2.4), implies

$$\mathcal{I}_R(\pi_1^*\mathcal{F}, E^1) = \mathcal{I}_R(\pi_1^*\mathcal{G}, E^1)$$

for any point R in E^1 . This gives a contradiction with the hypothesis over the foliations \mathcal{F} and \mathcal{G} .

The condition over the Camacho–Sad indices of the foliations in the previous lemma is related with the notion of collinear point that will be introduced in Section 4.

2.3. Equisingularity data and dual graph of plane curves

In this subsection we will fix some notations concerning the equisingularity data of a plane curve $C = \bigcup_{i=1}^{r} C_i$ in $(\mathbb{C}^2, 0)$ (see Appendix A.1 for more details). Given an irreducible component C_i of C, we will denote $n^i = \nu_0(C_i)$ the multiplicity of C_i at the origin, $\{\beta_0^i, \beta_1^i, \ldots, \beta_{g_i}^i\}$ the characteristic exponents of C_i and $\{(m_l^i, n_l^i)\}_{l=1}^{g_l}$ the Puiseux pairs of C_i .

Let us denote $\pi_C : X_C \to (\mathbb{C}^2, 0)$ the minimal reduction of singularities of the curve C. The dual graph G(C) is constructed as follows: each irreducible component E of the exceptional divisor $\pi_C^{-1}(0)$ is represented by a vertex which we also call E (we identify a divisor and its associated vertex in the dual graph). Two vertices are joined by an edge if and only if the associated divisors intersect. Each irreducible component of C is represented by an arrow joined to the only divisor which meets the strict transform of C by π_C . We can give a weight to each vertex E of G(C) equal to the self-intersection of the divisor $E \subset X_C$ and this weighted dual graph is equivalent to the equisingularity data of C.

If we denote by E^1 the irreducible component of $\pi_C^{-1}(0)$ corresponding to the divisor obtained by the blow-up of the origin, we can give an orientation to the graph G(C) beginning from the first divisor E^1 . The geodesic of a divisor E is the path which joins the first divisor E^1 with the divisor E. The geodesic of a curve is the geodesic of the divisor that meets the strict transform of the curve. Thus, there is a partial order in the set of vertices of G(C) given by E < E' if and only if the geodesic of E' goes through E. A maximal divisor in G(C) will be a maximal element in the set of vertices of G(C) with this partial order. Given a divisor E of G(C), we denote by I_E the set of indices $i \in \{1, 2, ..., r\}$ such that E belongs to the geodesic of the curve C_i .

Given a vertex E of G(C), we define the number b_E^C in the following way: $b_E^C + 1$ is the valence of E if $E \neq E^1$ and $b_{E^1}^C$ is the valence of E^1 in G(C) (recall that the valence of a divisor E in G(C) is the number of arrows and edges attached to E in G(C)). Given a divisor E of G(C), we say that Eis a bifurcation divisor of G(C) if $b_E^C \ge 2$ and a terminal divisor of G(C)if $b_E^C = 0$. A dead arc is a path which joins a bifurcation divisor with a terminal one without going through other bifurcation divisor. We denote by B(C) the set of bifurcation divisors of G(C). If there is no confusion with the curve C we are working with, we will denote $b_E = b_E^C$ for any divisor E in G(C).

Given an irreducible component E of $\pi_C^{-1}(0)$, we denote by $\pi_E : X_E \to (\mathbb{C}^2, 0)$ the morphism reduction of π_C to E (see [10]), that is, the morphism which verifies that

- the morphism π_C factorizes as $\pi_C = \pi_E \circ \pi'_E$ where π_E and π'_E are composition of punctual blow-ups;
- the divisor E is the strict transform by π'_E of an irreducible component E_{red} of $\pi_E^{-1}(0)$ and $E_{\text{red}} \subset X_E$ is the only component of $\pi_E^{-1}(0)$ with self-intersection equal to -1.

We will denote by $\pi_E^* C$ the strict transform of C by the morphism π_E . The points $\pi_E^* C \cap E_{\text{red}}$ are called *infinitely near points* of C in E.

Remark 2.4. — If C is a curve with only non-singular irreducible components and E is an irreducible component of $\pi_C^{-1}(0)$, then the number of infinitely near points of π_E^*C in E_{red} is equal to b_E . That is, the cardinal of the set $\pi_E^*C \cap E_{\text{red}}$ coincides with b_E .

3. Logarithmic foliations

Consider a germ of plane curve $C = \bigcup_{i=1}^{r} C_i$ in $(\mathbb{C}^2, 0)$. Take $f \in \mathbb{C}\{x, y\}$ such that C = (f = 0) and let us write $f = f_1 \cdots f_r$ with $f_i \in \mathbb{C}\{x, y\}$ irreducible. Given $\lambda = (\lambda_1, \ldots, \lambda_r) \in \mathbb{C}^r$, we can consider the logarithmic foliation \mathcal{L}^C_{λ} defined by

(3.1)
$$f_1 \cdots f_r \sum_{i=1}^r \lambda_i \frac{\mathrm{d}f_i}{f_i} = 0.$$

The logarithmic foliation $\mathcal{L}_{\lambda}^{C}$ belongs to \mathbb{G}_{C} provided that λ avoids certain rational resonances. Each generalized curve foliation $\mathcal{F} \in \mathbb{G}_{C}$ has a logarithmic model $\mathcal{L}_{\lambda}^{C}$, that is, a logarithmic foliation such that the Camacho–Sad indices of \mathcal{F} and $\mathcal{L}_{\lambda}^{C}$ coincide along the reduction of singularities (see [9]); note that \mathcal{F} and $\mathcal{L}_{\lambda}^{C}$ have the same separatrices and the same minimal reduction of singularities. Moreover, the logarithmic model of \mathcal{F} is unique once a reduced equation of the separatrices is fixed. Thus, for each foliation $\mathcal{F} \in \mathbb{G}_C$, we denote by $\lambda(\mathcal{F})$ the exponent vector of the logarithmic model of \mathcal{F} and we denote $\mathbb{G}_{C,\lambda}$ the set of foliations $\mathcal{F} \in \mathbb{G}_C$ such that $\lambda(\mathcal{F}) = \lambda$. A particular case of logarithmic foliation is the "hamiltonian" foliation defined by df = 0 which corresponds to $\lambda = (1, \ldots, 1)$; this foliation coincides with the foliation \mathcal{G}_f used in Section 2.

Let us fix some notations concerning logarithmic foliations that will be used in the sequel. Assume that the curve $C = \bigcup_{i=1}^{r} C_i$ has only nonsingular irreducible components and consider a non-dicritical logarithmic foliation $\mathcal{L}^{C}_{\lambda}$ given by (3.1). Let $\pi_{C} : X_{C} \to (\mathbb{C}^{2}, 0)$ be the minimal reduction of singularities of C, take E an irreducible component of $\pi_{C}^{-1}(0)$ and consider $\pi_{E} : X_{E} \to (\mathbb{C}^{2}, 0)$ the morphism reduction of π_{C} to E (see Subsection 2.3). Given an irreducible component C_{j} of C and a divisor Fof $\pi_{C}^{-1}(0)$, we denote $\varepsilon_{F}^{C_{j}} = 1$ if the geodesic of C_{j} contains the divisor Fand $\varepsilon_{F}^{C_{j}} = 0$ otherwise, that is,

$$\varepsilon_F^{C_j} = \begin{cases} 1, & \text{if } j \in I_F, \\ 0, & \text{if } j \notin I_F. \end{cases}$$

The residue of the logarithmic foliation $\mathcal{L}_{\lambda}^{C}$ along the divisor E is given by

(3.2)
$$\kappa_E(\mathcal{L}^C_{\lambda}) = \sum_{j=1}^r \lambda_j \sum_{E' \leqslant E} \varepsilon_{E'}^{C_j},$$

where $E' \leq E$ means all divisors in G(C) which are in the geodesic of E (including E itself). Note that $\kappa_E(\mathcal{L}^C_{\lambda}) = \sum_{j=1}^r \lambda_j m_E^{C_j}$ where $m_E^{C_j}$ is the multiplicity of $f_j \circ \pi_E$ along the divisor E (see [27, 28]).

Let $\{R_1^E, R_2^E, \ldots, R_{b_E}^E\}$ be the set of points $\pi_E^* C \cap E_{\text{red}}$ where we denote $b_E = b_E^C$ and put $I_{R_l^E}^C = \{i \in \{1, \ldots, r\} : \pi_E^* C_i \cap E_{\text{red}} = \{R_l^E\}\}$ for $l = 1, 2, \ldots, b_E$, that is, $i \in I_{R_l^E}^C$ if E belongs to the geodesic of the curve C_i in G(C) and R_i^E is an infinitely near point of C_i . With the notations introduced in Subsection 2.3, we have that $I_E = \bigcup_{l=1}^{b_E} I_{R_l^E}^C$.

An easy computation shows that the Camacho–Sad index of $\pi_E^* \mathcal{L}_{\lambda}^C$ relative to E_{red} at a point R_l^E is given by

(3.3)
$$\mathcal{I}_{R_l^E}(\pi_E^*\mathcal{L}_{\lambda}^C, E_{\mathrm{red}}) = -\frac{\sum_{i \in I_{R_l^E}} \lambda_i}{\kappa_E(\mathcal{L}_{\lambda}^C)}$$

(see [9, Section 4]).

TOME 75 (2025), FASCICULE 1

In Appendix A we include a subsection where we explain the behaviour of the above invariants associated to logarithmic foliations after ramification (see Subsection A.3).

4. Collinear and non-collinear points and divisors

In this section we will introduce some notations and definitions in order to describe properties of the jacobian curve that will be given in Section 5.

Let C and D be two plane curves in $(\mathbb{C}^2, 0)$ without common branches and assume that the curve $Z = C \cup D$ has only non-singular irreducible components. Take now $\mathcal{F} \in \mathbb{G}_C$ and $\mathcal{G} \in \mathbb{G}_D$ and let $\pi_Z : X_Z \to (\mathbb{C}^2, 0)$ be the minimal reduction of singularities of Z which gives a common reduction of singularities of \mathcal{F} and \mathcal{G} . Recall that, given an irreducible component Eof $\pi_Z^{-1}(0)$, we denote by $\pi_E : X_E \to (\mathbb{C}^2, 0)$ the morphism reduction of π_Z to E (see Section 2.3).

Remark 4.1. — With the above assumptions about Z, if v(E) = p (see Appendix A.1), then the morphism π_E is a composition of p punctual blowups

$$(\mathbb{C}^2, 0) \xleftarrow{\sigma_1} (X_1, P_1) \xleftarrow{\sigma_2} \cdots \xleftarrow{\sigma_{p-1}} (X_{p-1}, P_{p-1}) \xleftarrow{\sigma_p} X_p = X_E.$$

Moreover, if (x, y) are coordinates in $(\mathbb{C}^2, 0)$, there is a change of coordinates $(x, y) = (\bar{x}, \bar{y} + \varepsilon_E(\bar{x}))$, with $\varepsilon_E(x) = a_1x + \cdots + a_{p-1}x^{p-1}$, such that the blow-up σ_j is given by $x_{j-1} = x_j$, $y_{j-1} = x_jy_j$, for $j = 1, 2, \ldots, p$, where (x_j, y_j) are coordinates centered at P_j and $(x_0, y_0) = (\bar{x}, \bar{y})$. We say that (\bar{x}, \bar{y}) are coordinates in $(\mathbb{C}^2, 0)$ adapted to E. Note that in these coordinates, if (x_p, y_p) are coordinates in the first chart of E_{red} we have that $\pi_E(x_p, y_p) = (x_p, x_p^p y_p)$ and $E_{\text{red}} = (x_p = 0)$.

In this section, we will denote $b_E = b_E^Z$. Let $\{R_1^E, R_2^E, \ldots, R_{b_E}^E\}$ be the infinitely near points of Z in E, that is, $\pi_E^* Z \cap E_{\text{red}} = \{R_1^E, R_2^E, \ldots, R_{b_E}^E\}$. Note that these points are the union of the singular points of $\pi_E^* \mathcal{F}$ and $\pi_E^* \mathcal{G}$ in the first chart of E_{red} (the singular points of the foliations which do not correspond to a corner of the divisor). We denote

$$\Delta_E^{\mathcal{F},\mathcal{G}}(R_i^E) = \mathcal{I}_{R_i^E}(\pi_E^*\mathcal{G}, E_{\mathrm{red}}) - \mathcal{I}_{R_i^E}(\pi_E^*\mathcal{F}, E_{\mathrm{red}})$$

where $\mathcal{I}_{R_i^E}(\pi_E^*\mathcal{F}, E_{\text{red}})$ is the Camacho–Sad index of $\pi_E^*\mathcal{F}$ relative to E_{red} at the point R_i^E (see definition given in (2.3)). We will denote $\Delta_E(R_i^E) = \Delta_E^{\mathcal{F},\mathcal{G}}(R_i^E)$ if it is clear the foliations \mathcal{F} and \mathcal{G} we are working with.

In view of the notations given in [19, 20] for curves, we introduce the following definitions for foliations:

DEFINITION 4.2. — We say that an infinitely near point R_l^E of Z is a collinear point for the foliations \mathcal{F} and \mathcal{G} in E if $\Delta_E(R_l^E) = 0$; otherwise we say that R_l^E is a non-collinear point.

We say that a divisor E is collinear (for the foliations \mathcal{F} and \mathcal{G}) if $\Delta_E(R_l^E) = 0$ for all $l = 1, \ldots, b_E$; otherwise E is called a non-collinear divisor. A divisor E is called purely non-collinear if $\Delta_E(R_l^E) \neq 0$ for each $l = 1, \ldots, b_E$.

We denote by $\operatorname{Col}(E)$ the set of collinear points of E and by $\operatorname{NCol}(E)$ the set of non-collinear points (for the foliations \mathcal{F} and \mathcal{G}). It is clear that $\operatorname{Col}(E) \cup \operatorname{NCol}(E) = \{R_1^E, R_2^E, \ldots, R_{b_F}^E\}.$

Remark 4.3. — Note that if E is a maximal bifurcation divisor (with the partial order in G(C) given in Subsection 2.3), then E is purely noncollinear. This follows from the fact that, if E is a maximal bifurcation divisor, then each infinitely near point R_l^E of Z in E is in the geodesic of only one irreducible component of Z and hence it is a singular point for only one of the foliations $\pi_E^* \mathcal{F}$ or $\pi_E^* \mathcal{G}$. Moreover, it is a simple singularity. In fact, we have that

$$\Delta_E(R_l^E) = \begin{cases} -\mathcal{I}_{R_l^E}(\pi_E^*\mathcal{F}, E_{\text{red}}), & \text{if } R_l^E \in \pi_E^*C \cap E, \\ \mathcal{I}_{R_l^E}(\pi_E^*\mathcal{G}, E_{\text{red}}), & \text{if } R_l^E \in \pi_E^*D \cap E. \end{cases}$$

If $R_l^E \in \pi_E^* C \cap E$, we have that $\mathcal{I}_{R_l^E}(\pi_E^* \mathcal{F}, E_{\text{red}}) \neq 0$ since \mathcal{F} is a generalized curve foliation and R_l^E is a simple singularity of $\pi_E^* \mathcal{F}$ (similarly when $R_l^E \in \pi_E^* D \cap E$). Consequently, $\Delta_E(R_l^E) \neq 0$ for each $l = 1, \ldots, b_E$.

Although the definition of $\Delta_E(R_l^E)$ seems different to the one given by Kuo and Parusiński in [19, 20], we will show in Subsection 7.1 that both definitions coincide in the case of curves.

Take coordinates (x_p, y_p) in the first chart of E_{red} such that $\pi_E(x_p, y_p) = (x_p, x_p^p y_p), E_{\text{red}} = (x_p = 0)$ and assume that $R_l^E = (0, c_l^E), l = 1, 2, \dots, b_E$, in these coordinates. We define the rational function $\mathcal{M}_E(z) = \mathcal{M}_E^{\mathcal{F},\mathcal{G}}(z)$ associated to the divisor E for the foliations \mathcal{F} and \mathcal{G} by

(4.1)
$$\mathcal{M}_E(z) = \sum_{l=1}^{b_E} \frac{\Delta_E(R_l^E)}{z - c_l^E}.$$

Remark 4.4. — Note that although $\Delta_E^{\mathcal{F},\mathcal{G}}(R_l^E)$ and $\mathcal{M}_E^{\mathcal{F},\mathcal{G}}(z)$ depend on the foliations \mathcal{F} and \mathcal{G} , we have that

$$\Delta_E^{\mathcal{F},\mathcal{G}}(R_l^E) = \Delta_E^{\mathcal{L}_\lambda^C,\mathcal{L}_\mu^D}(R_l^E); \qquad \mathcal{M}_E^{\mathcal{F},\mathcal{G}}(z) = \mathcal{M}_E^{\mathcal{L}_\lambda^C,\mathcal{L}_\mu^D}(z)$$

provided that $\mathcal{F} \in \mathbb{G}_{C,\lambda}$ and $\mathcal{G} \in \mathbb{G}_{D,\mu}$.

TOME 75 (2025), FASCICULE 1

Remark 4.5. — Observe that if E is a non-collinear divisor, then $\mathcal{M}_E(z) \neq 0$.

Let $M(E) = \{Q_1^E, \ldots, Q_{s_E}^E\}$ be the set of points of E_{red} given by $Q_l^E = (0, q_l)$ in coordinates (x_p, y_p) where $\{q_1, \ldots, q_{s_E}\}$ is the set of zeros of $\mathcal{M}_E(z)$. We denote by $t_{Q_l^E}$ the multiplicity of q_l as a zero of $\mathcal{M}_E(z)$ and $t(E) = \sum_{l=1}^{s_E} t_{Q_l^E}$ the degree of the numerator of the rational function $\mathcal{M}_E(z)$. We put $t_P = 0$ for any $P \in E \setminus \mathcal{M}(E)$. Note that it can happen that $\mathcal{M}(E) = \emptyset$ (see Example in [20, p. 584]).

LEMMA 4.6. — If $NCol(E) \neq \emptyset$, that is, E is a non-collinear divisor, then we have that

(4.2)
$$\operatorname{NCol}(E) \cap M(E) = \emptyset$$
 and $\#\operatorname{NCol}(E) \ge 1 + \sum_{P \in M(E)} t_P = 1 + t(E).$

Moreover, if $\sum_{R_l^E \in \text{NCol}(E)} \Delta_E(R_l^E) \neq 0$, then we have that

$$\sharp \operatorname{NCol}(E) = 1 + \sum_{P \in M(E)} t_P.$$

Proof. — With the notations above, we can write $\mathcal{M}_E(z)$ as follows

$$\mathcal{M}_E(z) = \sum_{R_l^E \in \mathrm{NCol}(E)} \frac{\Delta_E(R_l^E)}{z - c_l^E}.$$

Thus, the set of zeros of $\mathcal{M}_E(z)$ is given by the roots of the polynomial

(4.3)
$$\sum_{\substack{R_l^E \in \mathrm{NCol}(E) \\ R_j^E \in \mathrm{NCol}(E)}} \Delta_E(R_l^E) \prod_{\substack{j \text{ with } j \neq l \\ R_j^E \in \mathrm{NCol}(E)}} (z - c_j^E).$$

Consequently, if $z = c_{l_0}^E$ is a zero of $\mathcal{M}_E(z)$, then

$$\Delta_E(R_{l_0}^E) \prod_{\substack{R_j^E \in \operatorname{NCol}(E)\\ j \neq l_0}} (c_{l_0}^E - c_j^E) = 0$$

which implies that $\Delta_E(R_{l_0}^E) = 0$ and hence $R_{l_0}^E \notin \operatorname{NCol}(E)$. Moreover, the degree of the polynomial given in (4.3) is $\leq \sharp \operatorname{NCol}(E) - 1$; the equality is attained when $\sum_{R_l^E \in \operatorname{NCol}(E)} \Delta_E(R_l^E) \neq 0$. Thus we have the statements of the lemma.

Remark 4.7. — Observe that we can have that $\sum_{R_l^E \in \text{NCol}(E)} \Delta_E(R_l^E) = 0$ even if E is a purely non-collinear divisor. This can happen for instance when $E = E^1$ is a bifurcation divisor since in this situation $\sum_{i=1}^{b_E} \mathcal{I}_{R_i^E}(\pi_E^*\mathcal{F}, E_{\text{red}}) = \sum_{i=1}^{b_E} \mathcal{I}_{R_i^E}(\pi_E^*\mathcal{G}, E_{\text{red}}) = -1$ and this implies $\sum_{R_l^E \in \text{NCol}(E)} \Delta_E(R_l^E) = 0$ (see also Remark 4.11 and Corollary 4.12).

ANNALES DE L'INSTITUT FOURIER

Remark 4.8. — Note that it can happen that $\operatorname{Col}(E) \cap M(E) \neq \emptyset$. With the notations of Section 3, consider the foliations $\mathcal{F} = \mathcal{L}_{\lambda}^{C}$ and $\mathcal{G} = \mathcal{L}_{\mu}^{D}$ where

$$C = (f = 0), \quad f(x, y) = (y - x)(y + x^2)(y - x^2)(y + 2x^2), \quad \lambda = (1, 1, 1, 3)$$

$$\begin{split} D &= (g=0), \quad g(x,y) = (y+x)(y+x^2+x^3)(y-x^2+x^3), \quad \mu = (3,3,1). \\ \text{Take } Z &= C \cup D. \text{ Consider the morphism } \sigma = \pi_1 \circ \pi_2 \text{ where } \pi_1 : X_1 \rightarrow (\mathbb{C}^2,0) \text{ is the blow-up of the origin, } E_1 = \pi_1^{-1}(0) \text{ and } \pi_2 : X_2 \rightarrow (X_1,P_1) \\ \text{is the blow-up of the origin } P_1 \text{ of the first chart of } E_1 \text{ and } E_2 = \pi_2^{-1}(P_1). \\ \text{Taking coordinates } (x_2,y_2) \text{ in the first chart of } E_2 \text{ such that } \sigma(x_2,y_2) = (x_2,x_2^2y_2) \text{ we have that } \sigma^*Z \cap E_2 = \{R_1^{E_2},R_2^{E_2},R_3^{E_2}\} \text{ where } R_1^{E_2} = (0,-1), \\ R_2^{E_2} &= (0,1) \text{ and } R_3^{E_2} = (0,-2). \text{ A simple computation shows that} \end{split}$$

$$\Delta_E(R_1^{E_2}) = -\frac{2}{11}; \quad \Delta_E(R_2^{E_2}) = 0; \quad \Delta_E(R_3^{E_2}) = \frac{3}{11}$$

Thus $Col(E_2) = \{R_2^{E_2}\}$ and $NCol(E_2) = \{R_1^{E_2}, R_3^{E_2}\}$. Moreover, we have that

$$\mathcal{M}_{E_2}(z) = -\frac{2}{11}\frac{1}{(z+1)} + \frac{3}{11}\frac{1}{(z+2)} = \frac{z-1}{11(z+2)(z+1)}$$

which implies that $M(E_2) = \{R_2^{E_2}\}.$

Given a non-collinear divisor E and a point $P \in E_{red}$, we define

(4.4)
$$\tau_E(P) = \begin{cases} t_P, & \text{if } P \in M(E), \\ -1, & \text{if } P \in \text{NCol}(E) \\ 0, & \text{otherwise.} \end{cases}$$

Note that $\sum_{P \in E_{\text{red}}} \tau_E(P) = t(E) - \# \operatorname{NCol}(E)$ which is a negative integer (it is the degree of the rational function $\mathcal{M}_E(z)$).

4.1. Collinear and non-collinear infinitely near points

Let us explain now the behaviour of collinear (resp. non-collinear) infinitely near points by blowing-up. Recall that here we denote $b_E = b_E^Z$ for any divisor E in G(Z).

LEMMA 4.9. — Let E and E' be two consecutive divisors in G(Z) with E < E' and $b_{E'} = 1$. We can write $\pi_{E'} = \pi_E \circ \sigma$ where $\sigma : X_{E'} \to (X_E, P)$ is the blow-up with center at a point $P \in E_{\text{red}}$. Let Q be the point $\pi_{E'}^* Z \cap E'_{\text{red}}$.

If P is a collinear point (resp. a non-collinear point) for the foliations \mathcal{F} and \mathcal{G} in E, then Q is a collinear point (resp. non-collinear point) for E'.

Nuria CORRAL

Proof. — Let us denote by \widetilde{E}_{red} the strict transform of E_{red} by σ and $\widetilde{P} = \widetilde{E}_{red} \cap E'_{red}$. Given any singular foliation \mathcal{F} , the Camacho–Sad indices verify the following equalities (see [4])

$$\mathcal{I}_{\tilde{P}}(\pi_{E'}^*\mathcal{F}, E_{\mathrm{red}}) = \mathcal{I}_P(\pi_E^*\mathcal{F}, E_{\mathrm{red}}) - 1$$
$$\mathcal{I}_{\tilde{P}}(\pi_{E'}^*\mathcal{F}, E'_{\mathrm{red}}) + \mathcal{I}_Q(\pi_{E'}^*\mathcal{F}, E'_{\mathrm{red}}) = -1.$$

Moreover, since \mathcal{F} is a generalized curve foliation, then \widetilde{P} is a simple singularity for $\pi_{E'}^* \mathcal{F}$ and hence we have that

$$\mathcal{I}_{\tilde{P}}(\pi_{E'}^*\mathcal{F}, \widetilde{E}_{\mathrm{red}}) \cdot \mathcal{I}_{\tilde{P}}(\pi_{E'}^*\mathcal{F}, E'_{\mathrm{red}}) = 1.$$

Then the index $\mathcal{I}_Q(\pi_{E'}^*\mathcal{F}, E'_{red})$ can be computed as

$$\mathcal{I}_Q(\pi_{E'}^*\mathcal{F}, E_{\mathrm{red}}') = -1 - \frac{1}{\mathcal{I}_P(\pi_E^*\mathcal{F}, E_{\mathrm{red}}) - 1} = -\frac{\mathcal{I}_P(\pi_E^*\mathcal{F}, E_{\mathrm{red}})}{\mathcal{I}_P(\pi_E^*\mathcal{F}, E_{\mathrm{red}}) - 1}$$

Thus, using the expression above for the foliations \mathcal{F} and \mathcal{G} , we have that

$$\Delta_{E'}(Q) = \begin{vmatrix} 1 & \mathcal{I}_Q(\pi_{E'}^*\mathcal{F}, E'_{\text{red}}) \\ 1 & \mathcal{I}_Q(\pi_{E'}^*\mathcal{G}, E'_{\text{red}}) \end{vmatrix}$$
$$= \frac{\Delta_E(P)}{(\mathcal{I}_P(\pi_E^*\mathcal{F}, E_{\text{red}}) - 1)(\mathcal{I}_P(\pi_E^*\mathcal{G}, E_{\text{red}}) - 1)}$$

and then $\Delta_{E'}(Q) = 0$ if and only if $\Delta_E(P) = 0$. This gives the result. \Box

Consider now E and E' two consecutive bifurcation divisors in G(Z), that is, there is a chain of consecutive divisors

$$E_0 = E < E_1 < \dots < E_{k-1} < E_k = E'$$

with $b_{E_l} = 1$ for l = 1, ..., k - 1 and the morphism $\pi_{E'} = \pi_E \circ \sigma$ where $\sigma : X_{E'} \to (X_E, P)$ is a composition of k punctual blow-ups

$$(4.5) \qquad (X_E, P) \stackrel{\sigma_1}{\longleftarrow} (X_{E_1}, P_1) \stackrel{\sigma_2}{\longleftarrow} \cdots \stackrel{\sigma_{k-1}}{\longleftarrow} (X_{E_{k-1}}, P_{k-1}) \stackrel{\sigma_k}{\longleftarrow} X_{E'}.$$

If E and E' are two consecutive bifurcation divisors as above, we say that E' arises from E at P and we denote $E <_P E'$.

As a consequence of Lemma 4.9, we have that if P is a collinear point (resp. a non-collinear point) for \mathcal{F} and \mathcal{G} relative to E, then P_l is a collinear point (resp. non-collinear point) relative to E_l for $l = 1, \ldots, k-1$. Moreover, we have that

COROLLARY 4.10. — Let *E* be the first bifurcation divisor in G(Z). We can write $\pi_E = \sigma_1 \circ \sigma_2 \circ \cdots \circ \sigma_k$ as a composition of *k* punctual blow-ups

$$(\mathbb{C}^2, 0) \xleftarrow{\sigma_1} (X_1, P_1) \xleftarrow{\sigma_2} \cdots \xleftarrow{\sigma_{k-1}} (X_{k-1}, P_{k-1}) \xleftarrow{\sigma_k} X_k = X_E.$$

We denote $E_i = \sigma_i^{-1}(P_{i-1})$ with $P_0 = 0$. Then all the divisors E_i , $1 \leq i \leq k-1$, are collinear.

ANNALES DE L'INSTITUT FOURIER

Proof. — Since $b_{E_l} = 1$ for $l = 1, \ldots, k - 1$, it is enough to prove that P_1 is a collinear point for \mathcal{F} and \mathcal{G} relative to $E_1 = E^1$. But this is a consequence of the fact that $I_{P_1}(\sigma_1^*\mathcal{F}, E_{1, \text{red}}) = I_{P_1}(\sigma_1^*\mathcal{G}, E_{1, \text{red}}) = -1$. Thus the result follows straightforward.

Remark 4.11. — Note that, if E is the first bifurcation divisor, the properties of the Camacho–Sad index imply that

$$\sum_{l=1}^{b_E} \Delta_E(R_l^E) = 0$$

where $\pi_{E}^{*}Z \cap E_{\text{red}} = \{R_{1}^{E}, \dots, R_{b_{E}}^{E}\}.$

The above equality also holds in the following context:

COROLLARY 4.12. — Let E and E' be two consecutive bifurcation divisors in G(Z) such that E' arises from E at P. If P is a collinear point, then

$$\sum_{l=1}^{b_{E'}} \Delta_{E'}(R_l^{E'}) = 0$$

where $\pi_{E'}^* Z \cap E'_{\text{red}} = \{R_1^{E'}, \dots, R_{b_{E'}}^{E'}\}.$

Proof. — As we have explained before we have that $\pi_{E'} = \pi_E \circ \sigma$ where $\sigma : X_{E'} \to (X_E, P)$ is a composition of k punctual blow-ups

$$(X_E, P) \xleftarrow{\sigma_1} (X_{E_1}, P_1) \xleftarrow{\sigma_2} \cdots \xleftarrow{\sigma_{k-1}} (X_{E_{k-1}}, P_{k-1}) \xleftarrow{\sigma_k} X_{E'}.$$

We denote $E_i = \sigma_i^{-1}(P_{i-1})$ with $P_0 = P$ and we have that $b_{E_i} = 1$ for $i = 1, \ldots, k-1$. Since P is a collinear point, then $\mathcal{I}_P(\pi_E^*\mathcal{F}, E_{\text{red}}) = \mathcal{I}_P(\pi_E^*\mathcal{G}, E_{\text{red}})$ and the properties of the Camacho–Sad indices imply that $\mathcal{I}_{P_i}(\pi_{E_i}^*\mathcal{F}, F_{i,\text{red}}) = \mathcal{I}_{P_i}(\pi_{E_i}^*\mathcal{G}, E_{i,\text{red}})$ for $i = 1, \ldots, k-1$. Consequently, we have that

$$\sum_{l=1}^{b_{E'}} \mathcal{I}_{R_l^{E'}}(\pi_{E'}^*\mathcal{F}, E_{\text{red}}') = \sum_{l=1}^{b_{E'}} \mathcal{I}_{R_l^{E'}}(\pi_{E'}^*\mathcal{G}, E_{\text{red}}')$$

which is equivalent to $\sum_{l=1}^{b_{E'}} \Delta_{E'}(R_l^{E'}) = 0.$

4.2. Weighted initial forms and non-collinear divisors

Let us introduce the following notation in order to describe the relationship between the Newton polygon and the infinitely near points of a curve (see Subsection 2.2 and also [10]). From now on we will always assume that we choose coordinates (x, y) such that x = 0 is not tangent to the curve

 $Z = C \cup D$ union of the separatrices of \mathcal{F} and \mathcal{G} . This will imply that the first side of the Newton polygons $\mathcal{N}(\mathcal{F}; x, y)$ and $\mathcal{N}(\mathcal{G}; x, y)$ has slope greater or equal to -1.

Assume that Z is a curve with only non-singular irreducible components and consider $\pi_Z : X_Z \to (\mathbb{C}^2, 0)$ its minimal reduction of singularities. Take any divisor E of $\pi_Z^{-1}(0)$ with v(E) = p and consider $\pi_E : X_E \to (\mathbb{C}^2, 0)$ the morphism reduction of π_Z to E. With the notations introduced in Remark 4.1, if (x, y) are coordinates adapted to E, the points $\pi_E^*Z \cap E_{\text{red}}$ are determined by $\text{In}_p(h; x, y)$ where h = 0 is a reduced equation of the curve Z. More precisely, if we take (x_p, y_p) coordinates in the first chart of E_{red} with $\pi_E(x_p, y_p) = (x_p, x_p^p y_p)$ and $E_{\text{red}} = (x_p = 0)$, thus the points of $\pi_E^*Z \cap E_{\text{red}}$ are given by $x_p = 0$ and $\sum_{i+pj=k} h_{ij} y_p^j = 0$ where $\text{In}_p(h; x, y) = \sum_{i+pj=k} h_{ij} x^i y^j$.

We are interested in determine the points $\pi_E^* \mathcal{J}_{\mathcal{F},\mathcal{G}} \cap E_{\text{red}}$, thus if v(E) = p and (x, y) are coordinates adapted to E, we would like to determine $\ln_p(J; x, y)$ where J(x, y) = 0 is an equation of the jacobian curve. Next result proves that the initial form $\ln_p(J; x, y)$ is determined by the initial forms $\ln_p(\omega), \ln_p(\eta)$ of the 1-forms defining the foliations \mathcal{F} and \mathcal{G} provided that the divisor E is non-collinear.

LEMMA 4.13. — Let *E* be an irreducible component of $\pi_Z^{-1}(0)$ with v(E) = p and take (x, y) coordinates adapted to *E*. If *E* is a non-collinear divisor, then

 $\operatorname{In}_p(\omega) \wedge \operatorname{In}_p(\eta) \not\equiv 0,$

where $\operatorname{In}_p(\omega) = \operatorname{In}_p(\omega; x, y)$ and $\operatorname{In}_p(\eta) = \operatorname{In}_p(\eta; x, y)$, and hence

$$\operatorname{In}_p(J; x, y) = J_p(x, y)$$

with $\operatorname{In}_p(\omega) \wedge \operatorname{In}_p(\eta) = J_p(x, y) \, \mathrm{d}x \wedge \, \mathrm{d}y.$

Proof. — Take an irreducible component E of $\pi_Z^{-1}(0)$ with v(E) = p and let (x, y) be coordinates adapted to E. Assume that $\ln_p(\omega) \wedge \ln_p(\eta) \equiv 0$, that is, if we write

$$In_p(\omega) = A_I(x, y) dx + B_I(x, y) dy$$

$$In_p(\eta) = P_I(x, y) dx + Q_I(x, y) dy$$

then

(4.6)
$$A_I(x,y)Q_I(x,y) - B_I(x,y)P_I(x,y) \equiv 0.$$

Note that $(A_I, B_I) \neq (0, 0)$ and $(P_I, Q_I) \neq (0, 0)$.

Take coordinates (x_p, y_p) in the first chart of E_{red} such that $\pi_E(x_p, y_p) = (x_p, x_p^p y_p), E_{\text{red}} = (x_p = 0)$ and assume that $R_l^E = (0, c_l^E)$, for $l = 1, \ldots, b_E$, in these coordinates where $\pi_E^* Z \cap E_{\text{red}} = \{R_1^E, \ldots, R_{b_E}^E\}$. Let ω^E and η^E be the strict transforms of ω and η by π_E with

$$\begin{split} \omega^E &= A^E(x_p, y_p) \, \mathrm{d} x_p + x_p B^E(x_p, y_p) \, \mathrm{d} y_p, \\ \eta^E &= P^E(x_p, y_p) \, \mathrm{d} x_p + x_p Q^E(x_p, y_p) \, \mathrm{d} y_p. \end{split}$$

From the definition of the Camacho–Sad index we have that

$$\mathcal{I}_{R_l^E}(\pi_E^*\mathcal{F}, E_{\mathrm{red}}) = -\operatorname{Res}_{y=c_l^E} \frac{B^E(0, y)}{A^E(0, y)},$$
$$\mathcal{I}_{R_l^E}(\pi_E^*\mathcal{G}, E_{\mathrm{red}}) = -\operatorname{Res}_{y=c_l^E} \frac{Q^E(0, y)}{P^E(0, y)}.$$

Note that $A^{E}(0, y) = A_{I}(1, y) + pyB_{I}(1, y), B^{E}(0, y) = B_{I}(1, y), P^{E}(0, y) = P_{I}(1, y) + pyQ_{I}(1, y)$ and $Q^{E}(0, y) = Q_{I}(1, y)$. Thus, the equality given in (4.6) implies

$$\frac{B^E(0,y)}{A^E(0,y)} = \frac{Q^E(0,y)}{P^E(0,y)}$$

and hence $\mathcal{I}_{R_l^E}(\pi_E^*\mathcal{F}, E_{\text{red}}) = \mathcal{I}_{R_l^E}(\pi_E^*\mathcal{G}, E_{\text{red}})$ for $l = 1, \ldots, b_E$, in contradiction with the fact that the divisor E is non-collinear.

Observe that the result above is true for the first divisor E^1 although the curves C and D have singular irreducible components, and hence we have that Lemma 2.3 can be obtained as a consequence of the previous result since the conditions over the foliations \mathcal{F} and \mathcal{G} in Lemma 2.3 imply that E^1 is a non-collinear divisor.

Moreover, next result shows that given $\mathcal{F}, \widetilde{\mathcal{F}} \in \mathbb{G}_{C,\lambda}$ and $\mathcal{G}, \widetilde{\mathcal{G}} \in \mathbb{G}_{D,\mu}$, we have that

$$\operatorname{In}_p(J_{\mathcal{F},\mathcal{G}};x,y) = \operatorname{In}_p(J_{\tilde{\mathcal{F}},\tilde{\mathcal{G}}};x,y)$$

provided that E is a non-collinear divisor with v(E) = p, where (x, y) are coordinates adapted to E and $J_{\mathcal{F},\mathcal{G}}(x, y) = 0$ and $J_{\tilde{\mathcal{F}},\tilde{\mathcal{G}}}(x, y) = 0$ are equations of the jacobian curves $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ and $\mathcal{J}_{\tilde{\mathcal{F}},\tilde{\mathcal{G}}}$ respectively. Given a foliation \mathcal{F} , we will denote by $\omega_{\mathcal{F}}$ a 1-form defining \mathcal{F} and $\operatorname{In}_p(\omega_{\mathcal{F}}) = \operatorname{In}_p(\omega_{\mathcal{F}}; x, y)$. Thus we have the following result

LEMMA 4.14. — Let E be an irreducible component of $\pi^{-1}(0)$ with v(E) = p and assume that (x, y) are coordinates adapted to E. Consider foliations $\mathcal{F}, \widetilde{\mathcal{F}} \in \mathbb{G}_{C,\lambda}$ and $\mathcal{G}, \widetilde{\mathcal{G}} \in \mathbb{G}_{D,\mu}$ then

$$\operatorname{In}_p(\omega_{\mathcal{F}}) = \operatorname{In}_p(\omega_{\tilde{\mathcal{F}}}); \qquad \operatorname{In}_p(\omega_{\mathcal{G}}) = \operatorname{In}_p(\omega_{\tilde{\mathcal{G}}}).$$

TOME 75 (2025), FASCICULE 1

Hence, if E is a non-collinear divisor, we have that

$$\operatorname{In}_p(J_{\mathcal{F},\mathcal{G}};x,y) = \operatorname{In}_p(J_{\tilde{\mathcal{F}},\tilde{\mathcal{G}}};x,y)$$

Proof. — Let us prove that $\ln_p(\omega_{\mathcal{F}}) = \ln_p(\omega_{\tilde{\mathcal{F}}})$. We can write

$$\begin{aligned} & \ln_p(\omega_{\mathcal{F}}) = A_I^{\mathcal{F}}(x, y) \, \mathrm{d}x + B_I^{\mathcal{F}}(x, y) \, \mathrm{d}y \\ & \ln_p(\omega_{\tilde{\mathcal{F}}}) = A_I^{\tilde{\mathcal{F}}}(x, y) \, \mathrm{d}x + B_I^{\tilde{\mathcal{F}}}(x, y) \, \mathrm{d}y. \end{aligned}$$

Take (x_p, y_p) coordinates in the first chart of E_{red} such that $\pi_E(x_p, y_p) = (x_p, x_p^p y_p)$ and $E_{\text{red}} = (x_p = 0)$. Let $\omega_{\mathcal{F}}^E$ and $\omega_{\tilde{\mathcal{F}}}^E$ be the strict transforms of $\omega_{\mathcal{F}}$ and $\omega_{\tilde{\mathcal{F}}}$ by π_E with

$$\begin{split} \omega_{\mathcal{F}}^E &= A_{\mathcal{F}}^E(x_p, y_p) \, \mathrm{d}x_p + x_p B_{\mathcal{F}}^E(x_p, y_p) \, \mathrm{d}y_p, \\ \omega_{\mathcal{F}}^E &= A_{\mathcal{F}}^E(x_p, y_p) \, \mathrm{d}x_p + x_p B_{\mathcal{F}}^E(x_p, y_p) \, \mathrm{d}y_p. \end{split}$$

Recall that we have that

$$\begin{split} A^{E}_{\mathcal{F}}(0,y) &= A^{\mathcal{F}}_{I}(1,y) + pyB^{\mathcal{F}}_{I}(1,y); \\ A^{E}_{\tilde{\mathcal{F}}}(0,y) &= A^{\tilde{\mathcal{F}}}_{I}(1,y) + pyB^{\tilde{\mathcal{F}}}_{I}(1,y); \\ \end{split} \qquad \begin{array}{l} B^{E}_{\mathcal{F}}(0,y) &= B^{\mathcal{F}}_{I}(1,y) \\ B^{\tilde{\mathcal{F}}}_{\tilde{\mathcal{F}}}(0,y) &= B^{\tilde{\mathcal{F}}}_{I}(1,y) \end{array}$$

and that the Camacho–Sad indices coincide for \mathcal{F} and $\widetilde{\mathcal{F}}$, that is,

$$\mathcal{I}_{P_l^E}(\pi_E^*\mathcal{F}, E_{\mathrm{red}}) = \mathcal{I}_{P_l^E}(\pi_E^*\widetilde{\mathcal{F}}, E_{\mathrm{red}}), \qquad \text{for } l = 1, 2, \dots, k,$$

where $\pi_E^* C \cap E_{\text{red}} = \{P_1^E, \dots, P_k^E\}$. Since C has only non-singular irreducible components, if we write $P_l^E = (0, d_l^E)$ in coordinates (x_p, y_p) and denote $m_l^C = \nu_{P_l^E}(\pi_E^* C)$, we have that

$$A_{\mathcal{F}}^{E}(0,y) = A_{\tilde{\mathcal{F}}}^{E}(0,y) = \prod_{l=1}^{k} (y - d_{l}^{E})^{m_{l}^{C}}$$

up to divide $\omega_{\mathcal{F}}^{E}$ and $\omega_{\mathcal{F}}^{E}$ by a constant. Moreover, if we consider (x_{l}, y_{l}) coordinates centered at P_{l}^{E} with $x_{l} = x_{p}$ and $y_{l} = y_{p} - d_{l}^{E}$, the equality of the Newton polygons $\mathcal{N}(\pi_{E}^{*}\mathcal{F}; x_{l}, y_{l}), \mathcal{N}(\pi_{E}^{*}\widetilde{\mathcal{F}}; x_{l}, y_{l})$ and $\mathcal{N}(\pi_{E}^{*}C; x_{l}, y_{l})$ implies

$$\operatorname{ord}_{y=d_l^E}(B_{\mathcal{F}}^E(0,y)) \geqslant m_l^C - 1 \qquad \operatorname{ord}_{y=d_l^E}(B_{\tilde{\mathcal{F}}}^E(0,y)) \geqslant m_l^C - 1$$

(see [11, Lemma 1]) and we can write $B_{\mathcal{F}}^E(0, y) = \prod_{l=1}^k (y - d_l^E)^{m_l^C - 1} \widetilde{B}_{\mathcal{F}}^E(y)$, $B_{\tilde{\mathcal{F}}}^E(0, y) = \prod_{l=1}^k (y - d_l^E)^{m_l^C - 1} \widetilde{B}_{\tilde{\mathcal{F}}}^E(y)$. Thus, from the definition of the

ANNALES DE L'INSTITUT FOURIER

Camacho–Sad index given in (2.3), we have that

$$\begin{split} \mathcal{I}_{P_l^E}(\pi_E^*\mathcal{F}, E_{\mathrm{red}}) &= -\operatorname{Res}_{y=d_l^E} \frac{B_{\mathcal{F}}^E(0, y)}{A_{\mathcal{F}}^E(0, y)} = -\operatorname{Res}_{y=d_l^E} \frac{B_{\mathcal{F}}^E(y)}{\prod_{l=1}^k (y - d_l^E)},\\ \mathcal{I}_{P_l^E}(\pi_E^*\widetilde{\mathcal{F}}, E_{\mathrm{red}}) &= -\operatorname{Res}_{y=d_l^E} \frac{B_{\tilde{\mathcal{F}}}^E(0, y)}{A_{\tilde{\mathcal{F}}}^E(0, y)} = -\operatorname{Res}_{y=d_l^E} \frac{\widetilde{B}_{\tilde{\mathcal{F}}}^E(y)}{\prod_{l=1}^k (y - d_l^E)}. \end{split}$$

The equality of the Camacho–Sad indices

$$\mathcal{I}_{P_l^E}(\pi_E^*\mathcal{F}, E_{\mathrm{red}}) = \mathcal{I}_{P_l^E}(\pi_E^*\mathcal{F}, E_{\mathrm{red}}),$$

for l = 1, ..., k, implies $\widetilde{B}_{\mathcal{F}}^E(y) = \widetilde{B}_{\tilde{\mathcal{F}}}^E(y)$ and hence $\operatorname{In}_p(\omega_{\mathcal{F}}) = \operatorname{In}_p(\omega_{\tilde{\mathcal{F}}})$. Finally, if E is a non-collinear divisor, the equality

$$\operatorname{In}_p(J_{\mathcal{F},\mathcal{G}};x,y) = \operatorname{In}_p(J_{\tilde{\mathcal{F}},\tilde{\mathcal{G}}};x,y)$$

is a direct consequence of Lemma 4.13.

Remark 4.15. — Note that x = 0 can be a branch of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ although x = 0 is not tangent to the curve Z. Let us consider the foliations \mathcal{F} and \mathcal{G} given by $\omega = 0$ and $\eta = 0$ with

$$\omega = (xy - 6x^2) dx + (y^2 - 6xy + 10x^2) dy$$
$$\eta = -6x^5 dx + 3y^2 dy.$$

Thus $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ is given by J(x,y) = 0 with

$$J(x,y) = 3x(y^3 - 6xy^2 + 2x^4y^2 - 12x^5y + 20x^6).$$

In this example, if we consider the blow-up $\pi_1 : X_1 \to (\mathbb{C}^2, 0)$ of the origin, the first divisor E^1 is non-collinear. Thus the result of Lemma 4.13 above holds: we have that $\operatorname{In}_1(\omega) = \omega$, $\operatorname{In}_1(\eta) = 3y^2 \, dy$ and hence $\operatorname{In}_1(J) = 3xy^2(y - 6x)$.

Note that the rational function $\mathcal{M}_{E^1}(z)$ is given by

$$\mathcal{M}_{E^1}(z) = -\frac{z-6}{z(z-1)(z-2)(z-3)}$$

which determines the branch $J_{nc}^{E^1}$ whose tangent cone is given by y-6x=0 but we cannot determine the branch x=0 of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ (see statement of Theorem 5.8).

The strategy to prove the results about the jacobian curve is based on the properties that share a foliation and its logarithmic model. Next lemma will allow to follow this strategy.

LEMMA 4.16. — Consider foliations $\mathcal{F}, \mathcal{L}_{\lambda}^{C} \in \mathbb{G}_{C,\lambda}$ and $\mathcal{G}, \mathcal{L}_{\mu}^{D} \in \mathbb{G}_{D,\mu}$. Let $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ be the jacobian curve of \mathcal{F} and \mathcal{G} and $\mathcal{J}_{\lambda,\mu}$ the jacobian curve of $\mathcal{L}_{\lambda}^{C}$ and \mathcal{L}_{μ}^{D} . Let E be an irreducible component of $\pi_{Z}^{-1}(0)$. If E is a non-collinear divisor, we have that

$$\pi_E^* \mathcal{J}_{\mathcal{F},\mathcal{G}} \cap E_{\mathrm{red}} = \pi_E^* \mathcal{J}_{\lambda,\mu} \cap E_{\mathrm{red}}$$

and the multiplicities satisfy that

$$\nu_P(\pi_E^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \nu_P(\pi_E^*\mathcal{J}_{\lambda,\mu})$$

at each point $P \in \pi_E^* \mathcal{J}_{\mathcal{F},\mathcal{G}} \cap E_{\mathrm{red}}$.

Proof. — Let *E* be an irreducible component of $\pi_Z^{-1}(0)$ with v(E) = p and take (x, y) coordinates adapted to *E*. The result is a direct consequence of the equality

$$\operatorname{In}_p(J_{\mathcal{F},\mathcal{G}};x,y) = \operatorname{In}_p(J_{\lambda,\mu};x,y)$$

given in Lemma 4.14 provided that E is a non-collinear divisor, where jacobian curves $\mathcal{J}_{\mathcal{F},\mathcal{G}}$, $\mathcal{J}_{\lambda,\mu}$ are given by $J_{\mathcal{F},\mathcal{G}}(x,y) = 0$ and $J_{\lambda,\mu}(x,y) = 0$ respectively.

5. Properties of the jacobian curve

Let us consider two singular foliations \mathcal{F} and \mathcal{G} in $(\mathbb{C}^2, 0)$ defined by the 1-forms $\omega = 0$ and $\eta = 0$ with $\omega = A(x, y) dx + B(x, y) dy$ and $\eta = P(x, y) dx + Q(x, y) dy$. Recall that the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ is defined by J(x, y) = 0 where

$$J(x,y) = A(x,y)Q(x,y) - B(x,y)P(x,y).$$

Next remark shows that the jacobian curve behaves well by a change of coordinates.

Remark 5.1. — If $F : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ is a change of coordinates with $F = (F_1, F_2)$, the Jacobian curve of $F^*\mathcal{F}$ and $F^*\mathcal{G}$ is given by

$$\begin{vmatrix} A \circ F & B \circ F \\ P \circ F & Q \circ F \end{vmatrix} \cdot \begin{vmatrix} F_{1,x} & F_{1,y} \\ F_{2,x} & F_{2,y} \end{vmatrix} = 0.$$

Thus, the curve $\mathcal{J}_{F^*\mathcal{F},F^*\mathcal{G}}$ is defined by $J \circ F = 0$. Hence, $\mathcal{J}_{F^*\mathcal{F},F^*\mathcal{G}} = F^{-1}(\mathcal{J}_{\mathcal{F},\mathcal{G}})$.

In particular, we get that the analytic type of the jacobian curve of the foliations \mathcal{F} and \mathcal{G} is an invariant of the analytic type of the pair of foliations \mathcal{F} and \mathcal{G} .

Assume that $\mathcal{F} \in \mathbb{G}_{C,\lambda}$ and $\mathcal{G} \in \mathbb{G}_{D,\mu}$ where $C = \bigcup_{i=1}^{r} C_i$ and $D = \bigcup_{i=1}^{s} D_i$ are two plane curves in $(\mathbb{C}^2, 0)$ without common irreducible components. In this section, we will assume that the curve $Z = C \cup D$ has only non-singular irreducible components and we consider $\pi_Z : X_Z \to (\mathbb{C}^2, 0)$ the minimal reduction of singularities of Z. Note that since Z has only non-singular irreducible components, the centers of the blow-ups to obtain π_Z are all free infinitely near points of Z. The general case will be treated in Section 6.

In Section 4 we have introduced all the notions we need to state the results concerning the properties of the jacobian curve. If we want to compute the infinitely near points of the jacobian curve of two foliations, Lemma 4.16 will allow to do computations for the jacobian curve of two logarithmic foliations and then get the result for the jacobian curve of two generalized curve foliations. The first result gives the multiplicity of the jacobian curve at a point in the reduction of singularities in terms of the multiplicities of the curves C and D. In particular, we obtain that all infinitely near points of the jacobian curve in the first chart of a divisor E of $\pi_Z^{-1}(0)$ are either infinitely near points of Z or a point in M(E). Note that x = 0 can be tangent to the jacobian curve although it is not tangent to Z and we cannot control this with the rational function $\mathcal{M}_E(z)$ (see Remark 4.15). More precisely, given a divisor E, we can fix coordinates (x, y) adapted to E (see Remark 4.1) and we can denote by $E_{\rm red}^*$ the points in the first chart of $E_{\rm red}$. Then, we have

THEOREM 5.2. — Let *E* be an irreducible component of $\pi_Z^{-1}(0)$ and assume that *E* is a non-collinear divisor. Given any $P \in E^*_{red}$, we have that

$$\nu_P(\pi_E^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \nu_P(\pi_E^*C) + \nu_P(\pi_E^*D) + \tau_E(P).$$

In particular, if $P \in E_{\text{red}}^*$ with $\nu_P(\pi_E^* \mathcal{J}_{\mathcal{F},\mathcal{G}}) > 0$, then P is an infinitely near point of Z or a point in M(E).

Proof. — We prove here the result for logarithmic foliations. The general case, when \mathcal{F} and \mathcal{G} are not necessarily logarithmic foliations is consequence of Lemma 4.16.

Consider the logarithmic foliations $\mathcal{L}_{\lambda}^{C}$ and \mathcal{L}_{μ}^{D} given by $\omega_{\lambda} = 0$ and $\eta_{\mu} = 0$ with

$$\omega_{\lambda} = \prod_{i=1}^{r} (y - \alpha_i(x)) \sum_{i=1}^{r} \lambda_i \frac{\mathrm{d}(y - \alpha_i(x))}{y - \alpha_i(x)}$$
$$\eta_{\mu} = \prod_{i=1}^{s} (y - \beta_i(x)) \sum_{i=1}^{s} \mu_i \frac{\mathrm{d}(y - \beta_i(x))}{y - \beta_i(x)}$$

TOME 75 (2025), FASCICULE 1

where the curve C_i is given by $y - \alpha_i(x) = 0$ with $\alpha_i(x) = \sum_{j=1}^{\infty} a_j^i x^j \in \mathbb{C}\{x\}$ and the curve D_i is given by $y - \beta_i(x) = 0$ with $\beta_i(x) = \sum_{j=1}^{\infty} b_j^i x^j \in \mathbb{C}\{x\}$. Let us denote $\mathcal{J}_{\lambda,\mu}$ the jacobian curve of \mathcal{L}_{λ}^C and \mathcal{L}_{μ}^D which is defined by $J_{\lambda,\mu}(x,y) = 0$ with

$$J_{\lambda,\mu}(x,y) = A_{\lambda}(x,y)Q_{\mu}(x,y) - B_{\lambda}(x,y)P_{\mu}(x,y)$$

where we write $\omega_{\lambda} = A_{\lambda}(x, y) dx + B_{\lambda}(x, y) dy$ and $\eta_{\mu} = P_{\mu}(x, y) dx + Q_{\mu}(x, y) dy$. More precisely, we can write

(5.1)
$$J_{\lambda,\mu}(x,y) = f(x,y)g(x,y) \begin{vmatrix} -\sum_{i=1}^{r} \lambda_i \frac{\alpha'_i(x)}{y - \alpha_i(x)} & \sum_{i=1}^{r} \frac{\lambda_i}{y - \alpha_i(x)} \\ -\sum_{i=1}^{s} \mu_i \frac{\beta'_i(x)}{y - \beta_i(x)} & \sum_{i=1}^{s} \frac{\mu_i}{y - \beta_i(x)} \end{vmatrix}$$

where $f(x, y) = \prod_{i=1}^{r} (y - \alpha_i(x))$ and $g(x, y) = \prod_{i=1}^{s} (y - \beta_i(x))$ are equations of the curves C and D respectively.

Let $\pi_Z : X_Z \to (\mathbb{C}^2, 0)$ be the minimal reduction of singularities of Z. Take E a bifurcation divisor of G(Z) with v(E) = p and consider $\pi_E : X_E \to (\mathbb{C}^2, 0)$ the reduction of π_Z to E. Since the jacobian curve behaves well by a change of coordinates (see Remark 5.1), we can assume that the coordinates (x, y) are adapted to E. Take (x_p, y_p) coordinates in the first chart of $E_{\text{red}} \subset X_E$ such that $\pi_E(x_p, y_p) = (x_p, x_p^p y_p)$ and $E_{\text{red}} = (x_p = 0)$. Let us compute the strict transform of $\mathcal{J}_{\lambda,\mu}$ by π_E .

Let us denote $I = \{1, \ldots, r\}, J = \{1, \ldots, s\}, I^E = \{i \in I : E \text{ belongs to the geodesic of } C_i\}$ and $J^E = \{j \in J : E \text{ belongs to the geodesic of } D_i\}$. We can write

$$\omega_{\lambda} = f(x,y) \left[\sum_{i \in I \smallsetminus I^{E}} \lambda_{i} \frac{-\alpha_{i}'(x) \, \mathrm{d}x + \mathrm{d}y}{y - \alpha_{i}(x)} + \sum_{i \in I^{E}} \lambda_{i} \frac{-\alpha_{i}'(x) \, \mathrm{d}x + \mathrm{d}y}{y - \alpha_{i}(x)} \right]$$
$$\eta_{\mu} = g(x,y) \left[\sum_{i \in J \smallsetminus J^{E}} \mu_{i} \frac{-\beta_{i}'(x) \, \mathrm{d}x + \mathrm{d}y}{y - \beta_{i}(x)} + \sum_{i \in J^{E}} \mu_{i} \frac{-\beta_{i}'(x) \, \mathrm{d}x + \mathrm{d}y}{y - \beta_{i}(x)} \right]$$

and hence, the jacobian curve $\mathcal{J}_{\lambda,\mu}$ is given by $J_{\lambda,\mu}(x,y) = 0$ with

$$J_{\lambda,\mu}(x,y) = f(x,y)g(x,y)M(x,y)$$

ANNALES DE L'INSTITUT FOURIER

where

$$M(x,y) = \begin{vmatrix} \sum_{i \in I \setminus I^E} \lambda_i \frac{-\alpha'_i(x)}{y - \alpha_i(x)} + \sum_{i \in I^E} \lambda_i \frac{-\alpha'_i(x)}{y - \alpha_i(x)} & \sum_{i \in I \setminus I^E} \frac{\lambda_i}{y - \alpha_i(x)} + \sum_{i \in I^E} \frac{\lambda_i}{y - \alpha_i(x)} \\ \sum_{i \in J \setminus J^E} \mu_i \frac{-\beta'_i(x)}{y - \beta_i(x)} + \sum_{i \in J^E} \mu_i \frac{-\beta'_i(x)}{y - \beta_i(x)} & \sum_{i \in J \setminus J^E} \frac{\mu_i}{y - \beta_i(x)} + \sum_{i \in J^E} \frac{\mu_i}{y - \beta_i(x)} \end{vmatrix}$$

Since v(E) = p and (x, y) are coordinates adapted to E, we have that

$$\operatorname{ord}_{x}(\alpha_{i}(x)) \geq p \quad \text{if} \quad i \in I^{E}; \quad \operatorname{ord}_{x}(\alpha_{i}(x)) = n_{i}$$

Thus, $\alpha_i(x) = \sum_{j \ge p} a_j^i x^j$ if $i \in I^E$ and $\beta_i(x) = \sum_{j \ge p} b_j^i x^j$ if $i \in J^E$, but $\alpha_i(x) = \sum_{j \ge n_i} a_j^i x^j$ with $n_i < p$ if $i \in I \smallsetminus I^E$ and $\beta_i(x) = \sum_{j \ge o_i} b_j^i x^j$ with $o_i < p$ if $i \in J \smallsetminus J^E$. Then, $J_{\lambda,\mu}(x_p, x_p^p y_p)$ is given by

$$J_{\lambda,\mu}(x_p, x_p^p y_p) = f(x_p, x_p^p y_p)g(x_p, x_p^p y_p)M_E^*(x_p, y_p)$$

with

$$M_E^*(x_p, y_p) = \frac{1}{x_p^{p+1}} M_E(x_p, y_p)$$

where

$$M_{E}(x_{p}, y_{p}) = \begin{vmatrix} \sum_{i \in I \smallsetminus I^{E}} \lambda_{i} \frac{-n_{i}a_{n_{i}}^{i} + x_{p}(\cdots)}{x_{p}^{p-n_{i}}y_{p} - a_{n_{i}}^{i} + x_{p}(\cdots)} & \sum_{i \in I \smallsetminus I^{E}} \frac{\lambda_{i}x_{p}^{p-n_{i}}}{x_{p}^{p-n_{i}}y_{p} - a_{n_{i}}^{i} + x_{p}(\cdots)} \\ + \sum_{i \in I^{E}} \lambda_{i} \frac{-pa_{p}^{i} + x_{p}(\cdots)}{y_{p} - a_{p}^{i} + x_{p}(\cdots)} & + \sum_{i \in I^{E}} \frac{\lambda_{i}}{y_{p} - a_{p}^{i} + x_{p}(\cdots)} \\ \sum_{i \in J \smallsetminus J^{E}} \mu_{i} \frac{-o_{i}b_{o_{i}}^{i} + x_{p}(\cdots)}{x_{p}^{p-o_{i}}y_{p} - b_{o_{i}}^{i} + x_{p}(\cdots)} & \sum_{i \in J \smallsetminus J^{E}} \frac{\mu_{i}x_{p}^{p-o_{i}}}{x_{p}^{p-o_{i}}y_{p} - b_{o_{i}}^{i} + x_{p}(\cdots)} \\ + \sum_{i \in J^{E}} \mu_{i} \frac{-pb_{p}^{i} + x_{p}(\cdots)}{y_{p} - b_{p}^{i} + x_{p}(\cdots)} & + \sum_{i \in J^{E}} \frac{\mu_{i}}{y_{p} - b_{p}^{i} + x_{p}(\cdots)} \end{vmatrix}$$

If $M_E(0, y_p) \neq 0$, then the points $\pi_E^* \mathcal{J}_{\lambda,\mu} \cap E_{\text{red}}$, in the first chart of E_{red} , are given by $x_p = 0$ and $J_E(y_p) = 0$ where

$$J_E(y_p) = f(0, y_p)\widetilde{g}(0, y_p)M_E(0, y_p).$$

Let $\{R_1^E, \ldots, R_{b_E}^E\}$ be the union of the singular points of $\pi_E^* \mathcal{L}_{\lambda}^C$ and $\pi_E^* \mathcal{L}_{\mu}^D$ in the first chart of E_{red} , where $R_l^E = (0, c_l^E)$ in coordinates (x_p, y_p) . Note that $\{R_1^E, \ldots, R_{b_E}^E\} = \pi_E^* Z \cap E_{\text{red}}$. Denote $m_l^C = \nu_{R_l^E}(\pi_E^* C) =$ $\sharp \{i \in \{1, \ldots, r\} : \pi_E^* C_i \cap E_{\text{red}} = \{R_l^E\}\}$ and $m_l^D = \nu_{R_l^E}(\pi_E^* D) =$ $\sharp \{j \in \{1, \ldots, s\} : \pi_E^* D_j \cap E_{\text{red}} = \{R_l^E\}\}$ for $l = 1, 2, \ldots, b_E$.

TOME 75 (2025), FASCICULE 1

We have that

$$M_E(0, y_p) = \begin{vmatrix} \sum_{i \in I \smallsetminus I^E} \lambda_i n_i + \sum_{i \in I^E} \lambda_i \frac{-p a_p^i}{y_p - a_p^i} & \sum_{i \in I^E} \frac{\lambda_i}{y_p - a_p^i} \\ \sum_{i \in J \smallsetminus J^E} \mu_i o_i + \sum_{i \in J^E} \mu_i \frac{-p b_p^i}{y_p - b_p^i} & \sum_{i \in J^E} \frac{\mu_i}{y_p - b_p^i} \end{vmatrix}$$

$$= \begin{vmatrix} \sum_{i \in I \smallsetminus I^E} \lambda_i n_i + \sum_{i \in I^E} \lambda_i p & \sum_{i \in I^E} \frac{\lambda_i}{y_p - a_p^i} \\ \sum_{i \in J \smallsetminus J^E} \mu_i o_i + \sum_{i \in J^E} \mu_i p & \sum_{i \in J^E} \frac{\mu_i}{y_p - b_p^i} \end{vmatrix}$$
$$= -\kappa_E(\mathcal{L}^C_\lambda) \kappa_E(\mathcal{L}^D_\mu) \begin{vmatrix} 1 & \sum_{l=1}^{b_E} \frac{\mathcal{I}_{R_l^E}(\pi^*_E \mathcal{L}^C_\lambda, E_{\mathrm{red}})}{y_p - c_l^E} \\ 1 & \sum_{l=1}^{b_E} \frac{\mathcal{I}_{R_l^E}(\pi^*_E \mathcal{L}^D_\mu, E_{\mathrm{red}})}{y_p - c_l^E} \end{vmatrix}$$

where $\kappa_E(\mathcal{L}_{\lambda}^C) = \sum_{i \in I \setminus I^E} \lambda_i n_i + \sum_{i \in I^E} \lambda_i p$ and $\kappa_E(\mathcal{L}_{\mu}^D) = \sum_{i \in J \setminus J^E} \mu_i o_i + \sum_{i \in J^E} \mu_i p$ are the residues of the logarithmic foliations along the divisor E (see (3.2)) and we use the expression of the Camacho–Sad index for a logarithmic foliation given in (3.3). Consequently, we obtain that

$$M_E(0, y_p) = -\kappa_E(\mathcal{L}^C_\lambda)\kappa_E(\mathcal{L}^D_\mu)\mathcal{M}_E(y_p)$$

where $\mathcal{M}_E(z)$ is the rational function associated to the divisor E for the foliations \mathcal{L}^C_{λ} and \mathcal{L}^D_{μ} (see expression (4.1)) Then, the points $\pi^*_E \mathcal{J}_{\lambda,\mu} \cap E_{\text{red}}$, in the first chart of E_{red} , are given by $x_p = 0$ and

$$\prod_{i=1}^{b_E} (y_p - c_l^E)^{m_l^C + m_l^D} \mathcal{M}_E(y_p) = 0.$$

(note that the curve $C \cup D$ has only non-singular irreducible components). Let $\{q_1, \ldots, q_{s_E}\}$ be the set of zeros of $\mathcal{M}_E(z)$. For $l = 1, 2, \ldots, s_E$, put $Q_l^E = (0, q_l)$ and denote by $t_{Q_l^E}$ the multiplicity of q_l as a zero of $\mathcal{M}_E(z)$. Thus, the points in $\pi_E^* \mathcal{J}_{\lambda,\mu} \cap E_{\text{red}}$ belong to $\text{Col}(E) \cup \text{NCol}(E) \cup M(E)$. Moreover, the multiplicity of $\pi_E^* \mathcal{J}_{\lambda,\mu}$ at a point $P \in E_{\text{red}}$, in the first chart of E_{red} , is given by

$$\nu_P(\pi_E^*\mathcal{J}_{\lambda,\mu}) = \nu_P(\pi_E^*C) + \nu_P(\pi_E^*D) + \tau_E(P)$$

where $\tau_E(P)$ was defined by the expression (4.4).

254

Remark 5.3. — With the notations of the proof above, if the first divisor E^1 is non-collinear, we have that the tangent cone of $\mathcal{J}_{\mathcal{L}^C_{\lambda},\mathcal{L}^D_{\mu}}$ is given by $J_1(x,y) = 0$ where

$$J_{1}(x,y) = -\left(\sum_{i=1}^{r} \lambda_{i} a_{1}^{i} \prod_{j \neq i} (y - a_{1}^{j}x)\right) \left(\sum_{i=1}^{s} \mu_{i} \prod_{j \neq i} (y - b_{1}^{j}x)\right) + \sum_{i=1}^{r} \lambda_{i} \prod_{j \neq i} (y - a_{1}^{j}x) \sum_{i=1}^{s} \mu_{i} b_{1}^{i} \prod_{j \neq i} (y - b_{1}^{j}x).$$

Thus, x = 0 is not tangent to the jacobian curve $\mathcal{J}_{\mathcal{L}_{\lambda}^{C},\mathcal{L}_{u}^{D}}$ provided that

(5.2)
$$\kappa_{E^1}(\mathcal{L}^C_{\lambda}) \sum_{i=1}^s \mu_i b_1^i - \kappa_{E^1}(\mathcal{L}^D_{\mu}) \sum_{i=1}^r \lambda_i a_1^i \neq 0$$

where we recall that $\kappa_{E^1}(\mathcal{L}^C_{\lambda}) = \sum_{i=1}^r \lambda_i$ and $\kappa_{E^1}(\mathcal{L}^D_{\mu}) = \sum_{i=1}^s \mu_i$. By Lemma 4.14, the above remarks hold for the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ for any $\mathcal{F} \in \mathbb{G}_{C,\lambda}, \mathcal{G} \in \mathbb{G}_{D,\mu}$. In the example given in Remark 4.15 we have that $\sum_{i=1}^3 \lambda_i a_1^i = \sum_{i=1}^3 \mu_i b_1^i = 0$ and hence the condition in (5.2) does not hold whereas in the example given in Remark 4.8 condition in (5.2) holds and hence x = 0 is not tangent to the jacobian curve.

The reader can find in Appendix A.1 some definitions related to the equisingularity data of curves used in the statements of the following results.

Consider now two consecutive bifurcation divisors E and E' in G(Z) such that E' arises from E at P. As we have explained in Section 4, this means that there is a chain of consecutive divisors

$$E_0 = E < E_1 < \dots < E_{k-1} < E_k = E'$$

with $b_{E_l} = 1$ for l = 1, ..., k - 1 and the morphism $\pi_{E'} = \pi_E \circ \sigma$ where $\sigma: X_{E'} \to (X_E, P)$ is a composition of k punctual blow-ups

(5.3)
$$(X_E, P) \xleftarrow{\sigma_1} (X_{E_1}, P_1) \xleftarrow{\sigma_2} \cdots \xleftarrow{\sigma_{k-1}} (X_{E_{k-1}}, P_{k-1}) \xleftarrow{\sigma_k} X_{E'}.$$

Now we can explain the behaviour of the branches of the jacobian curve going through a non-collinear point. Next corollary states that the branches of the jacobian curve going through a non-collinear point P in a bifurcation divisor as above go through the points P_1, \ldots, P_{k-1} given in the sequence (5.3), that is, the divisor E' is in the geodesic of those branches of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ going through P in E_{red} . COROLLARY 5.4. — Let E and E' be two consecutive bifurcation divisors in G(Z) with $E <_P E'$. If $P \in NCol(E)$, we have that

$$\nu_P(\pi_E^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \sum_{Q \in E'_{\mathrm{red}}} \nu_Q(\pi_{E'}^*\mathcal{J}_{\mathcal{F},\mathcal{G}}).$$

In particular, we get that there is no irreducible component δ of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ such that $\pi_{E'}^*\delta$ is attached to some intermediate component E_i , $1 \leq i \leq k-1$, in the chain $E < E_1 < \cdots < E_{k-1} < E'$. Moreover,

(5.4)
$$1 + \sum_{Q \in M(E')} t_Q = \sharp \operatorname{NCol}(E').$$

Hence E' is non-collinear.

Remark 5.5. — Note that from the above result, we get that there is no irreducible component δ of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ such that

$$v(E) < \mathcal{C}(\delta, \gamma_{E'}) < v(E')$$

where $\gamma_{E'}$ is a E'-curvette.

Proof of Corollary 5.4. — Let E and E' be two consecutive bifurcation divisors in G(Z) with $E <_P E'$ and assume that $P \in \operatorname{NCol}(E)$, thus $\Delta_E(P) \neq 0$. By Theorem 5.2 we have that

$$\nu_P(\pi_E^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \nu_P(\pi_E^*C) + \nu_P(\pi_E^*D) - 1.$$

Recall that $E <_P E'$ implies the existence of a chain of consecutive divisors

$$E_0 = E < E_1 < \dots < E_{k-1} < E_k = E'$$

with $b_{E_l} = 1$ for l = 1, ..., k - 1 and the morphism $\pi_{E'} = \pi_E \circ \sigma$ where $\sigma: X_{E'} \to (X_E, P)$ is a composition of k punctual blow-ups

$$(X_E, P) \xleftarrow{\sigma_1} (X_{E_1}, P_1) \xleftarrow{\sigma_2} \cdots \xleftarrow{\sigma_{k-1}} (X_{E_{k-1}}, P_{k-1}) \xleftarrow{\sigma_k} X_{E'}.$$

Since P is non-collinear, then each P_i is non-collinear by Lemma 4.9, thus $\Delta_{E_i}(P_i) \neq 0$ and hence $M(E_i) = \emptyset$ for $i = 1, \ldots, k-1$. In particular, using again Theorem 5.2, we have that

$$\nu_{P_i}(\pi_{E_i}^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \nu_{P_i}(\pi_{E_i}^*C) + \nu_{P_i}(\pi_{E_i}^*D) - 1.$$

Since the curves C and D have only non-singular irreducible components, and P_i is the only infinitely near point of both curves in E_i , we have that $\nu_{P_i}(\pi_{E_i}^*C) = \nu_P(\pi_E^*C)$ and $\nu_{P_i}(\pi_{E_i}^*D) = \nu_P(\pi_E^*D)$ for all $i = 1, \ldots, k-1$. Consequently,

$$\nu_{P_i}(\pi_{E_i}^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \nu_P(\pi_E^*\mathcal{J}_{\mathcal{F},\mathcal{G}}), \quad \text{for } i = 1, \dots, k-1.$$

ANNALES DE L'INSTITUT FOURIER

256

Since E' is a bifurcation divisor, we get that

$$\nu_{P_{k-1}}(\pi^*_{E_{k-1}}\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \sum_{Q \in E'_{\mathrm{red}}} \nu_Q(\pi^*_{E'}\mathcal{J}_{\mathcal{F},\mathcal{G}}).$$

Hence, from all the equalities above, we deduce that

$$\nu_P(\pi_E^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \sum_{Q \in E'_{\mathrm{red}}} \nu_Q(\pi_{E'}^*\mathcal{J}_{\mathcal{F},\mathcal{G}}),$$

which proves the first statement of the corollary. Finally, in order to prove the equality given in (5.4), it is enough to show that

$$\sum_{R \in \operatorname{NCol}(E')} \Delta_{E'}(R) \neq 0$$

by Lemma 4.6. Let us assume that $\sum_{R \in \text{NCol}(E')} \Delta_{E'}(R) = 0$, which implies

$$\sum_{R \in \mathrm{NCol}(E')} \mathcal{I}_R(\pi_{E'}^* \mathcal{F}, E'_{\mathrm{red}}) = \sum_{R \in \mathrm{NCol}(E')} \mathcal{I}_R(\pi_{E'}^* \mathcal{G}, E'_{\mathrm{red}})$$

and, by the properties of the Camacho–Sad indices, we deduce that

$$\mathcal{I}_{\tilde{P}_{k-1}}(\pi_{E'}^*\mathcal{F}, E'_{\mathrm{red}}) = \mathcal{I}_{\tilde{P}_{k-1}}(\pi_{E'}^*\mathcal{G}, E'_{\mathrm{red}})$$

where we denote $\widetilde{P}_{k-1} = \widetilde{E}_{k-1, \text{red}} \cap E'_{\text{red}}$ and $\widetilde{E}_{k-1, \text{red}}$ is the strict transform of $E_{k-1, \text{red}}$ by σ_k . Since \mathcal{F} and \mathcal{G} are generalized curve foliations, then \widetilde{P}_{k-1} is a simple singularity for $\pi^*_{E'}\mathcal{F}$ and $\pi^*_{E'}\mathcal{G}$ and hence we have that

$$\begin{split} \mathcal{I}_{\tilde{P}_{k-1}}(\pi_{E'}^*\mathcal{F},E_{\mathrm{red}}')\cdot\mathcal{I}_{\tilde{P}_{k-1}}(\pi_{E'}^*\mathcal{F},E_{k-1,\mathrm{red}}) &= 1\\ \mathcal{I}_{\tilde{P}_{k-1}}(\pi_{E'}^*\mathcal{G},E_{\mathrm{red}}')\cdot\mathcal{I}_{\tilde{P}_{k-1}}(\pi_{E'}^*\mathcal{G},\widetilde{E}_{k-1,\mathrm{red}}) &= 1. \end{split}$$

Consequently, given that

$$\begin{split} \mathcal{I}_{P_{k-1}}(\pi^*_{E_{k-1}}\mathcal{F}, E_{k-1, \text{red}}) &= \mathcal{I}_{\tilde{P}_{k-1}}(\pi^*_{E'}\mathcal{F}, \tilde{E}_{k-1, \text{red}}) + 1\\ &= \frac{1}{\mathcal{I}_{\tilde{P}_{k-1}}(\pi^*_{E'}\mathcal{F}, E'_{\text{red}})} + 1,\\ \mathcal{I}_{P_{k-1}}(\pi^*_{E_{k-1}}\mathcal{G}, E_{k-1, \text{red}}) &= \mathcal{I}_{\tilde{P}_{k-1}}(\pi^*_{E'}\mathcal{G}, \tilde{E}_{k-1, \text{red}}) + 1\\ &= \frac{1}{\mathcal{I}_{\tilde{P}_{k-1}}(\pi^*_{E'}\mathcal{G}, E'_{\text{red}})} + 1, \end{split}$$

we obtain that $\mathcal{I}_{P_{k-1}}(\pi^*_{F_{k-1}}\mathcal{F}, E_{k-1, \text{red}}) = \mathcal{I}_{P_{k-1}}(\pi^*_{F_{k-1}}\mathcal{G}, E_{k-1, \text{red}})$. This implies that $\Delta_{E_{k-1}}(P_{k-1}) = 0$ which is not possible since P_{k-1} is a non-collinear point by Lemma 4.9. This ends the proof.

In order to explain the behaviour of the branches of the jacobian curve going through a collinear point, we introduce the following definition. DEFINITION 5.6. — Let E be a bifurcation divisor of G(Z) and take P a collinear point of E. We say that a set of non-collinear bifurcation divisors $\{E_1, \ldots, E_u\}$ is a (non-collinear) cover of E at P if the following conditions hold:

- (i) E is in the geodesic of each E_l ;
- (ii) if $\{E_1^l, \ldots, E_{r(l)}^l\}$ is the set of all bifurcation divisors in the geodesic of E_l with

$$E <_P E_1^l < \dots < E_{r(l)}^l < E_l$$

then either r(l) = 0 or else each E_i^l is collinear;

(iii) if Z_j is an irreducible component of Z with $\pi_E^* Z_j \cap E_{\text{red}} = \{P\}$, then there exists a divisor E_l in the cover such that $\pi_{E_l}^* Z_j \cap E_l \neq \emptyset$, that is, there is a divisor E_l in the cover which is in the geodesic of Z_j .

Given a collinear point P of E, there is a unique cover of E at P. We can find it as follows: take an irreducible component Z_j of Z with $\pi_E^* Z_j \cap E_{\text{red}} =$ $\{P\}$. Let E' be the consecutive bifurcation divisor to E with $E <_P E'$ belonging to the geodesic of Z_j . If E' is non-collinear, then E' is one of the bifurcation divisors in the cover of E at P, otherwise we repeat the process above with the following bifurcation divisor in the geodesic of Z_j . Since the maximal bifurcation divisors are non-collinear (see Remark 4.3), we will always find a non-collinear divisor in the geodesic of Z_j verifying condition (iii) in the above definition.

THEOREM 5.7. — Consider a non-collinear bifurcation divisor E of G(Z) and a collinear point P of E. Take a cover $\{E_1, \ldots, E_u\}$ of E at P. Then

$$\nu_P(\pi_E^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) - \sum_{l=1}^u \sum_{Q \in E_{l,\mathrm{red}}} \nu_Q(\pi_{E_l}^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = t_P + \sum_{l=1}^u (\sharp \operatorname{NCol}(E_l) - t(E_l)).$$

Consequently, there is a curve J_P^E composed by irreducible components of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ such that, if δ is a branch of J_P^E ,

• $\pi_E^* \delta \cap E_{\text{red}} = \{P\},\$

• $\mathcal{C}(\delta, \gamma_{E_l}) < v(E_l)$ for l = 1, ..., u, where γ_{E_l} is any E_l -curvette.

Moreover, we have that

$$\nu_0(J_P^E) = t_P + \sum_{l=1}^u (\sharp \operatorname{NCol}(E_l) - t(E_l)).$$

ANNALES DE L'INSTITUT FOURIER

Proof. — Let E be a non-collinear bifurcation divisor of G(Z) and a point $P \in \text{Col}(E)$. Consider a cover $\{E_1, \ldots, E_u\}$ of E at P. By Theorem 5.2, we have that

$$\nu_P(\pi_E^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \nu_P(\pi_E^*C) + \nu_P(\pi_E^*D) + t_P$$
$$\sum_{l=1}^u \sum_{Q \in E_{l,\mathrm{red}}} \nu_Q(\pi_{E_l}^*\mathcal{J}_{\mathcal{F},\mathcal{G}}) = \sum_{l=1}^u \sum_{Q \in E_{l,\mathrm{red}}} (\nu_Q(\pi_{E_l}^*C) + \nu_Q(\pi_{E_l}^*D) + \tau_{E_l}(Q)).$$

By the properties of a cover given in Definition 5.6, we have that

$$\nu_P(\pi_E^*C) + \nu_P(\pi_E^*D) = \sum_{l=1}^u \sum_{Q \in E_{l, \text{red}}} (\nu_Q(\pi_{E_l}^*C) + \nu_Q(\pi_{E_l}^*D))$$

and the result is straightforward.

The results above allow to give a decomposition of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ into bunches of branches in the sense of the decomposition theorem of polar curves. Recall that given a divisor E of $\pi_Z^{-1}(0)$, we denote by $\pi_E : X_E \to (\mathbb{C}^2, 0)$ the morphism reduction of π_Z to E and we write $\pi_Z = \pi_E \circ \pi'_E$. Let B(Z)be the set of bifurcation divisors of G(Z). Given any $E \in B(Z)$ which is a non-collinear divisor for \mathcal{F} and \mathcal{G} , we define J_{nc}^E as the union of the branches ξ of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ such that

• $\pi_E^* \xi \cap \pi_E^* Z = \emptyset$,

• if
$$E' < E$$
, then $\pi_E^* \xi \cap \pi'_E(E') = \emptyset$,

• if E' > E, then $\pi_{E'}^* \xi \cap \overline{E'_{red}} = \emptyset$.

Moreover, given a non-collinear divisor E, we denote $J_c^E = \bigcup_{P \in \operatorname{Col}(E)} J_P^E$ (with $J_c^E = \emptyset$ if $\operatorname{Col}(E) = \emptyset$).

Thus, the previous results allow us to give a decomposition of

$$\mathcal{J}_{\mathcal{F},\mathcal{G}} = J^* \cup \left(\bigcup_{E \in B_N(Z)} J^E\right)$$

(see below for the precise statement) such there is a certain control of the topology of the irreducible components of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ obtained from the data of the foliations \mathcal{F} and \mathcal{G} provided that the component of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ is attached either to a non-collinear divisor or to a chain of collinear divisors which are between two non-collinear bifurcation divisors. The irreducible components corresponding to J^* are the one attached to "isolated" collinear divisors for which no control is possible.

Given a non-collinear bifurcation divisor E of G(Z), we denote

$$t^*(E) = \sum_{Q \in M(E) \smallsetminus \operatorname{Col}(E)} t_Q.$$

TOME 75 (2025), FASCICULE 1

 \square

that is, the number of zeros of $\mathcal{M}_E(z)$ (counting with multiplicities) which do not correspond to collinear points. Then, we can state the properties of the decomposition of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ as follows:

THEOREM 5.8. — Consider $\mathcal{F} \in \mathbb{G}_C$ and $\mathcal{G} \in \mathbb{G}_D$ such that $Z = C \cup D$ is a curve with only non-singular irreducible components. Let $B_N(Z)$ be the set of non-collinear bifurcation divisors of G(Z). Then there is a unique decomposition $\mathcal{J}_{\mathcal{F},\mathcal{G}} = J^* \cup (\bigcup_{E \in B_{\mathcal{N}}(Z)} J^E)$ where $J^E = J^E_{nc} \cup J^E_c$ with the following properties

- (i) $\nu_0(J_{nc}^E) = t^*(E)$. In particular, $\nu_0(J_{nc}^E) \leq \sharp \operatorname{NCol}(E) 1 \leq b_E 1$.
- (ii) $\pi_E^* J_{nc}^E \cap \pi_E^* Z = \emptyset.$
- (iii) if E' < E, then $\pi_E^* J_{nc}^E \cap \pi'_E(E') = \emptyset$.

- (iv) if E' > E, then $\pi_{E'}^* J_{nc}^E \cap E'_{red} = \emptyset$. (v) if δ is a branch of J_c^E , then $\pi_E^* \delta \cap E_{red}$ is a point in $\operatorname{Col}(E)$. (vi) $\nu_0(J_c^E) = \sum_{P \in C(E)} (t_P + \sum_{l=1}^{u(P)} (\sharp \operatorname{NCol}(E_l^P) t(E_l^P)))$ where $\{E_1^P, \ldots, E_{u(P)}^P\}$ is a cover of E at P.

Moreover, if E is a purely non-collinear divisor with

$$\sum_{R_l^E \in \mathrm{NCol}(E)} \Delta_E(R_l^E) \neq 0,$$

then

(5.5)
$$\nu_0(J^E) = \nu_0(J^E_{nc}) = b_E - 1.$$

Proof. — We have that

$$\nu_0(J_{nc}^E) = \sum_{P \in M(E) \smallsetminus \operatorname{Col}(E)} \nu_P(\pi_E^* \mathcal{J}_{\mathcal{F},\mathcal{G}}) = \sum_{P \in M(E) \smallsetminus \operatorname{Col}(E)} t_P = t^*(E)$$
$$\leqslant \sum_{P \in M(E)} t_P \leqslant \sharp \operatorname{NCol}(E) - 1 \leqslant b_E - 1$$

where we have used the inequality given in (4.2) and the fact that $\sharp \operatorname{NCol}(E) \leq b_E$. This gives the first statement of the theorem.

Moreover, if E is a purely non-collinear divisor, then $\operatorname{Col}(E) = \emptyset$, $J^E =$ J_{nc}^E and $\sharp \operatorname{NCol}(E) = b_E$. In addition, when $\sum_{R^E \in \operatorname{NCol}(E)} \Delta_E(R_l^E) \neq 0$ we have that $\sum_{P \in M(E)} t_P = \# \operatorname{NCol}(E) - 1$ by Lemma 4.6. Consequently, we deduce that

$$\nu_0(J^E) = \sum_{P \in M(E)} \nu_P(\pi_E^* \mathcal{J}_{\mathcal{F},\mathcal{G}}) = \sum_{P \in M(E)} t_P = \sharp \operatorname{NCol}(E) - 1 = b_E - 1$$

and we obtain expression (5.5).

ANNALES DE L'INSTITUT FOURIER

Properties (ii), (iii) and (iv) are consequence of the definition of J_{nc}^E . Properties (v) and (vi) follow directly from the definition of J_c^E and Theorem 5.7.

Note that the properties of J_{nc}^E can be stated in terms of coincidences as follows: if δ is an irreducible component of J_{nc}^E (with E a non-collinear bifurcation divisor) and Z_i is an irreducible component of $Z = C \cup D$, then

$$\mathcal{C}(\delta, Z_i) = \begin{cases} v(E), & \text{if } E \text{ is in the geodesic of } Z_i, \\ \mathcal{C}(\gamma_E, Z_i), & \text{otherwise.} \end{cases}$$

where γ_E is any *E*-curvette. Observe that $v(E) = \mathcal{C}(\gamma_E, Z_i)$ when *E* is in the geodesic of Z_i and γ_E does not intersect *E* at the points $\pi_E^* Z \cap E$.

Next result determines the intersection multiplicity of J_{nc}^E with the curves of separatrices C and D of the foliations \mathcal{F} and \mathcal{G} .

COROLLARY 5.9. — If E is a non-collinear bifurcation divisor, then

$$(J_{nc}^{E}, C)_{0} = \nu_{E}(C) \cdot t^{*}(E); \qquad (J_{nc}^{E}, D)_{0} = \nu_{E}(D) \cdot t^{*}(E)$$

where $\nu_E(C) = (C, \gamma_E)_0$ and $\nu_E(D) = (D, \gamma_E)_0$ with γ_E any *E*-curvette.

Proof. — Let E be a non-collinear bifurcation divisor of G(Z) and let γ_E be any E-curvette which does not intersect E at the points $\pi_E^* Z \cap E$. By the properties of J_{nc}^E given in Theorem 5.8, we have that if δ is a branch of J_{nc}^E then

$$\mathcal{C}(\delta, C_i) = \begin{cases} v(E), & \text{if } E \text{ is in the geodesic of } C_i, \\ \mathcal{C}(\gamma_E, C_i), & \text{othewise.} \end{cases}$$

Note that $C(\gamma_E, C_i) = v(E)$ if E is in the geodesic of C_i (that is, $i \in I_E$). Moreover, since γ_E and C_i are non-singular curves, we have that $C(\gamma_E, C_i) = (\gamma_E, C_i)_0$. Therefore, using the relationship between the coincidence and the intersection multiplicity of two branches given in Remark A.1, we have that

$$(\delta, C_i)_0 = \nu_0(\delta) \cdot (\gamma_E, C_i)_0$$

TOME 75 (2025), FASCICULE 1

for a branch δ of J_{nc}^E . Now, if we denote by $\mathcal{B}(J_{nc}^E)$ the set of branches of J_{nc}^E , we have that

$$(J_{nc}^{E}, C)_{0} = \sum_{i=1}^{r} (J_{nc}^{E}, C_{i})_{0} = \sum_{i=1}^{r} \sum_{\delta \in \mathcal{B}(J_{nc}^{E})} (\delta, C_{i})_{0}$$
$$= \sum_{i=1}^{r} \sum_{\delta \in \mathcal{B}(J_{nc}^{E})} \nu_{0}(\delta) \cdot (\gamma_{E}, C_{i})_{0} = \nu_{0}(J_{nc}^{E}) \sum_{i=1}^{r} (\gamma_{E}, C_{i})_{0}$$
$$= t^{*}(E) \cdot \nu_{E}(C).$$

As a consequence of the result above and Propositions B.1 and B.3 we obtain next corollary for non-dicritical generalized curve foliations which relates invariants of the foliations \mathcal{F} and \mathcal{G} , such as the Milnor numbers o the tangency orders, with data coming from the decomposition of the jacobian curve.

COROLLARY 5.10. — With the hypothesis and notations of Theorem 5.8, we get that

$$\sum_{E \in B_N(Z)} \nu_0(J_{nc}^E) \nu_E(C_i) \leq \mu_0(\mathcal{F}, C_i) + \tau_0(\mathcal{G}, C_i)$$

and

$$\sum_{E \in B_N(Z)} \nu_0(J_{nc}^E)(\nu_E(C) - \nu_E(D)) \leqslant \mu_0(\mathcal{F}) - \mu_0(\mathcal{G})$$

Proof. — We have just proved that $(J_{nc}^E, C_i)_0 = \nu_0(J_{nc}^E)\nu_E(C_i)$. Thus, using Proposition B.1, we get that

$$\sum_{E \in B_N(Z)} (J_{nc}^E, C_i)_0 = \sum_{E \in B_N(Z)} \nu_0(J_{nc}^E) \nu_E(C_i) \leqslant (\mathcal{J}_{\mathcal{F},\mathcal{G}}, C_i)_0$$
$$= \mu_0(\mathcal{F}, C_i) + \tau_0(\mathcal{G}, C_i).$$

Now, from Corollary 5.9 and Proposition B.3, we obtain that

$$\sum_{E \in B_N(Z)} \left((J_{nc}^E, C)_0 - (J_{nc}^E, D)_0) \right) = \sum_{E \in B_N(Z)} \nu_0(J_{nc}^E) (\nu_E(C) - \nu_E(D))$$
$$\leqslant (\mathcal{J}_{\mathcal{F},\mathcal{G}}, C)_0 - (\mathcal{J}_{\mathcal{F},\mathcal{G}}, D)_0$$
$$= \mu_0(\mathcal{F}) - \mu_0(\mathcal{G})$$

which gives the second inequality.

The general case of foliations with separatrices that can have singular irreducible components will be treated in next section.

6. General case

Consider two plane curves $C = \bigcup_{i=1}^{r} C_i$ and $D = \bigcup_{j=1}^{s} D_j$ which can have singular branches. Assume that C and D have no common irreducible components. Let $\rho : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ be a ramification given in coordinates by $\rho(u, v) = (u^n, v)$ such that the curve $\rho^{-1}Z$ has only non-singular irreducible components where $Z = C \cup D$. In this section we will denote \tilde{B} the curve $\rho^{-1}B$ for any plane curve B. See Appendix A for notations concerning ramifications.

Take \mathcal{F} and \mathcal{G} foliations with C and D as curve of separatrices respectively. Let us study the relationship between the curves $\widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}} = \rho^{-1}\mathcal{J}_{\mathcal{F},\mathcal{G}}$ and $\mathcal{J}_{\rho^*\mathcal{F},\rho^*\mathcal{G}}$.

Assume that the foliations \mathcal{F} and \mathcal{G} are given by $\omega = 0$ and $\eta = 0$ with

$$\omega = A(x, y) \,\mathrm{d}x + B(x, y) \,\mathrm{d}y; \quad \eta = P(x, y) \,\mathrm{d}x + Q(x, y) \,\mathrm{d}y,$$

then $\rho^* \mathcal{F}$ and $\rho^* \mathcal{G}$ are given by $\rho^* \omega = 0$ and $\rho^* \eta = 0$ where

$$\begin{split} \rho^* \omega &= A(u^n,v) n u^{n-1} \, \mathrm{d} u + B(u^n,v) \, \mathrm{d} v; \\ \rho^* \eta &= P(u^n,v) n u^{n-1} \, \mathrm{d} u + Q(u^n,v) \, \mathrm{d} v. \end{split}$$

Therefore, if we write J(x, y) = A(x, y)Q(x, y) - B(x, y)P(x, y), then the curve $\rho^{-1}\mathcal{J}_{\mathcal{F},\mathcal{G}}$ is given by $J(u^n, v) = 0$ whereas $\mathcal{J}_{\rho^*\mathcal{F},\rho^*\mathcal{G}}$ is given by $nu^{n-1}J(u^n, v) = 0$. Let us see (Corollary 6.2) that $\rho^{-1}\mathcal{J}_{\mathcal{F},\mathcal{G}} = \widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}}$ satisfies the statements of Theorem 5.2 with respect to $\rho^{-1}Z = \widetilde{Z}$.

Let $\pi_{\tilde{Z}} : \tilde{X} \to (\mathbb{C}^2, 0)$ be the minimal reduction of singularities of \tilde{Z} . We denote by \tilde{E} any irreducible component of $\pi_{\tilde{Z}}^{-1}(0)$ and by $\pi_{\tilde{E}} : \tilde{X}_{\tilde{E}} \to (\mathbb{C}^2, 0)$ the morphism reduction of $\pi_{\tilde{Z}}$ to \tilde{E} . Let us state some properties concerning the infinitely near points of $\tilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}}$ and $\mathcal{J}_{\rho^*\mathcal{F},\rho^*\mathcal{G}}$:

LEMMA 6.1. — Let \tilde{E} be an irreducible component of $\pi_{\tilde{Z}}^{-1}(0)$. We have that

$$\pi_{\tilde{E}}^* \widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}} \cap \widetilde{E}_{\mathrm{red}}^* = \pi_{\tilde{E}}^* \mathcal{J}_{\rho^* \mathcal{F},\rho^* \mathcal{G}} \cap \widetilde{E}_{\mathrm{red}}^*$$

where \widetilde{E}_{red}^* denote the points in the first chart of \widetilde{E}_{red} . Moreover,

$$\nu_P(\pi_{\tilde{E}}^*\widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}}) = \nu_P(\pi_{\tilde{E}}^*\mathcal{J}_{\rho^*\mathcal{F},\rho^*\mathcal{G}})$$

for each $P \in \pi_{\tilde{E}}^* \widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}} \cap \widetilde{E}_{\mathrm{red}}^*$.

Proof. — Take \widetilde{E} an irreducible component of $\pi_{\widetilde{Z}}^{-1}(0)$ with $v(\widetilde{E}) = p$ and assume that (u, v) are coordinates adapted to \widetilde{E} . If we denote $\widetilde{J}(u, v) = J(u^n, v)$, we have that

$$\operatorname{In}_p(nu^{n-1}\widetilde{J}; u, v) = nu^{n-1}\operatorname{In}_p(\widetilde{J}; u, v).$$

Write

$$\ln_p(\widetilde{J}; u, v) = \sum_{i+pj=k} h_{ij} u^i v^j.$$

Hence, if (u_p, v_p) are coordinates in the first chart of $\widetilde{E}_{red} \subset \widetilde{X}_{\tilde{E}}$ such that $\pi_{\tilde{E}}(u_p, v_p) = (u_p, u_p^p v_p)$ and $\widetilde{E}_{red} = (u_p = 0)$, then the points $\pi_{\tilde{E}}^* \widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}} \cap \widetilde{E}_{red}$, in the first chart of \widetilde{E}_{red} , are given by $u_p = 0$ and $\sum_{i+pj=k} h_{ij} v_p^j = 0$. This proves that

$$\pi_{\tilde{E}}^* \widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}} \cap \widetilde{E}_{\mathrm{red}}^* = \pi_{\tilde{E}}^* \mathcal{J}_{\rho^* \mathcal{F},\rho^* \mathcal{G}} \cap \widetilde{E}_{\mathrm{red}}^*$$

and that

$$\nu_P\left(\pi_{\tilde{E}}^*\widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}}\right) = \nu_P\left(\pi_{\tilde{E}}^*\mathcal{J}_{\rho^*\mathcal{F},\rho^*\mathcal{G}}\right)$$

 \square

for each $P \in \pi_{\tilde{E}}^* \widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}} \cap \widetilde{E}_{\mathrm{red}}^*$.

Hence, when \widetilde{E} is a non-collinear divisor for the foliations $\rho^* \mathcal{F}$ and $\rho^* \mathcal{G}$, the curve $\mathcal{J}_{\rho^* \mathcal{F}, \rho^* \mathcal{G}}$ satisfies Theorem 5.2 with respect to $\rho^{-1} Z = \widetilde{Z}$, and thanks to the previous lemma, we get the following result for $\rho^{-1} \mathcal{J}_{\mathcal{F}, \mathcal{G}} = \widetilde{\mathcal{J}}_{\mathcal{F}, \mathcal{G}}$:

COROLLARY 6.2. — Take \widetilde{E} an irreducible component of $\pi_{\widetilde{Z}}^{-1}(0)$ which is a non-collinear divisor for the foliations $\rho^* \mathcal{F}$ and $\rho^* \mathcal{G}$. Given any $P \in \widetilde{E}^*_{red}$, we have that

$$\nu_P\left(\pi_{\tilde{E}}^* \widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}}\right) = \nu_P\left(\pi_{\tilde{E}}^* \widetilde{C}\right) + \nu_P\left(\pi_{\tilde{E}}^* \widetilde{D}\right) + \tau_{\tilde{E}}(P).$$

In particular, if $P \in \widetilde{E}_{red}^*$ with $\nu_P(\pi_{\widetilde{E}}^* \widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}}) > 0$, then P is an infinitely near point of \widetilde{Z} or a point in $M(\widetilde{E})$.

Let E be a bifurcation divisor of G(Z) and consider $\widetilde{E}_l, \widetilde{E}_k$ two bifurcation divisors of $G(\widetilde{Z})$ associated to E. Recall that there is a bijection between the sets of points $\pi_{\widetilde{E}^l}^* \widetilde{Z} \cap \widetilde{E}_{\mathrm{red}}^l$ and $\pi_{\widetilde{E}^k}^* \widetilde{Z} \cap \widetilde{E}_{\mathrm{red}}^k$ given by the map $\rho_{l,k} : \widetilde{E}_{\mathrm{red}}^l \to \widetilde{E}_{\mathrm{red}}^k$ (see Appendix A). Thus we will denote by $\{R_1^{\widetilde{E}^l}, R_2^{\widetilde{E}^l}, \dots, R_{b_{\widetilde{E}^l}}^{\widetilde{E}^l}\}$ and $\{R_1^{\widetilde{E}^k}, R_2^{\widetilde{E}^k}, \dots, R_{b_{\widetilde{E}^k}}^{\widetilde{E}^k}\}$ the sets of points $\pi_{\widetilde{E}^l}^* \widetilde{Z} \cap \widetilde{E}_{\mathrm{red}}^l$ and $\pi_{\widetilde{E}^k}^* \widetilde{Z} \cap \widetilde{E}_{\mathrm{red}}^k$ respectively, with $R_t^{\widetilde{E}^k} = \rho_{l,k}(R_t^{\widetilde{E}^l})$ for $t = 1, 2, \dots, b_{\widetilde{E}^k}$. By the results in Appendix A.3 (see Proposition A.4 and (A.7)), we get that

$$\begin{split} \mathcal{I}_{R_{t}^{\tilde{E}^{l}}}\left(\pi_{\tilde{E}^{l}}^{*}\rho^{*}\mathcal{F}, \widetilde{E}_{\mathrm{red}}^{l}\right) &= \mathcal{I}_{R_{t}^{\tilde{E}^{k}}}\left(\pi_{\tilde{E}^{k}}^{*}\rho^{*}\mathcal{F}, \widetilde{E}_{\mathrm{red}}^{k}\right) \\ \mathcal{I}_{R_{t}^{\tilde{E}^{l}}}\left(\pi_{\tilde{E}^{l}}^{*}\rho^{*}\mathcal{G}, \widetilde{E}_{\mathrm{red}}^{l}\right) &= \mathcal{I}_{R_{t}^{\tilde{E}^{k}}}\left(\pi_{\tilde{E}^{k}}^{*}\rho^{*}\mathcal{G}, \widetilde{E}_{\mathrm{red}}^{k}\right) \end{split}$$

ANNALES DE L'INSTITUT FOURIER

264

which implies

$$\Delta_{\tilde{E}^l}\left(R_t^{\tilde{E}^l}\right) = \Delta_{\tilde{E}^k}\left(R_t^{\tilde{E}^k}\right) \quad \text{for} \quad t = 1, 2, \dots, b_{\tilde{E}^l},$$

with $\Delta_{\tilde{E}^l}(R_t^{\tilde{E}^l}) = \Delta_{\tilde{E}^l}^{\rho^* \mathcal{F}, \rho^* \mathcal{G}}(R_t^{\tilde{E}^l})$. Thus, \tilde{E}^l is collinear (resp. non-collinear) if and only if \tilde{E}^k is also collinear (resp. non-collinear). So we can introduce the following definition

DEFINITION 6.3. — We say that a bifurcation divisor E of G(Z) is collinear (resp. non-collinear) for the foliations \mathcal{F} and \mathcal{G} when any of its associated divisors \tilde{E}^l is collinear (resp. non-collinear) for the foliations $\rho^* \mathcal{F}$ and $\rho^* \mathcal{G}$.

Moreover, if $R_t^{\tilde{E}^l}, R_s^{\tilde{E}^l}$ are two points $\pi_{\tilde{E}^l}^* \widetilde{Z} \cap \widetilde{E}_{red}^l$ with $\rho_{\tilde{E}^l, E}(R_t^{\tilde{E}^l}) = \rho_{\tilde{E}^l, E}(R_s^{\tilde{E}^l})$ where $\rho_{\tilde{E}^l, E}: \widetilde{E}_{red}^l \to E_{red}$ is the ramification defined in appendix A, then

$$\Delta_{\tilde{E}^{l}}\left(R_{t}^{\tilde{E}^{l}}\right) = \Delta_{\tilde{E}^{k}}\left(R_{s}^{\tilde{E}^{l}}\right)$$

by (A.8) in Appendix A.3. Thus, we say that an infinitely near point R^E of Z in $E_{\rm red}$ is a collinear point (resp. non-collinear point) for the foliations \mathcal{F} and \mathcal{G} if, for any associated divisor \tilde{E}^l and any infinitely near point $R_t^{\tilde{E}^l}$ of $\rho^{-1}Z$ in $\tilde{E}_{\rm red}^l$ with $\rho_{\tilde{E}^l,E}(R_t^{\tilde{E}^l}) = R^E$, the point $R_t^{\tilde{E}^l}$ is collinear (resp. non-collinear) for the foliations $\rho^*\mathcal{F}$ and $\rho^*\mathcal{G}$. Given a bifurcation divisor E of G(Z), we denote by $\operatorname{Col}(E)$ the set of collinear points of E and $\operatorname{NCol}(E)$ the set of non-collinear points of E.

Corollary 6.2 and the results in Section 5 allow to give a decomposition of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$. By Theorem 5.8, we have a decomposition

$$\widetilde{\mathcal{J}}_{\mathcal{F},\mathcal{G}} = \widetilde{J}^* \cup \left(\bigcup_{\tilde{E} \in B_N(\tilde{Z})} J^{\tilde{E}}\right)$$

with $J^{\tilde{E}} = J_{nc}^{\tilde{E}} \cup J_c^{\tilde{E}}$. Given a non-collinear bifurcation divisor E of G(Z), we define $J^E = J_{nc}^E \cup J_c^E$ to be such that

$$\rho^{-1}J_{nc}^{E} = \bigcup_{l=1}^{\underline{n}_{E}} J_{nc}^{\tilde{E}^{l}}; \qquad \rho^{-1}J_{c}^{E} = \bigcup_{l=1}^{\underline{n}_{E}} J_{c}^{\tilde{E}^{l}}$$

where $\{\widetilde{E}^l\}_{l=1}^{\underline{n}_E}$ are the divisors of $G(\widetilde{Z})$ associated to E and J^* to be such that $\rho^{-1}J^* = \widetilde{J}^*$. Hence, we can state the main result of this paper

TOME 75 (2025), FASCICULE 1

THEOREM 6.4. — Let us write $Z = \bigcup_{i=1}^{r+s} Z_i$ with Z_i irreducible and denote by $B_N(Z)$ the set of non-collinear bifurcation divisors of G(Z). Then there is a decomposition

$$\mathcal{J}_{\mathcal{F},\mathcal{G}} = J^* \cup \left(\bigcup_{E \in B_N(Z)} J^E\right)$$

with $J^E = J^E_{nc} \cup J^E_c$ such that

- $\begin{array}{l} \text{h } J^{E} = J_{nc}^{E} \cup J_{c}^{E} \text{ such that} \\ \text{(i) } \nu_{0}(J_{nc}^{E}) \leqslant \begin{cases} \underline{n}_{E}n_{E}(b_{E}-1), & \text{if } E \text{ does not belong to} \\ & a \text{ dead arc,} \\ \underline{n}_{E}n_{E}(b_{E}-1) \underline{n}_{E}, & \text{otherwise.} \end{cases}$
- (ii) For each irreducible component δ of J_{nc}^E we have that
 - $C(\delta, Z_i) = v(E)$ if E belongs to the geodesic of Z_i ;
 - $\mathcal{C}(\delta, Z_i) = \mathcal{C}(Z_i, Z_i)$ if E belongs to the geodesic of Z_i but not to the one of Z_i .
- (iii) For each irreducible component δ of J_c^E , there exists an irreducible component Z_i of Z such that E belongs to its geodesic and

$$\mathcal{C}(\delta, Z_i) > v(E).$$

Moreover, if E' is the first non-collinear bifurcation divisor in the geodesic of Z_i after E, then

$$\mathcal{C}(\delta, Z_i) < v(E').$$

7. Jacobian curves of hamiltonian foliations and Polar curves of foliations

In this section we will explain how our results imply previous results concerning jacobian curves of two plane curves or polar curves of foliations.

7.1. Jacobian of two curves

In [19, 20], T.-C. Kuo and A. Parusińki consider the Jacobian f_xg_y – $f_y g_x = 0$ of a pair of germs of holomorphic functions f, g without common branches and give properties of its Puiseux series which they called *polar* roots of the Jacobian. They define a tree-model, noted T(f, g), which represents the Puiseux series of the curves C = (f = 0) and D = (g = 0)and the contact orders among these series. The tree-model T(f,g) is constructed as follows: it starts with an horizontal bar B_* called ground bar and a vertical segment on B_* called the main trunk of the tree. This trunk is marked with [p,q] where $p = \nu_0(C)$ and $q = \nu_0(D)$. Let $\{y_i^C(x)\}_{i=1}^{\nu_0(C)}$ and $\{y_i^D(x)\}_{i=1}^{\nu_0(D)}$ be the Puiseux series of C and D respectively, and denote $\{z_j(x)\}_{j=1}^N$ the set $\{y_i^C(x)\}_{i=1}^{\nu_0(C)} \cup \{y_i^D(x)\}_{i=1}^{\nu_0(D)}$ with $N = \nu_0(C) + \nu_0(D)$. Denote by $h_0 = \min\{\operatorname{ord}_x(z_i(x) - z_j(x)) : 1 \leq i, j \leq N\}$. A bar B_0 is drawn on top of the main trunk with $h(B_0) = h_0$ being the height of B_0 . The Puiseux series $\{z_j(x)\}$ are divided into equivalence classes (mod $B_0)$ by the following relation: $z_j(x) \sim_{B_0} z_k(x)$ if $\operatorname{ord}_x(z_j(x) - z_k(x)) > h_0$. Each equivalence class is represented by a vertical line, called trunk, drawn on the top of B_0 . Each trunk is marked by a bimultiplicity [s, t]where s (resp. t) denote the number of Puiseux series of C (resp. of D) in the equivalence class. The same construction is repeated recursively on each trunk. The construction finishes with trunks which have bimultiplicity [1,0] or [0,1] representing each Puiseux series of the curve $Z = C \cup D$.

Let us now consider the curve $\widetilde{Z} = \rho^{-1}Z$ where $\rho : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ is any Z-ramification given by $\rho(u, v) = (u^n, v)$ (see Appendix A). Since the branches of \widetilde{Z} are in bijection with the Puiseux series of Z and the valuations of the bifurcation divisors of $G(\widetilde{Z})$ represent the contact orders among these series, then the tree model above T(f,g) can be recovered from the dual graph of $G(\widetilde{Z})$: there is a bijection between the set of bars of T(f,g) which are not the ground bar and the bifurcation divisors $B(\widetilde{Z})$ of $G(\widetilde{Z})$. For instance, the first bar B_0 corresponds to the first bifurcation divisor \widetilde{E}_1 of $G(\widetilde{Z})$ and $h(B_0) = \frac{v(\widetilde{E}_1)}{n}$. The number of trunks on B_0 is equal to $b_{\widetilde{E}_1}$, that is, each trunk on B_0 correspond to an infinitely near point of \widetilde{Z} on $\widetilde{E}_{1,\text{red}}$. In particular, given a trunk with bimultiplicity [s,t] corresponding to a point $R_i^{\widetilde{E}_1}$, then $s = \nu_{R_i^{\widetilde{E}_1}}(\pi_{\widetilde{E}_1}^*C)$ and $t = \nu_{R_i^{\widetilde{E}_1}}(\pi_{\widetilde{E}_1}^*D)$.

We shall illustrate with an example the relationship between T(f,g) and $G(\tilde{Z})$. The following example corresponds to the Example 1.1 in [20].

Example 7.1. — Take positive integers d, f with d < f and non-zero constants A, B. Consider

$$\begin{split} f(x,y) &= (y+x)(y-x^{d+1}+Ax^{f+1})(y+x^{d+1}+Bx^{f+1})\\ g(x,y) &= (y-x)(y-x^{d+1}-Ax^{f+1})(y+x^{d+1}-Bx^{f+1}) \end{split}$$

and put C = (f = 0) and D = (g = 0). Since $Z = C \cup D$ has only nonsingular irreducible components, we do not need to consider a ramification. The dual graph G(Z) is given by



while the tree-model is given by



where we have indicated the branches of C and D corresponding to the terminal trunks. Thus, the bijection among the bars in T(f,g) and the bifurcation divisors in G(Z) is given by

 $B_0 \longleftrightarrow E_1; \qquad B_1 \longleftrightarrow E_{d+1}; \qquad B_2 \longleftrightarrow E'_{f+1}; \qquad B_3 \longleftrightarrow E_{f+1}$ with $h(B_0) = v(E_1) = 1, \ h(B_1) = v(E_{d+1}) = d+1, \ h(B_2) = h(B_3) = v(E_{f+1}) = v(E'_{f+1}) = f+1.$

Let us show that the notion of collinear point and collinear divisor given in Section 4 correspond to the ones given in [19, 20] thanks to the bijections explained above. Let B be a bar of T(f, g) and consider a Puiseux series $z_k(x)$ of Z which goes through B, this means, that

$$z_k(x) = z_B(x) + cx^{h(B)} + \cdots$$

where $z_B(x)$ depends only on the bar *B* and *c* is uniquely determined by $z_k(x)$. If *T* is a trunk which contains $z_k(x)$, then it is said that the trunk *T* grows on *B* at *c*. Let \tilde{E} be the bifurcation divisor of $G(\tilde{Z})$ corresponding to *B* and consider

$$v_k(u) = z_k(u^n) = z_B(u^n) + cu^{nh(B)} + \cdots$$

ANNALES DE L'INSTITUT FOURIER

Note that the curve given by $v - v_k(u) = 0$ is a branch of \widetilde{Z} such that \widetilde{E} belongs to its geodesic. This curve determines a unique point R in \widetilde{E}_{red} ; in this way we can establish the bijection among the trunks on B and the infinitely near points of \widetilde{Z} on \widetilde{E}_{red} .

Let now B be a bar of T(f,g) that corresponds to a bifurcation divisor \widetilde{E} of $G(\widetilde{Z})$ and T_k , $1 \leq k \leq b_{\widetilde{E}}$, be the set of trunks on B with bimultiplicity $[p_k, q_k]$ where the trunk T_i grows on B at c_i . Let us denote $\{R_1^{\widetilde{E}}, \ldots, R_{b_{\widetilde{E}}}^{\widetilde{E}}\}$ the set of infinitely near points of \widetilde{Z} in \widetilde{E}_{red} with $R_i^{\widetilde{E}}$ corresponding to the trunk T_i .

In [20], the authors define

$$\Delta_B(c_k) = \begin{vmatrix} \nu_f(B) & p_k \\ \nu_g(B) & q_k \end{vmatrix}, \qquad 1 \leqslant k \leqslant b_{\tilde{E}}$$

where $\nu_f(B) = \operatorname{ord}_x(f(x, z_B(x) + cx^{h(B)}))$ for $c \in \mathbb{C}$ generic (resp. $\nu_g(B)$), and the rational function associated to B as

$$\mathcal{M}_B(z) = \sum_{k=1}^{b_{\tilde{E}}} \frac{\Delta_B(c_k)}{z - c_k}.$$

Note that, if E is the bifurcation divisor of G(Z) such that \tilde{E} is associated to E, then the curve given by $y = z_B(x) + cx^{h(B)}$ is an E-curvette. Thus, taking into account Proposition 2.5.3 of [7] for instance, we get that

$$\nu_f(B) = \frac{(C, \gamma_E)_0}{m(E)}$$

with γ_E any *E*-curvette. Moreover, it is easy to verify that

$$\nu_f(B) = \frac{1}{n} \sum_{i=1}^{\nu_0(C)} (\gamma_i^C, \gamma_{\tilde{E}})_0$$

where γ_i^C is the curve given by $v - y_i^C(u^n) = 0$ and $\gamma_{\tilde{E}}$ is an \tilde{E} -curvette. Note that we can compute the intersection multiplicity $(\gamma_i^C, \gamma_{\tilde{E}})_0 = \sum_{\tilde{E}' \leqslant \tilde{E}} \varepsilon_{\tilde{E}'}^{\gamma_i^C}$ where the sum runs over all the divisors \tilde{E}' in $G(\tilde{C})$ in the geodesic of \tilde{E} and $\varepsilon_{\tilde{E}'}^{\gamma_i^C} = 1$ if the geodesic of γ_i^C contains the divisor \tilde{E}' and $\varepsilon_{\tilde{E}'}^{\gamma_i^C} = 0$ otherwise. Thus, with the notations given in Section 3, we have that

$$\nu_f(B) = \frac{1}{n} \kappa_{\tilde{E}}(\mathcal{L}^{\tilde{C}})$$

where $\mathcal{L}^{\tilde{C}} = \mathcal{G}_{\tilde{f}}$ is the logarithmic foliation in $\mathbb{G}_{\tilde{C}}$ with $\lambda = (1, 1, ..., 1)$, that is, the hamiltonian foliation defined by $d\tilde{f} = 0$ with $\tilde{f}(u, v) = f(u^n, v)$.

Moreover,

$$p_k = \nu_{R_k^{\tilde{E}}}(\pi_{\tilde{E}}^* \widetilde{C}), \qquad k = 1, \dots, b_{\tilde{E}}$$

and thus

$$\mathcal{I}_{R_k^{\tilde{E}}}(\pi_{\tilde{E}}^*\mathcal{L}^{\tilde{C}}, \widetilde{E}_{\mathrm{red}}) = -\frac{p_k}{n\nu_f(B)}, \qquad k = 1, \dots, b_{\tilde{E}}.$$

Consequently, with the notations introduced in Section 4, we have that

$$\Delta_B(c_k) = -\frac{1}{n} \Delta_{\tilde{E}}(R_k^{\tilde{E}}) \quad \text{and} \quad \mathcal{M}_{\tilde{E}}(z) = -n \mathcal{M}_B(z).$$

Thus the notions of collinear divisor and collinear point given in Section 4 correspond to the ones given in [20] for bars and points on them, and the results given in Section 5 imply some of the Theorems proved in [20].

7.2. Semiroots and Approximate roots

The notion of approximated root was introduced by Abhyankar and Moh in [1] where they proved the following result:

PROPOSITION 7.2. — Let A be an integral domain and $P(y) \in A[y]$ be a monic polynomial of degree d. If p is invertible in A and p divides d, then there exists a unique monic polynomial $Q(y) \in A[y]$ such that the degree of $P - Q^p$ is less than d - d/p.

The unique polynomial Q given by the previous proposition is called the p-th approximate root of P. Let us consider $f(x, y) \in \mathbb{C}\{x\}[y]$ an irreducible Weierstrass polynomial with characteristic exponents $\{\beta_0, \beta_1, \ldots, \beta_g\}$ and denote $e_k = \gcd(\beta_0, \beta_1, \ldots, \beta_k)$ for $k = 1, \ldots, g$. Thus e_k divides $\beta_0 = \deg_y f$. We will denote $f^{(k)}$ the e_k -approximate root of f and we call them the characteristic approximate roots of f. Next result ([1, Theorem 7.1]) gives the main properties of the characteristic approximate roots of f (see also [17, 29]):

PROPOSITION 7.3. — Let $f(x, y) \in \mathbb{C}\{x\}[y]$ be an irreducible Weierstrass polynomial with characteristic exponents $\{\beta_0, \beta_1, \ldots, \beta_g\}$. Then the characteristic approximate roots $f^{(k)}$ for $k = 0, 1, \ldots, g - 1$ verify:

- (i) The degree in y of $f^{(k)}$ is equal to β_0/e_k and $\mathcal{C}(f, f^{(k)}) = \beta_{k+1}/\beta_0$.
- (ii) The polynomial $f^{(k)}$ is irreducible with characteristic exponents $\{\beta_0/e_k, \beta_1/e_k, \dots, \beta_k/e_k\}.$

gl

ANNALES DE L'INSTITUT FOURIER

270

In [14], E. García Barroso and J. Gwoździewicz studied the jacobian curve of f and $f^{(k)}$ and they give a result concerning its factorization (see [14, Theorem 1]). In this section, we will prove that this result of factorization can be obtained as a consequence of Theorem 6.4.

Remark 7.4. — In [29], P. Popescu-Pampu proved that all polynomials in $\mathbb{C}\{x\}[y]$ satisfying condition (i) in the proposition above also verify condition (ii). Hence, given an irreducible Weierstrass polynomial $f(x, y) \in$ $\mathbb{C}\{x\}[y]$ with characteristic exponents $\{\beta_0, \beta_1, \ldots, \beta_g\}$, we can consider the monic polynomials in $\mathbb{C}\{x\}[y]$ satisfying condition (i) above which are called *k*-semiroots of f (see [29, Definition 6.4]). Since we only need the properties of characteristic approximate roots given in Proposition 7.3, in the rest of the section, we will denote by $f^{(k)}$ a *k*-semiroot of f, $0 \leq k \leq g - 1$.

Let C be the curve defined by f = 0 and denote $C^{(k)}$ the curve given by $f^{(k)} = 0$ with $0 \leq k \leq g-1$. Consider $\mathcal{F} \in \mathbb{G}_C$ and $\mathcal{F}^{(k)} \in \mathbb{G}_{C^{(k)}}$. Note that the minimal reduction of singularities $\pi_C : X_C \to (\mathbb{C}^2, 0)$ of the curve C gives also a reduction of singularities of $C \cup C^{(k)}$. There are g bifurcation divisors in G(C). The set of bifurcation divisors of G(C) will be denote by $\{E_1, \ldots, E_g\}$ with $v(E_i) = \frac{\beta_i}{\beta_0}$. Remark that the dual graph $G(C \cup C^{(k)})$ is given by (see [29] for instance):



Thus the sets of bifurcation divisors of G(C) and $G(C \cup C^{(k)})$ coincide. All bifurcation divisors of $G(C \cup C^{(k)})$ are Puiseux divisors for C while only E_1, \ldots, E_k are Puiseux divisors for $C^{(k)}$. Then we have

LEMMA 7.5. — The set of non-collinear bifurcation divisors of $G(C \cup C^{(k)})$ for the foliations \mathcal{F} and $\mathcal{F}^{(k)}$ is $\{E_{k+1}, \ldots, E_q\}$.

Proof. — Let $\{(m_1, n_1), \ldots, (m_g, n_g)\}$ be the Puiseux pairs of C, then we remind that $\beta_0 = \nu_0(C) = n_1 \cdots n_g$, $e_k = n_{k+1} \cdots n_g$ and $\beta_k/\beta_0 = m_k/n_1 \cdots n_k$ for $k = 1, \ldots, g$. Given a bifurcation divisor E_l of $G(C \cup C^{(k)})$, we have that $n_{E_l} = n_l$, $\underline{n}_{E_l} = n_1 \cdots n_{l-1} = \beta_0/e_{l-1}$ and $m(E_l) = \underline{n}_{E_l}n_E = n_1 \cdots n_l$. Consider now the ramification $\rho : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ given by $\rho(u, v) = (u^n, v)$ with $n = \beta_0$ and denote $\widetilde{C} = \rho^{-1}C$, $\widetilde{C}^{(k)} = \rho^{-1}C^{(k)}$. Take a bifurcation divisor E_l and let $\{\widetilde{E}_l^t\}_{t=1}^{n_{E_l}}$ be the set of bifurcation divisors of $G(\widetilde{C} \cup \widetilde{C}^{(k)})$ associated to E_l .

In the case l < k+1, we have that $\pi_{\tilde{E}_{l}^{t}}^{*} \widetilde{C} \cap \widetilde{E}_{l,\mathrm{red}}^{t} = \pi_{\tilde{E}_{l}^{t}}^{*} \widetilde{C}^{(k)} \cap \widetilde{E}_{l,\mathrm{red}}^{t}$ with $b_{\tilde{E}_{l}^{t}} = n_{l}$ in $G(\widetilde{C} \cup \widetilde{C}^{(k)})$. Let us denote $\pi_{\tilde{E}_{l}^{t}}^{*} \widetilde{C} \cap \widetilde{E}_{l,\mathrm{red}}^{t} = \{R_{1}^{\tilde{E}_{l}^{t}}, \ldots, R_{b_{\tilde{E}_{l}^{t}}}^{\tilde{E}_{l}^{t}}\}$. Using the equations given in Section 3 and Appendix A, the computation of the Camacho–Sad indices for the foliations $\rho^{*}\mathcal{F}$ and $\rho^{*}\mathcal{F}^{(k)}$ gives

(7.1)
$$\mathcal{I}_{R_{s}^{\tilde{E}_{l}^{t}}}(\pi_{\tilde{E}_{l}^{t}}^{*}\rho^{*}\mathcal{F}, \widetilde{E}_{l,\mathrm{red}}^{t}) = -\frac{n_{l+1}\cdots n_{g}}{\sum_{s=1}^{n}\sum_{\tilde{E}\leqslant\tilde{E}_{l}^{t}}\varepsilon_{\tilde{E}}^{\sigma_{s}}} \\ \mathcal{I}_{R_{s}^{\tilde{E}_{l}^{t}}}(\pi_{\tilde{E}_{l}^{t}}^{*}\rho^{*}\mathcal{F}^{(k)}, \widetilde{E}_{l,\mathrm{red}}^{t}) = -\frac{n_{l+1}\cdots n_{k}}{\sum_{s=1}^{n_{1}\cdots n_{k}}\sum_{\tilde{E}\leqslant\tilde{E}_{l}^{t}}\varepsilon_{\tilde{E}}^{\sigma_{s}^{(k)}}}$$

where $\widetilde{C} = \bigcup_{s=1}^{n} \sigma_s$ and $\widetilde{C}^{(k)} = \bigcup_{s=1}^{n_1 \cdots n_k} \sigma_s^{(k)}$. Hence, taking into account the results of Appendix A, we have

$$\mathcal{I}_{R_s^{\tilde{E}_l^t}}(\pi_{\tilde{E}_l^t}^*\rho^*\mathcal{F}, \tilde{E}_{l,\mathrm{red}}^t) = \mathcal{I}_{R_s^{\tilde{E}_l^t}}(\pi_{\tilde{E}_l^t}^*\rho^*\mathcal{F}^{(k)}, \tilde{E}_{l,\mathrm{red}}^t), \qquad s = 1, \dots, b_{\tilde{E}_l^t}$$

and consequently, $\Delta_{\tilde{E}_l^t}(R_s^{\tilde{E}_l^t}) = 0$, $s = 1, \ldots, b_{\tilde{E}_l^t}$, for the foliations $\rho^* \mathcal{F}$ and $\rho^* \mathcal{F}^{(k)}$. This proves that the bifurcation divisors E_l of $G(C \cup C^{(k)})$ with l < k + 1 are collinear for \mathcal{F} and $\mathcal{F}^{(k)}$.

Consider now the bifurcation divisor E_{k+1} of $G(C \cup C^{(k)})$ and let $\{\widetilde{E}_{k+1}^t\}_{t=1}^{\underline{n}_{E_{k+1}}}$ be the set of bifurcation divisors of $G(\widetilde{C} \cup \widetilde{C}^{(k)})$ associated to E_{k+1} . Although the curve $\pi_{E_{k+1}}^*C^{(k)}$ does not intersect $E_{k+1,\text{red}}$, the curve $\pi_{\widetilde{E}_{k+1}^t}^*\widetilde{C}^{(k)}$ intersects $\widetilde{E}_{k+1,\text{red}}^t$ in one point for each $t = 1, \ldots, \underline{n}_{E_{k+1}}$ which is different from the n_{k+1} points where $\pi_{\widetilde{E}_{k+1}^t}^*\widetilde{C}$ intersects $\widetilde{E}_{k+1,\text{red}}^t$. Note that $b_{E_{k+1}} = 2$ in $G(C \cup C^{(k)})$ and, by (A.2), $b_{\widetilde{E}_{k+1}^t} = n_{k+1} + 1$ in $G(\widetilde{C} \cup \widetilde{C}^{(k)})$. Let $\{R_1^{\widetilde{E}_{k+1}^t}, \ldots, R_{b_{\widetilde{E}_{k+1}^t}}^{\widetilde{E}_{k+1}^t}\}$ be the set of points $(\pi_{\widetilde{E}_{k+1}^t}^*\widetilde{C} \cap \widetilde{E}_{k+1,\text{red}}^t) \cup (\pi_{\widetilde{E}_{k+1}^t}^*\widetilde{C}^{(k)} \cap \widetilde{E}_{k+1,\text{red}}^t)$ in the previous case, and prove that $\Delta_{\widetilde{E}_{k+1}^t}(R_s^{\widetilde{E}_{k+1}^t}) \neq 0$, $s = 1, \ldots, b_{\widetilde{E}_{k+1}^t}$, for the foliations $\rho^*\mathcal{F}$ and $\rho^*\mathcal{F}^{(k)}$. Consequently E_{k+1} is a non-collinear divisor for \mathcal{F} and $\mathcal{F}^{(k)}$.

ANNALES DE L'INSTITUT FOURIER

However, we have that

(7.2)
$$\sum_{s=1}^{b_{\tilde{E}_{k+1}^t}} \Delta_{\tilde{E}_{k+1}^t} (R_s^{\tilde{E}_{k+1}^t}) = 0.$$

In fact, the divisor \widetilde{E}_{k+1}^t arises from one of the divisors \widetilde{E}_k^r at one of the points $R_t^{\widetilde{E}_k^r}$ of the set $\pi_{\widetilde{E}_k^r}^* \widetilde{C} \cap \widetilde{E}_{k,\mathrm{red}}^r$. Since \widetilde{E}_k^r is a collinear divisor, then $R_t^{\widetilde{E}_k^r}$ is a collinear point and (7.2) follows from Corollary 4.12.

Consider now a bifurcation divisor E_l of $G(C \cup C^{(k)})$ with l > k + 1. In this case, we have that the curve $\pi_{E_l}^* C^{(k)}$ does not intersect $E_{l,\text{red}}$, the curve $\pi_{\tilde{E}_l^t}^* \tilde{C}^{(k)}$ does not intersect $E_{l,\text{red}}$, the curve $\pi_{\tilde{E}_l^t}^* \tilde{C}^{(k)}$ does not intersect $\tilde{E}_{l,\text{red}}^t$ and $b_{\tilde{E}_l^t} = n_l$ in $G(\tilde{C} \cup \tilde{C}^{(k)})$ (see (A.2)). Let us denote $\{R_1^{\tilde{E}_l^t}, \ldots, R_{b_{\tilde{E}_l^t}}^{\tilde{E}_l^t}\}$ the set of points $\pi_{\tilde{E}_l^t}^* \tilde{C} \cap \tilde{E}_{l,\text{red}}^t = \pi_{\tilde{E}_l^t}^* (\tilde{C} \cup \tilde{C}^{(k)}) \cap \tilde{E}_{l,\text{red}}^t$. With the notations above, for $s \in \{1, \ldots, b_{\tilde{E}_l^t}\}$, we have that $\mathcal{I}_{R_s^{\tilde{E}_l^t}}(\pi_{\tilde{E}_l^t}^* \rho^* \mathcal{F}, \tilde{E}_{l,\text{red}}^t)$ is given by (7.1) while $\mathcal{I}_{R_s^{\tilde{E}_l^t}}(\pi_{\tilde{E}_l^t}^* \rho^* \mathcal{F}^{(k)}, \tilde{E}_{l,\text{red}}^t) = 0$ since the points $R_s^{\tilde{E}_l^t}$ are non-singular points for $\rho^* \mathcal{F}^{(k)}$. This implies $\Delta_{\tilde{E}_l^t}(R_s^{\tilde{E}_l^t}) \neq 0, 1 \leqslant s \leqslant b_{\tilde{E}_l^t}$, and hence E_l is a non-collinear divisor for the foliations $\rho^* \mathcal{F}$ and $\rho^* \mathcal{F}^{(k)}$. Moreover, we have that

$$\sum_{s=1}^{b_{\tilde{E}_{l}^{t}}} \Delta_{\tilde{E}_{l}^{t}}(R_{s}^{\tilde{E}_{l}^{t}}) = -\sum_{s=1}^{b_{\tilde{E}_{l}^{t}}} \mathcal{I}_{R_{s}^{\tilde{E}_{l}^{t}}}(\pi_{\tilde{E}_{l}^{t}}^{*}\rho^{*}\mathcal{F}, \widetilde{E}_{l,\mathrm{red}}^{t}) = 1 + \mathcal{I}_{Q}(\pi_{\tilde{E}_{l}^{t}}^{*}\rho^{*}\mathcal{F}, \widetilde{E}_{l,\mathrm{red}}^{t})$$

where Q is the only singular point of $\pi_{\tilde{E}_{l}^{t}}^{*}\rho^{*}\mathcal{F}$ in \tilde{E}_{l}^{t} different from the points $R_{s}^{\tilde{E}_{l}^{t}}$. By Proposition 4.4 in [9], we know that $\mathcal{I}_{Q}(\pi_{\tilde{E}_{l}^{t}}^{*}\rho^{*}\mathcal{F}, \tilde{E}_{l,\mathrm{red}}^{t}) \neq -1$ and hence

$$\sum_{s=1}^{b_{\tilde{E}_l}^t} \Delta_{\tilde{E}_l^t}(R_s^{\tilde{E}_l^t}) \neq 0.$$

Thus, by Theorem 6.4 there is a decomposition

$$\mathcal{J}_{\mathcal{F},\mathcal{F}^{(k)}} = J^* \cup \left(\bigcup_{i=k+1}^g J^i\right)$$

where $J^i = J_{nc}^{E_i}$, such that

- (i) $\nu_0(J^{k+1}) < n_1 \cdots n_{k+1}$.
- (ii) $\nu_0(J^i) = n_1 \cdots n_{i-1}(n_i 1)$ for $k + 2 \le i \le g$.
- (iii) if γ is a branch of J^i , $k+1 \leq i \leq g$, we have that $\mathcal{C}(\gamma, C) = \frac{\beta_i}{\beta_0}$.
- (iv) if γ is a branch of J^* , then $\mathcal{C}(\gamma, C) < \frac{\beta_{k+1}}{\beta_0}$.

Note that γ is a branch of J^* if it is not a branch of any of the curves J^i , $k+1 \leq i \leq g$, and hence, $\pi_C^* \gamma$ intersects a component E of the exceptional divisor $\pi_C^{-1}(0)$ which appears in the reduction of singularities of C before than E_{k+1} . Consequently we have that $\mathcal{C}(\gamma, C) < v(E_{k+1}) = \frac{\beta_{k+1}}{\beta_0}$ which gives property (iv).

Let us prove that $J^{k+1} = \emptyset$. From Section 6 we have that $\rho^{-1}J^{k+1} = \bigcup_{t=1}^{n_1 \cdots n_k} J_{nc}^{\tilde{E}_{k+1}^t}$ with the notations of the proof of Lemma 7.5. Let us compute $\mathcal{M}_{\tilde{E}_{k+1}^t}(z)$ for any $t \in \{1, \ldots, n_1 \cdots n_k\}$. To simplify notations, let us denote $\tilde{b} = b_{\tilde{E}_{k+1}^t} = n_{k+1} + 1$, $\tilde{R}_s = R_s^{\tilde{E}_{k+1}^t}$ and $\Delta(\tilde{R}_l) = \Delta_{\tilde{E}_{k+1}^t}(\tilde{R}_l)$. Thus, we have that

$$\mathcal{M}_{\tilde{E}_{k+1}^t}(z) = \sum_{s=1}^{\tilde{b}-1} \frac{\Delta(\tilde{R}_s)}{z-\xi^s} + \frac{\Delta(\tilde{R}_{\tilde{b}})}{z}$$

where ξ is a primitive n_{k+1} -root of a value $a = a^{\tilde{E}_{k+1}^t}$ determined by the Puiseux parametrizations of C. From the proof of Lemma 7.5, we obtain that $\Delta(\tilde{R}_s) = \Delta(\tilde{R}_t)$ for any $s, t \in \{1, \ldots, \tilde{b} - 1\}$. Thus, taking into account (7.2), we get that

$$\mathcal{M}_{\tilde{E}_{k+1}^{t}}(z) = \Delta(\tilde{R}_{1}) \frac{\sum_{s=1}^{b-1} \prod_{\substack{t=1\\t \neq s}}^{n_{k+1}} (z-\xi^{t})}{z^{n_{k+1}} - a} + \frac{\Delta(\tilde{R}_{\tilde{b}})}{z}$$
$$= \Delta(\tilde{R}_{1}) \frac{n_{k+1} z^{n_{k+1}-1}}{z^{n_{k+1}} - a} + \frac{\Delta(\tilde{R}_{\tilde{b}})}{z} = \frac{-a\Delta(\tilde{R}_{\tilde{b}})}{z(z^{n_{k+1}} - a)}$$

where $a\Delta(\widetilde{R}_{\tilde{b}}) \neq 0$. Hence, $J_{nc}^{\tilde{E}_{k+1}^{t}} = \emptyset$ for all $t = 1, \ldots, n_1 \cdots n_k$ and consequently $J^{k+1} = \emptyset$.

COROLLARY 7.6. — Let C be an irreducible curve and $C^{(k)}$ the curve given by the k-characteristic approximate root (or by a k-semiroot) with $0 \leq k \leq g - 1$. Consider $\mathcal{F} \in \mathbb{G}_C$ and $\mathcal{F}^{(k)} \in \mathbb{G}_{C^{(k)}}$. Thus, the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{F}^{(k)}}$ has a decomposition

$$\mathcal{J}_{\mathcal{F},\mathcal{F}^{(k)}} = J^* \cup \left(\bigcup_{i=k+2}^g J^i\right)$$

such that

- (i) $\nu_0(J^i) = n_1 \cdots n_{i-1}(n_i 1)$ for $k + 2 \le i \le g$.
- (ii) if γ is a branch of J^i , $k+2 \leq i \leq g$, we have that $\mathcal{C}(\gamma, C) = \frac{\beta_i}{\beta_0}$.
- (iii) if γ is a branch of J^* , then $\mathcal{C}(\gamma, C) < \frac{\beta_{k+1}}{\beta_0}$.

In particular, the result above implies the result of E. García Barroso and J. Gwoździewicz ([14, Theorem 1]) concerning the jacobian curve of a plane curve and its characteristic approximate roots.

Moreover, in [31], it is considered the jacobian curve $\mathcal{J}_{\mathcal{F},\mathcal{G}^{(k)}}$ of a foliation \mathcal{F} with an irreducible separatrix f = 0 and the hamiltonian foliation $\mathcal{G}^{(k)}$ defined by $df^{(k)} = 0$ with $f^{(k)}$ a characteristic approximate root of f. Corollary 7.6 also implies the main result of N. E. Saravia in [31, Theorem 4.1] concerning factorization of $\mathcal{J}_{\mathcal{F},\mathcal{G}^{(k)}}$.

Remark 7.7. — Note that E^1 is always a collinear divisor for the foliations \mathcal{F} and $\mathcal{F}^{(k)}$, hence the hypothesis of Lemma 2.3 are not satisfied and the multiplicity of $\mathcal{J}_{\mathcal{F},\mathcal{F}^{(k)}}$ can be greater than $\nu_0(\mathcal{F}) + \nu_0(\mathcal{F}^{(k)}) =$ $\nu_0(C) + \nu_0(C^{(k)}) - 2$ as showed in the examples given in [14] or [31].

7.3. Polar curves of foliations

Given a germ of foliation \mathcal{F} in $(\mathbb{C}^2, 0)$, a polar curve of \mathcal{F} corresponds to the jacobian curve of \mathcal{F} and a non-singular foliation \mathcal{G} . If we are interested in the topological properties of a generic polar curve of \mathcal{F} , it is enough to consider a generic curve $\mathcal{P}_{[a;b]}^{\mathcal{F}}$ in the family of curves given by

$$\omega \wedge (a \,\mathrm{d} y - b \,\mathrm{d} x) = 0$$

where $\omega = 0$ is a 1-form defining \mathcal{F} and $[a:b] \in \mathbb{P}^1_{\mathbb{C}}$ (see [9, Section 2]). When \mathcal{F} is a hamiltonian foliation given by df = 0 we recover the notion of polar curve of a plane curve. As we mention in the introduction, polar curves play an important role in the study of singularities of plane curves and also of foliations. There is a result, known as "decomposition theorem", which describes the minimal topological properties of the generic polar curve of a plane curve C in terms of the topological type of the curve C(see [25] for the case of C irreducible; [13] for C with several branches). In the case of foliations, the decomposition theorem also holds for the generic polar curve of a generalized curve foliation \mathcal{F} with an irreducible separatrix (see [30]). In the general case of a generalized curve foliation \mathcal{F} whose curve of separatrices is not irreducible, the decomposition theorem for its generic polar curve only holds under some conditions on the foliation \mathcal{F} (see [9]). Let us see that all these results can be recovered from the results in this paper. In particular, we show that we can prove Theorems 5.1 and 6.1 in [9] which give the decomposition theorem for the polar curve of a generalized curve foliation \mathcal{F} and hence we get all the other results concerning decompositions theorems.

Let \mathcal{F} be a generalized curve foliation in $(\mathbb{C}^2, 0)$ with C as curve of separatrices and denote by $\mathcal{P}^{\mathcal{F}}$ a generic polar curve of \mathcal{F} . We can assume that $\mathcal{P}^{\mathcal{F}} = \mathcal{J}_{\mathcal{F},\mathcal{G}}$ where \mathcal{G} is a non-singular foliation. Note that the curve of separatrices D of \mathcal{G} is a non-singular irreducible plane curve. Let us assume first that C has only non-singular irreducible components, all of them different from D, and take the notations of Section 4. Thus the minimal reduction of singularities $\pi_C : X_C \to (\mathbb{C}^2, 0)$ is also the minimal reduction of singularities of $Z = C \cup D$. Note that the dual graph G(Z) is obtained from G(C) adding an arrow to the first divisor E^1 which represents the curve D. Hence, if we denote b_E^Z , b_E^C the number associated to a divisor E in G(Z)or G(C) respectively, as defined in Subsection 2.3, then $b_{E^1}^Z = b_{E^1}^C + 1$ and $b_E^Z = b_E^C$ otherwise.

Consider E an irreducible component of the exceptional divisor $\pi_C^{-1}(0)$. If $E = E^1$ is the divisor which appears after the blow-up of the origin, then $\pi_{E^1}^* D \cap E_{\text{red}}^1 = \{Q\}$ and $Q \notin \pi_{E^1}^* C \cap E_{\text{red}}^1$. Thus, for $R \in E_{\text{red}}^1$ we have that

$$\Delta_{E^1}^{\mathcal{F},\mathcal{G}}(R) = \begin{cases} \mathcal{I}_Q(\pi_{E^1}^*\mathcal{G}, E_{\mathrm{red}}^1), & \text{if } R = Q, \\ -\mathcal{I}_R(\pi_{E^1}^*\mathcal{F}, E_{\mathrm{red}}^1), & \text{otherwise.} \end{cases}$$

with $\mathcal{I}_Q(\pi_{E^1}^*\mathcal{G}, E_{\mathrm{red}}^1) = -1$. If $E \neq E^1$, then $\pi_E^*Z \cap E_{\mathrm{red}} = \pi_E^*C \cap E_{\mathrm{red}}$ and then $\Delta_E^{\mathcal{F},\mathcal{G}}(R) = -\mathcal{I}_R(\pi_E^*\mathcal{F}, E_{\mathrm{red}})$ for any $R \in E_{\mathrm{red}}$.

With the hypothesis above and the notations of Section 4, we have that

LEMMA 7.8. — The following conditions are equivalent:

- (i) There is no corner in $\pi_C^{-1}(0)$ such that $\pi_C^* \mathcal{F}$ has Camacho–Sad index equal to -1.
- (ii) All the components of the exceptional divisor $\pi_C^{-1}(0)$ are purely non-collinear.

Proof. — Assume that (i) holds and that there is a component E of the exceptional divisor which is not purely non-collinear, that is, there is a singular point $R \in E_{\text{red}}$ of $\pi_E^* \mathcal{F}$ with $\mathcal{I}_R(\pi_E^* \mathcal{F}, E_{\text{red}}) = 0$. Then R is not a simple singularity for $\pi_E^* \mathcal{F}$ and hence, if $\sigma : X_{E'} \to X_E$ is the blow-up with center in R, and we denote by $\widetilde{E}_{\text{red}}$ the strict transform of E_{red} by σ , then we have that $\mathcal{I}_{\tilde{R}}(\pi_{\tilde{E}}^* \mathcal{F}, \widetilde{E}_{\text{red}}) = -1$ where $\widetilde{R} = E'_{\text{red}} \cap \widetilde{E}_{\text{red}}$. Thus we get a corner in $\pi_C^{-1}(0)$ with Camacho–Sad index equal to -1.

Conversely, assume now that all the components of the exceptional divisor $\pi_C^{-1}(0)$ are purely non-collinear and there is a corner $\widetilde{R} = E_{k-1} \cap E_k$ with $I_{\widetilde{R}}(\pi_C^*\mathcal{F}, E_{k-1}) = -1$. Consider the morphism $\pi_{E_{k-1}} : X_{E_{k-1}} \to (\mathbb{C}^2, 0)$ and take the point $R \in E_{k-1,\text{red}}$ that we have to blow-up to obtain the divisor E_k . Thus, by the properties of the Camacho–Sad, we have that $\mathcal{I}_R(\pi^*_{E_{k-1,\mathrm{red}}}\mathcal{F}, E_{k-1,\mathrm{red}}) = 0$ but this contradicts that E_{k-1} is purely noncollinear.

In particular, let us see that Theorem 6.1 in [9] is consequence of Theorem 5.2. Assume that the logarithmic model of \mathcal{F} is non resonant, this implies condition (i) in the lemma above (by [9, Proposition 4.4]) and hence all the divisors in G(Z) are non-collinear. Note that E^1 is always a bifurcation divisor in G(Z) and we have that $\sum_{R \in E^1} \Delta_{E^1}(R) = 0$ by Remark 4.11. If we write $\pi_{E^1}^* C \cap E_{\mathrm{red}}^1 = \{R_1^{E^1}, \ldots, R_{b_{E^1}}^{E^1}\}$ and $\pi_{E^1}^* D \cap E_{\mathrm{red}}^1 = \{Q\}$ with $R_l^{E^1} = (0, c_l^{E^1}), \ l = 1, \ldots, b_{E^1}^C$, and Q = (0, d) in coordinates in the first chart of E_{red}^1 , then the set of zeros of $\mathcal{M}_{E^1}(z)$ are given by the roots of the polynomial

$$\prod_{j=1}^{b_{E^1}^C} (z - c_j^{E^1}) + (z - d) \sum_{l=1}^{b_{E^1}^C} \mathcal{I}_{R_l^{E^1}}(\pi_{E^1}^* \mathcal{F}, E_{\text{red}}^1) \prod_{j \neq l} (z - c_j^{E^1})$$

which has multiplicity equal to $b_{E^1}^C - 1$ provided that

$$\sum_{j=1}^{b_{E^1}^C} c_j^{E^1} + d \sum_{l=1}^{b_{E^1}^C} \mathcal{I}_{R_l^{E^1}}(\pi_{E^1}^*\mathcal{F}, E_{\mathrm{red}}^1) \neq 0.$$

Note that we can assume that this condition holds since we are consider a generic polar curve.

Consider a component E of the exceptional divisor $\pi_C^{-1}(0)$. We have that $\operatorname{NCol}(E) = \pi_E^* C \cap E_{\operatorname{red}}$ if $E \neq E^1$ and $\operatorname{NCol}(E^1) = (\pi_{E^1}^* C \cap E_{\operatorname{red}}^1) \cup (\pi_{E^1}^* D \cap E_{\operatorname{red}}^1)$. Given a point P in $\pi_E^* C \cap E_{\operatorname{red}}$, by Theorem 5.2, we have that

$$\nu_P(\pi_E^*\mathcal{P}^\mathcal{F}) = \nu_P(\pi_E^*C) - 1$$

and hence we have [9, Theorem 6.1]. If we take the point Q given by $\pi_{E^1}^* D \cap E^1_{\text{red}}$, we have that $\nu_Q(\pi_{E^1}^* \mathcal{P}^{\mathcal{F}}) = \nu_Q(\pi_{E^1}^* D) - 1 = 0$. Thus, by Theorem 5.8 we obtain that $\mathcal{P}^{\mathcal{F}} = \bigcup_{E \in B(C)} J^E$ with the following properties

(i) $\nu_0(J^E) = b_E^C - 1$,

(ii)
$$\pi_E^* J^E \cap \pi_E^* C = \emptyset$$
,

- (iii) if E' < E, then $\pi_E^* J^E \cap \pi'_E(E') = \emptyset$,
- (iv) if E' > E, then $\pi_{E'}^* J^E \cap E'_{\text{red}} = \emptyset$,

which in particular implies the decomposition of the generic polar curve given in Corollary 6.2 of [9] for C with non-irreducible components. Thus the decomposition in the general case ([9, Theorem 5.1]) follows from Theorem 6.4.

Appendix A. Equisingularity data and Ramification

The aim of this appendix is to explain the behaviour of plane curves and their invariants under the action of a ramification. Although some of these results can be found in [10, Appendix B], we include them here for completeness.

A.1. Equisingularity data

In Subsection 2.3 we have introduced some notations concerning equisingularity of plane curves that will be used in the sequel. This appendix completes Subsection 2.3 with more notations related with equisingularity data that have already been used to prove some results or that will be useful in order to describe the effect of ramification over a plane curve.

Recall that $\pi_C : X_C \to (\mathbb{C}^2, 0)$ is the minimal reduction of singularities of a curve $C = \bigcup_{i=1}^r C_i$. Given an irreducible component E of $\pi_C^{-1}(0)$, a curvette $\tilde{\gamma}$ of the divisor E is a non-singular curve transversal to E at a non-singular point of $\pi_C^{-1}(0)$. The projection $\gamma = \pi_C(\tilde{\gamma})$ is a germ of plane curve in $(\mathbb{C}^2, 0)$ and we say that γ is an E-curvette. We denote by m(E) the multiplicity at the origin of any E-curvette and by v(E) the coincidence $\mathcal{C}(\gamma_E, \gamma'_E)$ of two E-curvettes γ_E, γ'_E which cut E in different points. Note that v(E) < v(E') if E < E'. Recall that the coincidence $\mathcal{C}(\gamma, \delta)$ between two irreducible curves γ and δ is defined as

(A.1)
$$\mathcal{C}(\gamma, \delta) = \sup_{\substack{1 \leq i \leq \nu_0(\gamma) \\ 1 \leq j \leq \nu_0(\delta)}} \{ \operatorname{ord}_x(y_i^{\gamma}(x) - y_j^{\delta}(x)) \}$$

where $\{y_i^{\gamma}(x)\}_{i=1}^{\nu_0(\gamma)}, \{y_j^{\delta}(x)\}_{j=1}^{\nu_0(\delta)}$ are the Puiseux series of γ and δ respectively.

Remark A.1. — Note that the coincidence $C(\gamma, \delta)$ between two irreducible curves γ and δ and the intersection multiplicity $(\gamma, \delta)_0$ of both curves at the origin are related as follows (see Merle [25, Proposition 2.4]): if $\{\beta_0, \beta_1, \ldots, \beta_g\}$ are the characteristic exponents of γ and α is a rational number such that $\beta_q \leq \alpha < \beta_{q+1}$ ($\beta_{g+1} = \infty$), then the following statements are equivalent:

(i)
$$C(\gamma, \delta) = \frac{\alpha}{\nu_0(\gamma)}$$

(ii) $\frac{(\gamma, \delta)_0}{\nu_0(\delta)} = \frac{\overline{\beta}_q}{n_1 \cdots n_{q-1}} + \frac{\alpha - \beta_q}{n_1 \cdots n_q}$

where $\{(m_i, n_i)\}_{i=1}^g$ are the Puiseux pairs of γ $(n_0 = 1)$ and $\{\overline{\beta}_0, \overline{\beta}_1, \dots, \overline{\beta}_g\}$ is a minimal system of generators of the semigroup $S(\gamma)$ of γ .

Consider any curvette $\tilde{\gamma}_E$ of E, then $\pi'_E(\tilde{\gamma}_E)$ is also a curvette of $E_{\text{red}} \subset X_E$ and it is clear that $m(E) = m(E_{\text{red}})$ and $v(E) = v(E_{\text{red}})$. Let $\{\beta_0^E, \beta_1^E, \ldots, \beta_{g(E)}^E\}$ be the characteristic exponents of $\gamma_E = \pi_C(\tilde{\gamma}_E)$. Then we have that $m(E) = \beta_0^E = \nu_0(\gamma_E)$. There are two possibilities for the value of v(E):

- (i) either π_E is the minimal reduction of singularities of γ_E and then $v(E) = \beta_{q(E)}^E / \beta_0^E$. We say that E is a Puiseux divisor for π_C (or C);
- (ii) or π_E is obtained by blowing-up $q \ge 1$ times after the minimal reduction of singularities of γ_E and in this situation $v(E) = (\beta_{g(E)}^E + q)/\beta_0^E$. In this situation, if E is a bifurcation divisor, we say that E is a contact divisor for π_C (or C).

Moreover, a bifurcation divisor E can belong to a dead arc only if it is a Puiseux divisor.

Take *E* a bifurcation divisor of G(C) and let $\{(m_1^E, n_1^E), (m_2^E, n_2^E), \ldots, (m_{q(E)}^E, n_{q(E)}^E)\}$ be the Puiseux pairs of an *E*-curvette γ_E , we denote

 $n_E = \begin{cases} n_{g(E)}, & \text{if } E \text{ is a Puiseux divisor,} \\ 1, & \text{otherwise,} \end{cases}$

and $\underline{n}_E = m(E)/n_E$. Observe that, if E belongs to a dead arc with terminal divisor F, then $m(F) = \underline{n}_E$. We define k_E to be

$$k_E = \begin{cases} g(E) - 1, & \text{if } E \text{ is a Puiseux divisor,} \\ g(E), & \text{if } E \text{ is a contact divisor.} \end{cases}$$

Hence we have that $\underline{n}_E = n_1^E \cdots n_{k_E}^E$.

Remark A.2. — Let E be a bifurcation divisor of G(C) which is a Puiseux divisor for C and take any $i \in I_E$ (that is, E belongs to the geodesic of the curve C_i). We have two possibilities concerning v(E):

- either $v(E) = \beta_{k_E+1}^i / \beta_0^i$, then we say that E is a Puiseux divisor for C_i ;
- or v(E) corresponds to the coincidence of C_i with another branch of C, in this situation we say that E is a contact divisor for C_i .

Note that, if E is a Puiseux divisor for C, then it is a Puiseux divisor for at least one irreducible component C_i with $i \in I_E$, but it can be a contact divisor for other branches C_j with $j \in I_E$, $j \neq i$. Consider for instance the

curve $C = C_1 \cup C_2$ with $C_1 = (y^2 - x^3 = 0)$ and $C_2 = (y - x^2 = 0)$. The dual graph G(C) is given by



Thus the divisor E_3 is a Puiseux divisor for C and C_1 but it is a contact divisor for C_2 since $v(E_3) = 3/2 = \mathcal{C}(C_1, C_2)$.

Example A.3. — If we consider a semiroot $C^{(k)}$ of a curve C as in Subsection 7.2, then all the bifurcation divisors E_1, E_2, \ldots, E_g are Puiseux divisors for C; the divisors E_1, E_2, \ldots, E_k are also Puiseux divisors for $C^{(k)}$ but E_{k+1} is a contact divisor for $C^{(k)}$.

A.2. Ramification

Consider a plane curve $C = \bigcup_{i=1}^{r} C_i$ in $(\mathbb{C}^2, 0)$. Let $\rho : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ be any *C*-ramification, that is, ρ is transversal to C and $\widetilde{C} = \rho^{-1}C$ has only non-singular irreducible components. Consequently, if the ramification is given by $x = u^n, y = v$, then it is required that $n \equiv 0 \mod (n^1, n^2, \dots, n^r)$ to ensure that \widetilde{C} has only non-singular irreducible components where $n^i = \nu_0(C_i)$. Each curve $\widetilde{C}_i = \rho^{-1}C_i$ has exactly n^i irreducible components and the number of irreducible components of \widetilde{C} is equal to $\nu_0(C) = n^1 + \dots + n^r$.

More precisely, let $y^i(x) = \sum_{l \ge n^i} a_l^i x^{l/n^i}$ be a Puiseux series of C_i , thus all its Puiseux series are given by

$$y_j^i(x) = \sum_{l \ge n^i} a_l^i \varepsilon_i^{lj} x^{l/n^i} \quad \text{for } j = 1, 2, \dots, n^i,$$

where ε_i is a primitive n^i -root of the unity. Then $f_i(x, y) = \prod_{l=1}^{n^i} (y - y_l^i(x))$ is a reduced equation of C_i . If we put $v_j^i(u) = y_j^i(u^n)$, then $v_j^i(u) \in \mathbb{C}\{u\}$ since $n/n^i \in \mathbb{N}$. Hence the curve $\sigma_j^i = (v - v_j^i(u) = 0)$ is non-singular and it is one of the irreducible components of $\rho^{-1}C_i$. Thus an equation of $\rho^{-1}C_i$ is given by

$$g_i(u,v) = f_i(u^n,v) = \prod_{l=1}^{n^i} (v - v_l^i(u)).$$

In particular we have that the irreducible components $\{\sigma_j^i\}_{j=1}^{n^i}$ of $\rho^{-1}C_i$ are in bijection with the Puiseux series of C_i .

ANNALES DE L'INSTITUT FOURIER

It is well-known that the equisingularity type of a curve C is determined by the characteristic exponents $\{\beta_0^i, \beta_1^i, \ldots, \beta_{g_i}^i\}_{i=1}^r$ of its irreducible components and the intersection multiplicities $\{(C_i, C_j)_0\}_{i\neq j}$. In [10] it is proved that the equisingularity data of C can be recovered from the curve $\rho^{-1}C$.

Let us explain now the relationship between the dual graphs G(C) and $G(\widetilde{C})$ of the minimal reduction of singularities of C and \widetilde{C} respectively. Observe that, if \widetilde{E} and \widetilde{E}' are two consecutive vertices of $G(\widetilde{C})$ with $\widetilde{E} < \widetilde{E}'$, then $v(\widetilde{E}') = v(\widetilde{E}) + 1$. Thus, $G(\widetilde{C})$ is completely determined once we know the bifurcation divisors, the order relations among them and the number of edges which leave from each bifurcation divisor.

Let K_i be the geodesic in G(C) of a branch C_i of C and let \widetilde{K}_i be the sub-graph of $G(\widetilde{C})$ corresponding to the geodesics of the irreducible components $\{\sigma_l^i\}_{l=1}^{n^i}$ of $\rho^{-1}C_i$. Let us explain how to construct \widetilde{K}_i from K_i . Denote by $B(\widetilde{K}_i)$ and $B(K_i)$ the bifurcation vertices of \widetilde{K}_i and K_i respectively. We say that a vertex \widetilde{E} of $B(\widetilde{K}_i)$ is associated to a vertex Eof $B(K_i)$ if $v(\widetilde{E}) = nv(E)$. Note that there can be other bifurcation vertices in $G(\widetilde{C}) \smallsetminus B(\widetilde{K}_i)$ with valuation equal to nv(E) but they are not associated to E.

Take a vertex E of $B(K_i)$. Consider first the case of E being the first bifurcation divisor of $B(K_i)$ and take E' its consecutive vertex in $B(K_i)$. Then E has only one associated vertex \tilde{E} in $B(\tilde{K}_i)$ and there are two possibilities for the number of edges which leave from it:

- If E is a Puiseux divisor for C_i , then there are n_1^i edges which leave from \widetilde{E} in \widetilde{K}_i ; then E' has n_1^i associated vertices in $B(\widetilde{K}_i)$.

Note that, if E is a Puiseux divisor for C, then E is a Puiseux divisor for at least one irreducible component C_i but it can be a contact divisor for all the other irreducible components (see Remark A.2). Let E now be any vertex of $B(K_i)$ and assume that we know the part of \tilde{K}_i corresponding to the vertices of K_i with valuation $\leq v(E)$. Then there are $\underline{n}_E = n_1^i \cdots n_{k_E}^i$ vertices $\{\tilde{E}^l\}_{l=1}^{\underline{n}_E}$ associated to E and

- If E is a Puiseux divisor for C_i , then there are $n_{k_E+1}^i$ edges which leave from each vertex \tilde{E}_l in \tilde{K}_i .
- If E is a contact divisor for C_i, then there is only one edge which leaves from each vertex *Ẽ_l* in *K̃_i*.

Nuria CORRAL

The dual graph $G(\tilde{C})$ is constructed by gluing the graphs \tilde{K}_i . Thus we deduced that, if \tilde{E} is a divisor of $G(\tilde{C})$ associated to a divisor E of G(C), then

$$(A.2) \quad b_{\tilde{E}} = \begin{cases} b_E, & \text{if } E \text{ is a contact divisor for } C, \\ (b_E - 1)n_E, & \text{if } E \text{ is a Puiseux divisor for } C \\ & \text{which belongs to a dead arc,} \\ (b_E - 1)n_E + 1, & \text{if } E \text{ is a Puiseux divisor for } C \\ & \text{which does not belong to a dead arc.} \end{cases}$$

Thus all vertices in $G(\widetilde{C})$ associated to a divisor E of G(C) have the same valence. Moreover, if γ_E is an E-curvette of a bifurcation divisor E of G(C), the curve $\rho^{-1}\gamma_E$ has $m(E) = \underline{n}_E n_E$ irreducible components which are all non-singular and each divisor \widetilde{E}^l belongs to the geodesic of exactly n_E branches of $\rho^{-1}\gamma_E$ which are curvettes of \widetilde{E}^l in different points.

Observe that there are non-bifurcation divisors of G(C) without associated divisors in $G(\widetilde{C})$.

Due to the bijection between the Puiseux series of C_i and the irreducible components of $\rho^{-1}C_i$, we have that the choice of a vertex $\tilde{E}^l \in B(\tilde{K}_i)$ associated to a bifurcation divisor E is equivalent to the choice of a \underline{n}_E -th root ξ_l of the unity. This implies that the vertex \tilde{E}^l belongs to the geodesic of $e_E^i = n^i/\underline{n}_E$ irreducible components $\{\sigma_{lt}^i\}_{t=1}^{e_E^i}$ of $\rho^{-1}C_i$. Moreover, the curve σ_{lt}^i is given by $(v - \eta_{lt}^i(u) = 0)$ where

$$\eta_{lt}^i(u) = \sum_{s \ge n^i} a_s^i(\zeta_{ilt})^s u^{sn/n^i}, \text{ for } t = 1, \dots, e_E^i,$$

and $\{\zeta_{ilt}\}_{t=1}^{e_E^i}$ are the e_E^i -th roots of ξ_l . The cardinal of the set $\pi_{\tilde{E}^l}^* \tilde{C}_i \cap \tilde{E}_{red}^l$ is given by

$$\sharp(\pi_{\tilde{E}^l}^* \tilde{C}_i \cap \tilde{E}_{\mathrm{red}}^l) = \begin{cases} n_E, & \text{if } E \text{ is a Puiseux divisor for } C_i, \\ 1, & \text{if } E \text{ is a contact divisor for } C_i. \end{cases}$$

Furthermore, if E is a Puiseux divisor for C_i and P, Q are two different points in $\pi^*_{\tilde{E}^l} \tilde{C}_i \cap \tilde{E}^l_{red}$, we have that

(A.3)
$$\nu_P(\pi^*_{\tilde{E}^l}\widetilde{C}_i) = \nu_Q(\pi^*_{\tilde{E}^l}\widetilde{C}_i) = \frac{e^i_E}{n_E};$$

ANNALES DE L'INSTITUT FOURIER

282

and if E is a contact divisor for C_i and we denote P the only point in the set $\pi^*_{\tilde{E}^l} \tilde{C}_i \cap \tilde{E}^l_{red}$, then

(A.4)
$$\nu_P(\pi^*_{\tilde{E}^i}\tilde{C}_i) = e^i_E.$$

Consider now two divisors \widetilde{E}^l and \widetilde{E}^k associated to the same bifurcation divisor E of G(C), and let ξ_l and ξ_k be the \underline{n}_E -th roots of the unity corresponding to the divisors \widetilde{E}^l and \widetilde{E}^k . We can define a bijection $\rho_{l,k}: \widetilde{E}^l_{\rm red} \to \widetilde{E}^k_{\rm red}$ as follows: the map $\rho_{l,k}$ sends the "infinity point" of $\widetilde{E}^l_{\rm red}$ (that is, the origin of the second chart of $\widetilde{E}^l_{\rm red}$) into the "infinity point" of $\widetilde{E}^k_{\rm red}$. Given another point P of $\widetilde{E}^l_{\rm red}$, we take an \widetilde{E}^l -curvette $\gamma^P_{\widetilde{E}^l} = (v - \psi^P_{\widetilde{E}^l}(u) = 0)$ with

(A.5)
$$\psi_{\tilde{E}^{l}}^{P}(u) = \sum_{i=1}^{v(\tilde{E}^{l})-1} a_{i}^{\tilde{E}^{l}} u^{i} + a_{v(\tilde{E}^{l})}^{P} u^{v(\tilde{E}^{l})},$$

and such that $\pi_{\tilde{E}^l}^* \gamma_{\tilde{E}^l}^P \cap \tilde{E}_{red}^l = \{P\}$. We define $\rho_{l,k}(P)$ to be the point $\pi_{\tilde{E}^k}^* \gamma_{\tilde{E}^k}^{\rho_{l,k}(P)} \cap \tilde{E}_{red}^k$, where the curve $\gamma_{\tilde{E}^k}^{\rho_{l,k}(P)} = (v - \psi_{\tilde{E}^k}^{\rho_{l,k}(P)}(u) = 0)$ is given by

$$\psi_{\tilde{E}^{k}}^{\rho_{l,k}(P)}(u) = \sum_{i=1}^{v(\tilde{E}^{l})-1} a_{i}^{\tilde{E}^{l}} \left(\frac{\xi_{k}}{\xi_{l}}\right)^{i} u^{i} + a_{v(\tilde{E}^{l})}^{P} \left(\frac{\xi_{k}}{\xi_{l}}\right)^{v(\tilde{E}^{l})} u^{v(\tilde{E}^{l})}.$$

Note that $\gamma_{\widetilde{E}^k}^{\rho_{l,k}(P)}$ is an \widetilde{E}^k -curvette.

Recall also that given any bifurcation divisor E of G(C) or $E = E^1$ and any of its associated divisors \tilde{E}^l in $G(\tilde{C})$, there is a morphism $\rho_{\tilde{E}^l,E}$: $\tilde{E}^l_{\text{red}} \to E_{\text{red}}$ which is a ramification of order n_E (see [10, Lemma 8]). The map $\rho_{\tilde{E}^l,E}$ is defined as follows: $\rho_{\tilde{E}^l,E}$ sends the "infinity point" of \tilde{E}^l_{red} into the "infinity point" of E_{red} and the origin of the first chart of \tilde{E}^l_{red} is sent to the origin of the first chart of E_{red} . For any other point P of \tilde{E}^l_{red} , we can take an \tilde{E}^l -curvette $\gamma_{\tilde{E}^l}^P = (v - \psi_{\tilde{E}^l}^P(u) = 0)$ with $\pi^*_{\tilde{E}^l}\gamma_{\tilde{E}^l}^P \cap \tilde{E}^l_{\text{red}} = \{P\}$. Thus if $\psi_{\tilde{E}^l}^P(u)$ is given by (A.5), we can consider the E-curvette γ_E^P given by the Puiseux series

$$y^{P}(x) = \sum_{i=1}^{v(\tilde{E}^{l})-1} a_{i}^{\tilde{E}^{l}} x^{i/m(E)} + a_{v(\tilde{E}^{l})}^{P} x^{v(\tilde{E}^{l})/m(E)},$$

and $\rho_{\tilde{E}^l,E}(P)$ is the only point $\pi_E^* \gamma_E^P \cap E_{\text{red}}$. Observe that, if E is a bifurcation divisor in the geodesic of a branch C_i of C and \tilde{E}^l is any associated divisor to E in $G(\tilde{C})$, then the morphism $\rho_{\tilde{E}^l,E}$ maps all the points in $\pi_{\tilde{E}^l}^* \tilde{C}_i \cap \tilde{E}_{\text{red}}^l$ to the only point in $\pi_E^* C_i \cap E_{\text{red}}$.

Moreover, note that, if \tilde{E}^l and \tilde{E}^k are two divisors of $G(\tilde{C})$ associated to a bifurcation divisor E of G(C), then the following diagram



is commutative.

Finally remark that, if γ_{E_t} is a curvette of a terminal divisor E_t of a dead arc with bifurcation divisor E, then $\rho^{-1}\gamma_{E_t}$ is composed by $m(E_t) = \underline{n}_E$ non-singular irreducible components and each divisor \widetilde{E}^l belongs to the geodesic of exactly one branch of $\rho^{-1}\gamma_{E_t}$, where $\{\widetilde{E}^l\}_{l=1}^{\underline{n}_E}$ are the divisors associated to E in $G(\widetilde{C})$.

A.3. Logarithmic foliations and ramification

Consider the logarithmic foliation $\mathcal{L}_{\lambda}^{C}$ defined by

$$f_1 \cdots f_r \sum_{i=1}^r \lambda_i \frac{\mathrm{d}f_i}{f_i} = 0$$

with $\lambda = (\lambda_1, \ldots, \lambda_r) \in \mathbb{C}^r$ and $f_i \in \mathbb{C}\{x, y\}$ (see Section 3 for notations concerning logarithmic foliations). Let us see the behaviour of \mathcal{L}^C_{λ} after a ramification.

Consider the curve $C = \bigcup_{i=1}^{r} C_i$ with $C_i = (f_i = 0)$ and let $\rho : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ be any *C*-ramification, that is, ρ is transversal to *C* and the curve $\widetilde{C} = \rho^{-1}C$ has only non-singular irreducible components. We refer to Subsections 2.3, A.1 and A.2 for notations concerning equisingularity data of curves and ramifications.

We have that $\rho^* \mathcal{L}^C_{\lambda} = \mathcal{L}^{\tilde{C}}_{\lambda^*}$ with

$$\lambda^* = (\overbrace{\lambda_1, \dots, \lambda_1}^{n^1}, \dots, \overbrace{\lambda_r, \dots, \lambda_r}^{n^r})$$

where $n^i = \nu_0(C_i)$ for i = 1, ..., r. We put $\rho^{-1}C_i = \widetilde{C}_i = \{\sigma_t^i\}_{t=1}^{n^i}$ where each σ_t^i is an irreducible curve. Moreover, we have that logarithmic models behave well under ramification. More precisely,

PROPOSITION A.4 (see [9]). — Let $\mathcal{F} \in \mathbb{G}_C$ and \mathcal{L}^C_{λ} a logarithmic model of \mathcal{F} . If $\rho : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ is a *C*-ramification, then $\rho^* \mathcal{L}^C_{\lambda}$ is a logarithmic model of $\rho^* \mathcal{F}$.

Let $\pi_C : X_C \to (\mathbb{C}^2, 0)$ be the minimal reduction of singularities of Cand $\pi_{\tilde{C}} : X_{\tilde{C}} \to (\mathbb{C}^2, 0)$ be the minimal reduction of singularities of \tilde{C} . Take E a bifurcation divisor of G(C) and let \tilde{E}^l be any bifurcation divisor of $G(\tilde{C})$ associated to E. Given any $i \in I_E$ (that is, E is in the geodesic of C_i), there are e_E^i branches of $\rho^{-1}C_i$ such that \tilde{E}^l belongs to their geodesics where $e_E^i = n^i/\underline{n}_E$ and we have that the residue of the logarithmic foliation $\mathcal{L}_{\lambda^*}^{\tilde{C}}$ along the divisor \tilde{E}^l is given by

$$\kappa_{\tilde{E}^l}(\mathcal{L}_{\lambda^*}^{\tilde{C}}) = \sum_{i=1}^r \lambda_i \sum_{t=1}^{n^i} \sum_{\tilde{E} \leqslant \tilde{E}^l} \varepsilon_{\tilde{E}}^{\sigma_t^i}.$$

(see (3.2) for its definition). Let $\{R_1^{\tilde{E}^l}, R_2^{\tilde{E}^l}, \dots, R_{b_{\tilde{E}^l}}^{\tilde{E}^l}\}$ be the set of points $\pi_{\tilde{E}^l}^* \tilde{C} \cap \tilde{E}_{red}^l$ and put $m_{R_t^{\tilde{E}^l}}^i = \nu_{R_t^{\tilde{E}^l}}(\pi_{\tilde{E}^l}^* \tilde{C}_i)$ for $t = 1, 2, \dots, b_{\tilde{E}^l}$. Note that $m_{R_t^{\tilde{E}^l}}^i = \sharp\{s \in \{1, \dots, n^i\} : \pi_{\tilde{E}^l}^* \sigma_s^i \cap \tilde{E}_{red}^l = \{R_t^{\tilde{E}^l}\}\} = \frac{e_{\tilde{E}}^i}{n_E}$ (the last equality follows from equations (A.3) and (A.4) in Appendix A.2 where $m_{R_t^{\tilde{E}^l}}^i$ is also computed). With these notations we have that

(A.6)
$$\mathcal{I}_{R_{t}^{\tilde{E}^{l}}}(\pi_{\tilde{E}^{l}}^{*}\mathcal{L}_{\lambda^{*}}^{\tilde{C}}, \widetilde{E}_{\mathrm{red}}^{l}) = -\frac{\sum_{i \in I_{E}} \lambda_{i} m_{R_{t}^{\tilde{E}^{l}}}^{i}}{\kappa_{\tilde{E}^{l}}(\mathcal{L}_{\lambda^{*}}^{\tilde{C}})}.$$

Observe that if \widetilde{E}^{l} and \widetilde{E}^{k} are two bifurcation divisors of $G(\widetilde{C})$ associated to the same divisor E of G(C), we have that $\kappa_{\tilde{E}^{l}}(\mathcal{L}_{\lambda^{*}}^{\tilde{C}}) = \kappa_{\tilde{E}^{k}}(\mathcal{L}_{\lambda^{*}}^{\tilde{C}})$. Moreover, there is a bijection between the sets of points $\pi_{\tilde{E}^{l}}^{*}\widetilde{C} \cap \widetilde{E}_{\mathrm{red}}^{l}$ and $\pi_{\tilde{E}^{k}}^{*}\widetilde{C} \cap \widetilde{E}_{\mathrm{red}}^{k}$ induced by the map $\rho_{l,k} : \widetilde{E}_{\mathrm{red}}^{l} \to \widetilde{E}_{\mathrm{red}}^{k}$ (see Appendix A.2). Hence, if $\{R_{1}^{\tilde{E}^{k}}, R_{2}^{\tilde{E}^{k}}, \ldots, R_{b_{\tilde{E}^{k}}}^{\tilde{E}^{k}}\}$ is the set of points $\pi_{\tilde{E}^{k}}^{*}\widetilde{C} \cap \widetilde{E}_{\mathrm{red}}^{k}$ with $R_{t}^{\tilde{E}^{k}} = \rho_{l,k}(R_{t}^{\tilde{E}^{l}})$ for $t = 1, 2, \ldots, b_{\tilde{E}^{k}}$, we have that

(A.7)
$$\mathcal{I}_{R_t^{\tilde{E}^k}}(\pi_{\tilde{E}^k}^*\mathcal{L}_{\lambda^*}^{\tilde{C}}, \widetilde{E}_{\mathrm{red}}^k) = \mathcal{I}_{R_t^{\tilde{E}^l}}(\pi_{\tilde{E}^l}^*\mathcal{L}_{\lambda^*}^{\tilde{C}}, \widetilde{E}_{\mathrm{red}}^l), \quad t = 1, 2, \dots, b_{\tilde{E}^k}.$$

Moreover, if $R_t^{\tilde{E}^l}, R_s^{\tilde{E}^l}$ are two points in $\pi_{\tilde{E}^l}^* \widetilde{C} \cap \widetilde{E}_{red}^l$ with $\rho_{\tilde{E}^l, E}(R_t^{\tilde{E}^l}) = \rho_{\tilde{E}^l, E}(R_s^{\tilde{E}^l})$ where $\rho_{\tilde{E}^l, E}: \widetilde{E}_{red}^l \to E_{red}$ is the ramification defined in Appendix A.2, then

(A.8)
$$\mathcal{I}_{R_t^{\tilde{E}^l}}(\pi_{\tilde{E}^l}^*\mathcal{L}_{\lambda^*}^{\tilde{C}}, \widetilde{E}_{\mathrm{red}}^l) = \mathcal{I}_{R_s^{\tilde{E}^l}}(\pi_{\tilde{E}^l}^*\mathcal{L}_{\lambda^*}^{\tilde{C}}, \widetilde{E}_{\mathrm{red}}^l)$$

since $m_{R_t^{\tilde{E}l}}^i = m_{R_s^{\tilde{E}l}}^i = \frac{e_E^i}{n_E}$ for $i \in I_E$ by equations (A.3) and (A.4).

TOME 75 (2025), FASCICULE 1

Appendix B. Intersection multiplicities

We state now two results concerning the intersection multiplicity of the jacobian curve of two foliations either with a single separatrix of one of the foliations and with the curve of all separatrices. These intersection multiplicities are computed in terms of local invariants of the foliations (see Subsection 2.1 for notations). Consider two foliations \mathcal{F} and \mathcal{G} in (\mathbb{C}^2 , 0) and denote by $\mathcal{J}_{\mathcal{F},\mathcal{G}}$ the jacobian curve of \mathcal{F} and \mathcal{G} .

PROPOSITION B.1. — Assume that \mathcal{F} and \mathcal{G} have no common separatrix. If S is an irreducible separatrix of \mathcal{F} , we have that

$$(\mathcal{J}_{\mathcal{F},\mathcal{G}},S)_0 = \mu_0(\mathcal{F},S) + \tau_0(\mathcal{G},S).$$

Proof. — Let us write $\omega = A(x, y) dx + B(x, y) dy$ and $\eta = P(x, y) dx + Q(x, y) dy$ the 1-forms defining \mathcal{F} and \mathcal{G} respectively. Let $\gamma(t) = (x(t), y(t))$ be a parametrization of the curve S. We can assume, without loss of generality, that $x(t) \neq 0$ and thus $\dot{x}(t) \neq 0$. Since S is a separatrix of \mathcal{F} , then $A(\gamma(t))\dot{x}(t) + B(\gamma(t))\dot{y}(t) = 0$. Thus, we have that

$$\begin{aligned} (\mathcal{J}_{\mathcal{F},\mathcal{G}},S)_0 &= \operatorname{ord}_t \{ A(\gamma(t))Q(\gamma(t)) - B(\gamma(t))P(\gamma(t)) \} \\ &= \operatorname{ord}_t \left\{ \frac{-B(\gamma(t))\dot{y}(t)}{\dot{x}(t)}Q(\gamma(t)) - B(\gamma(t))P(\gamma(t)) \right\} \\ &= \operatorname{ord}_t(B(\gamma(t))) - (\operatorname{ord}_t(x(t)) - 1) \\ &+ \operatorname{ord}_t\{P(\gamma(t))\dot{x}(t) + Q(\gamma(t))\dot{y}(t)\} \\ &= \mu_0(\mathcal{F},S) + \tau_0(\mathcal{G},S) \end{aligned}$$

where the last equality comes from the expression of $\mu_0(\mathcal{F}, S)$ given in (2.1) and the definition of $\tau_0(\mathcal{G}, S)$ given in (2.2).

When \mathcal{G} is a non-singular foliation, we obtain Proposition 1 in [6] for the polar intersection number with respect to a branch of the curve of separatrices of \mathcal{F} . Note that, although in [6] it is assumed that the foliation \mathcal{F} is non-dicritical, the result is also true when \mathcal{F} is a dicritical foliation. Using property (iv) in Theorem 2.1, we get following consequence of the above result:

COROLLARY B.2. — If \mathcal{G} is a non-dicritical second type foliation and $S_{\mathcal{G}}$ is the curve of separatrices of \mathcal{G} , we have that

$$(\mathcal{J}_{\mathcal{F},\mathcal{G}},S)_0 = \mu_0(\mathcal{F},S) + (S_{\mathcal{G}},S)_0 - 1.$$

Next result gives a relationship among the intersection multiplicities of the jacobian curve with the curves of separatrices and the Milnor number of the foliations. PROPOSITION B.3. — Consider two non-dicritical second type foliations \mathcal{F} and \mathcal{G} without common separatrices. Thus

$$(\mathcal{J}_{\mathcal{F},\mathcal{G}},S_{\mathcal{F}})_0 - (\mathcal{J}_{\mathcal{F},\mathcal{G}},S_{\mathcal{G}})_0 = \mu_0(\mathcal{F}) - \mu_0(\mathcal{G}),$$

where $S_{\mathcal{F}}$, $S_{\mathcal{G}}$ are the curves of separatrices of \mathcal{F} and \mathcal{G} respectively.

Proof. — Let $\mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})$ be the set of irreducible components of $\mathcal{J}_{\mathcal{F},\mathcal{G}}$. Given any branch $\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})$, we denote by $\gamma_{\Gamma}(t) = (x_{\Gamma}(t), y_{\Gamma}(t))$ any primitive parametrization of Γ . Assume that the foliations \mathcal{F} and \mathcal{G} are defined by the 1-forms $\omega = A \, dx + B \, dy$ and $\eta = P \, dx + Q \, dy$ respectively. Thus we have that $A(\gamma_{\Gamma}(t))Q(\gamma_{\Gamma}(t)) - B(\gamma_{\Gamma}(t))P(\gamma_{\Gamma}(t)) = 0$. Since Γ is not a separatrix of \mathcal{G} , then either $Q(\gamma_{\Gamma}(t)) \neq 0$ or $P(\gamma_{\Gamma}(t)) \neq 0$. We will assume that $Q(\gamma_{\Gamma}(t)) \neq 0$. Let us compute the intersection multiplicity of $(\mathcal{J}_{\mathcal{F},\mathcal{G}}, S_{\mathcal{F}})_0$ taking into account property (iv) in Theorem 2.1:

$$\begin{split} (\mathcal{J}_{\mathcal{F},\mathcal{G}},S_{\mathcal{F}})_{0} &= \sum_{\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})} (\Gamma,S_{\mathcal{F}})_{0} = \sum_{\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})} (\tau_{0}(\mathcal{F},\Gamma)+1) \\ &= \sum_{\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})} (\operatorname{ord}_{t} \left\{ A(\gamma_{\Gamma}(t))\dot{x}_{\Gamma}(t) + B(\gamma_{\Gamma}(t))\dot{y}_{\Gamma}(t) \right\} + 1) \\ &= \sum_{\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})} \left(\operatorname{ord}_{t} \left\{ \frac{B(\gamma_{\Gamma}(t))P(\gamma_{\Gamma}(t))}{Q(\gamma_{\Gamma}(t))} \dot{x}_{\Gamma}(t) + B(\gamma_{\Gamma}(t))\dot{y}_{\Gamma}(t) \right\} + 1 \right) \\ &= \sum_{\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})} \operatorname{ord}_{t} \{B(\gamma_{\Gamma}(t))\} - \sum_{\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})} \operatorname{ord}_{t} \{Q(\gamma_{\Gamma}(t))\} \\ &+ \sum_{\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})} (\operatorname{ord}_{t} \{P(\gamma_{\Gamma}(t))\dot{x}_{\Gamma}(t) + Q(\gamma_{\Gamma}(t))\dot{y}_{\Gamma}(t)\} + 1) \\ &= \mu_{0}(\mathcal{F}) - \mu_{0}(\mathcal{G}) + \sum_{\Gamma \in \mathcal{B}(\mathcal{J}_{\mathcal{F},\mathcal{G}})} (\tau_{0}(\mathcal{G},\Gamma) + 1) \\ &= \mu_{0}(\mathcal{F}) - \mu_{0}(\mathcal{G}) + (\mathcal{J}_{\mathcal{F},\mathcal{G}},S_{\mathcal{G}})_{0}. \end{split}$$

BIBLIOGRAPHY

- S. S. ABHYANKAR & T. T. MOH, "Newton-Puiseux expansion and generalized Tschirnhausen transformation. I, II", J. Reine Angew. Math. 260 (1973), p. 47-83, ibid. 261 (1973), p. 29–54.
- [2] M. ALBERICH-CARRAMIÑANA & V. GONZÁLEZ-ALONSO, "Determining plane curve singularities from its polars", Adv. Math. 287 (2016), p. 788-822.
- [3] C. CAMACHO, A. LINS NETO & P. SAD, "Topological invariants and equidesingularization for holomorphic vector fields", J. Differ. Geom. 20 (1984), no. 1, p. 143-174.
- [4] C. CAMACHO & P. SAD, "Invariant varieties through singularities of holomorphic vector fields", Ann. Math. 115 (1982), no. 3, p. 579-595.

- [5] F. CANO, D. CERVEAU & J. DÉSERTI, Théorie élémentaire des feuilletages holomorphes singuliers, Collection Échelles, Belin, 2013.
- [6] F. CANO, N. CORRAL & R. MOL, "Local polar invariants for plane singular foliations", Expo. Math. 37 (2019), no. 2, p. 145-164.
- [7] E. CASAS-ALVERO, Singularities of plane curves, London Mathematical Society Lecture Note Series, vol. 276, Cambridge University Press, 2000, xvi+345 pages.
- [8] —, "Local geometry of planar analytic morphisms", Asian J. Math. 11 (2007), no. 3, p. 373-426.
- [9] N. CORRAL, "Sur la topologie des courbes polaires de certains feuilletages singuliers", Ann. Inst. Fourier 53 (2003), no. 3, p. 787-814.
- [10] _____, "Infinitesimal adjunction and polar curves", Bull. Braz. Math. Soc. (N.S.) 40 (2009), no. 2, p. 181-224.
- [11] ——, "Infinitesimal initial part of a singular foliation", An. Acad. Brasil. Ciênc. 81 (2009), no. 4, p. 633-640.
- [12] , "Polar pencil of curves and foliations", in Équations différentielles et singularités. En l'honneur de J. M. Aroca (F. Cano, F. Loray, J. J. Moralez-Ruiz, P. Sad & M. Spivakovsky, eds.), Astérisque, no. 323, Société Mathématique de France, 2009.
- [13] E. R. GARCÍA BARROSO, "Sur les courbes polaires d'une courbe plane réduite", Proc. Lond. Math. Soc. 81 (2000), no. 1, p. 1-28.
- [14] E. R. GARCÍA BARROSO & J. GWOŹDZIEWICZ, "On the approximate Jacobian Newton diagrams of an irreducible plane curve", J. Math. Soc. Japan 65 (2013), no. 1, p. 169-182.
- [15] Y. GENZMER & R. MOL, "Local polar invariants and the Poincaré problem in the dicritical case", J. Math. Soc. Japan 70 (2018), no. 4, p. 1419-1451.
- [16] O. GÓMEZ-MARTÍNEZ, "Foliaciones dicríticas en la realización de invariantes analíticos de curvas singulares", PhD Thesis, Universidad Nacional Autónoma de México, 2021.
- [17] J. GWOŹDZIEWICZ & A. PLOSKI, "On the approximate roots of polynomials", Ann. Pol. Math. 60 (1995), no. 3, p. 199-210.
- [18] A. HEFEZ, M. E. HERNANDES & M. F. H. IGLESIAS, "On the factorization of the polar of a plane branch", in *Singularities and foliations. geometry, topology and applications*, Springer Proceedings in Mathematics & Statistics, vol. 222, Springer, 2018, p. 347-362.
- [19] T.-C. KUO & A. PARUSIŃSKI, "On Puiseux roots of Jacobians", Proc. Japan Acad., Ser. A 78 (2002), no. 5, p. 55-59.
- [20] —, "Newton-Puiseux roots of Jacobian determinants", J. Algebr. Geom. 13 (2004), no. 3, p. 579-601.
- [21] D. T. LÊ, F. MICHEL & C. WEBER, "Sur le comportement des polaires associées aux germes de courbes planes", Compos. Math. 72 (1989), no. 1, p. 87-113.
- [22] J.-F. MATTEI & E. SALEM, "Modules formels locaux de feuilletages holomorphes", https://arxiv.org/abs/math/0402256, 2004.
- [23] H. MAUGENDRE, "Discriminant d'un germe (g, f): $(\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$ et quotients de contact dans la résolution de $f \cdot g$ ", Ann. Fac. Sci. Toulouse, Math. 7 (1998), no. 3, p. 497-525.
- [24] _____, "Discriminant of a germ Φ : ($\mathbf{C}^2, 0$) \rightarrow ($\mathbf{C}^2, 0$) and Seifert fibred manifolds", J. Lond. Math. Soc. **59** (1999), no. 1, p. 207-226.
- [25] M. MERLE, "Invariants polaires des courbes planes", Invent. Math. 41 (1977), no. 2, p. 103-111.

- [26] L. ORTIZ-BOBADILLA, E. ROSALES-GONZÁLEZ & S. M. VORONIN, "Rigidity theorems for generic holomorphic germs of dicritic foliations and vector fields in (C², 0)", Mosc. Math. J. 5 (2005), no. 1, p. 171-206.
- [27] E. PAUL, "Classification topologique des germes de formes logarithmiques génériques", Ann. Inst. Fourier 39 (1989), no. 4, p. 909-927.
- [28] _____, "Cycles évanescents d'une fonction de Liouville de type $f_1^{\lambda_1} \cdots f_p^{\lambda_p}$ ", Ann. Inst. Fourier **45** (1995), no. 1, p. 31-63.
- [29] P. POPESCU-PAMPU, "Approximate roots", in Valuation theory and its applications, Vol. II (Saskatoon, SK, 1999), Fields Institute Communications, vol. 33, American Mathematical Society, 2003, p. 285-321.
- [30] P. ROUILLÉ, "Théorème de Merle: cas des 1-formes de type courbes généralisées", Bol. Soc. Bras. Mat., Nova Sér. 30 (1999), no. 3, p. 293-314.
- [31] N. E. SARAVIA, "Curva polar de una foliación asociada a sus raíces aproximadas", PhD Thesis, Pontificia Universidad Católica del Perú, 2018.
- [32] A. SEIDENBERG, "Reduction of singularities of the differential equation A dy = B dx", Am. J. Math. **90** (1968), p. 248-269.

Manuscrit reçu le 12 juillet 2021, révisé le 21 décembre 2022, accepté le 15 mai 2023.

Nuria CORRAL Departamento de Matemáticas, Estadística y Computación, Universidad de Cantabria, Avda. de los Castros s/n, 39005 – Santander (Spain) nuria.corral@unican.es