

ANNALES DE L'institut fourier

Sébastien BOUCKSOM & Mattias JONSSON A non-Archimedean approach to K-stability, I: Metric geometry of spaces of test configurations and valuations Tome 75, n° 2 (2025), p. 829-927.

https://doi.org/10.5802/aif.3668

Article mis à disposition par ses auteurs selon les termes de la licence CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE [CC] BY-ND http://creativecommons.org/licenses/by-nd/3.0/fr/



Les Annales de l'Institut Fourier sont membres du Centre Mersenne pour l'édition scientifique ouverte www.centre-mersenne.org e-ISSN : 1777-5310

A NON-ARCHIMEDEAN APPROACH TO K-STABILITY, I: METRIC GEOMETRY OF SPACES OF TEST CONFIGURATIONS AND VALUATIONS

by Sébastien BOUCKSOM & Mattias JONSSON (*)

ABSTRACT. — For any polarized variety (X, L), we show that test configurations and, more generally, \mathbb{R} -test configurations (defined as finitely generated filtrations of the section ring) can be analyzed in terms of Fubini–Study functions on the Berkovich analytification of X with respect to the trivial absolute value on the ground field. Building on non-Archimedean pluripotential theory, we describe the (Hausdorff) completion of the space of test configurations, with respect to two natural pseudo-metrics, in terms of plurisubharmonic functions and measures of finite energy on the Berkovich space. We also describe the Hausdorff quotient of the space of all filtrations, and establish a 1–1 correspondence between divisorial norms and divisorial measures, both being determined in terms of finitely many divisorial valuations.

RÉSUMÉ. — Pour toute variété polarisée (X, L), nous montrons que les configurations test, et plus généralement les \mathbb{R} -configurations test (définies comme filtrations de type fini sur l'anneau des sections), peuvent être analysées en terme de fonctions de Fubini–Study sur l'analytifié de Berkovich de X pour la valuation triviale sur le corps de base. En s'appuyant sur la théorie du pluripotentiel non-archimédien, nous décrivons le complété (séparé) de l'espace des configurations test, relativement à deux pseudo-distances naturelles, en terme de fonctions plurisousharmoniques et de mesures d'énergie finie sur l'espace de Berkovich. Nous décrivons également le quotient séparé de l'espace de toutes les filtrations, et établissons une correspondance bijective entre normes divisorielles et mesures divisorielles, toutes deux déterminées par un nombre fini de valuations divisorielles.

Introduction

The notion of K-stability was introduced in complex differential geometry as a conjectural, and now partially confirmed, algebro-geometric criterion

 $K\!eywords\!:$ K-stability, filtrations, valuations, test configurations, non-Archimedean geometry.

²⁰²⁰ Mathematics Subject Classification: 32Q26, 32P05.

^(*) The first author was partially supported by the ANR grant GRACK. The second author was partially supported by NSF grants DMS-1600011, DMS-1900025, DMS-2154380, and the United States–Israel Binational Science Foundation.

for the existence of special Kähler metrics. Lately, it has also become a subject in its own respect. In a series of two (largely independent) papers of which this is the first, we show how global pluripotential theory over a trivially valued field, as developed in [18], can be used to study K-stability.

Let X be a projective variety (reduced and irreducible) of dimension $n \ge 1$ over an algebraically closed field k (assumed to be of characteristic 0 in this introduction), and L an ample Q-line bundle on X. The definition of K-stability of the polarized variety (X, L), as given by Donaldson [39], involves the sign of an invariant attached to (ample) test configurations for (X, L). As shown in [17, 18], test configurations can be alternatively understood in terms of (rational) Fubini–Study functions on the Berkovich analytification X^{an} , and uniform K-stability becomes a linear growth condition for the non-Archimedean K-energy on the set of such functions.

Filtrations of the section ring of (X, L) provide another, widely used description of test configurations; more precisely, the latter correspond to \mathbb{Z} filtrations of finite type [65, 68]. Recent works related to the Hamilton–Tian conjecture [8, 29, 38, 49] have emphasized the importance of considering more general \mathbb{R} -test configurations, defined as \mathbb{R} -filtrations of finite type, and one first objective of this paper is to show that these can again be understood as (real) Fubini–Study functions on X^{an} .

On the other hand, Chi Li's recent breakthrough on the Yau–Tian– Donaldson conjecture for cscK metrics [55] (based in part on the first version of the present paper) involves a stronger form of uniform K-stability, formulated as a linear growth condition for the K-energy on the space of functions of finite energy on X^{an} . The latter are obtained as limits of Fubini–Study functions, and are the central topic of pluripotential theory on X^{an} [18]. Building on the latter technology, the second objective of this paper is to show how functions and measures of finite energy can be used to describe the completion of the space of test configurations with respect to a natural metric, leading to a picture that is quite similar to the welldeveloped complex analytic case [32, 34].

From test configurations to Fubini–Study functions

Denote by $\mathcal{N}_{\mathbb{R}}$ the space of (decreasing, left-continuous, separated, exhaustive) filtrations of the section algebra $R^{(d)} = R(X, dL)$ for d sufficiently divisible. It is convenient to view these as norms $\chi \colon R^{(d)} \to \mathbb{R} \cup \{+\infty\}$, for which we use "additive" terminology, see Section 1.1. A norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is of finite type if the associated graded algebra is finitely generated. For

any subgroup $\Lambda \subset \mathbb{R}$, let $\mathcal{N}_{\Lambda} \subset \mathcal{N}_{\mathbb{R}}$ be the set of norms with values in $\Lambda \cup \{+\infty\}$, and denote by

$$\mathcal{T}_{\mathbb{R}} \subset \mathcal{N}_{\mathbb{R}} \quad ext{and} \quad \mathcal{T}_{\Lambda} \coloneqq \mathcal{T}_{\mathbb{R}} \cap \mathcal{N}_{\Lambda}$$

the subsets of norms of finite type.

As is by now well-known (see [17, 65, 68]), the Rees construction provides a 1–1 correspondence between $\mathcal{T}_{\mathbb{Z}}$ and the set of (ample) test configurations for (X, L). In line with [38], we view $\mathcal{T}_{\mathbb{R}}$ as the space of \mathbb{R} -test configurations. Any $\chi \in \mathcal{T}_{\mathbb{R}}$ lies in \mathcal{T}_{Λ} for some finitely generated subgroup $\Lambda \simeq \mathbb{Z}^r$, and χ can be geometrically realized as a $\mathbb{G}_{\mathrm{m}}^r$ -equivariant degeneration of (X, L)to a polarized scheme (see Section A.3), which is further reduced iff χ is homogeneous, in the sense that $\chi(s^d) = d\chi(s)$ for all $d \in \mathbb{N}$.

The space $\mathcal{N}_{\mathbb{R}}$ comes equipped with a non-decreasing family $(d_p)_{1 \leq p \leq \infty}$ of natural pseudo-metrics. By [14], the space $\mathcal{N}_{\mathbb{R}}(V)$ of norms on any finite dimensional vector space V is indeed endowed with a metric d_p for any $p \in [1, \infty]$, the distance between two norms being the ℓ^p -length of their relative spectrum, defined by joint diagonalization in some basis. For p = 2, this is the classical Tits metric of $\mathcal{N}_{\mathbb{R}}(V)$ as a Euclidean building, whose relevance to K-stability was already emphasized in [30, 61].

Any $\chi \in \mathcal{N}_{\mathbb{R}}$ restricts to a norm on $R_m := \mathrm{H}^0(X, mL)$ for all m sufficiently divisible, and we define the pseudometric d_p on $\mathcal{N}_{\mathbb{R}}$ by setting

$$\mathbf{d}_p(\chi, \chi') \coloneqq \limsup_m m^{-1} \mathbf{d}_p(\chi|_{R_m}, \chi'|_{R_m}),$$

where the limsup is actually a limit for $p < \infty$, by [13, 17, 27]. The L^p -norm of a test configuration in $\mathcal{T}_{\mathbb{Z}}$, as in [17, 39, 65], can be computed using d_p .

The pseudo-metric d_p is not a metric, even after restriction to \mathcal{T}_{Λ} , and our first main result describes the Hausdorff quotient of $(\mathcal{T}_{\Lambda}, d_p)$ as a natural space of functions on the Berkovich analytification of X (with respect to the trivial absolute value on k). Recall that the latter is a compact Hausdorff topological space X^{an} , whose elements are semivaluations v on X, i.e. \mathbb{R} valued valuations on the function field of some subvariety of X, trivial on k. The space X^{an} contains as a dense subset the space X^{div} of divisorial valuations on X, induced (up to scaling) by a prime divisor on a birational model of X.

For any $v \in X^{\mathrm{an}}$ and any section s of a line bundle on X, we can define $v(s) \in [0, +\infty]$ by trivializing the line bundle at the center of v, and setting $|s|(v) \coloneqq e^{-v(s)}$ defines a continuous function $|s|: X^{\mathrm{an}} \to [0, 1]$. Given a subgroup $\Lambda \subset \mathbb{R}$, a Λ -Fubini–Study function for L is a function

 $\varphi \in \mathbf{C}^0 = \mathbf{C}^0(X^{\mathrm{an}})$ of the form

$$\varphi = \frac{1}{m} \max_{j} \{ \log |s_j| + \lambda_j \},\$$

where $m \ge 1$ is sufficiently divisible, (s_j) is a finite set of sections of mL without common zeroes, and $\lambda_j \in \Lambda$.

The set $\mathcal{H}_{\Lambda} \subset C^0$ of Λ -Fubini–Study functions is stable under max and the action of Λ by translation, and satisfies $\mathcal{H}_{\Lambda} = \mathcal{H}_{\mathbb{Q}\Lambda}$. It is related to the space \mathcal{T}_{Λ} of Λ -test configurations by the *Fubini–Study operator*, a surjective map

$$FS: \mathcal{T}_{\Lambda} \longrightarrow \mathcal{H}_{\Lambda}$$

that associates to $\chi \in \mathcal{T}_{\Lambda}$ the Fubini–Study function

$$FS(\chi) = m^{-1} \max_{j} \{ \log |s_j| + \chi(s_j) \},\$$

where (s_j) is any χ -orthogonal basis of R_m for m sufficiently divisible. Viewed as a map from (usual) test configurations to Fubini–Study functions, FS: $\mathcal{T}_{\mathbb{Z}} \to \mathcal{H}_{\mathbb{Z}} = \mathcal{H}_{\mathbb{Q}}$ is compatible with the one constructed and studied in [17, 18] (see Appendix A).

THEOREM A. — For any polarized variety (X, L), any subgroup $\Lambda \subset \mathbb{R}$ and $p \in [1, \infty]$, the Fubini–Study operator identifies the Hausdorff quotient of the pseudo-metric space $(\mathcal{T}_{\Lambda}, d_p)$ with \mathcal{H}_{Λ} . For $p = \infty$, the induced metric d_{∞} on \mathcal{H}_{Λ} further coincides with the supnorm metric.

It is enough to prove this for $\Lambda = \mathbb{R}$. Let us first describe the case $p = \infty$. The restrictions $\chi|_{R_d}$ of any norm $\chi \in \mathcal{N}_{\mathbb{R}}$ generate a sequence of canonical approximants $\chi_d \in \mathcal{T}_{\mathbb{R}}$, which allows us to extend the Fubini–Study operator to a map

$$FS: \mathcal{N}_{\mathbb{R}} \longrightarrow \mathcal{L}^{\infty}$$

into the space of bounded functions on X^{an} , by setting $\operatorname{FS}(\chi) := \lim_d \operatorname{FS}(\chi_d)$. On the other hand, any $\varphi \in \mathcal{L}^{\infty}$ defines an *infimum norm* $\operatorname{IN}(\varphi) \in \mathcal{N}_{\mathbb{R}}$, the avatar of the usual supnorm $\sup_{X^{\operatorname{an}}} |s| e^{-m\varphi}$ on R_m in our additive terminology. This defines an operator

$$\operatorname{IN}\colon \mathcal{L}^\infty \longrightarrow \mathcal{N}^{\operatorname{hom}}_{\mathbb{R}}$$

into the space of homogeneous norms. Using standard but nontrivial results in non-Archimedean geometry, we show that:

• the composition IN \circ FS: $\mathcal{N}_{\mathbb{R}} \to \mathcal{N}_{\mathbb{R}}^{\text{hom}}$ coincides with the homogenization operator $\chi \mapsto \chi^{\text{hom}}$, where $\chi^{\text{hom}}(s) = \lim_{r \to \infty} r^{-1}\chi(s^r)$, which corresponds to the spectral radius construction in the usual "multiplicative" terminology;

- homogenization preserves the finite type condition, and hence maps $\mathcal{T}_{\mathbb{R}}$ onto $\mathcal{T}_{\mathbb{R}}^{hom} \coloneqq \mathcal{T}_{\mathbb{R}} \cap \mathcal{N}_{\mathbb{R}}^{hom}$;
- on $\mathcal{T}_{\mathbb{R}}$, both the Fubini–Study operator and the pseudo-metric d_{∞} factor through homogenization.

These results imply that FS: $(\mathcal{T}_{\mathbb{R}}, d_{\infty}) \to (\mathcal{H}_{\mathbb{R}}, d_{\infty})$ is a surjective isometry, which restricts to an isometric isomorphism $(\mathcal{T}_{\mathbb{R}}^{\text{hom}}, d_{\infty}) \simeq (\mathcal{H}_{\mathbb{R}}, d_{\infty})$, where d_{∞} on $\mathcal{H}_{\mathbb{R}}$ is the supnorm metric; this settles Theorem A for $p = \infty$.

For any $p \in [1, \infty)$, we have $d_1 \leq d_p \leq d_\infty$ as pseudo-metrics on $\mathcal{N}_{\mathbb{R}}$. By the previous step, the restriction of d_∞ to $\mathcal{T}_{\mathbb{R}}$ factors through the Fubini– Study operator. Thus $d_p \mid_{\mathcal{T}_{\mathbb{R}}}$ descends to a pseudo-metric on $\mathcal{H}_{\mathbb{R}}$, and Theorem A asserts that it is a metric, i.e. that it separates points. It is enough to prove this for p = 1, which is accomplished via an explicit expression for d_1 in terms of the Monge–Ampère energy, analogous to the known expression for the Darvas metric in the complex analytic case [31].

Our approach is based on the close relation of the d₁-pseudometric on $\mathcal{N}_{\mathbb{R}}$ to the volume of a norm $\chi \in \mathcal{N}_{\mathbb{R}}$, defined as the limit

$$\operatorname{vol}(\chi) = \lim_{m} m^{-1} \operatorname{vol}(\chi|_{R_m}) \in \mathbb{R},$$

where $\operatorname{vol}(\chi|_{R_m})$ is the barycenter of the spectrum of $\chi|_{R_m}$. Indeed, for all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, we have

$$d_1(\chi, \chi') = \operatorname{vol}(\chi) + \operatorname{vol}(\chi') - 2\operatorname{vol}(\chi \wedge \chi'),$$

with $\chi \wedge \chi' \in \mathcal{N}_{\mathbb{R}}$ the pointwise min of χ and χ' . When $\chi \in \mathcal{T}_{\mathbb{Z}}$ corresponds to a test configuration $(\mathcal{X}, \mathcal{L}) \to \mathbb{A}^1$, with canonical compactification $(\overline{\mathcal{X}}, \overline{\mathcal{L}}) \to \mathbb{P}^1$, it was proved in [17], using the Riemann–Roch formula, that

$$\operatorname{vol}(\chi) = \frac{(\bar{\mathcal{L}}^{n+1})}{(n+1)(L^n)},$$

where the right-hand side is also, by definition, the Monge–Ampère energy $E(\varphi)$ of $\varphi := FS(\chi) \in \mathcal{H}_{\mathbb{Q}}$. Setting $\widetilde{E}(\varphi) := \sup\{E(\psi) \mid \varphi \ge \psi \in \mathcal{H}_{\mathbb{Q}}\}$ defines the extended energy functional $\widetilde{E} \colon C^0 \to \mathbb{R}$, and an approximation argument based on Okounkov bodies leads to the key formula $vol(\chi) = \widetilde{E}(FS(\chi))$, which implies

$$d_1(\chi, \chi') = E(\varphi) + E(\varphi') - 2 \tilde{E}(\varphi \wedge \varphi')$$

for all $\chi, \chi' \in \mathcal{H}_{\mathbb{R}}$, where $\varphi = FS(\chi)$, $\varphi' = FS(\chi')$. The right-hand side thus defines a pseudo-metric d_1 on $\mathcal{H}_{\mathbb{R}}$, and a result of [18] allows us to show that it separates points, thereby finishing the proof of Theorem A. This formula also characterizes the metric d_1 on $\mathcal{H}_{\mathbb{R}}$ as the unique one such that $d_1(\varphi, \varphi') = \inf_{\varphi'' \leqslant \varphi, \varphi'} \{ d_1(\varphi, \varphi'') + d_1(\varphi'', \varphi') \}$ for all $\varphi, \varphi' \in \mathcal{H}_{\mathbb{R}}$, and $d_1(\varphi, \varphi') = E(\varphi) - E(\varphi')$ when $\varphi \geqslant \varphi'$.

Darvas metrics on functions and measures of finite energy

By Theorem A, the Hausdorff completion of $(\mathcal{T}_{\mathbb{R}}, \mathbf{d}_p)$ can be identified with the completion of the metric space $(\mathcal{H}_{\mathbb{R}}, \mathbf{d}_p)$. When $p = \infty$, this is simply the closure of $\mathcal{H}_{\mathbb{R}} \subset \mathbb{C}^0$ in the topology of uniform convergence, which is, by definition, the space CPSH of continuous *L*-psh functions (in line with [47, 70]).

For a norm $\chi \in \mathcal{N}_{\mathbb{R}}$, $\mathrm{FS}(\chi)$ lies in CPSH as soon as it is continuous (by Dini's lemma); we show that the set $\mathcal{N}_{\mathbb{R}}^{\mathrm{cont}} \subset \mathcal{N}_{\mathbb{R}}$ of such norms coincides with the d_{∞} -closure in $\mathcal{N}_{\mathbb{R}}$ of the set $\mathcal{T}_{\mathbb{Z}}$ of (ample) test configurations, and that it is a strict subset (except in the trivial case dim X = 0, see Section 2.5).

Our next goal is to describe the completion of $(\mathcal{H}_{\mathbb{R}}, d_1)$. The answer relies on global pluripotential theory over a trivially valued field, as developed in [18] (inspired in part by the discretely valued case studied in [15]). Let us briefly describe the salient points of this theory.

Inspired by the complex analytic case, we define an *L*-psh function $\varphi \colon X^{\mathrm{an}} \to \mathbb{R} \cup \{-\infty\}$ as an upper semicontinuous (usc) function that can be written as the limit of a decreasing sequence (or net) in $\mathcal{H}_{\mathbb{R}}$ (or $\mathcal{H}_{\mathbb{Z}} = \mathcal{H}_{\mathbb{Q}}$), excluding $\varphi \equiv -\infty$. Such functions are uniquely determined by their restrictions to $X^{\mathrm{div}} \subset X^{\mathrm{an}}$, which are further finite valued, and we equip the space PSH of *L*-psh functions with the topology of pointwise convergence on X^{div} . By Dini's Lemma, the space of continuous *L*-psh functions CPSH considered above can be described as CPSH = PSH $\cap \mathbb{C}^0$.

The Monge–Ampère energy E admits a unique usc, monotone increasing extension

E: PSH
$$\longrightarrow \mathbb{R} \cup \{-\infty\},\$$

given by $E(\varphi) = \inf \{ E(\psi) \mid \varphi \leq \psi \in CPSH \}$, and the space of *L*-psh functions of finite energy is defined as

$$\mathcal{E}^1 \coloneqq \{\mathrm{E} > -\infty\} \subset \mathrm{PSH}$$
 .

A function in \mathcal{E}^1 is thus a decreasing limit of functions in $\mathcal{H}_{\mathbb{Q}}$ with bounded energy. The space \mathcal{E}^1 is endowed with the *strong topology*, defined as the coarsest refinement of the subspace topology from PSH $\supset \mathcal{E}^1$ for which $E: \mathcal{E}^1 \to \mathbb{R}$ is continuous. Any decreasing net in \mathcal{E}^1 is strongly convergent, and $\mathcal{H}_{\mathbb{Q}}$ is thus dense in \mathcal{E}^1 in the strong topology. To each $\varphi \in \mathcal{E}^1$ is associated a Monge–Ampère measure $MA(\varphi)$, a (Radon) probability measure on X^{an} that integrates functions in \mathcal{E}^1 . When $\varphi \in \mathcal{H}_{\mathbb{Q}}$, $MA(\varphi)$ has finite support in X^{div} , and can be described using intersection numbers computed on the central fiber of an associated test configuration. The Monge–Ampère operator $\varphi \mapsto MA(\varphi)$ is continuous on \mathcal{E}^1 in the strong topology, and characterized as the derivative of E, i.e.

$$\frac{\mathrm{d}}{\mathrm{d}t}\bigg|_{t=0} \mathrm{E}\left((1-t)\varphi + t\psi\right) = \int_{X^{\mathrm{an}}} (\psi - \varphi) \operatorname{MA}(\varphi)$$

for all $\varphi, \psi \in \mathcal{E}^1$. It takes its values in the space \mathcal{M}^1 of measures of finite energy, i.e. Radon probability measures μ on X^{an} for which the Legendre transform

$$\mathbf{E}^{\vee}(\mu) \coloneqq \sup_{\varphi \in \mathcal{E}^1} \left\{ \mathbf{E}(\varphi) - \int \varphi \, \mathrm{d}\mu \right\} \in [0, +\infty]$$

is finite. In analogy to the complex analytic case [3], the variational approach of [18] shows that $\mu = MA(\varphi)$ with $\varphi \in \mathcal{E}^1$ iff φ achieves the supremum that defines $E^{\vee}(\mu)$.

The space \mathcal{M}^1 also comes with a strong topology, the coarsest refinement of the weak topology of measures such that $E^{\vee} \colon \mathcal{M}^1 \to \mathbb{R}_{\geq 0}$ is continuous. A key result of [18] shows that the Monge–Ampère operator induces a topological embedding with dense image

$$MA: \mathcal{E}^1/\mathbb{R} \hookrightarrow \mathcal{M}^1$$

(with respect to the strong topologies), which is further onto iff the envelope property holds for (X, L). The latter important property has several equivalent formulations, including the compactness of the quotient space PSH/ \mathbb{R} (a fundamental fact in the setting of compact complex manifolds); it is established when X is smooth, using multiplier ideals, and we conjecture that it holds as long as X is normal (or merely unibranch, which is in turn a necessary condition).

The Monge–Ampère operator naturally induces a map MA: $\mathcal{T}_{\mathbb{R}} \to \mathcal{M}^1$ by setting MA(χ) := MA(FS(χ)); as mentioned above, when $\chi \in \mathcal{T}_{\mathbb{Z}}$, the measure MA(χ) has finite support in X^{div} , and can be directly described in terms of intersection numbers on (the integral closure of) the test configuration corresponding to χ .

With these preliminaries in hand, we can now state:

THEOREM B. — For any polarized variety (X, L), the following holds:

(i) there exists a unique metric d_1 on \mathcal{E}^1 that defines the strong topology and extends the metric d_1 on $\mathcal{H}_{\mathbb{R}} \subset \mathcal{E}^1$;

- (ii) there exists a unique metric d_1 on \mathcal{M}^1 that defines the strong topology and induces the quotient metric of d_1 on $\mathcal{E}^1/\mathbb{R} \hookrightarrow \mathcal{M}^1$;
- (iii) the metric space (\mathcal{M}^1, d_1) is always complete, while (\mathcal{E}^1, d_1) is complete iff the envelope property holds for (X, L);
- (iv) the Monge–Ampère operator MA: $\mathcal{T}_{\mathbb{R}} \to \mathcal{M}^1$ uniquely extends to an isometry

$$MA: (\mathcal{N}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1) \longrightarrow (\mathcal{M}^1, d_1),$$

where \underline{d}_1 denotes the quotient pseudometric of d_1 .

In particular, the Monge–Ampère operator realizes \mathcal{M}^1 as the Hausdorff completion of $(\mathcal{T}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1)$, while \mathcal{E}^1 is the Hausdorff completion of $(\mathcal{T}_{\mathbb{R}}, \underline{d}_1)$ iff the envelope property holds, e.g. when X is smooth (see also [35] for an approach based on geodesic rays, when $k = \mathbb{C}$).

We call the metric d_1 on \mathcal{E}^1 the Darvas metric; its complex analytic analogue, introduced by T. Darvas [31], plays a crucial role in global pluripotential theory, and in particular in the variational approach to the Yau– Tian–Donaldson conjecture [4, 54, 55]. The space \mathcal{E}^1 is studied over more general non-Archimedean fields in [64], where it is shown that (\mathcal{E}^1, d_1) is a geodesic metric space (assuming the envelope property). In analogy with the complex analytic case [32, 34], we expect that, for any $p \in [1, \infty)$, the completion of $(\mathcal{H}_{\mathbb{R}}, d_p)$ can be identified with the space

$$\mathcal{E}^p \coloneqq \left\{ \varphi \in \mathcal{E}^1 \mid \varphi \in L^p(\mathrm{MA}(\varphi)) \right\},\$$

assuming the envelope property.

Among other things, the proof of Theorem B is based on a precise comparison between d_1 and quasi-metrics on \mathcal{E}^1 and \mathcal{M}^1 studied in [18], using estimates that ultimately derive from the Hodge Index Theorem. By construction, MA: $(\mathcal{T}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1) \rightarrow (\mathcal{M}^1, d_1)$ is an isometry, and (iv) is thus a consequence of (iii) and the d_1 -density of $\mathcal{T}_{\mathbb{Z}}$ in $\mathcal{N}_{\mathbb{R}}$, which we prove using Okounkov bodies (see Corollary 3.19).

If $\chi \in \mathcal{N}_{\mathbb{R}}^{\text{cont}}$ is a continuous norm, then $\mathrm{FS}(\chi) \in \mathrm{CPSH} \subset \mathcal{E}^1$, and $\mathrm{MA}(\chi) = \mathrm{MA}(\mathrm{FS}(\chi))$. If the envelope property holds for (X, L), then the usc regularization $\mathrm{FS}^*(\chi)$ lies in \mathcal{E}^1 for any norm $\chi \in \mathcal{N}_{\mathbb{R}}$, and $\mathrm{MA}(\chi) = \mathrm{MA}(\mathrm{FS}^*(\chi))$. In this case, we get a surjective isometry $\mathrm{FS}^* \colon (\mathcal{N}_{\mathbb{R}}, \mathrm{d}_1) \to (\mathcal{E}^{\infty}_{\uparrow}, \mathrm{d}_1)$, where $\mathcal{E}^{\infty}_{\uparrow}$ is the set of *L*-psh functions that are *regularizable from below*, i.e. limits in PSH of an increasing net in CPSH. This realizes $\mathcal{E}^{\infty}_{\uparrow}$ as the Hausdorff quotient of $\mathcal{N}_{\mathbb{R}}$. We emphasize, however, that (iv) is valid even without assuming the envelope property for (X, L).

Finally, we show that the functional $\chi \mapsto ||\chi|| \coloneqq E^{\vee}(MA(\chi))$ on $\mathcal{N}_{\mathbb{R}}$ extends (up to a normalization constant) the *minimum norm* of a test configuration in the sense of Dervan [36].

Divisorial norms and maximal norms

The set X^{val} of valuations on the function field of X, trivial on k, is a dense subset of X^{an} . Following [22] we say that $v \in X^{\text{val}}$ is of linear growth if there exists C > 0 such that $v(s) \leq Cm$ for all nonzero sections $s \in R_m = \mathrm{H}^0(X, mL)$ with m sufficiently divisible. In terms of pluripotential theory, the set $X^{\text{lin}} \subset X^{\text{val}}$ of valuations of linear growth coincides with the set of points $v \in X^{\text{an}}$ that are non-pluripolar, i.e. such that every $\varphi \in \mathrm{PSH}$ is finite at v; in particular, it contains the set X^{div} of divisorial valuations.

Any $v \in X^{\text{lin}}$ defines a (homogeneous) norm $\chi_v \in \mathcal{N}_{\mathbb{R}}^{\text{hom}}$, simply by setting $\chi_v(s) \coloneqq v(s)$. We say that a norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is divisorial if it is of the form $\chi = \min_i \{\chi_{v_i} + c_i\}$ for a finite set (v_i) in X^{div} and $c_i \in \mathbb{R}$. We denote by $\mathcal{N}_{\mathbb{R}}^{\text{div}}$ the set of divisorial norms, and by $\mathcal{N}_{\mathbb{Q}}^{\text{div}} \coloneqq \mathcal{N}_{\mathbb{R}}^{\text{div}} \cap \mathcal{N}_{\mathbb{Q}}$ the subset of rational divisorial norms, for which the c_i can be chosen in \mathbb{Q} . The latter contains the homogenization χ^{hom} of any ample test configuration $\chi \in \mathcal{T}_{\mathbb{Z}}$, and $\mathcal{N}_{\mathbb{Q}}^{\text{div}}$ can alternatively be described in terms of norms associated to (possibly non-ample) test configurations (see Theorem A.10).

On the other hand, we define a *divisorial measure* as a Radon probability measure μ on X^{an} with support a finite subset of X^{div} , i.e. $\mu = \sum_i m_i \delta_{v_i}$ for a finite subset (v_i) of X^{div} and $m_i \in \mathbb{R}_{>0}$ such that $\sum_i m_i = 1$. The set $\mathcal{M}^{\mathrm{div}} \subset \mathcal{M}^1$ of divisorial measures is thus the convex hull of the image of the canonical embedding $X^{\mathrm{div}} \hookrightarrow \mathcal{M}^1 \ v \mapsto \delta_v$. For any test configuration $\chi \in \mathcal{T}_{\mathbb{Z}}$, the norm χ^{hom} and the measure $\mathrm{MA}(\chi) = \mathrm{MA}(\chi^{\mathrm{hom}})$ are both divisorial. More generally, we show:

THEOREM C. — The Monge–Ampère operator induces an isometric isomorphism

MA:
$$(\mathcal{N}_{\mathbb{R}}^{\mathrm{div}}/\mathbb{R}, \underline{\mathbf{d}}_1) \xrightarrow{\sim} (\mathcal{M}^{\mathrm{div}}, \mathbf{\mathbf{d}}_1).$$

We emphasize that the envelope property is not assumed here. In the companion paper [20], divisorial measures are used to define the notion of *divisorial stability*, which implies (and is conjecturally equivalent to) uniform K-stability. Theorem C enables us to view divisorial stability as a condition on divisorial norms, and leads to the equivalence between divisorial stability and uniform K-stability with respect to norms/filtrations.

The proof of Theorem C is based on the variational approach to (non-Archimedean) Monge–Ampère equations developed in [18], recast in terms of norms.

Recall that the space of norms $\mathcal{N}_{\mathbb{R}}$ is equipped with pseudometrics $(d_p)_{p\in[1,\infty]}$, such that $d_1 \leq d_p \leq d_\infty$. For $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, the condition $d_p(\chi, \chi') = 0$ is independent of $p < \infty$; we say that χ and χ' are asymptotically equivalent when this holds. While d_∞ becomes a metric upon restriction to the space $\mathcal{N}_{\mathbb{R}}^{\text{hom}}$ of homogeneous norms, this is still not the case for d_p with $p < \infty$, and our next goal is to introduce a canonical maximal subspace on which d_p does become a metric.

To this end, we introduce the class $\mathcal{N}_{\mathbb{R}}^{\max} \subset \mathcal{N}_{\mathbb{R}}^{\hom}$ of maximal norms, of the form $\chi = \inf_{v \in X^{\operatorname{div}}} \{\chi_v + c_v\}$ for a bounded family of constants $(c_v)_{v \in X^{\operatorname{div}}}$. Any divisorial norm is maximal, and maximal norms can alternatively be characterized as decreasing limits of divisorial norms. We further show that any norm χ_v with $v \in X^{\operatorname{lin}}$ is maximal.

The following result accounts for the chosen terminology.

THEOREM D. — Any norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is asymptotically equivalent to a unique maximal norm $\chi^{\max} \in \mathcal{N}_{\mathbb{R}}^{\max}$, characterized as the largest norm in the asymptotic equivalence class of χ . In particular, for any $p \in [1, \infty)$, the restriction of the pseudometric d_p to $\mathcal{N}_{\mathbb{R}}^{\max}$ is a metric, and $\mathcal{N}_{\mathbb{R}}^{\max}$ is maximal in $\mathcal{N}_{\mathbb{R}}$ for this property.

To prove this result, we first construct a projection $\chi \mapsto \chi^{\max}$ onto $\mathcal{N}_{\mathbb{R}}^{\max}$, by setting $\chi^{\max} \coloneqq \inf_{v \in X^{\operatorname{div}}} \{\chi_v + \operatorname{FS}(\chi)(v)\}$, and show that $\chi^{\max} = \chi'^{\max}$ iff $\operatorname{FS}(\chi) = \operatorname{FS}(\chi')$ on X^{div} . Using Monge–Ampère estimates from [18], we show that this holds if $\chi \sim \chi'$. Conversely, we need to show $\chi \sim \chi^{\max}$. Since $\operatorname{FS}(\chi) = \sup_d \operatorname{FS}(\chi_d)$ is an envelope of *L*-psh functions, it follows from [18, 19] that $\operatorname{FS}(\chi) = \operatorname{FS}^*(\chi)$ on X^{div} , and

$$\operatorname{vol}(\chi) = \widetilde{\operatorname{E}}(\operatorname{FS}(\chi)) = \widetilde{\operatorname{E}}(\operatorname{FS}^{\star}(\chi)) \ge \operatorname{vol}(\chi^{\max}).$$

This yields the result, since $\chi \leq \chi^{\max}$ implies

$$d_1(\chi, \chi^{\max}) = \operatorname{vol}(\chi^{\max}) - \operatorname{vol}(\chi).$$

As before, Theorem D does not assume the envelope property, but the proof exploits it through the use of resolution of singularities, see [18, Theorem 5.20]. Note that closely related results were independently obtained in [6] in a more general local setting.

Valuations of linear growth

Finally we use the results above to study the structure of the space X^{lin} of valuations of linear growth, which we can endow with several metrics.

First, from the embedding $X^{\text{lin}} \hookrightarrow \mathcal{N}_{\mathbb{R}}$ given by $v \mapsto \chi_v$ we get a family of (pseudo)metrics d_p , $1 \leq p \leq \infty$. Denoting by $v_{\text{triv}} \in X^{\text{div}}$ the trivial valuation, we have in particular

$$d_{\infty}(v, v_{\text{triv}}) = T(v), \quad d_1(v, v_{\text{triv}}) = S(v)$$

where $S(v) \coloneqq \operatorname{vol}(\chi_v)$ is the expected vanishing order of L along v, widely used in relation to the stability threshold/ δ -invariant [5, 44, 53]. The invariant $d_p(v, v_{\text{triv}})$ with $v \in X^{\text{div}}$ also appears (under a slightly different guise) in [69].

Second, a valuation is of linear growth iff the Dirac mass δ_v is a measure of finite energy, and in fact we have

$$\mathbf{E}^{\vee}(\delta_v) = \mathbf{S}(v)$$

for any $v \in X^{\text{lin}}$, see Theorem 7.22. In particular, we have an embedding $X^{\text{lin}} \hookrightarrow \mathcal{M}^1$. Denote by $d_{\mathcal{M}^1}$ the pullback of the metric d_1 on \mathcal{M}^1 to X^{lin} .

COROLLARY E. — The pseudo-metric d_p on X^{lin} is an actual metric for $1 \leq p \leq \infty$. Further, the metrics $d_{\mathcal{M}^1}$ and d_p , $1 \leq p \leq \infty$, on X^{lin} are equivalent and complete, and they are independent of L up to bi-Lipschitz equivalence.

Completeness with respect to d_{∞} , as well as independence of L, was already observed in [18], and the key point is thus to show $d_{\infty} \leq C d_{\mathcal{M}^1}$, which is done by invoking inequalities involving Monge–Ampère integrals, as in the proof of Theorem A (see Section 7.6 for details).

In [20] we use the space \mathcal{M}^1 and its subspace \mathcal{M}^{div} to analyze K-stability. When X is a Fano variety, restricting to Dirac masses $\delta_v \in \mathcal{M}^1$, with v in X^{lin} or X^{div} , recovers the valuative criterion of K-stability of Fano varieties due to Fujita and Li [44, 53].

An interesting type of valuations $v \in X^{\text{lin}}$ are those for which the associated filtration χ_v is of finite type. If $v \in X^{\text{div}}$, this means v is "dreamy" in the sense of K. Fujita [44], associated to a test configuration with irreducible and reduced central fiber. While valuations $v \in X^{\text{lin}}$ with χ_v of finite type play a crucial role in recent work on K-stability of Fano varieties [7, 8, 9, 10, 49, 57], their role in the general polarized case is less clear (although see [37, 58]). The condition of χ_v being of finite type is quite subtle and in particular depends on the ample Q-line bundle L. For this reason we believe that it is useful to study K-stability using functionals on spaces such as X^{div} , X^{lin} , \mathcal{M}^{div} or \mathcal{M}^1 , without any finite type assumption.

Organization

After giving some background in Section 1, we study homogenization and the related Fubini–Study and infimum norm operators in Section 2, proving part of Theorem A. In Section 3 we make a spectral analysis of norms on the section ring of (X, L), building upon [14, 27]. After that we give additional background on non-Archimedean pluripotential theory from [18]; in particular we revisit the spaces used in Theorem B. In Section 5 we construct and study the Darvas metrics on \mathcal{E}^1 and \mathcal{M}^1 , and prove the remaining part of Theorem A as well as parts (i)–(iii) of Theorem B. The classes of divisorial and maximal norms are studied in Section 6, where we prove Theorem D and also consider the regularized Fubini–Study operator. In Section 7 we define the Monge–Ampère operator on general norms, and prove Theorem C as well as Theorem B(iv) and Corollary E. Finally, Appendix A revisits the relation between test configurations and Fubini–Study functions, and Appendix B provides some remarks on the toric case.

Notation and conventions

- We work over an algebraically closed field k, of arbitrary characteristic unless otherwise specified.
- For $x, y \in \mathbb{R}_+$, $x \leq y$ means $x \leq C_n y$ for a constant $C_n > 0$ only depending on n, and $x \approx y$ if $x \leq y$ and $y \leq x$. Here n will be the dimension of a fixed variety X over k.
- A pseudo-metric on a set Z is a function d: Z × Z → ℝ₊ that is symmetric, vanishes on the diagonal, and satisfies the triangle inequality. It is a metric if it further separates points.
- The Hausdorff quotient of a pseudo-metric space (Z, d) is the metric space (Z_H, d_H) where Z_H is the quotient of Z by the equivalence relation $x \sim y \Leftrightarrow d(x, y) = 0$, and d_H is the induced metric. The map $(Z, d) \to (Z_H, d_H)$ is the unique isometric map of (Z, d) onto a metric space, up to unique isomorphism.
- The Hausdorff completion of a pseudo-metric space (Z, d) is the complete metric space $(\widehat{Z}, \widehat{d})$ defined as the completion of the Hausdorff quotient (Z_H, d_H) . It comes with an isometric map $(Z, d) \rightarrow (\widehat{Z}, \widehat{d})$ with dense image, which is universal with respect to maps into complete metric spaces.

 A quasi-metric on Z is function d : Z × Z → ℝ₊ that is symmetric, vanishes precisely on the diagonal, and satisfies the quasi-triangle inequality

$$\varepsilon d(x,y) \leq d(x,z) + d(z,y)$$

for some constant $\varepsilon > 0$. A quasi-metric space (Z, d) comes with a Hausdorff topology, and even a uniform structure. In particular, Cauchy sequences and completeness make sense for (Z, d). Such uniform structures have a countable basis of entourages, and are thus metrizable, by general theory.

- We use the standard abbreviations usc for "upper semicontinuous", *lsc* for "lower semicontinuous", *wlog* for "without loss of generality", and *iff* for "if and only if".
- A net is a family indexed by a directed set. On many occasions we shall consider nets (x_d) indexed by $d_0\mathbb{Z}_{\geq 1}$ for some $d_0 \geq 1$, and ordered by divisibility. Note that the sequence $(x_{m!})_{m\geq d_0}$ is cofinal in this net.

Acknowledgements

We thank E. Bedford, R. Berman, H. Blum, G. Codogni, T. Darvas, R. Dervan, A. Ducros, C. Favre, T. Hisamoto, C. Li, J. Poineau and M. Stevenson for fruitful discussions and useful comments.

1. Background

In the entire paper, (X, L) denotes a projective variety (reduced and irreducible) endowed with an ample \mathbb{Q} -line bundle. We review a number of basic facts about norms/filtrations and Berkovich analytification, referring for instance to [14, 18] for more details.

1.1. Norms on a vector space

As in [23] we will use "additive" terminology, so by a *norm* on a k-vector space V we mean a function $\chi: V \to \mathbb{R} \cup \{+\infty\}$ such that

- $\chi(v) = +\infty$ iff v = 0;
- $\chi(av) = \chi(v)$ for $a \in k^{\times}$ and $v \in V$; and
- $\chi(v+w) \ge \min\{\chi(v), \chi(w)\}$ for all $v, w \in V$.

Note that $\|\cdot\|_{\chi} \coloneqq e^{-\chi(\cdot)}$ is then a non-Archimedean norm on V with respect to the trivial absolute value on k in the usual ("multiplicative") sense [11]. Setting

$$F^{\lambda}V \coloneqq \{v \in V \mid \chi(v) \ge \lambda\}, \quad \chi(v) \coloneqq \max\{\lambda \in \mathbb{R} \mid v \in F^{\lambda}V\}$$

for $\lambda \in \mathbb{R}$ yields a 1–1 correspondence between norms on V and (nonincreasing, left-continuous, exhaustive and separated) filtrations of V. We also write $F^{>\lambda}V := \bigcup_{\lambda'>\lambda} F^{\lambda'}V = \{\chi > \lambda\}$, and define the associated graded space as the \mathbb{R} -graded vector space

$$\operatorname{gr}_{\chi} V \coloneqq \bigoplus_{\lambda \in \mathbb{R}} F^{\lambda} V / F^{>\lambda} V.$$

Each norm χ on V turns it into a (Hausdorff) topological vector space, in which $(F^{m\varepsilon}V)_{m\in\mathbb{N}}$ forms a countable basis of (open and closed) neighborhood of 0, for any $\varepsilon > 0$. The normed space (V, χ) admits a *completion* \hat{V} , a complete topological vector space containing V as a dense subspace, whose topology is defined by a (unique) norm on \hat{V} extending χ . The inclusion $V \hookrightarrow \hat{V}$ induces an isomorphism

(1.1)
$$\operatorname{gr}_{\chi} V \xrightarrow{\sim} \operatorname{gr}_{\chi} \widehat{V}.$$

We denote by $\mathcal{N}_{\mathbb{R}}(V)$ the set of norms on V. It has a distinguished element χ_{triv} , the trivial norm, such that $\chi_{\text{triv}}(v) = 0$ for all $v \neq 0$, and it admits a scaling action by $\mathbb{R}_{>0}$ and a partial ordering defined by $\chi \leq \chi'$ iff $\chi(v) \leq \chi'(v)$ for all v. Any two elements $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}(V)$ admit an infimum $\chi \wedge \chi' \in \mathcal{N}_{\mathbb{R}}(V)$, defined pointwise by

$$(\chi \wedge \chi')(v) \coloneqq \min\{\chi(v), \chi'(v)\}.$$

For any subgroup $\Lambda \subset \mathbb{R}$, we denote by $\mathcal{N}_{\Lambda}(V)$ the set of norms with values in $\Lambda \cup \{+\infty\}$. Thus

$$\{\chi_{\mathrm{triv}}\} = \mathcal{N}_{\{0\}}(V) \subset \mathcal{N}_{\Lambda}(V) \subset \mathcal{N}_{\mathbb{R}}(V).$$

A norm $\chi \in \mathcal{N}_{\mathbb{R}}(V)$ lies in $\mathcal{N}_{\Lambda}(V)$ iff the \mathbb{R} -grading of $\operatorname{gr}_{\chi} V$ reduces to a Λ -grading.

Assume now that V is finite dimensional. Any norm χ on V admits an orthogonal basis (e_i) , i.e. a basis of V such that

$$\chi\left(\sum_{i} a_{i} e_{i}\right) = \min_{a_{i} \neq 0} \chi(e_{i})$$

ANNALES DE L'INSTITUT FOURIER

for all $a_i \in k$. Up to reordering, an orthogonal basis is simply a compatible basis for the flag of linear subspaces underlying the filtration defined by χ , and elementary linear algebra thus implies that any two norms χ, χ' on V admit a joint orthogonal basis.

In particular, all norms on V are equivalent, which means, in our additive terminology, that $\chi - \chi'$ is a bounded function on $V \setminus \{0\}$ for all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}(V)$. The classical Goldman–Iwahori metric on $\mathcal{N}_{\mathbb{R}}(V)$ is defined by

(1.2)
$$d_{\infty}(\chi,\chi') = \sup_{v \in V \smallsetminus \{0\}} |\chi(v) - \chi'(v)|,$$

where the supremum is achieved among the elements of any joint orthogonal basis for χ and χ' . For later use, note that

(1.3)
$$\chi \leq \chi' \leq \chi'' \Longrightarrow d_{\infty}(\chi, \chi') \leq d_{\infty}(\chi, \chi'').$$

The metric space $(\mathcal{N}_{\mathbb{R}}(V), d_{\infty})$ is complete, but not locally compact as soon as dim $V \ge 2$. Note also that $\mathcal{N}_{\mathbb{Z}}(V)$ is a closed, discrete subset of $\mathcal{N}_{\mathbb{R}}(V)$, while $\mathcal{N}_{\mathbb{Q}}(V)$ is dense. For any $\chi \in \mathcal{N}_{\mathbb{R}}(V)$, we set

(1.4)
$$\lambda_{\min}(\chi) \coloneqq \min_{v \in V \smallsetminus \{0\}} \chi(v), \quad \lambda_{\max}(\chi) \coloneqq \max_{v \in V \smallsetminus \{0\}} \chi(v).$$

Thus

$$d_{\infty}(\chi, \chi_{triv}) = \max\{\lambda_{\max}(\chi), -\lambda_{\min}(\chi)\}.$$

Any norm χ on V induces a norm on the dual space and on all tensor powers, in such a way that the bases canonically induced by any given orthogonal basis of V remain orthogonal. If $\pi: V \to V'$ is a surjective linear map, then χ also induces a quotient norm χ' on V', such that $\chi'(v') =$ $\max{\{\chi(v) \mid \pi(v) = v'\}}$ for all $v' \in V'$.

1.2. Norms on a graded algebra

Let now $R = \bigoplus_{m \in \mathbb{N}} R_m$ be a graded k-algebra. It comes with an action of k^{\times} for which $a \cdot s = a^m s$ for $a \in k^{\times}$ and $s \in R_m$. We write $\mathcal{N}_{\mathbb{R}}(R)$ for the set of vector space norms $\chi \colon R \to \mathbb{R}$ that are

- superadditive, i.e. $\chi(fg) \ge \chi(f) + \chi(g)$ for $f, g \in R$;
- k^{\times} -invariant, i.e. $\chi(a \cdot f) = \chi(f)$ for $a \in k^{\times}$ and $f \in R$; this is equivalent to χ being compatible with the grading of R, that is, $\chi(\sum_m s_m) = \min_m \chi(s_m)$ where $s_m \in R_m$;
- linearly bounded, i.e. there exists C > 0 such that $|\chi| \leq Cm$ on $R_m \setminus \{0\}$ for all $m \geq 1$.

Norms in $\mathcal{N}_{\mathbb{R}}(R)$ are in 1–1 correspondence with graded, linearly bounded filtrations of R as in [13, 68]. Each $\chi \in \mathcal{N}_{\mathbb{R}}(R)$ defines a graded algebra

$$\operatorname{gr}_{\chi} R = \bigoplus_{m \in \mathbb{N}} \operatorname{gr}_{\chi} R_m = \bigoplus_{(m,\lambda) \in \mathbb{N} \times \mathbb{R}} F^{\lambda} R_m / F^{>\lambda} R_m.$$

LEMMA 1.1. — A norm $\chi \in \mathcal{N}_{\mathbb{R}}(R)$ is a valuation on R, i.e. it satisfies $\chi(fg) = \chi(f) + \chi(g)$ for all $f, g \in R$, iff $\operatorname{gr}_{\chi} R$ is an integral domain.

The set $\mathcal{N}_{\mathbb{R}}(R) \hookrightarrow \prod_m \mathcal{N}_{\mathbb{R}}(R_m)$ is stable under the scaling action of $\mathbb{R}_{>0}$ and infima; it further admits an additive action of \mathbb{R} , denoted by $(c, \chi) \mapsto \chi + c$, such that

(1.5)
$$(\chi + c)(s) \coloneqq \chi(s) + cm \text{ for } s \in R_m.$$

For any subgroup $\Lambda \subset \mathbb{R}$, denote by $\mathcal{N}_{\Lambda}(R) \subset \mathcal{N}_{\mathbb{R}}(R)$ the set of norms with values in $\Lambda \cup \{+\infty\}$. Norms in $\mathcal{N}_{\mathbb{Z}}(R)$ and $\mathcal{N}_{\mathbb{Q}}(R)$ will be called *integral* and *rational*, respectively. Integral norms are in 1–1 correspondence with \mathbb{Z} -filtrations, as considered in [65].

For any norm $\chi \in \mathcal{N}_{\mathbb{R}}(R)$, the round-down $|\chi| \in \mathcal{N}_{\mathbb{Z}}(R)$, defined by

(1.6)
$$\lfloor \chi \rfloor(s) \coloneqq \lfloor \chi(s) \rfloor, \quad s \in R_m \setminus \{0\},$$

is an integral norm.

Example 1.2. — Consider the algebra $k[z] = k[z_1, \ldots, z_n]$ of polynomials in *n* variables, with the usual grading. For each $\xi \in \mathbb{R}^n$, the monomial valuation

(1.7)
$$\chi_{\xi}\left(\sum_{\alpha\in\mathbb{N}^n}c_{\alpha}z^{\alpha}\right) = \min_{c_{\alpha}\neq 0}\langle\alpha,\xi\rangle = \min_{\alpha}\{v_{\mathrm{triv}}(c_{\alpha}) + \langle\alpha,\xi\rangle\}$$

defines a norm on the graded algebra k[z]. The completion of $(k[z], \chi_{\xi})$ is the algebra $k\{z;\xi\}$ of formal power series $\sum_{\alpha} c_{\alpha} z^{\alpha} \in k[\![z]\!]$ such that $\lim_{\alpha} (v_{\text{triv}}(c_{\alpha}) + \langle \alpha, \xi \rangle) = +\infty$, whose norm is still defined by (1.7). In multiplicative notation, $k\{z;\xi\}$ is the polydisc algebra $k\{r^{-1}z\}$, with $r_j = e^{-\xi_j}$, a building block of Berkovich spaces [1, 2].

From now on, we assume that R is finitely generated, so that each graded piece R_m is finite dimensional.

DEFINITION 1.3. — We say that a norm $\chi \in \mathcal{N}_{\mathbb{R}}(R)$ is generated in degree 1 if R is generated in degree 1 and, for any $m \ge 1$, the restriction $\chi|_{R_m}$ is the quotient norm of $S^m(\chi|_{R_1})$ under the canonical surjective map $S^m R_1 \to R_m$.

Concretely, χ is generated in degree 1 iff, given a χ -orthogonal basis (s_i) of R_1 , any $s \in R_m$ can be written as $s = \sum_{|\alpha|=m} c_\alpha \prod_i s_i^{\alpha_i}$ with $c_\alpha \in k$ and $\chi(s) = \min_{c_\alpha \neq 0} \sum_i \alpha_i \chi(s_i)$.

LEMMA 1.4. — For any subgroup $\Lambda \subset \mathbb{R}$ and $\chi \in \mathcal{N}_{\Lambda}(R)$, the following conditions are equivalent:

- (i) χ is generated in degree 1;
- (ii) $\operatorname{gr}_{\chi} R = \bigoplus_{m \in \mathbb{N}} \operatorname{gr}_{\chi} R_m$ is generated in degree 1;
- (iii) there exists $\xi \in \Lambda^N$ and a surjective map of graded k-algebras $\pi: k[z_1, \ldots, z_N] \to R$ with respect to which χ is the quotient norm of χ_{ξ} as in Example 1.2.

When this holds, we further have $\chi \in \mathcal{N}_{\Lambda'}(R)$ for some finitely generated subgroup $\Lambda' \subset \Lambda$.

Proof. — Assume (i). Choose a χ -orthogonal basis $(s_i)_{1 \leq i \leq N}$ of R_1 , and set $\xi_i := \chi(s_i)$. As noted above, any $s \in R_m \setminus \{0\}$ can be written as $s = \sum_{|\alpha|=m} c_\alpha \prod_i s_i^{\alpha_i}$ with $\chi(s) = \min_{c_\alpha \neq 0} \sum_i \alpha_i \xi_i$. This already yields the final assertion, with $\Lambda' := \sum_i \mathbb{Z}\xi_i$.

Define A as the set of α achieving $\min_{c_{\alpha}\neq 0} \sum_{i} \alpha_{i}\xi_{i} = \chi(s)$ and set $s' := \sum_{\alpha \in A} c_{\alpha} \prod_{i} s_{i}^{\alpha_{i}}$. Then $s - s' \in F^{>\chi(s)}R_{m}$, so s = s' in $\operatorname{gr}_{\chi} R_{m}$. This shows that $S^{m} \operatorname{gr}_{\chi} R_{1} \to \operatorname{gr}_{\chi} R_{m}$ is surjective, and hence (i) \Rightarrow (ii). If we define $\pi \colon k[z] \to R$ by $\pi(z_{i}) = s_{i}$, then it is clear that χ is the quotient norm of χ_{ξ} with $\xi = (\xi_{i})$, hence (i) \Rightarrow (ii).

Conversely, any quotient of a norm generated in degree 1 is plainly generated in degree 1 as well; hence (iii) \Rightarrow (i). Assume now (ii), and pick again a χ -orthogonal basis (s_i) of R_1 . Each $s \in R_m \setminus \{0\}$ can then be written as $s = \sum_{|\alpha|=m} a_{\alpha} \prod_i s_i^{\alpha_i} + s'$ where $a_{\alpha} \in k, \ \chi(s) = \sum_i \alpha_i \chi(s_i)$ for all α and $s' \in F^{>\chi(s)}R_m$. Repeating the procedure with s' in place of s and using the fact that $\lambda \mapsto F^{\lambda}R_m$ jumps only finitely many times (by finite-dimensionality of R_m), we end up with a decomposition $s = \sum_{|\alpha|=m} c_{\alpha} \prod_i s_i^{\alpha_i}$ such that $\chi(s) = \min_{c_{\alpha}\neq 0} \sum_i \alpha_i \chi(s_i)$. This proves that χ is generated in degree 1, thus (ii) \Rightarrow (i).

1.3. Norms on section rings

Recall that L is an ample \mathbb{Q} -line bundle on a projective variety X. For any $d \in \mathbb{N}$ such that dL is an actual line bundle, we write $R_d := \mathrm{H}^0(X, dL)$, and denote by

$$R^{(d)} = R(X, dL) = \bigoplus_{m \in \mathbb{N}} R_{md}$$

the *d*-th Veronese algebra, i.e. the section ring of dL; it is generated in degree 1 for all *d* sufficiently divisible, since *L* is ample. When *d* divides d', we have a restriction map $\mathcal{N}_{\mathbb{R}}(R^{(d)}) \to \mathcal{N}_{\mathbb{R}}(R^{(d')})$, and we set

(1.8)
$$\mathcal{N}_{\mathbb{R}} = \mathcal{N}_{\mathbb{R}}(X, L) \coloneqq \varinjlim_{d} \mathcal{N}_{\mathbb{R}}(R^{(d)}).$$

The set $\mathcal{N}_{\mathbb{R}}$ inherits a partial order with finite infima, and commuting actions of $\mathbb{R}_{>0}$ (by scaling) and \mathbb{R} (by translation).

An element $\chi \in \mathcal{N}_{\mathbb{R}}$ is represented by a norm on some $R^{(d)}$, two such norms being identified if they coincide on some further Veronese subalgebra; for convenience, we simply refer to χ as a norm. For all m sufficiently divisible, we denote by $\chi|_{R_m} \in \mathcal{N}_{\mathbb{R}}(R_m)$ the restriction of χ to R_m .

Remark 1.5. — To define $\chi|_{R_m}$, one needs to choose a representative of χ as a norm on some $R^{(d)}$. But any other choice leads to the same norms $\chi|_{R_m} \in \mathcal{N}_{\mathbb{R}}(R_m)$ for *m* sufficiently divisible, and the choice of representative can thus safely be ignored.

For any subgroup $\Lambda \subset \mathbb{R}$, we similarly introduce

$$\mathcal{N}_{\Lambda} \coloneqq \varinjlim_{d} \mathcal{N}_{\Lambda}(R^{(d)})$$

It can be identified with the set of $\chi \in \mathcal{N}_{\mathbb{R}}$ such that $\chi(R_m \setminus \{0\}) \subset \Lambda$ for m sufficiently divisible. Note that \mathcal{N}_{Λ} is invariant under the scaling action of $\{t \in \mathbb{R}_{>0} \mid t\Lambda \subset \Lambda\}$ and the translation action of the divisible group $\mathbb{Q}\Lambda \subset \mathbb{R}$, by (1.5).

Example 1.6. — Any (not necessarily ample) test configuration $(\mathcal{X}, \mathcal{L})/\mathbb{A}^1$ defines a norm $\chi_{\mathcal{L}} \in \mathcal{N}_{\mathbb{Z}}$ (see Section A.1). In this case, the translation action by $c \in \mathbb{Q}$ corresponds to twisting \mathcal{L} by $c\mathcal{X}_0$, while the scaling action by $d \in \mathbb{Z}_{>0}$ corresponds to the base change $\mathbb{A}^1 \to \mathbb{A}^1$ given by $z \mapsto z^d$.

The Goldman–Iwahori metric (1.2) induces a pseudo-metric d_{∞} on $\mathcal{N}_{\mathbb{R}}$ by setting

(1.9)
$$d_{\infty}(\chi,\chi') \coloneqq \limsup_{m} m^{-1} d_{\infty}(\chi|_{R_{m}},\chi'|_{R_{m}}) \in \mathbb{R}_{\geq 0}.$$

The limsup is taken with respect to the partial ordering on $\mathbb{Z}_{>0}$ by divisibility, and it is finite, by linear boundedness of χ, χ' . This pseudo-metric is not a metric (see however Proposition 2.8):

Example 1.7. — Pick any norm $\chi \in \mathcal{N}_{\mathbb{R}}$, with round-down $\lfloor \chi \rfloor \in \mathcal{N}_{\mathbb{Z}}$, see (1.6). For *m* sufficiently divisible, we then have $d_{\infty}(\chi|_{R_m}, \lfloor \chi \rfloor|_{R_m}) \leq 1$, and hence $d_{\infty}(\chi, \lfloor \chi \rfloor) = 0$. In particular, $\mathcal{N}_{\mathbb{Z}}$ is dense in $\mathcal{N}_{\mathbb{R}}$ in the d_{∞} topology. We also introduce

(1.10)
$$\lambda_{\max}(\chi) \coloneqq \lim_{m} m^{-1} \lambda_{\max}(\chi|_{R_m}),$$

where $\lambda_{\max}(\chi|_{R_m})$ is defined by (1.4) and the limit exists and is finite because $m^{-1}\lambda_{\max}(\chi|_{R_m})$ is increasing with respect to divisibility, and bounded by linear boundedness of χ . Note that

(1.11)
$$\chi \ge \chi_{\text{triv}} \Longrightarrow d_{\infty}(\chi, \chi_{\text{triv}}) = \lambda_{\max}(\chi)$$

1.4. \mathbb{R} -test configurations

DEFINITION 1.8. — We say that a norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is of finite type if it is represented by a norm on some $R^{(d)}$ whose associated graded algebra $\operatorname{gr}_{\chi} R^{(d)}$ is of finite type.

Equivalently, a norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is of finite type iff it is represented by a norm on some $R^{(d)}$ that is generated in degree 1, by Lemma 1.4. We denote by

 $\mathcal{T}_{\mathbb{R}} \subset \mathcal{N}_{\mathbb{R}}$

the set of such norms. In line with [38, 49], we interpret the elements of $\mathcal{T}_{\mathbb{R}}$ as \mathbb{R} -test configurations. This is justified by the Rees construction, which sets up a 1–1 correspondence between the subset

$$\mathcal{T}_{\mathbb{Z}} \coloneqq \mathcal{N}_{\mathbb{Z}} \cap \mathcal{T}_{\mathbb{R}}$$

of \mathbb{Z} -valued norms in $\mathcal{T}_{\mathbb{R}}$ and the set of (usual) ample test configurations for (X, L) (see Appendix A). For any $\chi \in \mathcal{N}_{\mathbb{Z}}$, note further that

(1.12)
$$\chi \in \mathcal{T}_{\mathbb{Z}} \iff \bigoplus_{\lambda \in \mathbb{Z}} F^{\lambda} R^{(d)}$$
 of finite type over k for d sufficiently divisible.

More generally, for any subgroup $\Lambda \subset \mathbb{R}$ we set

$$\mathcal{T}_{\Lambda} := \mathcal{N}_{\Lambda} \cap \mathcal{T}_{\mathbb{R}}.$$

As above, \mathcal{T}_{Λ} is invariant under the scaling action of $\{t \in \mathbb{R}_{>0} \mid t\Lambda \subset \Lambda\}$ and the translation action of $\mathbb{Q}\Lambda$. In particular, $\mathcal{T}_{\mathbb{Z}}$ is invariant under translation by \mathbb{Q} . It is also easy to see that

(1.13)
$$\chi \in \mathcal{T}_{\Lambda} \Longrightarrow \lambda_{\max}(\chi) \in \mathbb{Q}\Lambda.$$

By Lemma 1.4, we have

$$\mathcal{T}_{\mathbb{R}} = \bigcup_{\Lambda \subset \mathbb{R} \text{ finitely generated}} \mathcal{T}_{\Lambda}.$$

The central fiber of an \mathbb{R} -test configuration $\chi \in \mathcal{T}_{\mathbb{R}}$ is defined as the polarized scheme

(1.14)
$$(\mathcal{X}_0, \mathcal{L}_0) \coloneqq \left(\operatorname{Proj}\left(\operatorname{gr}_{\chi} R^{(d)}\right), d^{-1} \mathcal{O}(1) \right),$$

for $d \ge 1$ sufficiently divisible. If $\chi \in \mathcal{T}_{\Lambda}$ with $\Lambda \simeq \mathbb{Z}^r$ finitely generated, the Λ -grading of $\operatorname{gr}_{\chi} R^{(d)}$ provides a $\mathbb{G}_{\mathrm{m}}^r$ -action on $(\mathcal{X}_0, \mathcal{L}_0)$.

The smallest value of r is called the rank of χ ; it is equal to 1 iff χ is a usual test configuration, up to scaling.

Remark 1.9. — For an \mathbb{R} -test configuration $\chi \in \mathcal{T}_{\mathbb{R}}$, there does not generally exist a smallest subgroup $\Lambda \subset \mathbb{R}$ such that $\chi \in \mathcal{T}_{\Lambda}$, because the subgroup $\Lambda_m \subset \mathbb{R}$ generated by the values of $\chi|_{R_m}$ need not stabilize for m sufficiently divisible. However, the associated \mathbb{Q} -vector space $\mathbb{Q}\Lambda_m$ does stabilize, its dimension being the rank of χ .

Example 1.10. — Extending Example 1.2, suppose that (X, L) is acted upon by a torus $T = \mathbb{G}_{\mathrm{m}}^r$. Then each $\xi \in \mathbb{R}^r$ defines a norm $\chi = \chi_{\xi} \in \mathcal{N}_{\mathbb{R}}$, given by

$$\chi(s) \coloneqq \min\left\{ \langle \alpha, \xi \rangle \mid \alpha \in \mathbb{Z}^r, \, s_\alpha \neq 0 \right\}$$

for $s \in R_m$ with *m* sufficiently divisible, where $s = \sum_{\alpha \in \mathbb{Z}^r} s_\alpha$ is the weight decomposition. This norm satisfies

$$\operatorname{gr}_{\chi} R^{(d)} \simeq \bigoplus_{\lambda \in \mathbb{R}} \left(\bigoplus_{\alpha \in M, \, \langle \alpha, \xi \rangle = \lambda} R^{(d)}_{\alpha} \right) = R^{(d)},$$

which shows that $\chi \in \mathcal{T}_{\mathbb{R}}$ is of finite type, with central fiber isomorphic to (X, L). Further, χ lies in \mathcal{T}_{Λ} for the finitely generated subgroup $\Lambda = \sum_{i} \mathbb{Z}\xi_{i}$.

Example 1.11. — Pick an embedding $X \hookrightarrow \mathbb{P}^N$ in a projective space such that $\mathcal{O}(1)|_X = dL$ for some $d \ge 1$, and suppose we are given an action of a torus $T = \mathbb{G}_{\mathrm{m}}^r$ on $(\mathbb{P}^N, \mathcal{O}(1))$. By Example 1.10, each $\xi \in \mathbb{R}^r$ defines a norm on $R(\mathbb{P}^N, \mathcal{O}(1))$, generated in degree 1, which restricts to a norm in $\mathcal{T}_{\mathbb{R}}$. By Lemma 1.4, every element of $\mathcal{T}_{\mathbb{R}}$ conversely arises in this way (compare [49, Lemma 2.10]).

Following [49, 50], one can use Example 1.11 to provide a geometric realization of \mathbb{R} -test configurations as equivariant polarized families over a toric base (see Section A.3 for a brief discussion).

DEFINITION 1.12. — We define the canonical approximants of a norm $\chi \in \mathcal{N}_{\mathbb{R}}$ as the sequence $\chi_d \in \mathcal{T}_{\mathbb{R}}$ defined for $d \in \mathbb{Z}_{\geq 1}$ sufficiently divisible by letting χ_d be the (class of the) norm on $\mathbb{R}^{(d)}$ generated in degree 1 by χ_d .

If d divides d' then $\chi_d \leq \chi_{d'} \leq \chi$. As in Remark 1.5, this construction is not entirely canonical, as it depends on the choice of a representative of χ , but this can be ignored as any other choice leads to the same approximants χ_d for d sufficiently divisible.

A norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is of finite type iff $\chi = \chi_d$ for all sufficiently divisible d. Note also that

$$\chi \in \mathcal{N}_{\Lambda} \Longrightarrow \chi_d \in \mathcal{T}_{\Lambda}$$

for any subgroup $\Lambda \subset \mathbb{R}$.

1.5. The Berkovich analytification

By a valuation on X we mean a real-valued valuation $v: k(X)^{\times} \to \mathbb{R}$, trivial on k. We denote by X^{val} the space of valuations, endowed with the topology of pointwise convergence on $k(X)^{\times}$. The trivial valuation $v_{\text{triv}} \in X^{\text{val}}$ is defined by $v_{\text{triv}}(f) = 0$ for all $f \in k(X)^{\times}$.

By [1], the space X^{val} admits a natural compactification X^{an} , which as a set equals $X^{\text{an}} = \coprod Y^{\text{val}}$ with Y ranging over all (closed) subvarieties of X. We somewhat imprecisely refer to the points on X^{an} as semivaluations on X. The support of a semivaluation in $Y^{\text{val}} \subset X^{\text{an}}$ is the subvariety Y.

By the valuative criterion of properness, each valuation $v \in X^{\text{val}}$ admits a center $c_X(v) \in X$, characterized as the unique (scheme) point $\xi \in X$ such that $v \ge 0$ on the local ring $\mathcal{O}_{X,\xi}$ and v > 0 on its maximal ideal. This applies to semivaluations as well, replacing X with a subvariety, and thus defines a map $c_X \colon X^{\text{an}} \to X$ (which turns out to be anticontinuous, i.e. the preimage of an open subset is closed).

The space X^{an} comes with a natural action of $\mathbb{R}_{>0}$ by scaling $(t, v) \mapsto tv$. This induces an action $(t, \varphi) \mapsto t \cdot \varphi$ on functions φ on X^{an} by setting

(1.15)
$$(t \cdot \varphi)(v) \coloneqq t\varphi(t^{-1}v),$$

whose fixed points are functions that are homogeneous, i.e. $\varphi(tv) = t\varphi(v)$ for all $t \in \mathbb{R}_{>0}$ and $v \in X^{\mathrm{an}}$.

The set X^{an} is also endowed with a partial order relation, for which $v \ge v'$ iff $c_X(v)$ is a specialization of $c_X(v')$ and $v \ge v'$ pointwise on the local ring at $c_X(v)$. The trivial valuation satisfies $v \ge v_{\text{triv}}$ for all $v \in X^{\text{an}}$.

A (rational) divisorial valuation v on X is a valuation of the form $v = t \operatorname{ord}_E$, where E is a prime divisor on a normal, projective birational model $X' \to X$ and $t \in \mathbb{Q}_{>0}$. The center $c_X(v)$ is then the generic point of the image of E in X. For convenience, we also count the trivial valuation v_{triv} as divisorial, i.e. we allow t = 0 above. The set X^{div} of divisorial valuations is dense in X^{an} (see for instance [18, Theorem 2.14]).

1.6. Semivaluations and line bundles

A semivaluation $v \in X^{\operatorname{an}}$ can be naturally evaluated on a section $s \in \mathrm{H}^0(X, M)$ of any line bundle M on X, by defining v(s) as the value of v on the local function corresponding to s in any local trivialization of M at $c_X(v)$. Thus $v(s) \in [0, +\infty], v(s) > 0$ iff s vanishes at $c_X(v)$, and $v(s) = \infty$ iff s vanishes along the support of v. Further, $v \in X^{\operatorname{val}}$ iff $v(s) < +\infty$ for all $s \in \mathrm{H}^0(X, M) \smallsetminus \{0\}$ and all line bundles M. We define a continuous function $|s|: X^{\operatorname{an}} \to [0, 1]$ by setting

$$(1.16) |s|(v) \coloneqq \exp(-v(s)).$$

Now suppose L is an (ample) line bundle. The \mathbb{Z} -grading of R = R(X, L) defines an action of \mathbb{G}_m on the affine cone $Y \coloneqq \operatorname{Spec} R$, which comes with a natural surjective \mathbb{G}_m -invariant morphism $\pi \colon Y \smallsetminus \{o\} \to X$, where the vertex o of Y is the point defined by the maximal ideal $\bigoplus_{m>0} R_m$. For any $\xi \in X$, the fiber $\pi^{-1}(\xi)$ contains a unique \mathbb{G}_m -invariant point defined by the homogeneous prime ideal generated by all sections $s \in R_m, m \ge 1$ that vanish at ξ .

By general properties of the analytification functor in [1], the \mathbb{G}_{m} -action on Y induces an action of $\mathbb{G}_{m}(k) = k^{\times}$ on Y^{an} , and π induces a surjective k^{\times} -invariant map $\pi^{\mathrm{an}} \colon Y^{\mathrm{an}} \smallsetminus \{w_o\} \to X^{\mathrm{an}}$, where $w_o \in Y^{\mathrm{an}}$ is the trivial semivaluation with support o, which satisfies $w_o = +\infty$ on $\bigoplus_{m>0} R_m$. A semivaluation $w \in Y^{\mathrm{an}}$ is k^{\times} -invariant iff $w(\sum_m s_m) = \min_m w(s_m)$, where $s_m \in R_m$.

It is easy to see [53, Section 4.2] that if $v \in X^{\mathrm{an}}$, then the set of k^{\times} -invariant points in $(\pi^{\mathrm{an}})^{-1}(v)$ is of the form $\{w_{v,c}\}_{c\in\mathbb{R}}$, where $w_{v,c}$ is defined by

(1.17)
$$w_{v,c}(s) = \min_{m} \{v(s_m) + cm\}$$
 for any $s = \sum_{m} s_m \in R$

and where the value $v(s_m)$ is defined at the top of this section. Note that $w_{v,c}$ is centered at the vertex o iff c > 0.

1.7. Valuations of linear growth and dreamy valuations

Following [22], we define the maximal vanishing order of (multisections of) L at $v \in X^{\text{an}}$ as

(1.18)
$$\mathbf{T}(v) \coloneqq \mathbf{T}_L(v) = \sup m^{-1} v(s) \in [0, +\infty],$$

where the supremum is over m sufficiently divisible and $s \in R_m \setminus \{0\}$. We say that v has linear growth if $T(v) < +\infty$; this notion is independent of the ample \mathbb{Q} -line bundle L. The set $X^{\text{lin}} \subset X^{\text{an}}$ of valuations of linear growth satisfies

$$X^{\operatorname{div}} \subset X^{\operatorname{lin}} \subset X^{\operatorname{val}}.$$

Further, setting

(1.19)
$$d_{\infty}(v,w) \coloneqq \sup m^{-1}|v(s) - w(s)|,$$

where the supremum is again over m sufficiently divisible and $s \in R_m \setminus \{0\}$, defines a metric on X^{lin} such that $(X^{\text{lin}}, d_{\infty})$ is complete (see [18, Section 11.3]). We refer to the d_{∞} -topology of X^{lin} as the strong topology.

Example 1.13. — If $\pi: X' \to X$ is a proper birational morphism, with X' normal, and $E \subset X'$ is a prime divisor which is Q-Cartier, then $T_L(E)$ coincides with the pseudoeffective threshold $\sup\{t \ge 0 \mid \pi^*L - tE \in \operatorname{Psef}(X')\}$ (see [22, Theorem 2.24]).

Any $v \in X^{\text{lin}}$ defines a norm $\chi_v \in \mathcal{N}_{\mathbb{R}}$, given by $\chi_v(s) \coloneqq v(s)$ for $s \in R_m$ with *m* sufficiently divisible. It satisfies $\lambda_{\max}(\chi_v) = T(v)$ (see (1.10)). Further, the map

$$X^{\operatorname{lin}} \longrightarrow \mathcal{N}_{\mathbb{R}}, \quad v \longmapsto \chi_v$$

is injective, because the function field of X coincides with the homogeneous fraction field of $R^{(d)}$ for any d sufficiently divisible.

For any $v \in X^{\text{lin}}$ and $c \in \mathbb{R}$, the norm $\chi_v + c$ can be viewed as a valuation on the affine cone Spec $R^{(d)}$ for d sufficiently divisible; it coincides with $w_{v,c}$ in the notation of (1.17). By Lemma 1.1, such norms are characterized as follows.

LEMMA 1.14. — A norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is of the form $\chi = \chi_v + c$ with $v \in X^{\text{lin}}$ and $c \in \mathbb{R}$ iff $\operatorname{gr}_{\chi} R^{(d)}$ is an integral domain for some (or any) sufficiently divisible d.

When $\chi \in \mathcal{T}_{\mathbb{R}}$ is of finite type, the latter condition means that the corresponding central fiber \mathcal{X}_0 is reduced and irreducible, see (1.14).

Example 1.15. — Suppose that a torus T acts on (X, L). By Example 1.10, each $\xi \in N_{\mathbb{R}}$ defines a norm $\chi_{\xi} \in \mathcal{T}_{\mathbb{R}}$ whose associated central fiber $\mathcal{X}_0 \simeq X$ is integral. By Lemma 1.14, χ_{ξ} thus determines a valuation $v_{\xi} \in X^{\text{lin}}$, which only depends on the T-action on X, and can be obtained by the "action" of $\xi \in N_{\mathbb{R}} \subset T^{\text{an}}$ on $v_{\text{triv}} \in X^{\text{an}}$ in the sense of "peaked points" (see [1, Section 5.2]).

In the terminology of [44], a divisorial valuation $v \in X^{\text{div}}$ such that χ_v is of finite type is called *dreamy* (with respect to L).

Example 1.16. — Assume X is normal and $E \subset X$ is a Q-Cartier prime divisor. If $v \coloneqq \operatorname{ord}_E$ is dreamy with respect to L, then the pseudoeffective threshold

$$\sup\{t \ge 0 \mid L - tE \in \operatorname{Psef}(X)\} = \operatorname{T}_L(v) = \lambda_{\max}(\chi_v)$$

is necessarily rational (cf. Example 1.13 and (1.13)). Examples with an irrational threshold are well-known (e.g. when X is an abelian surface of Picard number at least 2), and therefore provide simple examples of non-dreamy valuations.

The next result generates examples of divisorial valuations that are not dreamy for any polarization of X.

LEMMA 1.17. — Pick a dreamy valuation $v \in X^{\text{div}}$ (with respect to a given ample \mathbb{Q} -line bundle L), and assume that v is centered at a closed point $p \in X(k)$, with valuation ideals

$$\mathfrak{a}_m \coloneqq \{ f \in \mathcal{O}_{X,p} \mid v(f) \ge m \}.$$

Then the Rees algebra $\bigoplus_{m \in \mathbb{N}} \mathfrak{a}_m$ is of finite type over $\mathcal{O}_{X,p}$.

In particular, the (local) volume of v

$$\operatorname{vol}(v) = \lim_{m \to \infty} \frac{n!}{m^m} \operatorname{dim}(\mathcal{O}_{X,p}/\mathfrak{a}_m)$$

must be rational (see [41]).

Proof. — After replacing v with a multiple, we may assume that v is \mathbb{Z} -valued, and hence that χ_v is a \mathbb{Z} -filtration. By (1.12), the bigraded k-algebra $\bigoplus_{(\lambda,m)\in\mathbb{Z}\times\mathbb{N}}F^{\lambda}R_{dm}$ is finitely generated over k for d sufficiently divisible, and hence so is the graded subalgebra $\bigoplus_{m\in\mathbb{N}}F^mR_{dm}$.

On the other hand, by [22, Lemma 2.17], we can find $d \ge 1$ sufficiently divisible such that $\mathcal{O}_X(mdL) \otimes \mathfrak{a}_m$ is globally generated for all $m \in \mathbb{N}$. Since $\mathrm{H}^0(X, \mathcal{O}_X(mdL) \otimes \mathfrak{a}_m) = F^m R_{dm}$, we infer that $\bigoplus_{m \in \mathbb{N}} \mathfrak{a}_m$ is of finite type over $\mathcal{O}_{X,p}$.

Example 1.18. — Assume $k = \mathbb{C}$, dim $X \ge 4$, and pick a smooth point $p \in X(k)$. By [51], we can find a divisorial valuation $v \in X^{\text{div}}$ centered at p such that vol(v) is irrational. By Lemma 1.17, v is not dreamy with respect to any ample \mathbb{Q} -line bundle L on X.

1.8. Fubini–Study functions

A Fubini–Study function (for L) is a function $\varphi \in \mathbf{C}^0 = \mathbf{C}^0(X^{\mathrm{an}})$ of the form

(1.20)
$$\varphi = \frac{1}{m} \max_{j} \{ \log |s_j| + \lambda_j \},$$

with $m \ge 1$ such that mL is a (globally generated) line bundle, (s_j) a finite set of R_m without common zeros, and $\lambda_j \in \mathbb{R}$. Recall that (1.20) means $\varphi(v) = \frac{1}{m} \max_j \{-v(s_j) + \lambda_j\}$ for all $v \in X^{\mathrm{an}}$, see (1.16).

Remark 1.19. — The function φ defines a continuous metric $|\cdot|e^{-m\varphi}$ on the Berkovich analytification of mL. This metric is the pullback of a standard (non-Archimedean) Fubini–Study (or Weil) metric on $\mathcal{O}(1)$ under the morphism $X \to \mathbb{P}^N$ defined by $(s_j)_{0 \leq j \leq N}$, which explains the chosen terminology.

If the λ_j in (1.20) can be chosen in a subgroup $\Lambda \subset \mathbb{R}$, we say that φ is a Λ -Fubini–Study function, and write $\mathcal{H}_{\Lambda} = \mathcal{H}_{\Lambda}(L) \subset \mathbb{C}^0$ for the set of such functions. Thus

$$\{0\} = \mathcal{H}_{\{0\}} \subset \mathcal{H}_{\Lambda} \subset \mathcal{H}_{\mathbb{R}}.$$

Note that

(1.21)
$$\mathcal{H}_{\Lambda} = \mathcal{H}_{\mathbb{Q}\Lambda}$$

and $\mathcal{H}_{\Lambda}(dL) = d\mathcal{H}_{\Lambda}(L)$ for any $d \in \mathbb{Q}_{>0}$. The set \mathcal{H}_{Λ} is stable under finite max and under the action of $\mathbb{Q}\Lambda$ by translation.

Recall the action (1.15) of $\mathbb{R}_{>0}$ on functions on X^{an} . If φ is given by (1.20) and $t \in \mathbb{R}_{>0}$, then

$$t \cdot \varphi = \frac{1}{m} \max_{j} \{ \log |s_j| + t\lambda_j \}.$$

Thus $\mathcal{H}_{\mathbb{R}}$ is stable under the action of $\mathbb{R}_{>0}$, while \mathcal{H}_{Λ} is stable under the action of the stabilizer $\{t \in \mathbb{R}_{>0} \mid t\Lambda \subset \Lambda\}$. In particular, $\mathcal{H}_{\mathbb{Q}}$ is stable under the action of $\mathbb{Q}_{>0}$.

2. Homogenization and the Fubini–Study operator

In this section we study the homogenization of a norm, and the related Fubini–Study and infimum norm operators. We show that homogenization preserves norms of finite type, establish a 1–1 correspondence between homogeneous norms of finite type and Fubini–Study functions, and we prove Theorem A in the case $p = \infty$.

In what follows, \mathcal{L}^{∞} denotes the space of bounded functions $\varphi \colon X^{\mathrm{an}} \to \mathbb{R}$, endowed with its usual supnorm metric $\mathrm{d}_{\infty}(\varphi, \varphi') \coloneqq \sup_{X^{\mathrm{an}}} |\varphi - \varphi'|$.

2.1. Homogenization

In this section, $R = \bigoplus_{m \in \mathbb{N}} R_m$ denotes any reduced graded k-algebra.

DEFINITION 2.1. — We say that a norm $\chi \in \mathcal{N}_{\mathbb{R}}(R)$ is homogeneous if $\chi(f^d) = d\chi(f)$ for all $f \in R$ and $d \in \mathbb{N}$.

In multiplicative terminology, this means that $\|\cdot\|_{\chi} = e^{-\chi}$ is powermultiplicative, see [11]. It is easy to see that a norm $\chi \in \mathcal{N}_{\mathbb{R}}(R)$ is homogeneous iff the associated graded algebra $\operatorname{gr}_{\chi} R$ is reduced. We denote by

$$\mathcal{N}^{\mathrm{hom}}_{\mathbb{R}}(R) \subset \mathcal{N}_{\mathbb{R}}(R)$$

the set of homogeneous norms on R. For any Veronese subalgebra $R^{(d)} = \bigoplus_{m \in \mathbb{N}} R_{dm}, d \ge 1$, the restriction map $\mathcal{N}_{\mathbb{R}}(R) \to \mathcal{N}_{\mathbb{R}}(R^{(d)})$ induces a bijection

(2.1)
$$\mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}(R) \xrightarrow{\sim} \mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}(R^{(d)}).$$

Any norm $\chi \in \mathcal{N}_{\mathbb{R}}(R)$ is dominated by a minimal homogeneous norm, namely its homogenization χ^{hom} , defined by

(2.2)
$$\chi^{\text{hom}}(f) \coloneqq \sup_{d \ge 1} \frac{1}{d} \chi(f^d) = \lim_{d \to \infty} \frac{1}{d} \chi(f^d),$$

where the second equality holds by superadditivity of $d \mapsto \chi(f^d)$ and Fekete's Lemma. It is indeed easy to check that (2.2) defines a vector space norm on R that is superadditive, k^{\times} -invariant, linearly bounded and homogeneous, i.e. an element $\chi^{\text{hom}} \in \mathcal{N}_{\mathbb{R}}^{\text{hom}}(R)$.

Remark 2.2. — Note that $\|\cdot\|_{\chi^{\text{hom}}} = e^{-\chi^{\text{hom}}(\cdot)}$ is the spectral radius (semi)norm of $\|\cdot\|_{\chi}$ in the (multiplicative) terminology of [1].

Using standard but nontrivial results on k-affinoid algebras, we prove:

THEOREM 2.3. — Let $\chi \in \mathcal{N}_{\mathbb{R}}(R)$ be a norm generated in degree 1, with homogenization χ^{hom} . Then:

- (i) there exists C > 0 such that $\chi(f) \leq \chi^{\text{hom}}(f) \leq \chi(f) + C$ for all $f \in R$;
- (ii) the k-algebra $\operatorname{gr}_{\chi^{\text{hom}}} R$ is finitely generated;
- (iii) if $\chi \in \mathcal{N}_{\Lambda}(R)$ for a subgroup $\Lambda \subset \mathbb{R}$, then $\chi^{\text{hom}} \in \mathcal{N}_{\mathbb{Q}\Lambda}(R)$.

Proof. — Pick a surjective map of graded algebras $\pi: k[z] = k[z_1, \ldots, z_N] \to R$ and $\xi \in \mathbb{R}^N$ such that χ is the quotient norm of χ_{ξ} (see Lemma 1.4). As in Example 1.2, the completion of k[z] with respect to χ_{ξ} is the polydisc algebra $k\{z;\xi\}$, and π induces a surjection $k\{z;\xi\} \to \hat{R}$ onto the completion of R, whose norm is the quotient of the norm χ_{ξ} of $k\{z;\xi\}$.

As a consequence, \widehat{R} is a k-affinoid algebra in the sense of [1], corresponding geometrically to the affinoid domain $Y^{\mathrm{an}} \cap \mathbb{D}(r)$ of the Berkovich analytification $Y^{\mathrm{an}} \hookrightarrow \mathbb{A}^{N,\mathrm{an}}$ of the affine cone $Y \coloneqq \operatorname{Spec} R \hookrightarrow \mathbb{A}^N = \operatorname{Spec} k[z]$, where $\mathbb{D}(r) \subset \mathbb{A}^{N,\mathrm{an}}$ is the closed polydisc of polyradius $r = (\mathrm{e}^{-\xi_1}, \ldots, \mathrm{e}^{-\xi_N})$.

Since R is assumed to be reduced, it follows from the non-Archimedean GAGA principle that \hat{R} is reduced as well (see [40, Théorème 3.3]), and (i) is now a consequence of [1, Proposition 2.1.4(ii)], which states (in multiplicative terminology) that the spectral radius (semi)norm of any reduced k-affinoid algebra is equivalent to the given norm.

Next, note that $\operatorname{gr}_{\chi^{\operatorname{hom}}} \widehat{R}$ coincides, by definition, with the graded reduction of \widehat{R} in the sense of Temkin [66, Section 3]. By [66, Proposition 3.1], the surjection $k\{z;\xi\} \to \widehat{R}$ therefore induces a finite morphism $\operatorname{gr}_{\chi_{\xi}} k\{z;\xi\} \to$ $\operatorname{gr}_{\chi^{\operatorname{hom}}} \widehat{R}$. Now we have $\operatorname{gr}_{\chi_{\xi}} k\{z;\xi\} \simeq \operatorname{gr}_{\chi_{\xi}} k[z]$ and $\operatorname{gr}_{\chi^{\operatorname{hom}}} \widehat{R} \simeq \operatorname{gr}_{\chi^{\operatorname{hom}}} R$, see (1.1). Thus $\operatorname{gr}_{\chi^{\operatorname{hom}}} R$ is finite over $\operatorname{gr}_{\chi_{\xi}} k[z] \simeq k[T]$, which yields (ii).

Finally suppose that $\chi \in \mathcal{N}_{\Lambda}(R)$ for a subgroup $\Lambda \subset \mathbb{R}$. In this case, we can choose $\xi \in \Lambda^N$, so $k\{T;\xi\}$ and \widehat{R} are both Λ -strict k-affinoid algebras. By [67, 3.1.2.1 (iv)], we thus have $\chi^{\text{hom}}(\widehat{R} \setminus \{0\}) \subset \mathbb{Q}\Lambda$, which proves (iii).

2.2. Homogenization of norms on section rings

Returning to the setting of a polarized variety (X, L) and its space of norms $\mathcal{N}_{\mathbb{R}}$ (see Section 1.3), we introduce:

DEFINITION 2.4. — Consider a norm $\chi \in \mathcal{N}_{\mathbb{R}}$. Then:

- (i) we say that χ is homogeneous if it admits a homogeneous representative on $R^{(d)} = R(X, dL)$ for some d;
- (ii) we define the homogenization of χ as the norm $\chi^{\text{hom}} \in \mathcal{N}_{\mathbb{R}}$ induced by the homogenization of any representative of χ .

By (2.2), χ^{hom} is well-defined; it satisfies $\chi^{\text{hom}} \ge \chi$, and is characterized as the smallest homogeneous norm with this property.

Example 2.5. — For any $\chi \in \mathcal{N}_{\mathbb{R}}$ we have $\chi^{\text{hom}} = (\lfloor \chi \rfloor)^{\text{hom}}$. This is indeed a direct consequence of (2.2).

We denote by

 $\mathcal{N}^{\mathrm{hom}}_{\mathbb{R}} \subset \mathcal{N}_{\mathbb{R}}$

the subset of homogeneous norms. By (2.1), we have

(2.3)
$$\mathcal{N}_{\mathbb{R}}^{\mathrm{hom}} \simeq \mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}(R^{(d)})$$

for any d such that dL is a line bundle. In other words, a homogeneous norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is well-defined on $R_m = \mathrm{H}^0(X, mL)$ for any m such that mLis a line bundle.

Remark 2.6. — An element of $\chi \in \mathcal{N}_{\mathbb{R}}$ is a norm on $\mathbb{R}^{(d)}$ for some sufficiently divisible d that depends on χ , so in general it does not make sense to talk about pointwise convergence of sequences or nets in $\mathcal{N}_{\mathbb{R}}$. By (2.3), it does however make sense when the norms are homogeneous, as they are then defined on $\mathbb{R}^{(d)}$ any fixed d such that dL is an honest line bundle.

The subset $\mathcal{N}_{\mathbb{R}}^{\text{hom}}$ is stable under minima, and under the scaling action of $\mathbb{R}_{>0}$ and the additive action of \mathbb{R} . For any subgroup $\Lambda \subset \mathbb{R}$, we set

$$\mathcal{N}^{\mathrm{hom}}_{\Lambda}\coloneqq\mathcal{N}_{\Lambda}\cap\mathcal{N}^{\mathrm{hom}}_{\mathbb{R}}$$

Recall that $\mathcal{N}_{\mathbb{R}}$ is equipped with a pseudo-metric d_{∞} , see (1.9). Using (2.2), it is straightforward to check:

LEMMA 2.7. — Homogenization $\chi \mapsto \chi^{\text{hom}}$ defines a projection $\mathcal{N}_{\mathbb{R}} \twoheadrightarrow \mathcal{N}_{\mathbb{R}}^{\text{hom}}$ which is equivariant for the actions of $\mathbb{R}_{>0}$ and \mathbb{R} , commutes with minima, and satisfies

$$d_{\infty}(\chi^{\text{hom}},\chi'^{\text{hom}}) \leqslant d_{\infty}(\chi,\chi'), \quad \lambda_{\max}(\chi^{\text{hom}}) = \lambda_{\max}(\chi)$$

for all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$.

The restriction of d_{∞} to $\mathcal{N}_{\mathbb{R}}^{\text{hom}}$ is further well-behaved:

PROPOSITION 2.8. — The restriction of d_{∞} to $\mathcal{N}_{\mathbb{R}}^{hom}$ is a metric. Furthermore, the metric space $(\mathcal{N}_{\mathbb{R}}^{hom}, d_{\infty})$ is complete, and contains $\mathcal{N}_{\mathbb{Z}}^{hom}$ as a closed subset.

Note that $\mathcal{N}_{\mathbb{Z}}^{\text{hom}}$ is always a strict subset of $\mathcal{N}_{\mathbb{R}}^{\text{hom}}$, thanks to the scaling action of $\mathbb{R}_{>0}$. In contrast, recall that $\mathcal{N}_{\mathbb{Z}}$ is d_{∞} -dense in $\mathcal{N}_{\mathbb{R}}$ (see Example 1.7).

LEMMA 2.9. — Pick $d \ge 1$ such that dL is an honest line bundle, and view d_{∞} as a pseudo-metric on $\mathcal{N}^{\text{hom}}(R^{(d)})$ via (2.3). For all $\chi, \chi' \in \mathcal{N}^{\text{hom}}(R^{(d)})$ we then have

$$\mathrm{d}_{\infty}(\chi,\chi') = \sup_{m \ge 1} \frac{1}{md} \,\mathrm{d}_{\infty}(\chi|_{R_{md}},\chi'|_{R_{md}}).$$

Proof. — By homogeneity of χ, χ' , we have for all $m, l \ge 1$

$$d_{\infty}(\chi|_{R_{md}},\chi'|_{R_{md}}) = \sup_{s \in R_{md} \smallsetminus \{0\}} |\chi(s) - \chi'(s)|$$

= $l^{-1} \sup_{s \in R_{lmd} \smallsetminus \{0\}} |\chi(s^{l}) - \chi'(s^{l})|$
 $\leq l^{-1} \sup_{t \in R_{lmd} \smallsetminus \{0\}} |\chi(t) - \chi'(t)|$
= $l^{-1} d_{\infty}(\chi|_{R_{lmd}},\chi'|_{R_{lmd}}).$

Thus $m \mapsto \frac{1}{md} d_{\infty}(\chi|_{R_{md}}, \chi'|_{R_{md}})$ is increasing with respect to divisibility, and (1.9) yields the result (recall that the limsup in the latter formula is understood with respect to the divisibility order).

Proof of Proposition 2.8. — Pick d as in Lemma 2.9. For each $m \ge 1$, $(\mathcal{N}_{\mathbb{R}}(R_{md}), d_{\infty})$ is a complete metric space, in which $\mathcal{N}_{\mathbb{Z}}(R_{md})$ sits as a closed subspace. This implies that $\mathcal{N}_{\mathbb{R}}(R^{(d)}) \hookrightarrow \prod_{m \ge 1} \mathcal{N}_{\mathbb{R}}(R_{md})$ is complete with respect to the metric

$$\widetilde{\mathrm{d}}_{\infty}(\chi,\chi') \coloneqq \sup_{m \geqslant 1} \frac{1}{md} \, \mathrm{d}_{\infty}(\chi|_{R_{md}},\chi'|_{R_{md}}),$$

and that $\mathcal{N}_{\mathbb{Z}}(R^{(d)}) \hookrightarrow \prod_{m \ge 1} \mathcal{N}_{\mathbb{Z}}(R_{md})$ is closed. It is also clear that $\mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}(R^{(d)})$ is closed in $\mathcal{N}_{\mathbb{R}}(R^{(d)})$ with respect to \widetilde{d}_{∞} , so the result now follows from Lemma 2.9.

Note that Lemma 2.9 ensures compatibility of the d_{∞} -metrics on $\mathcal{N}_{\mathbb{R}}^{\text{hom}}$ and X^{lin} (see (1.19)):

COROLLARY 2.10. — The map $v \mapsto \chi_v$ defines an isometric embedding $(X^{\text{lin}}, \mathbf{d}_{\infty}) \hookrightarrow (\mathcal{N}_{\mathbb{R}}^{\text{hom}}, \mathbf{d}_{\infty})$, i.e. $\mathbf{d}_{\infty}(v, w) = \mathbf{d}_{\infty}(\chi_v, \chi_w)$ for all $v, w \in X^{\text{lin}}$.

For any subgroup $\Lambda \subset \mathbb{R}$, we denote by

$$\mathcal{T}^{\mathrm{hom}}_{\Lambda} \coloneqq \mathcal{T}_{\Lambda} \cap \mathcal{N}^{\mathrm{hom}}_{\mathbb{R}} = \mathcal{T}_{\mathbb{R}} \cap \mathcal{N}^{\mathrm{hom}}_{\Lambda}$$

the set of homogeneous Λ -valued norms of finite type. As a straightforward consequence of Theorem 2.3, we get:

LEMMA 2.11. — For any $\chi \in \mathcal{T}_{\mathbb{R}}$, the following holds:

- (i) $\chi^{\text{hom}} \in \mathcal{T}^{\text{hom}}_{\mathbb{R}}$;
- (ii) for *m* sufficiently divisible, $d_{\infty}(\chi|_{R_m}, \chi^{\text{hom}}|_{R_m})$ is bounded, and hence $d_{\infty}(\chi, \chi^{\text{hom}}) = 0$;
- (iii) for any subgroup $\Lambda \subset \mathbb{R}, \chi \in \mathcal{T}_{\Lambda} \Longrightarrow \chi^{\text{hom}} \in \mathcal{T}_{\mathbb{Q}\Lambda}^{\text{hom}}$.

As we shall see, homogenization in fact maps \mathcal{T}_{Λ} onto $\mathcal{T}_{\mathbb{Q}\Lambda}^{\text{hom}}$ (cf. Corollary 2.18). For $\Lambda = \mathbb{Z}$, the homogenization map $\mathcal{T}_{\mathbb{Z}} \twoheadrightarrow \mathcal{T}_{\mathbb{Q}}^{\text{hom}}$ is closely related to integral closure (see Appendix A for a detailed discussion).

2.3. The Fubini–Study operator

Assume first that L is a globally generated line bundle. To any norm χ on $R_1 = \mathrm{H}^0(X, L)$, we associate a function on X^{an} by setting

(2.4)
$$\operatorname{FS}_{L}(\chi) \coloneqq \sup_{s \in R_{1} \smallsetminus \{0\}} \{ \log |s| + \chi(s) \},$$

i.e. $FS_L(\chi)(v) = \sup_{s \in R_1 \setminus \{0\}} \{-v(s) + \chi(s)\}$ for $v \in X^{an}$. Given a χ -orthogonal basis (s_i) of R_1 , one easily checks that

(2.5)
$$\operatorname{FS}_{L}(\chi) = \max_{i} \{ \log |s_{i}| + \chi(s_{i}) \} \in \mathcal{H}_{\mathbb{R}},$$

see [14, Lemma 7.17]. This implies

(2.6)
$$\lambda_{\max}(\chi) = \sup_{X^{\mathrm{an}}} \mathrm{FS}_L(\chi) = \mathrm{FS}_L(\chi)(v_{\mathrm{triv}}),$$

as well as

(2.7)
$$\chi \in \mathcal{N}_{\Lambda}(R_1) \Longrightarrow \mathrm{FS}_L(\chi) \in \mathcal{H}_{\Lambda}.$$

for any subgroup $\Lambda \subset \mathbb{R}$.

LEMMA 2.12. — Assume that L is a line bundle, and let χ be a norm on R = R(X, L). For each $m \ge 1$ we then have

$$\operatorname{FS}_{mL}(\chi|_{R_m}) \ge m \operatorname{FS}_L(\chi|_{R_1}),$$

and equality holds if χ is generated in degree 1.

Proof. — For each $s \in R_1 \setminus \{0\}$ we have $\chi(s^m) \ge m\chi(s)$, and the inequality follows, by (2.4). Assume that χ is generated in degree 1. To get equality, we need to show

$$v(s) \ge \chi(s) - m \operatorname{FS}_L(\chi_1)(v)$$

ANNALES DE L'INSTITUT FOURIER

for all $s \in R_m \setminus \{0\}$ and $v \in X^{\mathrm{an}}$. To see this, pick an orthogonal basis (s_i) of R_1 , and write $s = \sum_{|\alpha|=m} c_{\alpha} \prod_i s_i^{\alpha_i}$ with $\chi(s) = \min_{\alpha} \sum_i \alpha_i \chi(s_i)$ for some α with $c_{\alpha} \neq 0$. Then $\mathrm{FS}_L(\chi)(v) = \max_i \{\chi(s_i) - v(s_i)\}$, and hence

$$v(s) \ge \min_{c_{\alpha} \neq 0} \sum_{i} \alpha_{i} v(s_{i})$$
$$\ge \min_{c_{\alpha} \neq 0} \sum_{i} \alpha_{i} \left(\chi(s_{i}) - \mathrm{FS}_{L}(\chi_{1})(v) \right) = \chi(s) - m \, \mathrm{FS}_{L}(\chi_{1})(v),$$

which concludes the proof.

Returning to the general case of a \mathbb{Q} -line bundle, pick $\chi \in \mathcal{N}_{\mathbb{R}}$, and set

$$\operatorname{FS}_m(\chi) \coloneqq m^{-1} \operatorname{FS}_{mL}(\chi|_{R_m}) \in \mathcal{H}_{\mathbb{R}}$$

for m sufficiently divisible. By Lemma 2.12, $FS_m(\chi)$ is an increasing function of m with respect to divisibility, and is further uniformly bounded, by linear boundedness of χ . We may thus introduce:

DEFINITION 2.13. — The Fubini–Study operator FS: $\mathcal{N}_{\mathbb{R}} \to \mathcal{L}^{\infty}$ takes a norm $\chi \in \mathcal{N}_{\mathbb{R}}$ to the bounded function $FS(\chi): X^{an} \to \mathbb{R}$ defined as the pointwise limit

$$FS(\chi) \coloneqq \lim_{m} FS_m(\chi) = \sup_{m} FS_m(\chi).$$

Recall that \mathcal{L}^{∞} denotes the space of bounded functions on X^{an} . The bounded function $\mathrm{FS}(\chi)$ is lsc (lower semicontinuous), being a supremum of continuous functions; it is however not continuous in general (see Theorem 2.19 below). By (2.6) we have

(2.8)
$$\lambda_{\max}(\chi) = FS(\chi)(v_{\text{triv}}).$$

Note also that the canonical approximants $\chi_d \in \mathcal{T}_{\mathbb{R}}$ of any norm $\chi \in \mathcal{N}_{\mathbb{R}}$ satisfy

(2.9)
$$FS(\chi_d) = FS_d(\chi) \in \mathcal{H}_{\mathbb{R}}$$

for all *d* sufficiently divisible, by Lemma 2.12. In particular, if $\chi \in \mathcal{T}_{\mathbb{R}}$ is of finite type then the limit in Definition 2.13 is stationary.

The Fubini–Study operator FS: $\mathcal{N}_{\mathbb{R}} \to \mathcal{L}^{\infty}$ is increasing with respect to the partial orderings on $\mathcal{N}_{\mathbb{R}}$ and \mathcal{L}^{∞} , and equivariant under the actions of \mathbb{R} and $\mathbb{R}_{>0}$. It is also easily seen to be 1-Lipschitz with respect to the d_{∞} -(pseudo)metrics, i.e.

(2.10)
$$d_{\infty}(FS(\chi), FS(\chi')) \leq d_{\infty}(\chi, \chi') \text{ for all } \chi, \chi' \in \mathcal{N}_{\mathbb{R}}.$$

As we see below, equality holds when χ, χ' are homogeneous (see Corollary 2.17).

The next result shows that FS is invariant under homogenization.

 \square

PROPOSITION 2.14. — For any $\chi \in \mathcal{N}_{\mathbb{R}}$ we have $FS(\chi) = FS(\chi^{\text{hom}})$.

Proof. — That $FS(\chi) \leq FS(\chi^{\text{hom}})$ is clear, since $\chi \leq \chi^{\text{hom}}$. Now let $v \in X^{\text{an}}$ and $\varepsilon > 0$. Successively pick $m \geq 1$ sufficiently divisible such that $FS(\chi^{\text{hom}})(v) \leq FS_m(\chi^{\text{hom}})(v) + \varepsilon$, then $s \in R_m \setminus \{0\}$ such that $mFS_m(\chi^{\text{hom}})(v) \leq \chi^{\text{hom}}(s) - v(s) + m\varepsilon$, and finally $d \geq 1$ such that $\chi^{\text{hom}}(s) \leq \frac{1}{d}\chi(s^d) + m\varepsilon$, see (2.2). Then

$$FS(\chi^{hom})(v) \leq \frac{1}{md}(\chi(s^d) - v(s^d)) + 3\varepsilon \leq FS_{md}(\chi)(v) + 3\varepsilon \leq FS(\chi)(v) + 3\varepsilon,$$

completing the proof.

The Fubini–Study operator relates norms of finite type and Fubini–Study functions, as follows:

PROPOSITION 2.15. — For any subgroup $\Lambda \subset \mathbb{R}$, we have

$$\mathrm{FS}(\mathcal{T}_{\Lambda}) = \mathrm{FS}(\mathcal{T}_{\mathbb{Q}\Lambda}) = \mathrm{FS}(\mathcal{T}_{\mathbb{Q}\Lambda}^{\mathrm{hom}}) = \mathcal{H}_{\Lambda} = \mathcal{H}_{\mathbb{Q}\Lambda}$$

As we shall see, the map FS: $\mathcal{T}_{\mathbb{Q}\Lambda}^{\text{hom}} \to \mathcal{H}_{\Lambda}$ is further 1–1 (see Corollary 2.18).

Proof. — If $\chi \in \mathcal{T}_{\Lambda}$ then $\chi = \chi_m$ for m sufficiently divisible, and hence $FS(\chi) = FS(\chi_m) = FS_m(\chi) \in \mathcal{H}_{\Lambda}$ by (2.7). Conversely, pick $\varphi \in \mathcal{H}_{\Lambda}$, and write $\varphi = d^{-1} \max_i \{ \log |s_i| + \lambda_i \}$ with $d \ge 1$, a finite family $(s_i)_{1 \le i \le N}$ in R_d without common zeros, and $\lambda_i \in \Lambda$. After enlarging the family (s_i) and choosing the corresponding $\lambda_i \ll 0$ in Λ , we may further assume that (s_i) spans R_d . Consider the surjective map $k^N \to R_d$ that takes the canonical basis (e_i) to (s_i) . Denote by χ_0 the norm on k^N that is diagonal in (e_i) , with $\chi_0(e_i) = \lambda_i$, and let $\chi_d \in \mathcal{N}_{\mathbb{R}}(R_d)$ be the quotient norm. It is then easy to check (see [14, Lemma 7.17]) that

$$\varphi = d^{-1} \max_{j} \{ \log |s_j| + \lambda_j \} = d^{-1} \operatorname{FS}_{dL}(\chi_d).$$

By Lemma 2.12, the norm $\chi \in \mathcal{N}_{\Lambda}(R^{(d)})$ generated in degree 1 by χ_d satisfies $FS(\chi) = \varphi$, which proves $FS(\mathcal{T}_{\Lambda}) = \mathcal{H}_{\Lambda}$. By Proposition 2.14 and Lemma 2.11, we infer

$$\mathcal{H}_{\Lambda} = \mathrm{FS}(\mathcal{T}_{\Lambda}) \subset \mathrm{FS}(\mathcal{T}_{\mathbb{Q}\Lambda}^{\mathrm{hom}}) \subset \mathcal{H}_{\mathbb{Q}\Lambda},$$
which concludes the proof since $\mathcal{H}_{\mathbb{Q}\Lambda} = \mathcal{H}_{\Lambda}$, see (1.21).

2.4. The infimum norm and homogenization

Next we define an operator

IN:
$$\mathcal{L}^{\infty} \longrightarrow \mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}$$

ANNALES DE L'INSTITUT FOURIER

that to a bounded function φ on X^{an} attaches a homogeneous norm, the infimum norm $\chi = \text{IN}(\varphi)$. For any $m \in \mathbb{N}$ such that mL is a line bundle, it is defined on R_m by

(2.11)
$$\chi(s) \coloneqq \inf_{v \in X^{\mathrm{an}}} \{ v(s) + m\varphi(v) \} = \inf_{X^{\mathrm{an}}} \{ m\varphi - \log |s| \}.$$

In "multiplicative" notation, this is simply the usual supremum norm

$$\|s\|_{\varphi} = \sup_{X^{\mathrm{an}}} |s| \,\mathrm{e}^{-m\varphi},$$

compare for instance [14, Section 6]. The operator $IN = IN_L$ is increasing, and equivariant for the actions of \mathbb{R} and $\mathbb{R}_{>0}$, i.e.

(2.12)
$$\operatorname{IN}(\varphi + c) = \operatorname{IN}(\varphi) + c, \quad \operatorname{IN}(t \cdot \varphi) = t \operatorname{IN}(\varphi)$$

for $\varphi \in \mathcal{L}^{\infty}$, $c \in \mathbb{R}$ and $t \in \mathbb{R}_{>0}$. For any $\varphi, \varphi' \in \mathcal{L}^{\infty}$, it is also easy to see that

(2.13)
$$\operatorname{IN}(\varphi \wedge \varphi') = \operatorname{IN}(\varphi) \wedge \operatorname{IN}(\varphi'),$$

(2.14)
$$d_{\infty}(IN(\varphi), IN(\varphi')) \leq d_{\infty}(\varphi, \varphi')$$

(2.15)
$$\operatorname{IN}(\varphi) = \operatorname{IN}(\varphi_{\star}),$$

where $\varphi_{\star} \in \mathcal{L}^{\infty}$ denotes the *lsc regularization* of φ , i.e. the largest *lsc* function such that $\varphi_{\star} \leq \varphi$. The following key result relates homogenization and infimum norms:

THEOREM 2.16. — For any $\chi \in \mathcal{N}_{\mathbb{R}}$, we have $IN(FS(\chi)) = \chi^{hom}$.

COROLLARY 2.17. — For all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$ we have $d_{\infty}(FS(\chi), FS(\chi')) = d_{\infty}(\chi^{\text{hom}}, \chi'^{\text{hom}})$. In particular, the Fubini–Study operator defines an isometric embedding of complete metric spaces

$$FS\colon (\mathcal{N}^{\hom}_{\mathbb{R}}, d_{\infty}) \hookrightarrow (\mathcal{L}^{\infty}, d_{\infty}).$$

Recall that d_{∞} also denotes the supnorm metric on the space \mathcal{L}^{∞} of bounded functions on X^{an} .

The following result settles the case $p = \infty$ of Theorem A in the introduction:

COROLLARY 2.18. — For any subgroup $\Lambda \subset \mathbb{R}$, the Fubini–Study operator defines a surjective isometry FS: $(\mathcal{T}_{\Lambda}, d_{\infty}) \twoheadrightarrow (\mathcal{H}_{\Lambda}, d_{\infty})$, which factors as

- a surjective isometry $(\mathcal{T}_{\Lambda}, d_{\infty}) \twoheadrightarrow (\mathcal{T}_{\mathbb{Q}\Lambda}^{hom}, d_{\infty})$ defined by homogenization;
- an isomorphism FS: $(\mathcal{T}_{\mathbb{Q}\Lambda}^{hom}, d_{\infty}) \xrightarrow{\sim} (\mathcal{H}_{\Lambda}, d_{\infty})$, with inverse IN: $(\mathcal{H}_{\Lambda}, d_{\infty}) \xrightarrow{\sim} (\mathcal{T}_{\mathbb{Q}\Lambda}^{hom}, d_{\infty})$.

For any homogeneous norm $\chi \in \mathcal{N}_{\mathbb{R}}^{\text{hom}}$, we further have $FS(\chi) \in \mathcal{H}_{\Lambda} \Leftrightarrow \chi \in \mathcal{T}_{\mathbb{Q}\Lambda}$.

The last equivalence fails in general for non-homogeneous norms, see Example 2.22

The proof of Theorem 2.16 uses the Berkovich maximum modulus principle as well as the remarks in Section 1.6.

Proof of Theorem 2.16. — After passing to a multiple, we may assume that L is a line bundle and χ is a norm on R = R(X, L). Let M be the Berkovich spectrum of the normed ring (R, χ) , i.e. the set of semivaluations $w: R \to \mathbb{R} \cup \{+\infty\}$ such that $w \ge \chi$. Geometrically, M sits as a compact subset of the analytification Y^{an} of the affine cone Y = Spec R, and is obtained as the image of the unit disc bundle in the total space of L^{\vee} , i.e. the blowup of $o \in Y$ (compare [42]).

Since homogenization corresponds to the spectral radius seminorm construction (see Remark 2.2), the Berkovich maximum modulus principle [1, Theorem 1.3.1] (applied to the completion of (R, χ)) yields $\chi^{\text{hom}}(f) = \min_{w \in M} w(f)$ for any $f \in R$, where the infimum is attained by compactness of M. In particular, for any $s \in R_m \setminus \{0\}, m \ge 1$, we have

$$\chi^{\hom}(s) = \min_{w \in M} w(s).$$

Let M^{inv} be the set of k^{\times} -invariant semivaluations in M. We have a projection $p: M \to M^{inv}$ defined by

$$p(w)\left(\sum_{m} s_{m}\right) = \min_{m} w(s_{m})$$

where $s_m \in R_m$. Thus $p(w) \leq w$, so in the formula for $\chi^{\text{hom}}(s)$, it suffices to take the infimum over $w \in M^{\text{inv}}$. As in Section 1.6 consider the projection $\pi^{\text{an}}: Y^{\text{an}} \setminus \{w_o\} \to X^{\text{an}}$, where w_o is the trivial semivaluation at the vertex of the cone; this satisfies $w_o(f) = +\infty$ for $f \in \bigoplus_{m>0} R_m$ and $w_o(f) = 0$ for $f \in R \setminus \bigoplus_{m>0} R_m$. For any $v \in X^{\text{an}}$, the set of k^{\times} -invariant points in $(\pi^{\text{an}})^{-1}(v)$ is of the form $\{w_{v,c}\}_{c\in\mathbb{R}}$, where $w_{v,c}$ is the unique k^{\times} -invariant point such that

$$w_{v,c}(s) = v(s) + cm$$

for $s \in R_m$, $m \ge 1$, see (1.17).

It follows that $M^{\text{inv}} \subset M \subset Y^{\text{an}}$ is the set of semivaluations $w_{v,c}$, where $v \in X^{\text{an}}$ and $v(s) + mc \ge \chi(s)$ for all $s \in R_m$, $m \ge 1$. Note that this condition on c means precisely that $c \ge FS(\chi)(v)$. Altogether, this means

that if $s \in R_m$, $m \ge 1$, then

$$\chi^{\text{hom}}(s) = \inf\{v(s) + mc \mid v \in X^{\text{an}}, c \ge \text{FS}(\chi)(v)\}$$
$$= \inf\{v(s) + m \text{FS}(\chi)(v) \mid v \in X^{\text{an}}\}$$
$$= \text{IN}(\text{FS}(\chi))(s),$$

which completes the proof.

Proof of Corollary 2.17. — By Proposition 2.14, we may assume that χ, χ' are homogeneous. By Theorem 2.16, (2.14) and (2.10), we then have $d_{\infty}(\chi,\chi') = d_{\infty}(IN(FS(\chi)), IN(FS(\chi'))) \leqslant d_{\infty}(FS(\chi), FS(\chi')) \leqslant d_{\infty}(\chi,\chi').$ \square

Thus equality holds everywhere, and the result follows.

Proof of Corollary 2.18. — By Corollary 2.17 and Lemma 2.11, the Fubini–Study operator defines a surjective isometry FS: $(\mathcal{T}_{\Lambda}, d_{\infty}) \twoheadrightarrow (\mathcal{H}_{\Lambda}, d_{\infty}),$ which factors as homogenization followed by FS: $(\mathcal{T}_{\mathbb{Q}\Lambda}^{\mathrm{hom}}, \mathrm{d}_{\infty}) \xrightarrow{\sim} (\mathcal{H}_{\Lambda}, \mathrm{d}_{\infty})$, whose inverse is necessarily given by IN, by Theorem 2.16. This implies that homogenization defines a surjective isometry $(\mathcal{T}_{\Lambda}, d_{\infty}) \twoheadrightarrow (\mathcal{T}_{\mathbb{Q}\Lambda}^{hom}, d_{\infty}).$

The final assertion follows, by injectivity of FS on $\mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}.$ \square

2.5. Continuous norms

Building on the previous results, we are now in a position to characterize the d_{∞} -closure of the set $\mathcal{T}_{\mathbb{Z}}$ of test configurations, as follows.

THEOREM 2.19. — For any norm $\chi \in \mathcal{N}_{\mathbb{R}}$, the following are equivalent:

- (i) χ lies in the d_∞-closure of $\mathcal{T}_{\mathbb{Z}}$;
- (ii) χ lies in the d_∞-closure of $\mathcal{T}_{\mathbb{R}}$;
- (iii) the canonical approximants (χ_d) satisfy $d_{\infty}(\chi_d, \chi) \to 0$;
- (iv) $FS(\chi)$ is continuous;
- (v) $FS(\chi_d) \to FS(\chi)$ uniformly on X^{an} .

DEFINITION 2.20. — We say that $\chi \in \mathcal{N}_{\mathbb{R}}$ is continuous when the equivalent properties of Theorem 2.19 holds.

The set $\mathcal{N}_{\mathbb{R}}^{cont}$ of continuous norms is thus the d_{∞} -closure of $\mathcal{T}_{\mathbb{Z}}$ (or $\mathcal{T}_{\mathbb{R}}$); it is a strict subset of $\mathcal{N}_{\mathbb{R}}$ as soon as dim $X \ge 1$ (see Example 2.21 below).

Proof of Theorem 2.19. — We trivially have (i) \Rightarrow (ii). Assume (ii), and pick $\varepsilon > 0$. In view of (1.9), we can find $\chi' \in \mathcal{T}_{\mathbb{R}}$ and $d \ge 1$ such that

(2.16)
$$d_{\infty}(\chi|_{R_{md}},\chi'|_{R_{md}}) \leq md\varepsilon \text{ for all } m \geq 1.$$

 \square
Replacing d with a multiple, we can further assume $\chi'_d = \chi'$. For m = 1, (2.16) yields $\chi' \leq \chi + d\varepsilon$ on R_d , and hence $\chi' = \chi'_d \leq \chi_d + md\varepsilon$ on R_{md} for all $m \geq 1$. On the other hand, (2.16) yields $\chi \leq \chi' + md\varepsilon$ on R_{md} , which proves $\chi_d \leq \chi \leq \chi_d + 2md\varepsilon$ on R_{md} for all $m \geq 1$. This implies $d_{\infty}(\chi_d, \chi) \leq \varepsilon$. This proves (ii) \Rightarrow (iii), the converse being obvious since $\chi_d \in \mathcal{T}_{\mathbb{R}}$.

Assume (ii). To prove (i), we may replace χ with its round-down and assume $\chi \in \mathcal{N}_{\mathbb{Z}}$ (see Example 1.7). Its canonical approximants then satisfy $\chi_d \in \mathcal{T}_{\mathbb{Z}}$, and hence (ii) \Rightarrow (i), thanks to (iii).

Since $FS(\chi)$ is the pointwise limit of the increasing net of continuous functions $(FS(\chi_d))$, Dini's lemma yields (iv) \Leftrightarrow (v). Finally, we claim that

(2.17)
$$d_{\infty}(\chi_d, \chi) = d_{\infty}(FS(\chi_d), FS(\chi))$$

for d sufficiently divisible, which will show (iii) \Leftrightarrow (v). Note that

$$\chi_d \leq \chi \leq \chi^{\text{hom}} \Longrightarrow \mathrm{d}_{\infty}(\chi_d, \chi) \leq \mathrm{d}_{\infty}(\chi_d, \chi^{\text{hom}}),$$

see (1.3). Now $d_{\infty}(\chi_d, (\chi_d)^{\text{hom}})) = 0$ (see Lemma 2.11), and hence

$$d_{\infty}(\chi_d, \chi^{\text{hom}}) = d_{\infty}((\chi_d)^{\text{hom}}, \chi^{\text{hom}}) = d_{\infty}(\text{FS}(\chi_d), \text{FS}(\chi)).$$

by Corollary 2.17. This shows $d_{\infty}(\chi_d, \chi) \leq d_{\infty}(FS(\chi_d), FS(\chi))$, while the converse holds by (2.10). This shows (2.17), and concludes the proof. \Box

Example 2.21. — To each subvariety $Z \subsetneq X$ we associate a norm $\chi = \chi_Z \in \mathcal{N}_{\mathbb{Z}}^{\text{hom}}$ by setting, for each nonzero section $s \in R_m$ with m sufficiently divisible

$$\chi(s) = \begin{cases} m & \text{if } s|_Z \equiv 0, \\ 0 & \text{otherwise.} \end{cases}$$

We claim that χ is not continuous. Indeed, using the description of χ_d as a quotient norm, it is easy to check that any $s \in R_{dm}$ locally lies in I_Z^p with $p \coloneqq \chi_d(s) \in \mathbb{N}$. Choosing $s \in R_{dm} = \mathrm{H}^0(X, dmL)$ that locally belongs to I_Z but not I_Z^2 (which is possible for any *m* large enough, since *L* is ample), we get $\chi_d(s) = 1$, while $\chi(s) = dm$. This shows $\mathrm{d}_{\infty}(\chi|_{R_{dm}}, \chi_d|_{R_{dm}}) \geq dm-1$, and hence $\mathrm{d}_{\infty}(\chi, \chi_d) \geq 1$, which proves the claim.

Alternatively, one can show that $FS(\chi)$ is identically 1 on $X^{an} \setminus Z^{an}$, and 0 on Z^{an} , and hence is not continuous.

Example 2.22. — As a variant of Example 2.21, consider the norm $\chi \in \mathcal{N}_{\mathbb{R}}$ defined for $s \in R_m \setminus \{0\}$ by

$$\chi(s) = \begin{cases} m & \text{if } s|_Z \equiv 0, \\ m - \sqrt{m} & \text{otherwise,} \end{cases}$$

which is indeed a norm, by subadditivity of $m \mapsto \sqrt{m}$. Then $\chi^{\text{hom}} = \chi_{\text{triv}} + 1$, and hence $\text{FS}(\chi) = \text{FS}(\chi^{\text{hom}}) \equiv 1$. In particular, $\text{FS}(\chi) \in \mathcal{H}_{\mathbb{R}}$; however, χ is not of finite type. Indeed,

$$\mathbf{d}_{\infty}(\chi|_{R_m}, \chi^{\mathrm{hom}}|_{R_m}) = \sqrt{m}$$

is not bounded (see Lemma 2.11).

Remark 2.23. — It follows from Example 2.21 that the set $\mathcal{T}_{\mathbb{Z}}$ of test configurations is never d_{∞} -dense in $\mathcal{N}_{\mathbb{R}}$ when dim $X \ge 1$. In contrast, $\mathcal{T}_{\mathbb{Z}}$ is dense with respect to any of the weaker pseudometrics $d_p, p \in [1, \infty)$ to be introduced in Section 3 (see Corollary 3.19).

We next analyze the behavior of homogenization on continuous norms.

PROPOSITION 2.24. — For each $\chi \in \mathcal{N}_{\mathbb{R}}$, we have

$$\chi \in \mathcal{N}_{\mathbb{R}}^{\mathrm{cont}} \iff \chi^{\mathrm{hom}} \in \mathcal{N}_{\mathbb{R}}^{\mathrm{cont}} \Longrightarrow \mathrm{d}_{\infty}(\chi, \chi^{\mathrm{hom}}) = 0.$$

Further, homogenization induces a surjective isometry

$$(\mathcal{N}_{\mathbb{R}}^{\mathrm{cont}}, \mathrm{d}_{\infty}) \longrightarrow (\mathcal{N}_{\mathbb{R}}^{\mathrm{cont}, \mathrm{hom}}, \mathrm{d}_{\infty}).$$

Here $\mathcal{N}_{\mathbb{R}}^{\mathrm{cont,hom}} \coloneqq \mathcal{N}_{\mathbb{R}}^{\mathrm{cont}} \cap \mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}$ denotes the set of continuous homogeneous norms.

Proof. — The first equivalence follows from Proposition 2.14 and Theorem 2.19(iv). By Lemma 2.11, $d_{\infty}(\chi, \chi^{\text{hom}}) = 0$ holds on $\mathcal{T}_{\mathbb{R}}$. By d_{∞} continuity of homogenization (see Lemma 2.7), this extends to the d_{∞} closure $\mathcal{N}_{\mathbb{R}}^{\text{hom}}$. This proves the second implication, which in turn yields the
last point, by the triangle inequality.

When $\chi \in \mathcal{N}_{\mathbb{R}}$ is not continuous, the property $d_{\infty}(\chi, \chi^{\text{hom}}) = 0$ fails in general; in other words, homogenization is not a d_{∞} -isometry on the whole space $\mathcal{N}_{\mathbb{R}}$:

Example 2.25. — Pick a subvariety $Z \subsetneq X$, and set for $s \in R_m \setminus \{0\}$

$$\chi(s) = \begin{cases} m & \text{if } s \in I_Z^2, \\ m/2 & \text{if } s \in I_Z \setminus I_Z^2, \\ 0 & \text{if } s \notin I_Z. \end{cases}$$

As one easily checks, this defines a norm $\chi \in \mathcal{N}_{\mathbb{Q}}$, such that $\chi^{\text{hom}} = \chi_Z$ is the norm in Example 2.21. Further, $d_{\infty}(\chi|_{R_m}, \chi^{\text{hom}}|_{R_m}) = m/2$ for msufficiently divisible, and hence $d_{\infty}(\chi, \chi^{\text{hom}}) = 1/2$. Finally, recall from [18] that the space CPSH of continuous (bounded) *L*-psh functions on X^{an} can be described as the closure of $\mathcal{H}_{\mathbb{R}}$ (or, equivalently, $\mathcal{H}_{\mathbb{Q}} = \mathcal{H}_{\mathbb{Z}}$) with respect to uniform convergence (see also Section 4.1 below). We show:

THEOREM 2.26. — The Fubini–Study and infimum norm operators induce inverse isomorphic isometries FS: $(\mathcal{N}_{\mathbb{R}}^{\text{cont,hom}}, d_{\infty}) \xrightarrow{\sim} (\text{CPSH}, d_{\infty})$, IN: $(\text{CPSH}, d_{\infty}) \xrightarrow{\sim} (\mathcal{N}_{\mathbb{R}}^{\text{cont,hom}}, d_{\infty})$.

Proof. — By Corollary 2.17, the Fubini–Study operator defines an isometric embedding of complete metric spaces FS: $(\mathcal{N}_{\mathbb{R}}^{hom}, d_{\infty}) \hookrightarrow (\mathcal{L}^{\infty}, d_{\infty})$, which thus maps the closure of any subset onto the closure of its image. Now FS($\mathcal{T}_{\mathbb{R}}^{hom}$) = $\mathcal{H}_{\mathbb{R}}$ (see Proposition 2.15), where the closure of $\mathcal{T}_{\mathbb{R}}^{hom}$ in $(\mathcal{N}_{\mathbb{R}}^{hom}, d_{\infty})$ is $\mathcal{N}_{\mathbb{R}}^{cont,hom}$ (by Proposition 2.24) and the closure of $\mathcal{H}_{\mathbb{R}}$ in $(\mathcal{L}^{\infty}, d_{\infty})$ is CPSH. It follows that FS: $(\mathcal{N}_{\mathbb{R}}^{cont,hom}, d_{\infty}) \xrightarrow{\sim} (CPSH, d_{\infty})$ is an isometric isomorphism, whose inverse is necessarily given by IN, by Theorem 2.16.

2.6. The Fubini–Study envelope

As in [14, Section 7.5] and [18, Section 5.3], we define the Fubini–Study envelope of a bounded function $\varphi \in \mathcal{L}^{\infty}$ as the pointwise supremum

(2.18)
$$Q(\varphi) = Q_L(\varphi) \coloneqq \sup\{\psi \in \mathcal{H}_{\mathbb{R}} \mid \psi \leqslant \varphi\}.$$

Since any $\psi \in \mathcal{H}_{\mathbb{R}}$ is a uniform limit of functions in $\mathcal{H}_{\mathbb{Q}}$, one can replace $\mathcal{H}_{\mathbb{R}}$ with $\mathcal{H}_{\mathbb{O}}$ in this definition. We note that

(2.19)
$$Q_{dL}(d\varphi) = d Q_L(\varphi), \quad Q(t \cdot \varphi) = Q(\varphi), \quad Q(\varphi + c) = Q(\varphi) + c$$

for all $d \in \mathbb{Q}_{>0}$, $t \in \mathbb{R}_{>0}$, $c \in \mathbb{R}$, and refer to Section 4.4 for more information. We view the next result as a "dual" version of Proposition 2.14.

PROPOSITION 2.27. — For any $\varphi \in \mathcal{L}^{\infty}$ we have $IN(\varphi) = IN(Q(\varphi))$.

This is in fact a special case of [14, Lemma 7.23] but we repeat the simple argument for the convenience of the reader.

LEMMA 2.28. — If $\varphi \in \mathcal{L}^{\infty}$ and $s \in R_m$ with m sufficiently divisible, then $\log |s| \leq m\varphi$ iff $\log |s| \leq m Q(\varphi)$.

Proof. — We may assume m = 1. Since $Q(\varphi) \leq \varphi$, we only need to prove the direct implication. For $t \in \mathbb{R}$, set $\psi_t = \max\{\log |s|, -t\}$. Then $\psi_t \in \mathcal{H}_{\mathbb{R}}$, and $\psi_t \leq \varphi$ for $t \gg 0$ since φ is bounded. Thus $\psi_t \leq Q(\varphi)$ by the definition of Q, so $\log |s| \leq \psi_t \leq Q(\varphi)$. Proof of Proposition 2.27. — Pick $s \in R_m$ with m sufficiently divisible. We must prove that $\lambda := \inf_{X^{an}} (m\varphi - \log |s|)$ equals $\lambda' := \inf_{X^{an}} (m Q(\varphi) - \log |s|)$. Since $Q(\varphi) \leq \varphi$ we have $\lambda' \leq \lambda$. The reverse inequality follows from Lemma 2.28 applied to the bounded function $\varphi - m^{-1}\lambda$, together with (2.19).

We similarly have a dual version of Theorem 2.16:

PROPOSITION 2.29. — For any $\varphi \in \mathcal{L}^{\infty}$, we have $FS(IN(\varphi)) = Q(\varphi)$.

Proof. — After passing to a multiple, we may assume that L is a line bundle, so that $\chi := IN(\varphi)$ is a norm on R = R(X, L). For all $m \ge 1$ and $s \in R_m \setminus \{0\}$, we have $\log |s| \le m\varphi - \chi(s)$ on X^{an} , by definition of the infimum norm. By Lemma 2.28 and (2.19), this yields $\log |s| \le m Q(\varphi) - \chi(s)$, and hence

$$FS(\chi) = \sup\{m^{-1}(\log|s| + \chi(s)) \mid m \ge 1, s \in R_m \smallsetminus \{0\}\} \leqslant Q(\varphi).$$

For the reverse inequality, pick any $\psi \in \mathcal{H}_{\mathbb{R}}$ with $\psi \leq \varphi$. Since $FS(\mathcal{T}_{\mathbb{R}}) = \mathcal{H}_{\mathbb{R}}$ (see Proposition 2.15), there exists $m \ge 1$ and a norm χ' on R_m such that

$$m\psi = FS_{mL}(\chi') = \sup_{s \in R_m \setminus \{0\}} \{ \log |s| + \chi'(s) \}.$$

Since $\psi \leq \varphi$, this gives $\log |s| + \chi'(s) \leq m\varphi$, i.e. $\chi' \leq \chi|_{R_m}$ on R_m . As a result,

$$\psi = m^{-1} \operatorname{FS}_{mL}(\chi') \leqslant m^{-1} \operatorname{FS}_{mL}(\chi|_{R_m}) = \operatorname{FS}_m(\chi) \leqslant \operatorname{FS}(\chi),$$

which completes the proof.

Combining Theorem 2.16 and Proposition 2.29 with Propositions 2.14 and 2.27, we also obtain

COROLLARY 2.30. — We have $FS \circ IN \circ FS = FS \text{ on } \mathcal{N}_{\mathbb{R}}, \text{ and } IN \circ FS \circ IN = IN \text{ on } \mathcal{L}^{\infty}.$

3. Spectral analysis

In this section we define a volume function vol: $\mathcal{N}_{\mathbb{R}} \to \mathbb{R}$ as well as pseudo-metrics d_p , $p \in [1, \infty)$, on the space $\mathcal{N}_{\mathbb{R}}$ of norms on section rings of multiples of L. Much of the material is studied for more general non-Archimedean ground fields in [14, 27], but we present the details for the convenience of the reader.

 \square

3.1. The finite-dimensional case

We first describe the space $\mathcal{N}_{\mathbb{R}}(V)$ of non-Archimedean norms on a k-vector space V of dimension $N < \infty$, essentially following [14, 60].

Pick a norm $\chi \in \mathcal{N}_{\mathbb{R}}(V)$, and a χ -orthogonal basis $(e_j)_{1 \leq j \leq N}$ of V. After permutation, we may assume that the sequence $\lambda_j(\chi) \coloneqq \chi(e_j), j = 1, \ldots, N$ satisfies

$$\lambda_1(\chi) \ge \cdots \ge \lambda_N(\chi).$$

It is then independent of the choice of orthogonal basis and is called the spectrum of χ (i.e. the "jumping values" of the associated filtration in the terminology of [13]). In terms of (1.4) we have

$$\lambda_1(\chi) = \lambda_{\max}(\chi), \quad \lambda_N(\chi) = \lambda_{\min}(\chi).$$

The volume of χ is defined as the mean value⁽¹⁾ of its spectrum, i.e.

$$\operatorname{vol}(\chi) \coloneqq \frac{1}{N} \sum_{j} \lambda_j(\chi).$$

For any basis (e_j) of V we have $\operatorname{vol}(\chi) \ge N^{-1} \sum_j \chi(e_j)$ with equality iff (e_j) is χ -orthogonal.

More generally, any two norms χ, χ' admit a common orthogonal basis (e_i) . The relative spectrum of χ with respect to χ' is the sequence

$$\lambda_1(\chi,\chi') \ge \cdots \ge \lambda_N(\chi,\chi')$$

obtained by reordering $(\chi(e_j) - \chi'(e_j))_{1 \leq j \leq N}$, and the spectral measure of χ with respect to χ' is the corresponding probability measure

$$\sigma(\chi,\chi') \coloneqq \frac{1}{N} \sum_{j} \delta_{\lambda_j(\chi,\chi')}.$$

Its barycenter satisfies

(3.1)
$$\int_{\mathbb{R}} \lambda \, \mathrm{d}\sigma(\chi, \chi') = \frac{1}{N} \sum_{j} \lambda_{j}(\chi, \chi') = \mathrm{vol}(\chi) - \mathrm{vol}(\chi').$$

When $\chi' = \chi_{\text{triv}}$ is the trivial norm, we simply write

$$\sigma(\chi) = \sigma(\chi, \chi_{\text{triv}}) = \frac{1}{N} \sum_{j} \delta_{\lambda_i(\chi)},$$

 $^{^{(1)}}$ Note that a different normalization is used in [14, 16], where the volume is defined as the sum of the elements of the spectrum.

and call it the spectral measure of χ . In terms of the associated \mathbb{R} -filtration $F^{\lambda}V = \{\chi \ge \lambda\}$, we have

(3.2)
$$\sigma(\chi) = \frac{1}{N} \sum_{\lambda \in \mathbb{R}} \dim(F^{\lambda} V / F^{>\lambda} V) \,\delta_{\lambda}$$

To any basis $\mathbf{e} = (e_i)$ of V is associated an apartment $\mathbb{A}_{\mathbf{e}} \subset \mathcal{N}_{\mathbb{R}}(V)$, defined as the set of norms diagonalized in this basis. We then have a canonical parametrization

$$\iota_{\mathbf{e}} \colon \mathbb{R}^N \overset{\sim}{\longrightarrow} \mathbb{A}_{\mathbf{e}},$$

and a Gram–Schmidt retraction

$$\rho_{\mathbf{e}} \colon \mathcal{N}_{\mathbb{R}}(V) \longrightarrow \mathbb{A}_{\mathbf{e}}.$$

The map $\iota_{\mathbf{e}}$ sends $(\lambda_j) \in \mathbb{R}^N$ to the unique $\chi \in \mathbb{A}_{\mathbf{e}}$ such that $\chi(e_i) = \lambda_i$, while $\rho_{\mathbf{e}}$ sends a norm χ to the unique norm $\rho_{\mathbf{e}}(\chi) \in \mathbb{A}_{\mathbf{e}}$ such that

$$\rho_{\mathbf{e}}(\chi)(e_i) = \sup_{a \in k^N} \chi(e_i + \sum_{j < i} a_j e_j),$$

i.e. the norm induced, via the basis **e**, from the natural subquotient norm on the graded object of the complete flag defined by **e**. By additivity of the volume in exact sequences, we have

(3.3)
$$\operatorname{vol}(\rho_{\mathbf{e}}(\chi)) = \operatorname{vol}(\chi),$$

see [14, Lemma 2.12]. Each $\mathbb{A}_{\mathbf{e}}$ is trivially preserved by the translation action of \mathbb{R} , the scaling action by $\mathbb{R}_{>0}$, and by the operation $(\chi, \chi') \mapsto \chi \wedge \chi'$. Moreover,

$$\chi \leqslant \chi' \Longrightarrow \rho_{\mathbf{e}}(\chi) \leqslant \rho_{\mathbf{e}}(\chi').$$

3.2. Metrics on the space of norms

Generalizing the classical construction of the Tits metric on the Euclidean building $\mathcal{N}_{\mathbb{R}}(V)$ (see for instance [60, Section 2.2]), it is shown in [14, Theorem 3.1] that each \mathfrak{S}_N -invariant norm τ on \mathbb{R}^N induces a unique metric d_{τ} on $\mathcal{N}_{\mathbb{R}}(V)$ such that $\iota_{\mathbf{e}} : (\mathbb{R}^N, \tau) \hookrightarrow (\mathcal{N}_{\mathbb{R}}(V), d_{\tau})$ is an isometry for any basis **e**. It has the property that $\rho_{\mathbf{e}} : \mathcal{N}_{\mathbb{R}}(V) \to \mathbb{A}_{\mathbf{e}}$ is a contraction. All metrics on $\mathcal{N}_{\mathbb{R}}(V)$ obtained this way are equivalent. They turn $\mathcal{N}_{\mathbb{R}}(V)$ into a metric space that is complete, but not locally compact. In particular, for each $p \in [1, \infty]$ we define a metric d_p on $\mathcal{N}_{\mathbb{R}}(V)$ by setting for any two norms χ, χ' with relative spectrum $(\lambda_i) = (\lambda_i(\chi, \chi'))$

(3.4)
$$d_p(\chi,\chi') \coloneqq \left(N^{-1}\sum_i |\lambda_i|^p\right)^{1/p}$$

for $p \in [1, \infty)$, and

$$d_{\infty}(\chi, \chi') \coloneqq \max_{i} |\lambda_{i}|.$$

Thus $d_p(\chi, \chi')$ is the L^p -norm of the identity with respect to the spectral measure $\sigma(\chi, \chi')$. Note that

(3.5)
$$d_1 \leqslant d_p \leqslant d_\infty^{1-\frac{1}{p}} d_1^{\frac{1}{p}} \leqslant d_\infty$$

on $\mathcal{N}_{\mathbb{R}}(V)$ for $p \in (1, \infty)$. The metric d_2 is the Tits metric mentioned above, while d_{∞} coincides with the Goldman–Iwahori metric (1.2). Our main interest lies in the metric d_1 , which is closely related to the volume:

LEMMA 3.1. — For all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}(V)$ and $p \in [1, \infty)$ we have

(3.6)
$$d_p(\chi,\chi')^p = d_p(\chi,\chi\wedge\chi')^p + d_p(\chi\wedge\chi',\chi')^p.$$

For p = 1, we further have

(3.7)
$$d_1(\chi,\chi') = \operatorname{vol}(\chi) + \operatorname{vol}(\chi') - 2\operatorname{vol}(\chi \wedge \chi').$$

Proof. — The first assertion follows from the fact that the minimum $\chi \wedge \chi'$ of two norms in an apartment $\mathbb{A}_{\mathbf{e}} \simeq \mathbb{R}^N$ is computed componentwise, and the trivial identity

$$\sum_{i} |\lambda_i - \lambda'_i|^p = \sum_{i} |\lambda_i - \min\{\lambda_i, \lambda'_i\}|^p + \sum_{i} |\min\{\lambda_i, \lambda'_i\} - \lambda'_i|^p.$$

for all $\lambda, \lambda' \in \mathbb{R}^N$. On the other hand, it follows from (3.1) that $\chi \ge \chi' \Longrightarrow$ $d_1(\chi, \chi') = vol(\chi) - vol(\chi')$, and (3.7) follows.

The volume function is trivially 1-Lipschitz with respect to d_1 , i.e.

$$(3.8) |vol(\chi) - vol(\chi')| \leq d_1(\chi, \chi')$$

for all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}(V)$. This is also the case for the min operator:

LEMMA 3.2. — Let $\chi_i, \chi'_i, i = 1, 2$, be norms on V. Then

(3.9)
$$d_1(\chi_1 \wedge \chi_2, \chi'_1 \wedge \chi'_2) \leq d_1(\chi_1, \chi'_1) + d_1(\chi_2, \chi'_2).$$

Proof. — First assume $\chi_i \ge \chi'_i$, i = 1, 2. Pick a basis **e** such that $\chi'_1, \chi'_2 \in \mathbb{A}_{\mathbf{e}}$. Lemma 3.1 together with (3.3) show that replacing χ_i by $\rho_{\mathbf{e}}(\chi_i)$, i = 1, 2

does not change the right-hand side of (3.9). As for the left-hand side, $\rho_{\mathbf{e}} \colon \mathcal{N}_{\mathbb{R}}(V) \to \mathbb{A}_{\mathbf{e}}$ being order preserving implies

$$\rho_{\mathbf{e}}(\chi_1) \land \rho_{\mathbf{e}}(\chi_2) \ge \rho_{\mathbf{e}}(\chi_1 \land \chi_2) \ge \chi_1' \land \chi_2',$$

which shows that the left-hand side of (3.9) can only increase upon replacing χ_i by $\rho_{\mathbf{e}}(\chi_i)$, i = 1, 2, using again (3.3) and (3.7).

As a result, we may in fact assume that all four norms belong to $\mathbb{A}_{\mathbf{e}}$. Write $\chi_i(e_j) = \lambda_{i,j}$ and $\chi'_i(e_j) = \lambda'_{i,j}$ for $1 \leq j \leq N$ and i = 1, 2. Then $\lambda_{i,j} \geq \lambda'_{i,j}$ for all i, j, and we must prove that

$$\sum_{j} \lambda_{1,j} \wedge \lambda_{2,j} - \sum_{j} \lambda'_{1,j} \wedge \lambda'_{2,j} \leqslant \sum_{j} (\lambda_{1,j} - \lambda'_{1,j}) + \sum_{j} (\lambda_{2,j} - \lambda'_{2,j});$$

this is straightforward.

Finally consider arbitrary norms. Set $\chi''_i = \chi_i \wedge \chi'_i$ for i = 1, 2. By (3.6) we have

$$d_1(\chi_1 \land \chi_2, \chi'_1 \land \chi'_2) = d_1(\chi_1 \land \chi_2, \chi''_1 \land \chi''_2) + d_1(\chi'_1 \land \chi'_2, \chi''_1 \land \chi''_2)$$

and $\chi_i, \chi'_i \ge \chi''_i$, for i = 1, 2, so (3.9) follows from what precedes, together with (3.6).

3.3. Spectral measures and volume

Now we return to the setting of a projective variety X and an ample \mathbb{Q} line bundle L on X. The following equidistribution result is a special case of a result of Chen–Maclean [27], which deals with general non-Archimedean fields.

THEOREM 3.3. — For any two norms $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, the scaled spectral measures

$$(1/m)_{\star}\sigma(\chi|_{R_m},\chi'|_{R_m})$$

have uniformly bounded support, and they admit a weak limit.

The limit is taken with respect to the partial order by divisibility. If L is an actual line bundle and χ, χ' are norms on R(X, L), then the limit also exists as $m \to \infty$ in the usual total ordering.

DEFINITION 3.4. — For any $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, the spectral measure of χ with respect to χ' is the compactly supported (Borel) probability measure on \mathbb{R} defined as

$$\sigma(\chi,\chi') \coloneqq \lim_m (1/m)_\star \sigma(\chi|_{R_m},\chi'|_{R_m}).$$

The spectral measure of χ is $\sigma(\chi) \coloneqq \sigma(\chi, \chi_{\text{triv}})$, and the volume of χ is the barycenter

$$\operatorname{vol}(\chi) = \int_{\mathbb{R}} \lambda \, \mathrm{d}\sigma(\chi)$$

By (3.1), we have

(3.10)
$$\operatorname{vol}(\chi) = \lim_{m} m^{-1} \operatorname{vol}(\chi|_{R_m}).$$

Example 3.5. — For any $v \in X^{\text{lin}}$ with associated norm $\chi_v \in \mathcal{N}_{\mathbb{R}}^{\text{hom}}$, the spectrum of $\chi_v|_{R_m}$ is the vanishing sequence of R_m with respect to v as defined in [22], i.e. the (finite) set of values of v on nonzero elements of R_m , counted with multiplicity, and

(3.11)
$$S(v) = S_L(v) \coloneqq \operatorname{vol}(\chi_v)$$

coincides with the expected vanishing order of [5] (see also [59]).

The existence of the spectral measure $\sigma(\chi)$ (called the *limit measure* of the corresponding filtration in [17, Section 5.1]) follows from [13, Theorem A]. When $\chi \in \mathcal{T}_{\mathbb{Z}}$, the limit measure coincides with the Duistermaat–Heckman measure of the corresponding test configuration, see [17, Proposition 3.12].

As we shall see, a simple trick borrowed from [27] reduces the proof of Theorem 3.3 to this special case $\chi' = \chi_{\text{triv}}$.

Proof of Theorem 3.3. — The uniform boundedness part is a direct consequence of the linear boundedness condition that we impose on norms in $\mathcal{N}_{\mathbb{R}}(R)$. Set $N_m := \dim R_m$, denote by $(\lambda_{m,j})_{1 \leq j \leq N_m}$ the spectrum of $\chi|_{R_m}$ with respect to $\chi'|_{R_m}$, and set

$$\sigma_m \coloneqq (1/m)_\star \sigma(\chi|_{R_m}, \chi'|_{R_m}) = \frac{1}{N_m} \sum_{j=1}^{N_m} \delta_{m^{-1}\lambda_{m,j}}.$$

As is well-known, in order to prove convergence of σ_m , it suffices to show that

$$\int_{\mathbb{R}} \min\{\lambda, c\} \, \mathrm{d}\sigma_m = \frac{1}{mN_m} \sum_{j=1}^{N_m} \max\{\lambda_{m,j}, mc\}$$

converges for all $c \in \mathbb{R}$, see [27, Proposition 5.1]. But $(\min\{\lambda_{m,j}, cm\})_j$ is the spectrum of $\chi|_{R_m} \wedge (\chi'|_{R_m} + cm)$ with respect to $\chi'|_{R_m}$. Replacing χ with $\chi \wedge (\chi' + c)$ (where \mathbb{R} acts by translation according to (1.5)), we are reduced to proving that the barycenter

$$(mN_m)^{-1}\sum_j \lambda_{m,j}$$

ANNALES DE L'INSTITUT FOURIER

of the measure σ_m converges. Now, this barycenter is the difference of the barycenters of $(1/m)_{\star}\sigma(\chi|_{R_m})$ and $(1/m)_{\star}\sigma(\chi'|_{R_m})$, each of which admits a limit by [13, Theorem A], and we are done.

The proof of Theorem 3.3 shows that

(3.12)
$$\int \min\{\lambda, c\} \, \sigma(\chi, \chi')(\mathrm{d}\lambda) = \operatorname{vol}\left(\chi \wedge (\chi' + c)\right) - \operatorname{vol}(\chi')$$

for all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$ and $c \in \mathbb{R}$. Some further properties of the spectral measure $\sigma(\chi)$ are described by the following result.

THEOREM 3.6. — Pick $\chi \in \mathcal{N}_{\mathbb{R}}$, with associated filtration $F^{\lambda}R_m = \{s \in$ $R_m \mid \chi(s) \ge \lambda$. Then:

- (i) for each $\lambda \in \mathbb{R}$, dim $F^{m\lambda}R_m/\dim R_m$ admits a limit $\operatorname{vol}(\chi \ge \lambda) \in$ [0,1] as $m \to \infty$;
- (ii) the function $\lambda \mapsto \operatorname{vol}(\chi \geq \lambda)^{1/n}$ is positive and concave on $(-\infty, \lambda_{\max}(\chi))$, and vanishes on $(\lambda_{\max}(\chi), +\infty)$;
- (iii) $\sigma(\chi) = -\frac{d}{d\lambda} \operatorname{vol}(\chi \ge \lambda)$ in the sense of distributions;
- (iv) supp $\sigma(\chi) = [\lambda_{\min}(\chi), \lambda_{\max}(\chi)]$ with

(3.13)
$$\lambda_{\min}(\chi) \coloneqq \inf \left\{ \lambda \in \mathbb{R} \mid \operatorname{vol}(\chi \ge \lambda) < 1 \right\};$$

(v) for any $a \leq \lambda_{\min}(\chi)$, i.e. such that $\operatorname{vol}(\chi \geq a) = 1$, we have

$$\operatorname{vol}(\chi) = a + \int_{a}^{+\infty} \operatorname{vol}(\chi \ge \lambda) \, \mathrm{d}\lambda = a + \int_{a}^{\lambda_{\max}(\chi)} \operatorname{vol}(\chi \ge \lambda) \, \mathrm{d}\lambda.$$

Proof. — Properties (i)–(iv) are direct consequence of [13] (see [17, Section 3.1]. To see (v), set $b \coloneqq \lambda_{\max}(\chi), f(\lambda) \coloneqq \operatorname{vol}(\chi \ge \lambda)$, and pick a cut-off function $\theta \in C_c^{\infty}(\mathbb{R})$ such that $\theta \equiv 1$ on [a, b]. Since $\theta(\lambda)\lambda$ is smooth and compactly supported, (iii) and (iv) yield

$$\operatorname{vol}(\chi) = \int_{\mathbb{R}} \lambda \, \mathrm{d}\sigma(\chi) = -\int_{\mathbb{R}} \theta(\lambda)\lambda f'(\lambda) \, \mathrm{d}\lambda$$
$$= \int_{\mathbb{R}} (\theta(\lambda)\lambda)' f(\lambda) \, \mathrm{d}\lambda = \int_{\mathbb{R}} (\theta'(\lambda)\lambda + \theta(\lambda))f(\lambda) \, \mathrm{d}\lambda.$$

Since $f(\lambda) = 1$ for $\lambda \leq a$, $f(\lambda) = 0$ for $\lambda \geq b$ and $\theta(\lambda) = 1$ for $\lambda \geq a$, this is equal to

$$\int_{-\infty}^{a} (\theta'(\lambda)\lambda + \theta(\lambda)) \, \mathrm{d}\lambda + \int_{a}^{+\infty} f(\lambda) \, \mathrm{d}\lambda = a + \int_{a}^{+\infty} f(\lambda) \, \mathrm{d}\lambda = a + \int_{a}^{b} f(\lambda) \, \mathrm{d}\lambda,$$

by integration by parts. This proves (v).

by integration by parts. This proves (v).

The next result shows how spectral measures behave under operations on norms. It follows from elementary computations of spectra in joint orthogonal bases; the details are left to the reader.

PROPOSITION 3.7. — Let $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, and pick $c \in \mathbb{R}$, $t \in \mathbb{R}_{>0}$. Then:

- (i) $\sigma(\chi', \chi)$ is the pushforward of $\sigma(\chi, \chi')$ under $\lambda \mapsto -\lambda$;
- (ii) $\sigma(\chi + c, \chi')$ is the pushforward of $\sigma(\chi, \chi')$ under $\lambda \mapsto \lambda + c$;
- (iii) $\sigma(\chi, \chi \land \chi')$ is the pushforward of $\sigma(\chi, \chi')$ under $\lambda \mapsto \max\{\lambda, 0\}$;
- (iv) $\sigma(t\chi, t\chi')$ is the pushforward of $\sigma(\chi, \chi')$ under $\lambda \mapsto t\lambda$.

Remark 3.8. — In [8, Theorem 3.3] (which appeared after the first version of this article was posted), the authors construct a natural *joint spectral measure* $\rho(\chi, \chi')$ on \mathbb{R}^2 associated to any pair $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, that encodes the asymptotic behavior of the spectra of χ and χ' in jointly orthogonal bases. The spectral measure $\sigma(\chi, \chi')$ is the pushforward of $\rho(\chi, \chi')$ under the map $\mathbb{R}^2 \to \mathbb{R}$ given by $(\lambda, \lambda') \mapsto \lambda - \lambda'$.

3.4. The d_p -pseudometrics and asymptotic equivalence

Pick $p \in [1, \infty)$ and $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$. By definition, we have

$$\mathrm{d}_p(\chi|_{R_m},\chi'|_{R_m})^p = \int_{\mathbb{R}} |\lambda|^p \,\mathrm{d}\sigma(\chi|_{R_m},\chi'|_{R_m}).$$

Theorem 3.3 thus shows that the limit

$$\mathrm{d}_p(\chi,\chi') \coloneqq \lim_m m^{-1} \,\mathrm{d}_p(\chi|_{R_m},\chi'|_{R_m})$$

exists in $[0, +\infty)$, and coincides with the L^p -norm of the identity with respect to the spectral measure $\sigma(\chi, \chi')$, i.e.

(3.14)
$$d_p(\chi,\chi')^p = \int_{\mathbb{R}} |\lambda|^p \sigma(\chi,\chi').$$

It is clear that $(d_p)_{1 \leq p < \infty}$ is a non-decreasing family of pseudo-metrics on $\mathcal{N}_{\mathbb{R}}$. For p = 1, (3.7) further yields

(3.15)
$$d_1(\chi,\chi') = \operatorname{vol}(\chi) + \operatorname{vol}(\chi') - 2\operatorname{vol}(\chi \wedge \chi').$$

For any $p \in [1, \infty]$, we have $d_1 \leq d_p \leq d_\infty$. One also easily checks (using for instance (3.14) and Proposition 3.7)

(3.16)
$$d_p(t\chi, t\chi') = t d_p(\chi, \chi'),$$
$$d_p(\chi + c, \chi' + c) = d_p(\chi, \chi'),$$
$$d_p(\chi, \chi + c) = |c|$$

for all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}, t \in \mathbb{R}_{>0}, c \in \mathbb{R}$.

The pseudo-metric d_1 defines a (non-Hausdorff) topology on $\mathcal{N}_{\mathbb{R}}$, which is strictly coarser than the d_p -topology for any p > 1, when dim X > 0. However, (3.5) remains valid on $\mathcal{N}_{\mathbb{R}}$, and shows that the d_p -topologies with $p < \infty$ all agree on d_{∞} -bounded subsets of $\mathcal{N}_{\mathbb{R}}$. In particular, they share the same pairs of non-separated points, which gives rise to:

DEFINITION 3.9. — We say that two norms $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$ are asymptotically equivalent, and write $\chi \sim \chi'$, if the following equivalent conditions hold:

- (i) $d_1(\chi, \chi') = 0;$
- (ii) $d_p(\chi, \chi') = 0$ for all $p \in [1, \infty)$;
- (iii) $\sigma(\chi, \chi') = \delta_0$.

The equivalence between (i)–(iii) follows from (3.14). Since $d_1 \leq d_{\infty}$, we trivially have

$$d_{\infty}(\chi,\chi') = 0 \Longrightarrow \chi \sim \chi'.$$

The converse fails in general, and thus so does the analogue of (3.14) for the pseudo-metric d_{∞} (see, however, Corollary 6.26 below for the case of continuous norms).

Example 3.10. — Pick any subvariety $Z \subsetneq X$, and consider the norm $\chi = \chi_Z \in \mathcal{N}_{\mathbb{Z}}$ as in Example 2.21. Using (3.2), it is easy to see that $\sigma(\chi|_{R_m}) = \varepsilon_m \delta_0 + (1 - \varepsilon_m) \delta_m$ with

$$\varepsilon_m \coloneqq \dim \mathrm{H}^0(Z, mL) / \dim \mathrm{H}^0(X, mL) = O(1/m).$$

Thus $\sigma(\chi) = \lim_m (1/m)_\star \sigma(\chi|_{R_m}) = \delta_1$, and hence $\chi \sim \chi_{\text{triv}} + 1$ (see Proposition 3.7(ii)). On the other hand, since $\chi \neq \chi_{\text{triv}} + 1$ are both homogeneous, we have $d_{\infty}(\chi, \chi_{\text{triv}} + 1) > 0$ (see Proposition 2.8).

By (3.15), we have:

LEMMA 3.11. — If $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$ satisfy $\chi \ge \chi'$, then $\chi \sim \chi' \iff \operatorname{vol}(\chi) = \operatorname{vol}(\chi')$.

As we next show, spectral measures are continuous with respect to the d_1 -topology.

THEOREM 3.12. — Consider nets (χ_i) , (χ'_i) in $\mathcal{N}_{\mathbb{R}}$, converging respectively to $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$ in the d₁-topology. Then $\sigma(\chi_i, \chi'_i) \to \sigma(\chi, \chi')$ weakly.

Proof. — Set $\sigma_i \coloneqq \sigma(\chi_i, \chi'_i)$ and $\sigma \coloneqq \sigma(\chi, \chi')$. As in the proof of Theorem 3.3, it suffices to prove that

$$\int \min\{\lambda, c\} \,\sigma_i(\mathrm{d}\lambda) = \operatorname{vol}(\chi_i \wedge (\chi_i' + c)) - \operatorname{vol}(\chi_i')$$

converges to

$$\int \min\{\lambda, c\} \, \sigma(\mathrm{d}\lambda) = \operatorname{vol}(\chi \wedge (\chi' + c)) - \operatorname{vol}(\chi')$$

for all $c \in \mathbb{R}$, see (3.12). This follows immediately from the Lipschitz property of the volume and min operators, see (3.8) and Lemma 3.2.

COROLLARY 3.13. — For any $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, the quantities

$$\sigma(\chi, \chi'), \quad \lambda_{\max}(\chi) \quad and \quad \operatorname{vol}(\chi)$$

only depend on the asymptotic equivalence classes of χ, χ' . Further,

$$\chi_i \sim \chi'_i, i = 1, 2 \Longrightarrow \chi_1 \wedge \chi_2 \sim \chi'_1 \wedge \chi'_2.$$

Proof. — The first claim follows directly from Theorem 3.12. It implies the second one, as $\lambda_{\max}(\chi)$ can be reconstructed from $\sigma(\chi) = \sigma(\chi, \chi_{\text{triv}})$, by Theorem 3.6. Finally, the Lipschitz properties of the volume and the min operator (see (3.8) and Lemma 3.2) carry over to $\mathcal{N}_{\mathbb{R}}$, which takes care of the last two claims.

Following [43, 69], we finally show:

LEMMA 3.14. — Suppose $\chi \in \mathcal{N}_{\mathbb{R}}$ satisfies $\chi \ge \chi_{\text{triv}}$. For any $p \in [1, \infty)$, we then have:

(i)
$$d_p(\chi, \chi_{triv})^p = p \int_0^{+\infty} \lambda^{p-1} \operatorname{vol}(\chi \ge \lambda) d\lambda$$

 $= p \int_0^{\lambda_{\max}(\chi)} \lambda^{p-1} \operatorname{vol}(\chi \ge \lambda) d\lambda;$
(ii) $\lambda_{\max}(\chi) = d_{\infty}(\chi, \chi_{triv}) \leqslant C_{n,p} d_p(\chi, \chi_{triv}) \text{ with } C_{n,p} \coloneqq {\binom{n+p}{n}}^{1/p}.$

Note that $C_{n,p} \to 1$ as $p \to \infty$, and hence $d_p(\chi, \chi_{triv}) \to d_{\infty}(\chi, \chi_{triv})$.

Proof. — As in the proof of Theorem 3.6, set $b \coloneqq \lambda_{\max}(\chi)$, $f(\lambda) \coloneqq$ vol $(\chi \ge \lambda)$, and pick a smooth function $g \in C_c^{\infty}(\mathbb{R})$ such that $g(\lambda) = \lambda^p$ for $\lambda \in [0, b]$. Then $f(\lambda) = 1$ for $\lambda \le 0$, $f(\lambda) = 0$ for $\lambda > b$, and hence

$$d_p(\chi, \chi_{\text{triv}})^p = \int_0^b \lambda^p \, \mathrm{d}\sigma(\chi) = -\int_{\mathbb{R}} g(\lambda) f'(\lambda) \, \mathrm{d}\lambda = \int_{\mathbb{R}} g'(\lambda) f(\lambda) \, \mathrm{d}\lambda$$
$$= \int_{-\infty}^0 g'(\lambda) \mathrm{d}\lambda + \int_0^{+\infty} p \lambda^{p-1} f(\lambda) \, \mathrm{d}\lambda,$$

where $\int_{-\infty}^{0} g'(\lambda) d\lambda = g(0) = 0$. This proves (i).

The first equality in (ii) is (1.11). To prove the inequality, we argue as in [69, Section 5]. By Theorem 3.6, $f(\lambda)^{1/n}$ is concave on $(-\infty, b]$, and

hence $f(\lambda)^{1/n} \ge 1 - \lambda/b$ for $\lambda \in [0, b]$. Using (i), this yields

$$d_p(\chi, \chi_{\text{triv}})^p = p \int_0^{\lambda_{\max}} \lambda^{p-1} f(\lambda) \, d\lambda$$

$$\geq p \int_0^b \lambda^{p-1} \left(1 - \frac{\lambda}{b}\right)^n \, d\lambda$$

$$= b^p p \int_0^1 t^{p-1} (1-t)^n \, dt = \frac{p! n!}{(n+p)!} b^p,$$

and the result follows.

3.5. The space of norms modulo translation

For any $p \in [1, \infty]$, the additive action of \mathbb{R} on $\mathcal{N}_{\mathbb{R}}$ preserves the pseudometric d_p , which thus induces a quotient pseudo-metric \underline{d}_p on the space of norms modulo translation $\mathcal{N}_{\mathbb{R}}/\mathbb{R}$, such that

$$\underline{\mathbf{d}}_p(\chi, \chi') = \inf_{c \in \mathbb{R}} \mathbf{d}_p(\chi, \chi' + c)$$

for $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$. This supremum is actually achieved:

LEMMA 3.15. — For any $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, there exists $c \in \mathbb{R}$ such that $\underline{d}_p(\chi, \chi') = d_p(\chi, \chi' + c)$ and $|c| \leq 2 d_p(\chi, \chi')$.

In particular, $\underline{d}_p(\chi, \chi') = 0$ iff χ, χ' are asymptotically equivalent modulo translation, in the sense that $\chi \sim \chi' + c$ for some $c \in \mathbb{R}$ (which is then uniquely determined by $c = \operatorname{vol}(\chi) - \operatorname{vol}(\chi')$). When $\chi' = \chi_{\operatorname{triv}}$ we say that χ is asymptotically constant.

Proof. — By (3.16) , for all $c \in \mathbb{R}$ we have

$$|c| = d_p(\chi', \chi' + c) \leq d_p(\chi', \chi) + d_p(\chi, \chi' + c).$$

Thus $d_p(\chi, \chi' + c) \leq d_p(\chi, \chi') \Longrightarrow |c| \leq 2 d_p(\chi, \chi')$, and hence

$$\underline{\mathbf{d}}_p(\chi,\chi') = \inf \left\{ \mathbf{d}_p(\chi,\chi'+c) \mid c \in \mathbb{R}, \, |c| \leq 2 \, \mathbf{d}_p(\chi,\chi') \right\},\,$$

which is achieved by compactness and Lipschitz continuity of

$$c \mapsto \mathrm{d}_p(\chi, \chi' + c).$$

DEFINITION 3.16. — For each $p \in [1,\infty)$, we define the L^p -norm of $\chi \in \mathcal{N}_{\mathbb{R}}$ as

$$\|\chi\|_p \coloneqq \mathrm{d}_p(\chi, \chi_{\mathrm{triv}} + \mathrm{vol}(\chi)).$$

TOME 75 (2025), FASCICULE 2

This definition extends the notion in [39] of the L^p -norm of a test configuration, see [17, Remark 6.10]. Indeed, (3.14) yields

$$\|\chi\|_p^p = \int |\lambda - \bar{\lambda}|^p \,\mathrm{d}\sigma(\chi),$$

the *p*-th central moment of the spectral measure $\sigma(\chi)$, where $\overline{\lambda} = \int \lambda \, d\sigma(\chi) =$ vol (χ) is its barycenter. Note also that

$$\|\chi + c\|_p = \|\chi\|_p, \quad \|t\chi\|_p = t\|\chi\|_p$$

for $c \in \mathbb{R}$, $t \in \mathbb{R}_{>0}$.

PROPOSITION 3.17. — Given a norm $\chi \in \mathcal{N}_{\mathbb{R}}$, the following are equivalent:

- (i) χ is asymptotically constant;
- (ii) $\|\chi\|_p = 0$ for some $p \in [1, +\infty)$;
- (iii) $\|\chi\|_p = 0$ for all $p \in [1, +\infty)$.

Proof. — If $\chi \sim \chi_{\text{triv}} + c$ with $c \in \mathbb{R}$, then $c = \text{vol}(\chi)$, by (3.8). The rest is straightforward.

3.6. Convergence of the canonical approximants

The next result strengthens the approximation result proved in [13, Theorem 1.14] (see also [12, Théorème 3.15] and [30]). A version valid for arbitrary non-Archimedean fields is given in [63, Theorem 4.5.4] (which appeared after the first version of the current paper).

Recall that the canonical approximants $\chi_d \in \mathcal{T}_{\mathbb{R}}$ of a norm $\chi \in \mathcal{N}_{\mathbb{R}}$, which are defined for *d* sufficiently divisible, satisfy $\chi_d \leq \chi$ and form an increasing net with respect to divisibility.

THEOREM 3.18. — For any $\chi \in \mathcal{N}_{\mathbb{R}}$ and $p \in [1, \infty)$, we have $d_p(\chi_d, \chi) \to 0$.

Recall that the result holds for $p = \infty$ iff χ is continuous (see Theorem 2.19).

COROLLARY 3.19. — For any $p \in [1, \infty)$, the set $\mathcal{T}_{\mathbb{Z}}$ of test configurations is dense in $\mathcal{N}_{\mathbb{R}}$ in the d_p -topology.

Proof. — By Theorem 2.19, $\mathcal{T}_{\mathbb{Z}}$ is dense in $\mathcal{T}_{\mathbb{R}}$ for d_{∞} -topology, and hence also for the d_p -topology, since $d_p \leq d_{\infty}$. It therefore suffices to show that $\mathcal{T}_{\mathbb{R}}$ is d_p -dense in $\mathcal{N}_{\mathbb{R}}$, which follows from Theorem 3.18 since the canonical approximants of any norm lie in $\mathcal{T}_{\mathbb{R}}$. By d_1 -continuity of spectral measures (see Theorem 3.12) we also get:

COROLLARY 3.20. — For all
$$\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$$
 we have $\lim_{d} \sigma(\chi_{d}, \chi'_{d}) = \sigma(\chi, \chi')$.

Proof of Theorem 3.18. — Since $\chi_d \leq \chi$ is an increasing net, $d_{\infty}(\chi_d, \chi)$ is decreasing, and hence uniformly bounded. In view of (3.5), it is thus enough to show the result for p = 1, i.e.

$$d_1(\chi_d, \chi) = \operatorname{vol}(\chi) - \operatorname{vol}(\chi_d)$$

tends to 0. Since (χ_d) is d_{∞} -bounded, we can find a < b such that $\sigma(\chi)$ and each $\sigma(\chi_d)$ is has support in [a, b], and hence

$$\operatorname{vol}(\chi) - \operatorname{vol}(\chi_d) = \int_a^b \operatorname{vol}(\chi \ge \lambda) \, \mathrm{d}\lambda - \int_a^b \operatorname{vol}(\chi_d \ge \lambda) \, \mathrm{d}\lambda,$$

see Theorem 3.6. By dominated convergence, it will thus suffice to show $\lim_d \operatorname{vol}(\chi_d \ge \lambda) = \operatorname{vol}(\chi \ge \lambda)$ for each $\lambda \in \mathbb{R}$ fixed.

To see this, we may assume, after replacing L by a multiple, that R = R(X, L) is generated in degree 1 and that χ is defined on R. By definition (see Theorem 3.6(i)), we have

$$\operatorname{vol}(\chi \ge \lambda) = \frac{\operatorname{vol}(S)}{\operatorname{vol}(R)} = V^{-1} \operatorname{vol}(S),$$

where $S \subset R$ is the graded subalgebra with graded pieces $S_m := F^{m\lambda}R_m$ and $\operatorname{vol}(S) = \lim_{m \to \infty} \frac{n!}{m^n} \dim S_m$. Similarly,

$$\operatorname{vol}(\chi_d \ge \lambda) = (d^n V)^{-1} \operatorname{vol}(T(d))$$

where $T(d) \subset S^{(d)}$ is generated in degree 1 by $T(d)_1 = S_1^{(d)} = S_d$. The desired convergence now follows from Lemma 3.21 below.

LEMMA 3.21. — Let $S \subset R$ be a graded subalgebra, and suppose we are given, for each d divisible enough, a graded subalgebra $T(d) \subset S^{(d)}$ such that $T(d)_1 = S_1^{(d)} = S_d$. Then $\lim_d d^{-n} \operatorname{vol}(T(d)) = \operatorname{vol}(S)$.

Proof. — We use Okounkov bodies, following [12]. Set K := k(X), and pick a valuation $\nu \colon K^{\times} \to \mathbb{Z}^n$ of maximal rational rank, equal to n (e.g. associated to a flag of subvarieties as in [52]). Set $\Gamma_m := \nu(S_m \setminus \{0\})$ and $\Gamma(d)_m := \nu(T(d)_m \setminus \{0\})$ for $m \ge 1$. Let $\Delta(S)$ and $\Delta(T(d))$ be the closed convex hull inside \mathbb{R}^n of $\bigcup_m m^{-1}\Gamma_m$ and of $\bigcup_m m^{-1}\Gamma(d)_m$, respectively. Then $\operatorname{vol}(S) = n! \operatorname{vol}(\Delta(S))$ and $\operatorname{vol}(T(d)) = n! \operatorname{vol}(\Delta(T(d)))$, so it suffices to prove that $\lim_d \operatorname{vol}(d^{-1}\Delta(T(d))) = \operatorname{vol}(\Delta(S))$.

Since $T(d)_m \subset S_{dm}$, we get $\Gamma(d)_m \subset \Gamma_{dm}$ for all d, m, and hence $d^{-1}\Delta(T(d)) \subset \Delta(S)$ for all d. If $\operatorname{vol}(\Delta(S)) = 0$, we are done, so we may assume $\Delta(S)$ has nonempty interior. Pick compact subsets A and B of \mathbb{R}^n

with $A \in B \in \Delta(S)$. It suffices to prove that $d^{-1}\Delta(T(d)) \supset A$ for d sufficiently divisible. Now $d^{-1}\mathbb{Z}^n \cap B = d^{-1}\Gamma_d \cap B$, see [12, Lemme 1.13]. If Δ_d is the convex hull of $d^{-1}\Gamma_d$, it follows that $\Delta_d \supset A$. But $T(d)_1 = S_d$, so $\Gamma(d)_1 = \Gamma_d$, and hence $d^{-1}\Delta(T(d)) \supset \Delta_d \supset A$, which completes the proof.

4. Non-Archimedean pluripotential theory

In this section we summarize results from [18] that are relevant to our later purposes.

4.1. L-psh functions

An *L*-psh function $\varphi \colon X^{\mathrm{an}} \to [-\infty, +\infty)$ is defined as the pointwise limit of any decreasing net in $\mathcal{H}_{\mathbb{Q}}$ (or $\mathcal{H}_{\mathbb{R}}$), excluding $\varphi \equiv -\infty$. We denote by $\mathrm{PSH} = \mathrm{PSH}(L)$ the set of all *L*-psh functions. If $\varphi \in \mathrm{PSH}$, then $\varphi + c \in \mathrm{PSH}$ for all $c \in \mathbb{R}$. If $\varphi, \psi \in \mathrm{PSH}$, then $\max\{\varphi, \psi\} \in \mathrm{PSH}$. If $(\varphi_j)_j$ is a decreasing net in PSH, and φ is the pointwise limit of (φ_j) , then $\varphi \in \mathrm{PSH}$, or $\varphi \equiv$ $-\infty$. We can thus describe PSH as the smallest class of functions which is invariant under max, translation by a constant, decreasing limits, and contains all functions of the form $m^{-1} \log |s|$ with m sufficiently divisible and $s \in R_m \setminus \{0\}$.

By Dini's Lemma, the set

$$CPSH := PSH \cap C^0$$

of (bounded) continuous *L*-psh functions is the closure of $\mathcal{H}_{\mathbb{Q}}$ (or $\mathcal{H}_{\mathbb{R}}$) in C^{0} (in line with the definition of a semipositive (continuous) metric in [28, 47, 70]).

The set PSH is stable under convex combinations, and under the action $(t, \varphi) \mapsto t \cdot \varphi$ of $\mathbb{R}_{>0}$ on functions, see (1.15). If $v, v' \in X^{\mathrm{an}}$ and $v \leq v'$, then $\varphi(v) \geq \varphi(v')$ for all $\varphi \in \mathrm{PSH}$. In particular,

$$\sup \varphi \coloneqq \sup_{X^{\mathrm{an}}} \varphi = \varphi(v_{\mathrm{triv}})$$

for all $\varphi \in \text{PSH}$.

A subset $\Sigma \subset X^{an}$ is pluripolar if $\Sigma \subset \{\varphi = -\infty\}$ for some L-psh function φ . This condition is independent of the choice of ample Q-line bundle L, and Σ is nonpluripolar iff

(4.1)
$$T(\Sigma) \coloneqq \sup_{\varphi \in PSH} \left(\sup \varphi - \sup_{\Sigma} \varphi \right) \in [0, +\infty]$$

ANNALES DE L'INSTITUT FOURIER

is finite. If $\Sigma = \{v\}$ with $v \in X^{\operatorname{an}}$, then $\operatorname{T}(\{v\}) = \operatorname{T}(v)$ as defined in (1.18), and the set $X^{\operatorname{lin}} \subset X^{\operatorname{an}}$ of valuations of linear growth thus coincides with the set of non-pluripolar points $v \in X^{\operatorname{an}}$, i.e. such that every $\varphi \in \operatorname{PSH}$ is finite-valued on v.

Since every divisorial valuation has linear growth, *L*-psh functions are finite-valued on X^{div} . The restriction map PSH $\rightarrow \mathbb{R}^{X^{\text{div}}}$ is further injective [18, Corollary 4.23], and we endow PSH with the induced topology of pointwise convergence on X^{div} . This is in fact equivalent to pointwise convergence on X^{lin} [18, Theorem 11.4].

Note that since $\mathcal{H}_{\mathbb{R}}(dL) = d\mathcal{H}_{\mathbb{R}}(L)$ we have $\mathrm{PSH}(dL) = d\mathrm{PSH}(L)$ for any $d \in \mathbb{Q}_{>0}$. To study $\mathrm{PSH}(L)$ we may therefore in practice assume that L is an ample line bundle and that R(X, L) is generated in degree 1.

We refer to [18, Example 4.13] for a concrete description of L-psh functions on curves. See also Appendix B below for the toric case.

4.2. Monge–Ampère operator and energy on $\mathcal{H}_{\mathbb{O}}$

The mixed Monge–Ampère operator on $\mathcal{H}_{\mathbb{Z}} = \mathcal{H}_{\mathbb{Q}}$ associates to any tuple $(\varphi_1, \ldots, \varphi_n) \in \mathcal{H}_{\mathbb{Q}}^n$ a Radon probability measure $MA(\varphi_1, \ldots, \varphi_n)$, defined as follows. Pick integrally closed, semiample test configurations $(\mathcal{X}, \mathcal{L}_i)$ for (X, L) (with the same \mathcal{X}) such that $\varphi_i = \varphi_{\mathcal{L}_i}$, see Appendix A. Denoting by $\mathcal{X}_0 = \sum_i b_i E_i$ the irreducible decomposition of the central fiber, we then have

$$\mathrm{MA}(\varphi_1,\ldots,\varphi_n)=\sum_i b_i(\mathcal{L}_1|_{E_i}\cdot\ldots\cdot\mathcal{L}_n|_{E_i})\delta_{v_i},$$

where $v_i \in X^{\text{div}}$ is the divisorial valuation defined by E_i .

Following the strategy by Chambert-Loir in [25], the mixed Monge– Ampère operator admits a unique continuous extension to the space CPSH of *continuous L*-psh functions (with respect to uniform convergence), and this extension is in turn a special case of the with the general theory developed in [26].

LEMMA 4.1. — For any $\varphi_1, \ldots, \varphi_n \in \mathcal{H}_{\mathbb{R}}$, the support of $MA(\varphi_1, \ldots, \varphi_n)$ is a finite subset of X^{lin} .

Proof. — Set $\mu := \operatorname{MA}(\varphi_1, \ldots, \varphi_n)$ and $\Sigma := \operatorname{supp} \mu$. The finiteness of Σ is proved in [14, Example 8.11], as a consequence of [26, Proposition 6.9.2] and the invariance under ground field extension of the Chambert-Loir-Ducros construction. By [18, Proposition 7.21], we further have $\int \varphi \mu > -\infty$ for any $\varphi \in \operatorname{PSH}$. Since φ is bounded above, this implies that φ is finite at each $v \in \Sigma$; thus v is nonpluripolar, and hence $v \in X^{\operatorname{lin}}$.

We will use notation such as $MA(\varphi^{\langle j \rangle}, \psi^{\langle n-j \rangle})$, with j copies of φ and n-j copies of ψ , and write $MA(\varphi) = MA(\varphi^{\langle n \rangle})$. Thus $MA(0) = \delta_{v_{triv}}$.

The Monge–Ampère energy E: CPSH $\rightarrow \mathbb{R}$ is the primitive of the Monge–Ampère operator in the sense that

(4.2)
$$\frac{\mathrm{d}}{\mathrm{d}t}\Big|_{t=0} \mathrm{E}((1-t)\varphi + t\psi) = \int (\psi - \varphi) \,\mathrm{MA}(\varphi)$$

for $\varphi, \psi \in \text{CPSH}$, normalized by E(0) = 0. As such, E is monotone increasing, i.e. $\varphi \ge \psi \implies E(\varphi) \ge E(\psi)$. Integration along line segments yields

(4.3)
$$E(\varphi) - E(\psi) = \frac{1}{n+1} \sum_{j=0}^{n} \int_{X^{an}} (\varphi - \psi) \operatorname{MA}(\varphi^{(j)}, \psi^{(n-j)}),$$

for $\varphi, \psi \in CPSH$, and hence

$$\mathbf{E}(\varphi) = \frac{1}{n+1} \sum_{j=0}^{n} \int_{X^{\mathrm{an}}} \varphi \, \mathrm{MA}\left(\varphi^{\langle j \rangle}, 0^{\langle n-j \rangle}\right).$$

If $\varphi \in \mathcal{H}_{\mathbb{Z}} = \mathcal{H}_{\mathbb{Q}}$ is represented by a test configuration $(\mathcal{X}, \mathcal{L})$, then

(4.4)
$$\mathbf{E}(\varphi) = \frac{(\bar{\mathcal{L}}^{n+1})}{(n+1)(L^n)},$$

where $(\overline{\mathcal{X}}, \overline{\mathcal{L}}) \to \mathbb{P}^1$ is the canonical compactification of $(\mathcal{X}, \mathcal{L}) \to \mathbb{A}^1$.

The functional E is concave on CPSH, which amounts to

(4.5)
$$E(\varphi) - E(\psi) \leq \int (\varphi - \psi) MA(\psi)$$

for all $\varphi, \psi \in CPSH$, by (4.2). Combined with (4.3), this implies

(4.6)
$$\varphi \geqslant \psi \Longrightarrow \mathcal{E}(\varphi) - \mathcal{E}(\psi) \approx \int (\varphi - \psi) \operatorname{MA}(\psi).$$

In addition to E, we introduce the translation invariant functional

$$\mathbf{I}(\varphi,\psi) \coloneqq \int (\varphi - \psi) \left(\mathbf{MA}(\psi) - \mathbf{MA}(\varphi) \right) \ge 0,$$

which satisfies the quasi-triangle inequality

(4.7)
$$I(\varphi_1, \varphi_2) \lesssim I(\varphi_1, \varphi_3) + I(\varphi_3, \varphi_2).$$

We also set

$$I(\varphi) \coloneqq I(\varphi, 0) \coloneqq \sup \varphi - \int \varphi MA(\varphi).$$

The Monge–Ampère operator is homogeneous with respect to the action of $\mathbb{R}_{>0}$ on continuous *L*-psh functions φ , in the sense that MA $(t \cdot \varphi) = t_{\star}$ MA (φ) for all t > 0. Similarly, we have $\mathbf{E}(t \cdot \varphi) = t \mathbf{E}(\varphi)$, $\mathbf{I}(t \cdot \varphi) = t \mathbf{I}(\varphi)$.

ANNALES DE L'INSTITUT FOURIER

4.3. Functions and measures of finite energy

The Monge–Ampère energy admits a unique non-decreasing, usc extension E: PSH $\rightarrow \mathbb{R} \cup \{-\infty\}$, given for $\varphi \in PSH$ by

(4.8)
$$E(\varphi) := \inf \{ E(\psi) \mid \varphi \leqslant \psi \in CPSH \}.$$

We denote by

$$\mathcal{E}^1 \coloneqq \{\varphi \in \text{PSH} \mid E(\varphi) > -\infty\}$$

the set of *L*-psh functions of finite energy. In other words, functions in \mathcal{E}^1 are decreasing limits of nets $\varphi_i \in \mathcal{H}_{\mathbb{Q}}$ with energy $\mathcal{E}(\varphi_i)$ uniformly bounded below. In particular, $\text{CPSH} \subset \mathcal{E}^1$.

The weak topology of \mathcal{E}^1 is its subspace topology from PSH, and the strong topology on \mathcal{E}^1 is the coarsest refinement of the weak topology for which E becomes continuous.

For a decreasing or increasing net (φ_j) in \mathcal{E}^1 , strong and weak convergence coincide, i.e. $\varphi_j \to \varphi$ strongly in \mathcal{E}^1 iff $\varphi_j \to \varphi$ pointwise on X^{div} , see Example 12.2 and Theorem 12.5 in [18], respectively.

Denote by \mathcal{M} the space of Radon probability measures on X^{an} , endowed with the weak topology. The main point in introducing the strong topology is that the mixed Monge–Ampère operator MA, a priori only defined as a map (CPSH)^{*n*} $\rightarrow \mathcal{M}$, admits a (unique) extension $(\mathcal{E}^1)^n \rightarrow \mathcal{M}$ that is continuous in the strong topology on both sides.

Further,

$$(\varphi_0, \varphi_1, \dots, \varphi_n) \longmapsto \int \varphi_0 \operatorname{MA}(\varphi_1, \dots, \varphi_n)$$

is finite-valued and (strongly) continuous on tuples in \mathcal{E}^1 . In particular, the functional I from Section 4.2 extend continuously to \mathcal{E}^1 , and it induces a quasi-metric on \mathcal{E}^1/\mathbb{R} that defines the strong topology.

The energy of a probability measure $\mu \in \mathcal{M}$ on X^{an} is defined by

(4.9)
$$\mathbf{E}^{\vee}(\mu) \coloneqq \sup_{\varphi \in \mathcal{E}^1} \left\{ \mathbf{E}(\varphi) - \int \varphi \, \mathrm{d}\mu \right\},$$

where the supremum can be restricted to functions in $\mathcal{H}_{\mathbb{Q}}$, by approximation. This defines a convex, lsc function $E^{\vee} \colon \mathcal{M} \to [0, +\infty]$. We denote by

$$\mathcal{M}^1 \coloneqq \left\{ \mu \in \mathcal{M} \mid \mathbf{E}^{\vee}(\mu) < +\infty \right\}$$

the set of measures of finite energy. It comes with a strong topology, defined as the coarsest refinement of the weak topology of measures in which E^{\vee} is continuous. The topological space \mathcal{M}^1 does not depend on L.

By (4.5), for any $\varphi \in \mathcal{E}^1$, the measure $\mu = MA(\varphi)$ has finite energy, and φ achieves the supremum in (4.9), i.e.

(4.10)
$$\mathbf{E}^{\vee}(\mathbf{MA}(\varphi)) = \mathbf{E}(\varphi) - \int \varphi \ \mathbf{MA}(\varphi).$$

Conversely, a measure $\mu \in \mathcal{M}^1$ satisfies $\mu = MA(\varphi)$ with $\varphi \in \mathcal{E}^1$ iff φ achieves the supremum in (4.9). By a main result of [18], the Monge-Ampère operator induces a topological embedding with dense image

$$\mathrm{MA}\colon \mathcal{E}^1/\mathbb{R} \hookrightarrow \mathcal{M}^1,$$

with respect to the strong topology on both sides.

4.4. Envelopes

Consider a bounded-above family (φ_i) of *L*-psh functions, and set $\varphi \coloneqq \sup_i \varphi_i$. By definition, the usc regularization $\varphi^* \colon X^{\mathrm{an}} \to \mathbb{R} \cup \{-\infty\}$ is the smallest usc function such that $\varphi^* \ge \varphi$.

LEMMA 4.2. — The restriction of φ^* to X^{div} coincides with φ , and φ^* is the smallest usc function on X^{an} with this property.

Proof. — By [18, Theorem 5.6], points of X^{div} are non-negligible, which is a reformulation of the first assertion. Consider next a usc function $\psi: X^{\text{an}} \to \mathbb{R} \cup \{-\infty\}$ such that $\psi = \varphi$ on X^{div} . For each *i*, we then have $\varphi_i \leq \psi$ on X^{div} , and hence on X^{an} , by [18, Theorem 4.22]. Taking the supremum over *i* yields $\varphi \leq \psi$ on X^{an} , and hence $\varphi^* \leq \psi$, since ψ is usc.

We say that (X, L) has the envelope property if φ^* is L-psh for each bounded-above family of L-psh functions, using the above notation. It is proved in [18, Theorem 5.20] that the envelope property holds if X is smooth and k has characteristic zero, or in any characteristic if dim $X \leq 2$ [48]. For later use, we record:

LEMMA 4.3. — Let (φ_i) be a bounded-above, increasing net in \mathcal{E}^1 . Set $\varphi \coloneqq \sup_i \varphi_i$, and assume that φ^* is L-psh (e.g. L has the envelope property). Then $\varphi^* \in \mathcal{E}^1$ and $\varphi_i \to \varphi^*$ strongly in \mathcal{E}^1 .

Proof. — The first point holds because $\varphi^* \ge \varphi_i$, and hence $E(\varphi^*) \ge E(\varphi_i) > -\infty$. As recalled above, we have $\varphi^* = \varphi$ on X^{div} . Thus $\varphi_i \to \varphi^*$ pointwise on X^{div} , i.e. weakly in \mathcal{E}^1 , and hence strongly as well, since (φ_i) is an increasing net.

Given a function $\varphi \colon X^{\mathrm{an}} \to \mathbb{R} \cup \{\pm \infty\}$ we define the *psh envelope* pointwise as

$$\mathbf{P}(\varphi) \coloneqq \sup\{\psi \in \mathbf{PSH} \mid \psi \leqslant \varphi\}.$$

Note that the Fubini–Study envelope in (2.18) can be written

$$Q(\varphi) = \sup\{\psi \in \mathcal{H}_{\mathbb{Q}} \mid \psi \leqslant \varphi\} = \sup\{\psi \in CPSH \mid \psi \leqslant \varphi\}.$$

In both cases the convention $\sup \emptyset = -\infty$ applies. Clearly $Q(\varphi) \leq P(\varphi) \leq \varphi$, and either $\inf \varphi = -\infty \equiv Q(\varphi)$, or $Q(\varphi)$ is (finite-valued and) lsc. In the latter case,

(4.11)
$$Q(\varphi) = P(\varphi_{\star})$$

where φ_{\star} is the lsc regularization of φ (see [18, Lemma 5.19]). In particular, $Q(\varphi) = P(\varphi)$ when φ is continuous.

The functions $P(\varphi)$ and $Q(\varphi)$ are not psh in general. For any $\varphi \in C^0$, we have

$$P(\varphi) = Q(\varphi) \Longleftrightarrow P(\varphi) \in C^0 \Longleftrightarrow P(\varphi) \in PSH,$$

and these properties hold if (and only if) L has the envelope property. For the next result, see [18, Corollary 5.18].

LEMMA 4.4. — Assume that (X, L) has the envelope property, and consider a usc function $\varphi \colon X^{\mathrm{an}} \to \mathbb{R} \cup \{-\infty\}$. Then:

- (i) either $P(\varphi) \in PSH$ or $P(\varphi) \equiv -\infty$;
- (ii) if φ is the pointwise limit of a decreasing net of usc functions $\varphi_i \colon X^{\mathrm{an}} \to \mathbb{R} \cup \{-\infty\}$, then $\mathrm{P}(\varphi_i) \searrow \mathrm{P}(\varphi)$ pointwise on X^{an} .

Denote by $\mathcal{E}^{\infty} \subset \mathcal{E}^1$ the space of bounded *L*-psh functions. A function $\varphi \in \mathcal{E}^{\infty}$ is regularizable from below if there exists an increasing net $(\varphi_j)_j$ in CPSH that converges to φ in PSH (i.e. pointwise on X^{div}). Such a net can then be chosen in $\mathcal{H}_{\mathbb{Q}}$, and converges strongly to φ in \mathcal{E}^1 . We write

$$\mathcal{E}^\infty_\uparrow\subset\mathcal{E}^\infty$$

for the space of *L*-psh functions regularizable from below. If the envelope property holds, then a bounded function $\varphi \in \mathcal{E}^{\infty}$ lies in $\mathcal{E}^{\infty}_{\uparrow}$ iff its discontinuity locus $\{\varphi_{\star} < \varphi\}$ is pluripolar [18, Theorem 11.23].

Remark 4.5. — Assuming the envelope property, the inclusion $\mathcal{E}^{\infty}_{\uparrow} \subset \mathcal{E}^{\infty}$ is strict if $n \ge 1$, whereas CPSH $\subset \mathcal{E}^{\infty}_{\uparrow}$ is strict as soon as $n \ge 2$. See Examples 13.23 and 13.25 in [18].

For any bounded function $\varphi \in \mathcal{L}^{\infty}$, denote by $Q^*(\varphi) \coloneqq Q(\varphi)^*$ the use regularization of $Q(\varphi)$.

LEMMA 4.6. — Assume that (X, L) has the envelope property. Then:

- (i) $Q^*: \mathcal{L}^{\infty} \to \mathcal{L}^{\infty}$ is a projection operator onto $\mathcal{E}^{\infty}_{\uparrow}$;
- (ii) for all $\varphi, \psi \in \mathcal{E}^{\infty}_{\uparrow}$ we have $Q^{\star}(\varphi \wedge \psi) = P(\varphi \wedge \psi)$.

Proof. — (i) follows from [18, Theorem 13.24]. Pick $\varphi, \psi \in \mathcal{E}^{\infty}_{\uparrow}$. By (4.11) we have $Q(\varphi \land \psi) = P((\varphi \land \psi)_{\star}) = P(\varphi_{\star} \land \psi_{\star})$. Since φ, ψ are regularizable from below, their discontinuity locus is pluripolar, i.e. $\varphi_{\star} = \varphi, \psi_{\star} = \psi$, and hence $\varphi_{\star} \land \psi_{\star} = \varphi \land \psi$, outside a pluripolar set. By [18, Theorem 13.20], it follows that $P(\varphi_{\star} \land \psi_{\star})^{\star} = P(\varphi \land \psi)^{\star}$, which coincides with $P(\varphi \land \psi)$ since $\varphi \land \psi$ is usc (see Lemma 4.4). This proves (ii).

4.5. The extended energy

Recall from [18, Section 8]⁽²⁾ that the extended Monge–Ampère energy of an arbitrary function $\varphi \colon X^{\mathrm{an}} \to \mathbb{R} \cup \{\pm \infty\}$ is defined as

(4.12)
$$\widetilde{\mathcal{E}}(\varphi) \coloneqq \sup\{\mathcal{E}(\psi) \mid \psi \in \mathrm{PSH}, \psi \leqslant \varphi\} \in \mathbb{R} \cup \{\pm \infty\}.$$

Note that $\widetilde{E}(\varphi) = \widetilde{E}(P(\varphi))$, since any $\psi \in PSH$ satisfies $\psi \leq \varphi \Leftrightarrow \psi \leq P(\varphi)$. If $\varphi \colon X^{an} \to \mathbb{R} \cup \{+\infty\}$ is lsc (and hence bounded below), then $P(\varphi) = Q(\varphi)$, see (4.11), and hence

(4.13)
$$\widetilde{E}(\varphi) = \widetilde{E}(P(\varphi)) = \widetilde{E}(Q(\varphi)) = \sup\{E(\psi) \mid \psi \in \mathcal{H}_{\mathbb{R}}, \psi \leqslant \varphi\}.$$

A Dini-type argument (see [18, Proposition 8.3]) further yields:

LEMMA 4.7. — The functional $\varphi \mapsto \widetilde{E}(\varphi)$ is continuous along increasing nets of bounded-below lsc functions.

Following [19], we say that (X, L) has the weak envelope property if there exists a birational model $\pi: X' \to X$ and an ample Q-line bundle L'on X' such that $\pi^*L \leq L'$ and (X', L') has the envelope property. This is for instance the case whenever char k = 0, or if dim $X \leq 2$.

⁽²⁾ In loc. cit., the extended energy was simply denoted by $E(\varphi)$.

LEMMA 4.8. — Assume (X, L) has the weak envelope property, and pick any bounded-above family (φ_i) of L-psh functions. Set $\varphi \coloneqq \sup_i \varphi_i$. Then:

(i)
$$\varphi^* = \varphi$$
 on X^{lin} ;

(ii) if φ is further bounded below, then $\widetilde{E}(\varphi^*) = \widetilde{E}(\varphi)$.

Proof. — Point (i) means that each $v \in X^{\text{lin}}$ is non-negligible. Use the previous notation. Since π is birational, we have $\pi^{-1}(\{v\}) = \{v'\}$ with $v' \in X'^{\text{lin}}$. By [18, Lemma 5.4], it suffices to show that v' is non-negligible, and this follows from [18, Theorem 13.17], which applies because (X', L') has the envelope property. Finally, (ii) follows from [19, Theorem B], as the assumption guarantees that $P(\varphi) = \varphi$.

5. Darvas metrics

In this section, we study the metrics on the spaces $\mathcal{H}_{\mathbb{R}}$, \mathcal{E}^1 and \mathcal{M}^1 induced by the d₁-pseudometric of $\mathcal{N}_{\mathbb{R}}$, and prove the main part of Theorem B.

5.1. Volume vs. energy

The next result will be a key tool in what follows.

THEOREM 5.1. — For any $\chi \in \mathcal{N}_{\mathbb{R}}$ we have $\operatorname{vol}(\chi) = \widetilde{E}(FS(\chi))$.

Here $FS(\chi)$ is bounded and lsc, but not *L*-psh in general, and $E(FS(\chi))$ is its extended energy (see Section 4.5).

Proof. — Consider the round-down $\chi' := \lfloor \chi \rfloor \in \mathcal{N}_{\mathbb{Z}}$. Then $d_{\infty}(\chi, \chi') = 0$ (see Example 1.7), and hence $FS(\chi) = FS(\chi')$, $vol(\chi) = vol(\chi')$. As a result, we may and do assume $\chi \in \mathcal{N}_{\mathbb{Z}}$. By Theorem 3.18, the canonical approximants $\chi_d \in \mathcal{T}_{\mathbb{Z}}$ satisfy $vol(\chi_d) \to vol(\chi)$. On the other hand, $FS(\chi_d) = FS_d(\chi)$ increases pointwise to $FS(\chi)$ (see (2.9)), and hence $E(FS(\chi_d)) = \widetilde{E}(FS(\chi_d)) \to \widetilde{E}(FS(\chi))$, by Lemma 4.7.

We are thus reduced to the case $\chi \in \mathcal{T}_{\mathbb{Z}}$, which is a consequence of [17]. Indeed, χ corresponds to an ample test configuration $(\mathcal{X}, \mathcal{L})$ for (X, L) under the Rees correspondence (see Appendix A). By [17, Proposition 3.12], the spectral measure $\sigma(\chi)$ coincides with the Duistermaat–Heckman measure DH $(\mathcal{X}, \mathcal{L})$, and passing to the barycenters yields

$$\operatorname{vol}(\chi) = \frac{(\bar{\mathcal{L}}^{n+1})}{(n+1)(L^n)},$$

by [17, Lemma 7.3]. By (4.4), the right-hand side is also equal to $E(FS(\chi)) = \widetilde{E}(FS(\chi))$, and the result follows.

COROLLARY 5.2. — Any norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is asymptotically equivalent to its homogenization, i.e. $\chi \sim \chi^{\text{hom}}$.

Proof. — Since $\chi \leq \chi^{\text{hom}}$, it suffices to show $\text{vol}(\chi) = \text{vol}(\chi^{\text{hom}})$ (see Lemma 3.11). This follows from Theorem 5.1, since $\text{FS}(\chi) = \text{FS}(\chi^{\text{hom}})$ by Proposition 2.14. □

COROLLARY 5.3. — For any $\varphi \in \mathcal{L}^{\infty}$ we have $\operatorname{vol}(\operatorname{IN}(\varphi)) = \widetilde{\operatorname{E}}(\operatorname{Q}(\varphi)) = \widetilde{\operatorname{E}}(\varphi_{\star}).$

Proof. — Since $IN(\varphi_{\star}) = IN(\varphi)$ (see (2.15)), we may assume that φ is lsc, and hence $\widetilde{E}(\varphi) = \widetilde{E}(Q(\varphi))$, see (4.13). Now $FS(IN(\varphi)) = Q(\varphi)$, by Proposition 2.29, and we conclude by Theorem 5.1.

5.2. The Darvas metric on $\mathcal{H}_{\mathbb{R}}$

Recall from Corollary 2.18 that the operators

$$\mathrm{FS} \colon (\mathcal{T}_{\mathbb{R}}, \mathrm{d}_{\infty}) \longrightarrow (\mathcal{H}_{\mathbb{R}}, \mathrm{d}_{\infty}), \quad \mathrm{IN} \colon (\mathcal{H}_{\mathbb{R}}, \mathrm{d}_{\infty}) \hookrightarrow (\mathcal{T}_{\mathbb{R}}, \mathrm{d}_{\infty})$$

are isometries such that $FS \circ IN = id$.

For any $p \in [1, \infty]$, the pseudo-metric d_p on $\mathcal{T}_{\mathbb{R}} \subset \mathcal{N}_{\mathbb{R}}$ satisfies $d_p \leq d_{\infty}$; it is thus constant along the fibers of FS, and hence descends to a pseudometric d_p on $\mathcal{H}_{\mathbb{R}}$, such that

 $FS: (\mathcal{T}_{\mathbb{R}}, d_p) \longrightarrow (\mathcal{H}_{\mathbb{R}}, d_p), \quad IN: (\mathcal{H}_{\mathbb{R}}, d_p) \longmapsto (\mathcal{T}_{\mathbb{R}}, d_p)$

are isometries. Theorem A asserts that d_p is a metric on $\mathcal{H}_{\mathbb{R}}$. Since $d_p \ge d_1$, this follows from the following more precise result.

THEOREM 5.4. — The pseudo-metric d_1 on $\mathcal{H}_{\mathbb{R}}$ is a metric, uniquely characterized by

(5.1)
$$\varphi \ge \psi \Longrightarrow d_1(\varphi, \psi) = E(\varphi) - E(\psi);$$

(5.2)
$$d_1(\varphi, \psi) = \inf\{d_1(\varphi, \tau) + d_1(\tau, \psi) \mid \tau \in \mathcal{H}_{\mathbb{R}}, \tau \leq \varphi \land \psi\},\$$

for all $\varphi, \psi \in \mathcal{H}_{\mathbb{R}}$.

Proof. — We first prove that d_1 satisfies (5.1) and (5.2), which will take care of uniqueness. Pick $\varphi, \psi \in \mathcal{H}_{\mathbb{R}}$, and set $\chi := IN(\varphi), \chi' := IN(\psi)$. Then $FS(\chi) = \varphi$, $FS(\chi') = \psi$, and Theorem 5.1 implies $vol(\chi) = E(\varphi)$, $vol(\chi') = E(\psi)$. If $\varphi \ge \psi$, then $\chi \ge \chi'$, and (3.15) yields

$$d_1(\varphi, \psi) = d_1(\chi, \chi') = \operatorname{vol}(\chi) - \operatorname{vol}(\chi') = \operatorname{E}(\varphi) - \operatorname{E}(\psi),$$

ANNALES DE L'INSTITUT FOURIER

which proves (5.1). Next, pick $\varepsilon > 0$. Applying Theorem 3.18 to $\chi \wedge \chi'$ yields $\chi'' \in \mathcal{T}_{\mathbb{R}}$ such that $\chi'' \leq \chi \wedge \chi'$ and $d_1(\chi'', \chi \wedge \chi') \leq \varepsilon$. If we set $\tau := FS(\chi'') \in \mathcal{H}_{\mathbb{R}}$, then $\tau \leq \varphi, \psi$, and

$$d_1(\varphi, \tau) + d_1(\tau, \psi) = d_1(\chi, \chi'') + d_1(\chi'', \chi')$$

$$\leq d_1(\chi, \chi \land \chi') + d_1(\chi \land \chi', \chi') + 2\varepsilon$$

$$= d_1(\chi, \chi') + 2\varepsilon = d_1(\varphi, \psi) + 2\varepsilon,$$

where we have used (3.15). This proves (5.2). Assume now $d_1(\varphi, \psi) = 0$. By (5.1) and (5.2), there exists a sequence (τ_i) in $\mathcal{H}_{\mathbb{R}}$ such that $\tau_i \leq \varphi, \psi$ and $E(\tau_i) \to E(\varphi)$ and $E(\tau_i) \to E(\psi)$. By [18, Proposition 12.6], it follows that (τ_i) converges to both φ and ψ in \mathcal{E}^1 , and hence $\varphi = \psi$, since the topology is separated. This proves, as desired, that d_1 is a metric on $\mathcal{H}_{\mathbb{R}}$ (this conclusion alternatively follows from (5.7) below).

5.3. The Darvas metric on \mathcal{E}^1

We next prove that the metric d_1 on $\mathcal{H}_{\mathbb{R}}$ canonically extends to \mathcal{E}^1 , yielding an analogue in our context of the metric introduced by Darvas [31] in the complex analytic setting.

THEOREM 5.5. — There exists a unique metric d_1 on \mathcal{E}^1 that defines the strong topology and restricts to the previous metric d_1 on $\mathcal{H}_{\mathbb{R}}$. Further:

(i) for all $\varphi, \psi \in CPSH$, we have

(5.3)
$$d_1(\varphi, \psi) = E(\varphi) + E(\psi) - 2\widetilde{E}(P(\varphi \land \psi));$$

- (ii) the metric space (\$\mathcal{E}^1\$, d₁) is complete iff the envelope property holds for (\$X\$, \$L\$);
- (iii) if the envelope property holds, then (5.3) remains valid for all $\varphi, \psi \in \mathcal{E}^1$, and $P(\varphi \land \psi) \in \mathcal{E}^1$.

Recall that the envelope property holds whenever X is smooth and $\operatorname{char}(k) = 0$, and fails if X is not unibranch. We refer to the metric d_1 on \mathcal{E}^1 as the *Darvas metric*. By [64], (\mathcal{E}^1, d_1) is a geodesic metric space, assuming the envelope property.

Our strategy to extend d_1 to \mathcal{E}^1 is to compare it to the functional

$$\bar{\mathrm{I}}(\varphi,\psi) \coloneqq \mathrm{I}(\varphi,\psi) + |\mathrm{sup}\,\varphi - \mathrm{sup}\,\psi|.$$

It was indeed proven in [18, Section 12.1] that \overline{I} is a quasi-metric on \mathcal{E}^1 that defines the strong topology, and further satisfies

(5.4)
$$I(\varphi, \psi) = I(\varphi, \varphi \lor \psi) + I(\varphi \lor \psi, \psi).$$

 Set

$$\overline{\mathbf{I}}(\varphi) \coloneqq \overline{\mathbf{I}}(\varphi, 0), \quad \mathbf{d}_1(\varphi) \coloneqq \mathbf{d}_1(\varphi, 0).$$

By the quasi-triangle inequality, (5.4) implies

(5.5)
$$\overline{I}(\varphi \lor \psi) \lesssim \max{\{\overline{I}(\varphi), \overline{I}(\psi)\}}.$$

LEMMA 5.6. — The quasi-metrics d_1 and \overline{I} on $\mathcal{H}_{\mathbb{R}}$ are Hölder comparable, in the sense that

(5.6)
$$d_1(\varphi,\psi) \lesssim \bar{I}(\varphi,\psi)^{\alpha} \max\{\bar{I}(\varphi),\bar{I}(\psi)\}^{1-\alpha}$$

(5.7)
$$\bar{\mathbf{I}}(\varphi,\psi) \lesssim \mathbf{d}_1(\varphi,\psi)^{\alpha} \max\{\mathbf{d}_1(\varphi),\mathbf{d}_1(\psi)\}^{1-\alpha},$$

with $\alpha \coloneqq 1/2^n$. In particular, $d_1(\varphi) \approx \overline{I}(\varphi)$.

Proof. — First assume $\varphi \ge \psi$. Then (5.1) and (4.6) show that

$$d_1(\varphi, \psi) = E(\varphi) - E(\psi) \approx \int (\varphi - \psi) \operatorname{MA}(\psi),$$

and hence

$$I(\varphi, \psi) = \int (\varphi - \psi) (MA(\psi) - MA(\varphi)) \lesssim d_1(\varphi, \psi).$$

Now we can write

(5.8)
$$\bar{\mathbf{I}}(\varphi,\psi) = \int (\varphi-\psi) \, \mathbf{MA}(\psi) + \int (\varphi-\psi) (\mathbf{MA}(0) - \mathbf{MA}(\varphi)).$$

As a special case of [18, Lemma 7.30] we have the estimate

$$\left| \int (\varphi - \psi) (\mathrm{MA}(0) - \mathrm{MA}(\varphi)) \right| \lesssim \mathrm{I}(\varphi, \psi)^{\alpha} \max\{\mathrm{I}(\varphi), \mathrm{I}(\psi)\}^{1-\alpha}.$$

Since $I \leq I$, this yields on the one hand

$$\begin{split} d_{1}(\varphi,\psi) &\lesssim \bar{I}(\varphi,\psi) + I(\varphi,\psi)^{\alpha} \max\{\bar{I}(\varphi),\bar{I}(\psi)\}^{1-\alpha} \\ &\lesssim \bar{I}(\varphi,\psi)^{\alpha} \max\{\bar{I}(\varphi,\psi),\bar{I}(\varphi),\bar{I}(\psi)\}^{1-\alpha} \\ &\lesssim \bar{I}(\varphi,\psi)^{\alpha} \max\{\bar{I}(\varphi),\bar{I}(\psi)\}^{1-\alpha}. \end{split}$$

Since $I \leq d_1$, we get on the other hand

$$\begin{split} \bar{I}(\varphi,\psi) &\lesssim d_1(\varphi,\psi) + d_1(\varphi,\psi)^{\alpha} \max\{d_1(\varphi),d_1(\psi)\}^{1-\alpha} \\ &\lesssim d_1(\varphi,\psi)^{\alpha} \max\{d_1(\varphi),d_1(\psi)\}^{1-\alpha}. \end{split}$$

This proves (5.6), and (5.7) when $\varphi \ge \psi$.

Now consider arbitrary $\varphi, \psi \in \mathcal{H}_{\mathbb{R}}$. To prove (5.6), set $\sigma \coloneqq \varphi \lor \psi \in \mathcal{H}_{\mathbb{R}}$. From what precedes and (5.5), we have

$$d_1(\varphi,\sigma) \lesssim \overline{I}(\varphi,\sigma)^{\alpha} \max{\{\overline{I}(\varphi),\overline{I}(\psi)\}^{1-\alpha}}$$

ANNALES DE L'INSTITUT FOURIER

890

and

$$d_1(\psi, \sigma) \lesssim \bar{I}(\psi, \sigma)^{\alpha} \max\{\bar{I}(\varphi), \bar{I}(\psi)\}^{1-\alpha},$$

which together with the triangle inequality for d_1 yields (5.6).

The proof of (5.7) is similar. By (5.2) we can pick $\tau \in \mathcal{H}_{\mathbb{R}}$ with $\tau \leq \varphi, \psi$ such that $\max\{d_1(\tau, \varphi), d_1(\tau, \psi)\} \leq d_1(\varphi, \psi)$, and hence $d_1(\tau) \leq \max\{d_1(\varphi), d_1(\psi)\}$. Since $\tau \leq \varphi, \psi$, the first step yields

$$\bar{I}(\varphi,\tau) \lesssim d_1(\varphi,\tau)^{\alpha} \max\{d_1(\varphi),d_1(\tau)\}^{1-\alpha}$$

$$\lesssim d_1(\varphi,\psi)^{\alpha} \max\{d_1(\varphi),d_1(\psi)\}^{1-\alpha},$$

$$\bar{I}(\psi,\tau) \lesssim d_1(\psi,\tau)^{\alpha} \max\{d_1(\psi),d_1(\tau)\}^{1-\alpha}$$

$$\lesssim d_1(\varphi,\psi)^{\alpha} \max\{d_1(\varphi),d_1(\psi)\}^{1-\alpha},$$

and the quasi-triangle inequality for I yields (5.7).

Proof of Theorem 5.5. — Since $\mathcal{H}_{\mathbb{R}}$ is dense in \mathcal{E}^1 , uniqueness is clear. Given $\varphi, \psi \in \mathcal{E}^1$, pick sequences (φ_i) , (ψ_i) in $\mathcal{H}_{\mathbb{R}}$ converging strongly to φ and ψ , respectively (for example, we can use decreasing sequences). Thus $\lim_i \bar{I}(\varphi_i, \varphi) = \lim_i \bar{I}(\psi_i, \psi) = 0$. Using (5.6) this implies that $(d_1(\varphi_i, \psi_i))_i$ is a Cauchy sequence, so that $d_1(\varphi, \psi) \coloneqq \lim_j d_1(\varphi_j, \psi_j)$ exists. It is easy to see that it does not depend on the choice of sequence (φ_i) and (ψ_i) , and that the extension is a pseudo-metric on \mathcal{E}^1 . Further, the estimates of Lemma 5.6 still hold for $\varphi, \psi \in \mathcal{E}^1$. In particular $d_1(\varphi, \psi) = 0$ iff $\bar{I}(\varphi, \psi) = 0$ iff $\varphi = \psi$, so d_1 is a metric on \mathcal{E}^1 . These estimates also show that d_1 and \bar{I} share the same Cauchy sequences in \mathcal{E}^1 , so that (\mathcal{E}^1, d_1) is complete iff (\mathcal{E}^1, \bar{I}) is complete. By [18, Theorem 12.8], this is also equivalent to the envelope property for (X, L), which proves (ii).

Next, pick $\varphi, \psi \in \mathcal{H}_{\mathbb{R}}$. By (5.1) and (5.2), we have

$$d_{1}(\varphi, \psi) = \inf \left\{ d_{1}(\varphi, \tau) + d_{1}(\tau, \psi) \mid \tau \in \mathcal{H}_{\mathbb{R}}, \, \tau \leqslant \varphi \land \psi \right\}$$
$$= E(\varphi) + E(\psi) - 2 \sup \{ E(\psi) \mid \psi \in \mathcal{H}_{\mathbb{R}}, \, \psi \leqslant \varphi \land \psi \}$$
$$= E(\varphi) + E(\psi) - 2 \widetilde{E}(P(\varphi \land \psi)),$$

see (4.13). Since $d_1 \leq d_{\infty}$, all terms are continuous with respect to uniform convergence, and the identity therefore remains valid on CPSH, which yields (i).

Finally, assume that the envelope property holds. Pick $\varphi, \psi \in \mathcal{E}^1$, set $\rho \coloneqq P(\varphi \land \psi)$, and choose decreasing nets (φ_i) , (ψ_i) in $\mathcal{H}_{\mathbb{R}}$ converging to φ, ψ . By Lemma 4.4, we either have $\rho \in PSH$ or $\rho \equiv -\infty$, and $P(\varphi_i \land \psi_i)$ decreases pointwise to ρ . Since (5.3) holds for $\varphi_i, \psi_i \in \mathcal{H}_{\mathbb{R}}$, it also holds for

 φ, ψ , by continuity of E along decreasing nets. In particular, $E(\rho)$ is finite, and hence $\rho \in \mathcal{E}^1$. This proves (iii).

We also note the following useful Lipschitz property:

 $\text{Proposition 5.7.} \quad \text{ For all } \varphi, \psi \in \mathcal{E}^1 \text{ we have } |\operatorname{E}(\varphi) - \operatorname{E}(\psi)| \leqslant d_1(\varphi, \psi).$

Proof. — By continuity, we may assume $\varphi, \psi \in \mathcal{H}_{\mathbb{R}}$. Pick $\varepsilon > 0$ and $\tau \in \mathcal{H}_{\mathbb{R}}$ such that $\tau \leq \varphi \wedge \psi$ and $E(\varphi) + E(\psi) - 2E(\tau) \leq d_1(\varphi, \psi) + \varepsilon$, see (5.2). Then

$$\begin{aligned} |\mathrm{E}(\varphi) - \mathrm{E}(\psi)| &\leq |\mathrm{E}(\varphi) - \mathrm{E}(\tau)| + |\mathrm{E}(\tau) - \mathrm{E}(\psi)| = \mathrm{E}(\varphi) + \mathrm{E}(\psi) - 2\,\mathrm{E}(\tau) \\ &\leq \mathrm{d}_1(\varphi, \psi) + \varepsilon, \end{aligned}$$

 \square

and the result follows.

As in [33, Theorem 3.7], we next provide a comparison of the Darvas metric d_1 on \mathcal{E}^1 with the functional $I_1: \mathcal{E}^1 \times \mathcal{E}^1 \to \mathbb{R}_{\geq 0}$ defined by

$$I_1(\varphi,\psi) \coloneqq \int |\varphi - \psi|(MA(\varphi) + MA(\psi)).$$

This functional is obviously symmetric, and it separates points, as a consequence of the Domination Principle (see [18, Corollary 10.6]). For all $\varphi, \psi \in \mathcal{E}^1$, we further have

(5.9)
$$I_1(\varphi, \psi) = I_1(\varphi, \varphi \lor \psi) + I_1(\varphi \lor \psi, \psi)$$

As with I and \overline{I} , this follows from the Locality Principle (see [18, Theorem 7.40, Proposition 7.45]).

THEOREM 5.8. — For all $\varphi, \psi \in \mathcal{E}^1$ we have $d_1(\varphi, \psi) \approx I_1(\varphi, \psi)$.

The proof relies on the following analogue of [33, Lemma 3.8].

LEMMA 5.9. — If $\varphi, \psi \in \mathcal{E}^1$ and $\rho \coloneqq \frac{1}{2}(\varphi + \psi)$, then $d_1(\varphi, \psi) \approx d_1(\varphi, \rho) + d_1(\rho, \psi)$.

Proof. — By approximation, we may assume $\varphi, \psi \in \mathcal{H}_{\mathbb{R}}$. Pick any $\tau \in \mathcal{H}_{\mathbb{R}}$ with $\tau \leq \varphi \wedge \psi$. Then $\tau \leq \varphi, \psi, \rho$, and (5.1) yields

$$\begin{split} \mathbf{d}_{1}(\varphi,\rho) + \mathbf{d}_{1}(\rho,\psi) \\ &\leqslant \mathbf{d}_{1}(\varphi,\tau) + \mathbf{d}_{1}(\psi,\tau) + 2\,\mathbf{d}_{1}(\rho,\tau) \\ &= (\mathbf{E}(\varphi) - \mathbf{E}(\tau)) + (\mathbf{E}(\psi) - \mathbf{E}(\tau)) + 2(\mathbf{E}(\rho) - \mathbf{E}(\tau)) \\ &\approx \int (\varphi-\tau)\,\mathbf{M}\mathbf{A}(\tau) + \int (\psi-\tau)\,\mathbf{M}\mathbf{A}(\tau) + 2\int (\rho-\tau)\,\mathbf{M}\mathbf{A}(\tau) \\ &= 2\int (\varphi-\tau)\,\mathbf{M}\mathbf{A}(\tau) + 2\int (\psi-\tau)\,\mathbf{M}\mathbf{A}(\tau) \\ &\approx (\mathbf{E}(\varphi) - \mathbf{E}(\tau) + \mathbf{E}(\psi) - \mathbf{E}(\tau)) = \mathbf{d}_{1}(\varphi,\tau) + \mathbf{d}_{1}(\psi,\tau). \end{split}$$

ANNALES DE L'INSTITUT FOURIER

Here the first inequality is simply the triangle inequality for d_1 , whereas the third and fifth lines follow from (4.6). By (5.2), the infimum over τ of the right-hand side equals $d_1(\varphi, \psi)$ and we are done.

Proof of Theorem 5.8. — Since $d_1(\varphi, \psi)$ and $I_1(\varphi, \psi)$ are both continuous along decreasing nets, we may assume wlog $\varphi, \psi \in \mathcal{H}_{\mathbb{R}}$. Let us start by proving $d_1(\varphi, \psi) \leq I_1(\varphi, \psi)$. By (5.9) and the triangle inequality for d_1 , it suffices to consider the case $\varphi \geq \psi$. But in this case,(4.6) yields

$$d_1(\varphi, \psi) = E(\varphi) - E(\psi) \approx \int (\varphi - \psi) \operatorname{MA}(\psi) \leqslant I_1(\varphi, \psi).$$

It remains to prove $d_1(\varphi, \psi) \gtrsim I_1(\varphi, \psi)$. Set $\rho \coloneqq \frac{1}{2}(\varphi + \psi) \in \mathcal{H}_{\mathbb{R}}$, so that Lemma 5.9 gives $d_1(\varphi, \psi) \approx d_1(\varphi, \rho) + d_1(\rho, \psi)$. Pick $\varepsilon > 0$. By (5.2), we can find $\sigma, \tau \in \mathcal{H}_{\mathbb{R}}$ such that $\sigma \leq \varphi \land \rho, \tau \leq \rho \land \psi$, and

 $d_1(\varphi,\rho) \ge d_1(\varphi,\sigma) + d_1(\sigma,\rho) - \varepsilon \quad \text{and} \quad d_1(\rho,\psi) \ge d_1(\rho,\tau) + d_1(\tau,\psi) - \varepsilon,$ and hence

$$d_1(\varphi, \psi) \gtrsim d_1(\sigma, \rho) + d_1(\rho, \tau) - 2\varepsilon.$$

As $\sigma \leq \rho$, we have

$$d_1(\sigma,\rho) = E(\rho) - E(\sigma) \ge \int (\rho - \sigma) \operatorname{MA}(\rho) \ge 2^{-n} \int (\rho - \sigma) (\operatorname{MA}(\varphi) + \operatorname{MA}(\psi)),$$

where the last inequality follows by expanding $MA(\rho) = MA(\frac{1}{2}(\varphi + \psi))$. Combining this with the analogous lower bound on $d_1(\rho, \tau)$ yields

$$d_1(\varphi,\psi) \gtrsim \int (2\rho - \sigma - \tau) (MA(\varphi) + MA(\psi)) - 2\varepsilon$$

We conclude by noting that $2\rho - \sigma - \tau \ge \frac{1}{2}|\varphi - \psi|$ and letting $\varepsilon \to 0$. \Box

5.4. The Darvas metric on \mathcal{M}^1

By [18, Proposition 12.7], the Monge–Ampère operator induces a topological embedding with dense image

MA:
$$\mathcal{E}^1/\mathbb{R} \hookrightarrow \mathcal{M}^1$$
,

where \mathcal{E}^1 and \mathcal{M}^1 are both equipped with the strong topology. In particular, the quotient topology of \mathcal{E}^1/\mathbb{R} is Hausdorff. Since the action of \mathbb{R} on \mathcal{E}^1 by translation preserves d_1 , the topology of \mathcal{E}^1/\mathbb{R} is defined by the quotient metric

(5.10)
$$\underline{\mathbf{d}}_{1}(\varphi,\psi) = \inf_{c \in \mathbb{R}} \mathbf{d}_{1}(\varphi+c,\psi).$$

Note also that the isometric surjection FS: $(\mathcal{T}_{\mathbb{R}}, d_1) \twoheadrightarrow (\mathcal{H}_{\mathbb{R}}, d_1)$, being equivariant with respect to the action of \mathbb{R} , induces an isometric surjection

where \underline{d}_1 respectively denotes the restriction of the quotient metric on $\mathcal{N}_{\mathbb{R}}/\mathbb{R}$ and \mathcal{E}^1/\mathbb{R} .

As in the proof of Lemma 3.15, we have:

LEMMA 5.10. — For all $\varphi, \psi \in \mathcal{E}^1$, there exists $c \in \mathbb{R}$ such that $\underline{d}_1(\varphi, \psi) = d_1(\varphi + c, \psi)$ and $|c| \leq 2 d_1(\varphi, \psi) \lesssim \max\{\overline{I}(\varphi), \overline{I}(\psi)\}.$

This provides another reason why (5.10) defines a metric on \mathcal{E}^1/\mathbb{R} .

THEOREM 5.11. — There exists a unique metric d_1 on \mathcal{M}^1 that defines the strong topology and restricts to the quotient metric (5.10) on $\mathcal{E}^1/\mathbb{R} \to \mathcal{M}^1$. Furthermore, the metric space (\mathcal{M}^1, d_1) is complete.

Note that completeness this time holds with or without the envelope property, in contrast with Theorem 5.5. As with the latter, the proof is based on a comparison of \underline{d}_1 with the translation invariant functional I: $\mathcal{E}^1 \times \mathcal{E}^1 \to \mathbb{R}_{\geq 0}$.

LEMMA 5.12. — The quasi-metrics \underline{d}_1 and I on \mathcal{E}^1/\mathbb{R} are Hölder comparable, i.e.

(5.12) $\underline{\mathbf{d}}_{1}(\varphi,\psi) \lesssim \mathbf{I}(\varphi,\psi)^{\alpha} \max\{\mathbf{I}(\varphi),\mathbf{I}(\psi)\}^{1-\alpha},$

(5.13)
$$I(\varphi,\psi) \lesssim \underline{d}_1(\varphi,\psi)^{\alpha} \max\{\underline{d}_1(\varphi),\underline{d}_1(\psi)\}^{1-\alpha}$$

for all $\varphi, \psi \in \mathcal{E}^1$, with $\alpha \coloneqq 1/2^n$. In particular, $\underline{d}_1(\varphi) \approx I(\varphi)$.

Proof. — By translation invariance of I and \underline{d}_1 , we may assume $\sup \varphi = \sup \psi = 0$. Then (5.12) follows directly from (5.6), since $\underline{d}_1(\varphi, \psi) \leq \underline{d}_1(\varphi, \psi)$. By Lemma 5.10, we can find $c \in \mathbb{R}$ such that $\underline{d}_1(\varphi, \psi) = \underline{d}_1(\varphi + c, \psi)$ and $|c| \leq \max\{I(\varphi), I(\psi)\}$. By (5.7) we infer

(5.14)
$$I(\varphi,\psi) \leq \bar{I}(\varphi+c,\psi) \leq \underline{d}_1(\varphi,\psi)^{\alpha} \max\{I(\varphi),I(\psi)\}^{1-\alpha}.$$

In particular, $I(\varphi) \leq \underline{d}_1(\varphi)^{\alpha} I(\varphi)^{1-\alpha}$; hence $I(\varphi) \leq \underline{d}_1(\varphi)$, and (5.13) follows.

Proof of Theorem 5.11. — Since $\mathcal{E}^1/\mathbb{R} \hookrightarrow \mathcal{M}^1$ has dense image, uniqueness is clear. By [18, Theorem 10.12], the strong topology of \mathcal{M}^1 is defined by a certain quasi-metric I^{\vee} , that further satisfies $I^{\vee}(MA(\varphi), MA(\psi)) \approx$ $I(\varphi, \psi)$. Using the estimates of Lemma 5.12 and arguing just as in the proof of Theorem 5.5, we infer the existence of an extension of \underline{d}_1 to a pseudo-metric d_1 on \mathcal{M}^1 such that

(5.15)
$$d_1(\mu, \delta_{v_{\rm triv}}) \approx \mathbf{I}^{\vee}(\mu, \delta_{v_{\rm triv}}) \approx \mathbf{E}^{\vee}(\mu),$$

(5.16)
$$d_1(\mu, \mu') \lesssim I^{\vee}(\mu, \mu')^{\alpha} \max\{E^{\vee}(\mu), E^{\vee}(\mu')\}^{1-\alpha},$$

(5.17) $I^{\vee}(\mu,\mu') \lesssim d_1(\mu,\mu')^{\alpha} \max\{E^{\vee}(\mu)\}, E^{\vee}(\mu')\}^{1-\alpha},$

for all $\mu, \nu \in \mathcal{M}^1$ (recalling that MA(0) = $\delta_{v_{\text{triv}}}$). This shows that d_1 separates points, and hence is a metric on \mathcal{M}^1 , which further shares the same convergent and Cauchy sequences with I^{\vee} . It thus defines the strong topology of \mathcal{M}^1 , and (\mathcal{M}^1, d_1) is complete, because $(\mathcal{M}^1, I^{\vee})$ is complete by [18, Theorem 10.14].

Combining the above estimates with a key estimate for Monge–Ampère integrals from [18], we get the following Hölder continuity property:

THEOREM 5.13. — There exist $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}_{>0}$, only depending on n, such that $\sum_i \alpha_i = 1$ and

(5.18)
$$\left| \int |\varphi - \varphi'| \left(\mu - \mu' \right) \right| \lesssim \mathrm{d}_1(\varphi, \varphi')^{\alpha_1} \, \mathrm{d}_1(\mu, \mu')^{\alpha_2} M^{\alpha_3}$$

for all $\varphi, \varphi' \in \mathcal{E}^1$ and $\mu, \mu' \in \mathcal{M}^1$, where

$$M \coloneqq \max\{\mathbf{I}(\varphi), \mathbf{I}(\varphi'), \mathbf{E}^{\vee}(\mu), \mathbf{E}^{\vee}(\mu')\}.$$

Further, there exists $\alpha \in (0,1)$ only depending on n such that

(5.19)
$$\|\varphi - \varphi'\|_{L^1(\mu)} \lesssim d_1(\varphi, \varphi')^{\alpha} \max\{I(\varphi), I(\varphi'), E^{\vee}(\mu)\}^{1-\alpha}.$$

Proof. — By [18, Theorem 10.3], we have

$$\left| \int (\varphi - \varphi')(\mu - \mu') \right| \lesssim \mathbf{I}(\varphi, \varphi')^{\alpha} \, \mathbf{I}^{\vee}(\mu, \mu')^{\frac{1}{2}} M^{\frac{1}{2} - \alpha}$$

Injecting (5.13) and (5.17) yields

(5.20)
$$\left| \int (\varphi - \varphi')(\mu - \mu') \right| \lesssim \mathrm{d}_1(\varphi, \varphi')^{\alpha_1} \, \mathrm{d}_1(\mu, \mu')^{\alpha_2} M^{\alpha_3}$$

with $\alpha_1, \alpha_2, \alpha_3$ as above. Next, write

$$|\varphi - \varphi'| = 2(\tau - \varphi') + (\varphi' - \varphi)$$

with $\tau \coloneqq \varphi \lor \varphi' \in \mathcal{E}^1$. On the one hand, we have $I(\tau) \leq M$. On the other hand, Theorem 5.8 and (5.9) yield

$$d_1(\tau, \varphi') \approx I_1(\tau, \varphi') \leqslant I_1(\varphi, \varphi') \approx d_1(\varphi, \varphi').$$

TOME 75 (2025), FASCICULE 2

Applying (5.20) to τ, φ' and φ', φ now yields (5.18). To prove (5.19), set $\mu' = MA(\varphi)$. Then $E^{\vee}(\mu) \approx I(\varphi)$, and hence

$$\left| \int |\varphi - \varphi'| \left(\mu - \mathrm{MA}(\varphi) \right) \right| \lesssim \mathrm{d}_1(\varphi, \varphi')^{\alpha} \max\{\mathrm{I}(\varphi), \mathrm{I}(\varphi'), \mathrm{E}^{\vee}(\mu)\}^{1-\alpha}.$$

By Theorem 5.8, we have on the other hand $\int |\varphi - \varphi'| \operatorname{MA}(\varphi) \lesssim d_1(\varphi, \varphi')$, and summing up these estimates yields (5.19).

6. Divisorial and maximal norms

The restriction of the pseudometric d_1 to the subspace $\mathcal{N}_{\mathbb{R}}^{\text{hom}} \subset \mathcal{N}_{\mathbb{R}}$ of homogeneous norms is still not a metric unless dim X = 0, see Example 3.10. Here we study further subspaces on which d_1 does induce a metric.

One such subspace consists of divisorial norms, defined by finitely many divisorial valuations. These play an important role in the notion of divisorial stability introduced and studied in [20]. We then show that, at least in characteristic zero, there is a canonical maximal subspace of $\mathcal{N}_{\mathbb{R}}^{\text{hom}}$ on which d_1 is a metric. In particular, we prove Theorem D.

6.1. General infimum norms

The following construction generalizes the one in Section 2.4.

DEFINITION 6.1. — For any non-pluripolar set $\Sigma \subset X^{\mathrm{an}}$, and any bounded function $\varphi \colon \Sigma \to \mathbb{R}$, let $\mathrm{IN}_{\Sigma}(\varphi) \in \mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}$ denote the homogeneous norm defined for $s \in R_m$ by

(6.1)
$$\operatorname{IN}_{\Sigma}(\varphi)(s) = \inf_{v \in \Sigma} \{v(s) + m\varphi(v)\}.$$

Note that $\exp(-\operatorname{IN}_{\Sigma}(\varphi)(s))$ coincides with the more usual supnorm $\sup_{\Sigma} |s| e^{-m\varphi}$. The filtration corresponding to $\operatorname{IN}_{\Sigma}(\varphi)$ is given by

 $F^{\lambda}R_m = \{s \in R_m \mid v(s) + m\varphi(v) \ge \lambda \text{ for all } v \in \Sigma\}, \quad \lambda \in \mathbb{R}.$

The condition that Σ is non-pluripolar, which is equivalent to $T(\Sigma) < \infty$ (see (4.1)) and holds as soon as $\Sigma \cap X^{\text{lin}} \neq \emptyset$, guarantees that $\text{IN}_{\Sigma}(\varphi)$ is indeed a (linearly bounded) norm. More precisely:

LEMMA 6.2. — For any subset $\Sigma \subset X^{\mathrm{an}}$ and any bounded function $\varphi \colon \Sigma \to \mathbb{R}$, (6.1) defines a (linearly bounded) norm iff the closure $\overline{\Sigma} \subset X^{\mathrm{an}}$ is non-pluripolar. Further, we then have $\mathrm{T}(\overline{\Sigma}) = \lambda_{\max}(\mathrm{IN}_{\Sigma}(0))$.

Proof. — Since
$$\varphi$$
 is bounded, it is clear that $\operatorname{IN}_{\Sigma}(\varphi)$ is a norm iff
 $\operatorname{T}'(\Sigma) \coloneqq \sup \left\{ m^{-1} \inf_{v \in \Sigma} v(s) \mid s \in R_m \setminus \{0\} \text{ with } m \text{ sufficiently divisible} \right\}$

is finite. By continuity of $v \mapsto v(s)$ for any section s, we have $T'(\Sigma) = T'(\overline{\Sigma})$, and we may thus further assume that Σ is closed. It will then be enough to show that $T(\Sigma) = T'(\Sigma)$ (the case of a single point being [18, Lemma 4.46]). Note that $T'(\Sigma) = \sup_{\varphi}(-\sup_{\Sigma}\varphi)$ where φ runs over L-psh functions of the form $\varphi = m^{-1} \log |s|$ with $s \in R_m \setminus \{0\}$. Since $\sup_{X^{an}} \varphi = \varphi(v_{triv}) = 0$, we infer $T(\Sigma) \ge T'(\Sigma)$. Conversely, pick $\varphi \in PSH$. If $\varphi \in \mathcal{H}_{\mathbb{R}}$, then writing φ as in (1.20) yields $\sup_{X^{an}} \varphi \le \sup_{\Sigma} \varphi + T'(\Sigma)$. In the general case, write φ as the limit of a decreasing net in (φ_i) in $\mathcal{H}_{\mathbb{R}}$. Since $\sup_{X^{an}} \varphi_i = \varphi_i(v_{triv})$ converges to $\sup_{X^{an}} \varphi = \varphi(v_{triv})$, it suffices to show $\sup_{\Sigma} \varphi_i \to \sup_{\Sigma} \varphi$. As X^{an} , and hence Σ , are compact, we can find $v_i \in \Sigma$ such that $\varphi_i(v_i) =$ $\sup_{\Sigma} \varphi_i$, for each i. After passing to a subnet, we may further assume $v_i \to v \in \Sigma$. If $i \le j$ then $\varphi_i(v_j) \ge \varphi_j(v_j) = \sup_{\Sigma} \varphi_j$, and letting $j \to \infty$ yields $\varphi_i(v) \ge \lim_j \sup_{\Sigma} \varphi_j$. Since $\lim_i \varphi_i(v) = \varphi(v)$, we infer $\sup_{\Sigma} \varphi \ge$ $\varphi(v) \ge \lim_j \sup_{\Sigma} \varphi_j$, and the result follows. \Box

Remark 6.3. — Except in the trivial case dim X = 0, we can always find a pluripolar subset $\Sigma \subset X^{\text{an}}$ such that $\overline{\Sigma}$ is non-pluripolar. Indeed, the trivial valuation v_{triv} , which is non-pluripolar, lies in the closure of $X(k) \subset X^{\text{an}}$. By [62], v_{triv} thus lies in the closure of a countable subset $\Sigma \subset X(k)$, which is necessarily pluripolar (see [18, Lemma 4.37]).

For a fixed non-pluripolar subset $\Sigma \subset X^{\mathrm{an}}$, we write

$$\mathcal{N}^{\Sigma}_{\mathbb{R}} \subset \mathcal{N}^{\mathrm{hom}}_{\mathbb{R}}$$

for the set of norms $IN_{\Sigma}(\varphi)$, with φ ranging over bounded functions on Σ .

Example 6.4. — If $\varphi \colon X^{\mathrm{an}} \to \mathbb{R}$ is bounded, then $\mathrm{IN}_{X^{\mathrm{an}}}(\varphi) = \mathrm{IN}(\varphi)$, and Theorem 2.16 thus yields $\mathcal{N}_{\mathbb{R}}^{X^{\mathrm{an}}} = \mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}$.

A simple check shows that

$$\begin{split} \mathrm{IN}_{\Sigma}(\varphi \wedge \varphi') &= \mathrm{IN}_{\Sigma}(\varphi) \wedge \mathrm{IN}_{\Sigma}(\varphi'), \\ \mathrm{IN}_{\Sigma}(\varphi + c) &= \mathrm{IN}_{\Sigma}(\varphi) + c, \\ \mathrm{IN}_{t\Sigma}(t \cdot \varphi) &= t \, \mathrm{IN}_{\Sigma}(\varphi) \end{split}$$

and

(6.2)
$$d_{\infty}(IN_{\Sigma}(\varphi), IN_{\Sigma}(\varphi')) \leq \sup_{\Sigma} |\varphi - \varphi'|$$

for all bounded functions $\varphi, \varphi' \colon \Sigma \to \mathbb{R}, c \in \mathbb{R}$ and $t \in \mathbb{R}_{>0}$. Thus $\mathcal{N}_{\mathbb{R}}^{\Sigma}$ is invariant under the translation action of \mathbb{R} and under minima, and it is invariant under the scaling action of $\mathbb{R}_{>0}$ whenever Σ is.

PROPOSITION 6.5. — Pick a non-pluripolar subset $\Sigma \subset X^{an}$. Then:

- (i) each $\chi \in \mathcal{N}_{\mathbb{R}}^{\Sigma}$ satisfies $\chi = IN_{\Sigma}(\varphi)$ with $\varphi \coloneqq FS(\chi)|_{\Sigma}$, and φ is the smallest bounded function on Σ with this property;
- (ii) if $\Sigma \subset \Sigma' \subset X^{\mathrm{an}}$ then $\mathcal{N}_{\mathbb{R}}^{\Sigma} \subset \mathcal{N}_{\mathbb{R}}^{\Sigma'}$.
- (iii) if Σ is further dense in Σ' , then $IN_{\Sigma}(\varphi) = IN_{\Sigma'}(\varphi)$ for each bounded, usc function $\varphi \colon \Sigma' \to \mathbb{R}$.

Proof. — Pick a bounded function $\psi: \Sigma \to \mathbb{R}$ such that $\chi = \mathrm{IN}_{\Sigma}(\psi)$. For any $v \in \Sigma$ and any $s \in R_m \setminus \{0\}$ we have $\chi(s) \leq m^{-1}v(s) + \psi(v)$. On the one hand, this implies $\varphi(v) = \sup_s \{\chi(s) - m^{-1}v(s)\} \leq \psi(v)$ for any $v \in \Sigma$, and hence $\mathrm{IN}_{\Sigma}(\varphi) \leq \chi$. On the other hand, for any $s \in R_m \setminus \{0\}$ and any $v \in \Sigma$, we have $m^{-1}v(s) + \varphi(v) \geq \chi(s)$, so $\mathrm{IN}_{\Sigma}(\varphi) \geq \chi$. This proves (i).

To see (ii), pick $\chi \in \mathcal{N}_{\mathbb{R}}^{\Sigma}$, i.e. $\chi = \operatorname{IN}_{\Sigma}(\varphi)$ with $\varphi \colon \Sigma \to \mathbb{R}$ bounded. Pick C > 0 such that $\chi(s) \leq mC$ for $s \in R_m \setminus \{0\}$. We claim that χ coincides with $\chi' \coloneqq \operatorname{IN}_{\Sigma}(\varphi') \in \mathcal{N}_{\mathbb{R}}^{\Sigma'}$, where $\varphi' \colon \Sigma' \to \mathbb{R}$ is the extension of φ such that $\varphi' \equiv C$ on $\Sigma' \setminus \Sigma$. To see this, pick $s \in R_m \setminus \{0\}$ For each $v' \in \Sigma' \setminus \Sigma$ we have

$$v'(s) + m\varphi(v') \ge mC \ge \chi(s) = \inf_{v \in \Sigma} \{v(s) + m\varphi(v)\}$$

which yields, as desired, $\chi'(s) = \inf_{v \in \Sigma'} \{v(s) + m\varphi(v)\} = \inf_{v \in \Sigma} \{v(s) + m\varphi(v)\} = \chi(s).$

Finally, the inequality $\operatorname{IN}_{\Sigma}(\varphi) \geq \operatorname{IN}_{\Sigma'}(\varphi)$ in (iii) is trivial. Conversely, pick any $s \in R_m \setminus \{0\}$. Then $m^{-1}v(s) + \varphi(v) \geq \operatorname{IN}_{\Sigma'}(\varphi)(s)$ for all $v \in \Sigma$, and this inequality extends to Σ' as $v \mapsto m^{-1}v(s) + \varphi(v)$ is use on Σ' and $\Sigma \subset \Sigma'$ is dense. Thus $\operatorname{IN}_{\Sigma}(\varphi)(s) \leq \operatorname{IN}_{\Sigma'}(\varphi)(s)$, which proves (iii). \Box

COROLLARY 6.6. — Suppose $\Sigma \subset X^{\mathrm{an}}$ is non-pluripolar. If (χ_i) is a decreasing net in $\mathcal{N}_{\mathbb{R}}^{\Sigma}$ converging pointwise to $\chi \in \mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}$ (see Remark 2.6), then $\chi \in \mathcal{N}_{\mathbb{R}}^{\Sigma}$.

Proof. — Set $\varphi_i := FS(\chi_i)$. Then φ_i is a decreasing net of functions on X^{an} bounded below by $\varphi := FS(\chi)$. For any *i*, Proposition 6.5(i) and Example 6.4 imply

$$\chi_i = \mathrm{IN}_{\Sigma}(\varphi_i) \ge \mathrm{IN}_{\Sigma}(\varphi) \ge \mathrm{IN}(\varphi) = \chi.$$

Taking the infimum over *i* yields $\chi = IN_{\Sigma}(\varphi) \in \mathcal{N}_{\mathbb{R}}^{\Sigma}$.

Next we generalize the homogenization operator.

ANNALES DE L'INSTITUT FOURIER

DEFINITION 6.7. — For any non-pluripolar subset $\Sigma \subset X^{\mathrm{an}}$, we define an operator $P_{\Sigma} \colon \mathcal{N}_{\mathbb{R}} \to \mathcal{N}_{\mathbb{R}}^{\Sigma}$ by setting $P_{\Sigma}(\chi) \coloneqq \mathrm{IN}_{\Sigma}(\mathrm{FS}(\chi))$ for $\chi \in \mathcal{N}_{\mathbb{R}}$.

The map P_{Σ} is a projection, i.e. P_{Σ} is surjective and $P_{\Sigma} \circ P_{\Sigma} = P_{\Sigma}$, by Proposition 6.5(i); it is further 1-Lipschitz with respect to the d_{∞} pseudometric, by (2.10) and (6.2).

The map $P_{X^{an}}: \mathcal{N}_{\mathbb{R}} \to \mathcal{N}_{\mathbb{R}}^{X^{an}} = \mathcal{N}_{\mathbb{R}}^{hom}$ coincides with homogenization (see Theorem 2.16). Further $\Sigma \subset \Sigma' \Longrightarrow P_{\Sigma'}(\chi) \leq P_{\Sigma}(\chi)$. In particular, $\chi \leq \chi^{hom} \leq P_{\Sigma}(\chi)$, and $P_{\Sigma}(\chi)$ can be characterized as the smallest norm in $\mathcal{N}_{\Sigma}^{\Sigma}$ such that $\chi \leq P_{\Sigma}(\chi)$. A direct check further yields:

LEMMA 6.8. — Let (Σ_i) be an increasing net of non-pluripolar subsets of X^{an} , and set $\Sigma := \bigcup_i \Sigma_i$. Then P_{Σ_i} decreases pointwise to P_{Σ} on $\mathcal{N}_{\mathbb{R}}$.

For later use, we also note:

LEMMA 6.9. — For any non-pluripolar subset $\Sigma \subset X^{\mathrm{an}}$ and $\chi \in \mathcal{N}_{\mathbb{R}}$, we have $\mathrm{FS}(\mathrm{P}_{\Sigma}(\chi)) = \mathrm{FS}(\chi)$ on Σ .

Proof. — Set $\varphi := \operatorname{FS}(\chi)$. Since $\chi \leq \operatorname{P}_{\Sigma}(\chi)$, we have $\varphi \leq \operatorname{FS}(\operatorname{P}_{\Sigma}(\chi))$. Conversely, pick $v \in \Sigma$. For any $s \in R_m \setminus \{0\}$, we then have $\operatorname{P}_{\Sigma}(\chi)(s) = \operatorname{IN}_{\Sigma}(\varphi) \leq v(s) + m\varphi(v)$, and hence $\operatorname{FS}(\operatorname{P}_{\Sigma}(v)) = \sup_s \frac{1}{m}(\operatorname{P}_{\Sigma}(\chi)(s) - v(s)) \leq \varphi(v)$, which proves the result.

6.2. Divisorial norms and PL functions

In the next two subsections we consider two important cases of the construction above.

DEFINITION 6.10. — We define the set $\mathcal{N}_{\mathbb{R}}^{\mathrm{div}} \subset \mathcal{N}_{\mathbb{R}}^{\mathrm{hom}}$ of divisorial norms as the (increasing) union of $\mathcal{N}_{\mathbb{R}}^{\Sigma}$ over all finite subsets $\Sigma \subset X^{\mathrm{div}}$. The set of rational divisorial norms is

$$\mathcal{N}^{\mathrm{div}}_{\mathbb{Q}}\coloneqq\mathcal{N}^{\mathrm{div}}_{\mathbb{R}}\cap\mathcal{N}_{\mathbb{Q}}.$$

That the union is increasing follows from Proposition 6.5(ii). Also note that $\mathcal{N}_{\mathbb{R}}^{\text{div}}$ (resp. $\mathcal{N}_{\mathbb{Q}}^{\text{div}}$) is invariant under finite minima, under the scaling action by $\mathbb{Q}_{>0}$ and under the translation action by \mathbb{R} (resp. \mathbb{Q}).

Concretely, a norm χ is divisorial iff it can be written as

(6.3)
$$\chi = \max\{\chi_{v_i} + c_i\}$$

for a finite set of divisorial valuations (v_i) and $c_i \in \mathbb{R}$, and χ is rational iff the c_i can be chosen in \mathbb{Q} . Indeed:
LEMMA 6.11. — For any finite subset $\Sigma \subset X^{\text{div}}$, a norm χ lies in $\mathcal{N}_{\mathbb{Q}}^{\Sigma}$ iff it can be written $\chi = \text{IN}_{\Sigma}(\varphi)$ for some function $\varphi \colon \Sigma \to \mathbb{Q}$.

Proof. — The "if" part is clear. Conversely, assume $\chi \in \mathcal{N}_{\mathbb{Q}}^{\operatorname{div}}$, and write $\chi = \operatorname{IN}_{\Sigma}(\varphi)$ for some function $\varphi \colon \Sigma \to \mathbb{R}$ on a finite subset $\Sigma \subset X^{\operatorname{div}}$. Let $\Sigma' \coloneqq \{v \in \Sigma \mid \varphi(v) \in \mathbb{Q}\}$ and let $\varphi' \colon \Sigma \to \mathbb{Q}$ be any function such that $\varphi' \geqslant \varphi$ with equality on Σ' . Then $\chi' \coloneqq \operatorname{IN}_{\Sigma}(\varphi')$ equals χ . Indeed, $\chi' \geqslant \chi$, and if $s \in R_m \setminus \{0\}$, then $\chi(s) = \min_{v \in \Sigma} (m\varphi(v) - v(s))$. As $\chi(s) \in \mathbb{Q}$ and $v(s) \in \mathbb{Q}$ for every $v \in \Sigma$, the minimum cannot be attained on $\Sigma' \setminus \Sigma$, so $\chi(s) = \min_{v \in \Sigma'} \{m\varphi(v) - v(s)\} = \chi'(s)$.

Recall from [18] that the space $PL(X) \subset C^0(X)$ of piecewise linear functions $\varphi \colon X^{\mathrm{an}} \to \mathbb{R}$ is defined as the \mathbb{Q} -vector space spanned by $\mathcal{H}_{\mathbb{Q}}$. It is independent of the choice of L, stable under max and min, and is dense in $C^0(X)$ with respect to uniform convergence.

As we next show, rational divisorial norms arise precisely as infimum norms of PL functions:

THEOREM 6.12. — A norm $\chi \in \mathcal{N}_{\mathbb{R}}$ lies in $\mathcal{N}_{\mathbb{Q}}^{\text{div}}$ iff $\chi = \text{IN}(\varphi)$ with $\varphi \in \text{PL}(X)$.

COROLLARY 6.13. — Any rational homogeneous norm of finite type is divisorial, i.e. $\mathcal{T}_{\mathbb{Q}}^{\text{hom}} \subset \mathcal{N}_{\mathbb{Q}}^{\text{div}}$. In particular, the homogenization of any test configuration $\chi \in \mathcal{T}_{\mathbb{Z}}$ is a rational divisorial norm.

In contrast, $\mathcal{T}_{\mathbb{R}}^{\text{hom}}$ is generally not contained in $\mathcal{N}_{\mathbb{R}}^{\text{div}}$, see Example B.3. We refer to Appendix A (especially Theorem A.10) for a more detailed discussion of the relation between test configurations and rational divisorial norms.

COROLLARY 6.14. — The envelope property holds for (X, L) iff $\mathcal{N}_{\mathbb{R}}^{\text{div}} \subset \mathcal{N}_{\mathbb{R}}^{\text{cont}}$.

See Section 2.5 for the space $\mathcal{N}_{\mathbb{R}}^{\text{cont}}$ of continuous norms.

Example 6.15. — If X is a nodal curve, then the envelope property fails, and $\chi_v \in \mathcal{N}_{\mathbb{R}}^{\text{div}}$ is indeed not a continuous norm if v is a divisorial valuation with center at the node.

Proof of Theorem 6.12. — Assume first $\chi = IN(\varphi)$ with $\varphi \in PL(X)$. By [18, Lemma 4.26], there exists a finite subset $\Sigma \subset X^{div}$ such that $\sup_{X^{an}}(\psi - \varphi) = \max_{\Sigma}(\psi - \varphi)$ for all $\psi \in PSH(L)$. In particular, for any $s \in R_m \setminus \{0\}$ we have

$$\sup_{X^{\mathrm{an}}} (m^{-1} \log |s| - \varphi) = \max_{\Sigma} (m^{-1} \log |s| - \varphi),$$

i.e. $\chi(s) = \inf_{v \in X^{an}} \{v(s) + m\varphi(v)\} = \min_{v \in \Sigma} \{v(s) + m\varphi(v)\}$. This proves $\operatorname{IN}(\varphi) = \operatorname{IN}_{\Sigma}(\varphi)$, which lies in $\mathcal{N}_{\mathbb{Q}}^{\operatorname{div}}$ since PL functions take rational values on X^{div} .

Conversely, assume $\chi \in \mathcal{N}_{\mathbb{Q}}^{\mathrm{div}}$, i.e. $\chi \in \mathcal{N}_{\mathbb{Q}}^{\Sigma}$ for a finite subset $\Sigma \subset X^{\mathrm{div}}$. By [18, Lemma 2.12], Σ is contained in the set $\Sigma_{\mathfrak{a}}$ of Rees valuations of some flag ideal \mathfrak{a} of X; we may thus assume $\Sigma = \Sigma_{\mathfrak{a}}$, see Proposition 6.5(ii). By Lemma 6.11, we can write $\chi = \mathrm{IN}_{\Sigma}(\widetilde{\varphi})$ for some function $\widetilde{\varphi} \colon \Sigma \to \mathbb{Q}$. By [18, Lemma 2.28], there exists $\rho \in \mathrm{PL}^+(X)$ and $r \gg 1$ such that $\varphi \coloneqq r\varphi_{\mathfrak{a}} - \rho \in \mathrm{PL}(X)$ satisfies $\varphi = \widetilde{\varphi}$ on Σ , while [18, Lemma 2.12] shows that

$$\sup_{X^{\mathrm{an}}}(\psi - \varphi) = \max_{\Sigma}(\psi - \varphi)$$

for all $\psi \in \text{PL}^+(X)$, and hence also for all $\psi \in \text{PSH}(L)$ (as ψ can then be written as a decreasing limit of functions in $\mathcal{H}_{\mathbb{Q}} \subset \text{PL}^+(X)$). As above, this implies $\text{IN}(\varphi) = \chi$, which concludes the proof.

Proof of Corollary 6.13. — Any norm $\chi \in \mathcal{T}_{\mathbb{Q}}^{\text{hom}}$ satisfies $\chi = \text{IN}(\varphi)$ with $\varphi \coloneqq \text{FS}(\chi) \in \mathcal{H}_{\mathbb{Q}} \subset \text{PL}(X)$ (see Theorem 2.16 and Proposition 2.15). By Theorem 6.12, we thus have $\mathcal{T}_{\mathbb{Q}}^{\text{hom}} \subset \mathcal{N}_{\mathbb{Q}}^{\text{div}}$. The last point follows from Lemma 2.11.

Proof of Corollary 6.14. — By Theorem 6.12, $\mathcal{N}_{\mathbb{R}}^{\text{div}}$ is contained in $\mathcal{N}_{\mathbb{R}}^{\text{cont}}$ iff $\text{IN}(\varphi)$ is continuous for any $\varphi \in \text{PL}(X)$, i.e. $\text{FS}(\text{IN}(\varphi)) \in \text{C}^0(X)$ (see Theorem 2.19). Since φ is continuous, we have $\text{FS}(\text{IN}(\varphi)) = \text{Q}(\varphi) = \text{P}(\varphi)$ (see Proposition 2.29). Thus $\mathcal{N}_{\mathbb{R}}^{\text{div}} \subset \mathcal{N}_{\mathbb{R}}^{\text{cont}}$ holds iff $\text{P}(\varphi)$ is continuous for each $\varphi \in \text{PL}(X)$. By density of PL(X) in $\text{C}^0(X)$ and the Lipschitz property of P with respect to the supnorm, this is also equivalent to the continuity of $\text{P}(\varphi)$ for each $\varphi \in \text{C}^0(X)$, which holds in turn iff (X, L) has the envelope property (see [18, Lemma 5.17]).

6.3. Maximal norms and the regularized Fubini-Study operator

Specializing now the definitions of Section 6.1 to the whole set $\Sigma := X^{\text{div}}$, we introduce:

DEFINITION 6.16. — We say that a norm $\chi \in \mathcal{N}_{\mathbb{R}}$ is maximal if it lies in $\mathcal{N}_{\mathbb{R}}^{\max} \coloneqq \mathcal{N}_{\mathbb{R}}^{X^{\operatorname{div}}}$.

Explicitly, a norm is maximal iff it can be written as

$$\chi = \inf_{v \in X^{\mathrm{div}}} \{ \chi_v + c_v \}$$

for a bounded set of constants $(c_v)_{v \in X^{\text{div}}}$. The (slightly abusive) terminology will be justified by Corollary 6.25 below. By Proposition 6.5(ii), we have

$$\mathcal{N}^{\mathrm{div}}_{\mathbb{R}} \subset \mathcal{N}^{\mathrm{max}}_{\mathbb{R}} \subset \mathcal{N}^{\mathrm{hom}}_{\mathbb{R}},$$

both inclusions being strict (except in the trivial case dim X = 0), cf. Example 6.23 below.

For each $\chi \in \mathcal{N}_{\mathbb{R}}$ we set

$$\chi^{\max} \coloneqq \mathcal{P}_{X^{\operatorname{div}}}(\chi) = \operatorname{IN}_{X^{\operatorname{div}}}(\operatorname{FS}(\chi)).$$

Then $\chi \leq \chi^{\text{hom}} \leq \chi^{\text{max}}$, and χ^{max} is the smallest norm in $\mathcal{N}_{\mathbb{R}}^{\text{max}}$ such that $\chi \leq \chi^{\text{max}}$.

Before going further, recall from Section 2.3 that the Fubini–Study operator associates to any norm $\chi \in \mathcal{N}_{\mathbb{R}}$ with canonical approximants $\chi_d \in \mathcal{T}_{\mathbb{R}}$ a bounded, lsc function $FS(\chi): X^{an} \to \mathbb{R}$, such that $FS(\chi) = \sup_d FS(\chi_d)$ with $FS(\chi_d) \in \mathcal{H}_{\mathbb{R}}$. We denote by $FS^*(\chi) := FS(\chi)^*$ its usc regularization, which is thus a bounded usc function on X^{an} . The next result will be instrumental for what follows:

LEMMA 6.17. — For any norm $\chi \in \mathcal{N}_{\mathbb{R}}$, the following holds:

- (i) $FS^{\star}(\chi) = FS(\chi)$ on X^{div} ;
- (ii) if (X, L) has the weak envelope property, then $FS^{\star}(\chi) = FS(\chi)$ on X^{lin} , and $\widetilde{E}(FS^{\star}(\chi)) = \widetilde{E}(FS(\chi));$
- (iii) if $FS^*(\chi)$ is L-psh (e.g., if (X, L) has the envelope property), then $FS^*(\chi)$ lies in $\mathcal{E}^{\infty}_{\uparrow} \subset \mathcal{E}^1$, and $FS(\chi_d) \to FS^*(\chi)$ strongly in \mathcal{E}^1 .

We refer to Section 4.4 for the (weak) envelope property and the space $\mathcal{E}^{\infty}_{\uparrow}$ of psh functions approximable from below. Recall that the weak envelope property holds as soon as char k = 0, and that the envelope property then holds if X is further smooth.

Proof. — Since $FS(\chi) = \sup_d FS(\chi_d)$ with $FS(\chi_d)$ *L*-psh, (i) and (ii) respectively follow from Lemmas 4.2 and 4.8 (see Section 4.4). If (X, L) has the envelope property, then $FS^*(\chi)$ is *L*-psh, and the rest of (iii) follows from Lemma 4.3.

PROPOSITION 6.18. — For any $\chi \in \mathcal{N}_{\mathbb{R}}$, we have

- (i) $\chi^{\max} = IN(FS^{\star}(\chi));$
- (ii) χ is maximal iff $\chi = IN(\varphi)$ for some bounded usc function φ on X^{an} .

COROLLARY 6.19. — The space $\mathcal{N}_{\mathbb{R}}^{\max}$ is invariant under the scaling action by $\mathbb{R}_{>0}$, the translation action by \mathbb{R} , under finite minima, and under decreasing limits. Further, any $\chi \in \mathcal{N}_{\mathbb{R}}^{\max}$ can be written as the pointwise limit of a decreasing net in $\mathcal{N}_{\mathbb{R}}^{\text{div}}$.

COROLLARY 6.20. — Each continuous norm $\chi \in \mathcal{N}_{\mathbb{R}}^{\text{cont}}$ satisfies $\chi^{\text{hom}} = \chi^{\text{max}}$. In particular, every continuous homogeneous norm is maximal, i.e. $\mathcal{N}_{\mathbb{R}}^{\text{cont,hom}} \subset \mathcal{N}_{\mathbb{R}}^{\text{max}}$.

The first equality fails in general when χ is not continuous (see Example 6.23), and the last inclusion is strict in general (see Corollary 6.30 below).

COROLLARY 6.21. — If (X, L) has the weak envelope property, then $\mathcal{N}_{\mathbb{R}}^{\max} = \mathcal{N}_{\mathbb{R}}^{X^{\lim}}$; in particular, χ_v is then maximal for any $v \in X^{\lim}$.

Proof of Proposition 6.18. — Lemma 6.17(i) implies that $\chi^{\max} = P_{X^{\dim}}(FS(\chi))$ coincides with $IN_{X^{\dim}}(FS^*(\chi))$, which is also equal to $IN(FS^*(\chi))$, since $FS^*(\chi)$ is use on X^{an} and X^{div} is dense (see Proposition 6.5(iii)). This proves (i).

To see (ii), assume χ is maximal. By (i), we then have $\chi = \chi^{\max} = IN(FS^{\star}(\chi))$ where $FS^{\star}(\chi)$ is bounded and usc. Conversely, assume $\chi = IN(\varphi)$ with φ bounded and usc on X^{an} . By density of X^{div} , Proposition 6.5(iii) yields $\chi = IN_{X^{\operatorname{div}}}(\varphi)$, and hence $\chi \in \mathcal{N}_{\mathbb{R}}^{\max}$. This proves (ii). \Box

Proof of Corollary 6.19. — For any non-pluripolar $\Sigma \subset X^{\mathrm{an}}$, the space $\mathcal{N}_{\mathbb{R}}^{\Sigma}$ is invariant under the translation action by \mathbb{R} , under finite minima, and under decreasing limits, see Corollary 6.6. By Proposition 6.18(ii) and (2.12), $\mathcal{N}_{\mathbb{R}}^{\mathrm{max}}$ is further invariant under the scaling action of $\mathbb{R}_{>0}$ (even though X^{div} is only invariant under the scaling action of $\mathbb{Q}_{>0}$). The final assertion is an immediate consequence of Corollary 6.6 and Lemma 6.8.

Proof of Corollary 6.20. — If $\chi \in \mathcal{N}_{\mathbb{R}}^{\text{cont}}$ then $FS(\chi)$ is continuous (see Theorem 2.19), and hence $\chi^{\text{hom}} = IN(FS(\chi)) = \chi^{\text{max}}$, by Theorem 2.16 and Proposition 6.18(i).

Proof of Corollary 6.21. — Lemma 6.17(ii) implies that $P_{X^{\text{lin}}}(\chi) = IN_{X^{\text{lin}}}(FS(\chi))$ coincides with $IN_{X^{\text{lin}}}(FS^*(\chi))$. By Proposition 6.5(iii), this is also equal to $IN(FS^*(\chi))$, which is in turn equal to χ^{\max} , by Proposition 6.18(i). Thus $P_{X^{\text{lin}}}(\chi) = \chi^{\max}$, and hence $\chi \in \mathcal{N}_{\mathbb{R}}^{X^{\text{lin}}} \Leftrightarrow \chi \in \mathcal{N}_{\mathbb{R}}^{\max}$. \Box

We can now state the main result of this section:

THEOREM 6.22. — For all norms $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, we have $\chi \sim \chi' \Longrightarrow \chi^{\max} = \chi'^{\max}$. If (X, L) has the weak envelope property (e.g., if char k = 0), the converse implication holds.

Example 6.23. — For any subvariety $Z \subsetneq X$, the norm $\chi = \chi_Z \in \mathcal{N}_{\mathbb{Z}}^{\text{hom}}$ of Example 2.21 is not maximal. Indeed, χ is asymptotically equivalent to the maximal norm $\chi_{\text{triv}} + 1$ (see Example 3.10), and hence $\chi^{\text{max}} = \chi_{\text{triv}} + 1 \neq \chi$.

COROLLARY 6.24. — The restriction of d_1 to $\mathcal{N}_{\mathbb{R}}^{\max}$ a metric. If (X, L) has the weak envelope property, then $\mathcal{N}_{\mathbb{R}}^{\max}$ is further maximal for this property.

COROLLARY 6.25. — Assume (X, L) has the weak envelope property, and pick any norm $\chi \in \mathcal{N}_{\mathbb{R}}$. Then χ is maximal iff it is the largest norm in its asymptotic equivalence class.

COROLLARY 6.26. — If $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}^{\text{cont}}$ are continuous, then $\chi \sim \chi' \iff d_{\infty}(\chi, \chi') = 0.$

As a first step towards Theorem 6.22, we show:

LEMMA 6.27. — For all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$, the following are equivalent:

- (i) $\chi^{\max} = \chi'^{\max};$
- (ii) $FS(\chi) = FS(\chi')$ on X^{div} ;
- (iii) $FS^{\star}(\chi) = FS^{\star}(\chi')$ on X^{an} .

Proof. — Since $\chi^{\text{max}} = P_{X^{\text{div}}}(\chi)$, Lemma 6.9 yields $\text{FS}(\chi^{\text{max}}) = \text{FS}(\chi)$ on X^{div} , and similarly for χ' . This implies (i) ⇒ (ii), while Lemma 6.17(i) yields (ii) ⇔ (iii). Finally, (iii) ⇒ (i) follows from Proposition 6.18(i). □

LEMMA 6.28. — For all
$$\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$$
, we have
 $\chi \sim \chi' \iff \lim_{d} d_1(\mathrm{FS}(\chi_d), \mathrm{FS}(\chi'_d)) = 0 \Longrightarrow \mathrm{FS}(\chi) = \mathrm{FS}(\chi') \text{ on } X^{\mathrm{lin}}.$

Proof. — By construction of the d_1 -metric on $\mathcal{H}_{\mathbb{R}}$, $FS: (\mathcal{T}_{\mathbb{R}}, d_1) \to (\mathcal{H}_{\mathbb{R}}, d_1)$ is an isometry, and hence $d_1(\chi_d, \chi'_d) = d_1(FS(\chi_d), FS(\chi'_d))$ for all d sufficiently divisible. By Theorem 3.18, we have, on the other hand, $d_1(\chi_d, \chi'_d) \to d_1(\chi, \chi')$. This implies the first equivalence. For any $v \in X^{\text{lin}}$, the measure δ_v lies in \mathcal{M}^1 . By (5.19), we thus have

$$d_1(FS(\chi_d), FS(\chi'_d)) \longrightarrow 0 \Longrightarrow FS(\chi_d)(v) - FS(\chi'_d)(v) \longrightarrow 0$$

which yields the right-hand implication, since $FS(\chi_d) \to FS(\chi)$ and $FS(\chi'_d) \to FS(\chi')$ pointwise on X^{an} .

Proof of Theorem 6.22. — If $\chi \sim \chi'$, then $FS(\chi) = FS(\chi')$ on $X^{\text{lin}} \supset X^{\text{div}}$, by Lemma 6.28, and hence $\chi^{\text{max}} = \chi'^{\text{max}}$, by Lemma 6.27. Now assume the weak envelope property. To prove the converse implication, it suffices to show $\chi \sim \chi^{\text{max}}$ for any $\chi \in \mathcal{N}_{\mathbb{R}}$. Since $\chi \leq \chi^{\text{max}}$, this amounts to

 $\operatorname{vol}(\chi) = \operatorname{vol}(\chi^{\max})$ (see Lemma 3.11). By Theorem 5.1, we have $\operatorname{vol}(\chi) = \widetilde{E}(\varphi)$ with $\varphi := \operatorname{FS}(\chi)$. On the other hand, we have $\chi^{\max} = \operatorname{IN}(\varphi^*)$ (see Proposition 6.18(i)), and hence $\operatorname{vol}(\chi^{\max}) = \widetilde{E}((\varphi^*)_*)$, by Corollary 5.3. Since φ is lsc, we have $\varphi \leq (\varphi^*)_* \leq \varphi^*$, so by monotonicity of the energy, it suffices to prove that $\widetilde{E}(\varphi) = \widetilde{E}(\varphi^*)$, which follows from Lemma 6.17. \Box

Proof of Corollary 6.24. — That d_1 restricts to a metric on $\mathcal{N}_{\mathbb{R}}^{\max}$ is a direct consequence of Theorem 6.22. If d_1 is also a metric on a subset $\mathcal{N}' \subset \mathcal{N}_{\mathbb{R}}$ that contains $\mathcal{N}_{\mathbb{R}}^{\max}$, then any $\chi \in \mathcal{N}'$ satisfies $d_1(\chi, \chi^{\max}) = 0$, by Theorem 6.22, and hence $\chi = \chi^{\max} \in \mathcal{N}_{\mathbb{R}}^{\max}$. Thus $\mathcal{N}' = \mathcal{N}_{\mathbb{R}}^{\max}$.

Proof of Corollary 6.25. — Assume χ is maximal, and pick $\chi' \in \mathcal{N}_{\mathbb{R}}$ with $\chi \sim \chi'$. Then $\chi' \leq \chi'^{\max} = \chi^{\max} = \chi$, by Theorem 6.22, which proves that χ is the largest norm in its equivalence class. Conversely, this last property implies $\chi^{\max} \leq \chi$, since $\chi \sim \chi^{\max}$ by Theorem 6.22, and hence $\chi = \chi^{\max}$, i.e. $\chi \in \mathcal{N}_{\mathbb{R}}^{\max}$.

Proof of Corollary 6.26. — By Theorem 2.19, $FS(\chi)$ and $FS(\chi')$ are continuous, and are equal iff $d_{\infty}(\chi, \chi') = 0$. By Lemma 6.27, we thus have $\chi \sim \chi' \Rightarrow d_{\infty}(\chi, \chi') = 0$, while the converse trivially holds.

Assuming now the envelope property (e.g. X is smooth and char k = 0), we finally state:

THEOREM 6.29. — If (X, L) has the envelope property, then the regularized Fubini–Study operator defines a surjective isometry $\mathrm{FS}^* \colon (\mathcal{N}_{\mathbb{R}}, \mathrm{d}_1) \twoheadrightarrow$ $(\mathcal{E}^{\infty}_{\uparrow}, \mathrm{d}_1)$, which restricts to an isometric isomorphism $\mathrm{FS}^* \colon (\mathcal{N}^{\max}_{\mathbb{R}}, \mathrm{d}_1) \xrightarrow{\sim} (\mathcal{E}^{\infty}_{\uparrow}, \mathrm{d}_1)$ with inverse IN: $(\mathcal{E}^{\infty}_{\uparrow}, \mathrm{d}_1) \xrightarrow{\sim} (\mathcal{N}^{\max}_{\mathbb{R}}, \mathrm{d}_1)$.

Proof. — Since (X, L) has the envelope property, FS^{*}(χ) lies in $\mathcal{E}^{\infty}_{\uparrow}$ for each $\chi \in \mathcal{N}_{\mathbb{R}}$ (see Lemma 6.17(iii)). Since FS: ($\mathcal{T}_{\mathbb{R}}$, d₁) → ($\mathcal{H}_{\mathbb{R}}$, d₁) is an isometry, the canonical approximants $\chi_d, \chi'_d \in \mathcal{T}_{\mathbb{R}}$ satisfy d₁(χ_d, χ'_d) = d₁(FS(χ_d), FS(χ'_d)) for all *d* sufficiently divisible. Now, Theorem 3.18 implies on the one hand d₁(χ_d, χ'_d) → d₁(χ, χ'). On the other hand, Lemma 6.17(iii) implies d₁(FS(χ_d), FS(χ'_d)) → d₁(FS^{*}(χ), FS^{*}(χ')), since d₁ defines the strong topology of \mathcal{E}^1 (see Theorem 5.5). This proves that FS^{*}: ($\mathcal{N}_{\mathbb{R}}$, d₁) → ($\mathcal{E}^{\infty}_{\uparrow}$, d₁) is an isometry, whose restriction to $\mathcal{N}^{\max}_{\mathbb{R}}$ is necessarily injective, by Theorem 6.22. Conversely, pick $\varphi \in \mathcal{E}^{\infty}_{\uparrow}$. Then IN(φ) ∈ $\mathcal{N}^{\max}_{\mathbb{R}}$ (see Proposition 6.18(ii)). By Proposition 2.29 and Lemma 4.6(i), we further have FS^{*}(IN(φ)) = Q^{*}(φ) = φ . This shows that FS^{*}: ($\mathcal{N}^{\max}_{\mathbb{R}}$, d₁) $\xrightarrow{\rightarrow}$ ($\mathcal{E}^{\infty}_{\uparrow}$, d₁) is an isometric isomorphism, and the rest follows.

COROLLARY 6.30. — Assume (X, L) has the envelope property. Then

$$\mathcal{N}_{\mathbb{R}}^{\max} \subset \mathcal{N}_{\mathbb{R}}^{\text{cont}} \Longleftrightarrow \mathcal{E}^{\infty}_{\uparrow} = \text{CPSH} \Longleftrightarrow \dim X \leqslant 1.$$

Proof. — The first equivalence follows from Theorem 2.26 and Theorem 6.29, and the second one from [18, Example 13.25]).

7. The Monge–Ampère measure of a norm

Any ample test configuration for (X, L) defines a measure on X^{an} with finite support in X^{div} . This defines a Monge–Ampère operator MA: $\mathcal{T}_{\mathbb{Z}} \to \mathcal{M}^1$. Here we extend this operator to a map $\mathcal{N}_{\mathbb{R}} \to \mathcal{M}^1$ whose fibers consist of asymptotic equivalence classes modulo translation, thus completing the proof of Theorem B. The construction, which works even when in the absence of the envelope property, restricts to a homeomorphism between divisorial norms modulo translations and probability measures with finite support in X^{div} , thus proving Theorem C. We also extend Dervan's minimum norm functional from $\mathcal{T}_{\mathbb{Z}}$ to $\mathcal{N}_{\mathbb{R}}$.

7.1. Monge–Ampère measures of R-test configurations

We define the Monge–Ampère measure $MA(\chi) \in \mathcal{M}^1$ of an \mathbb{R} -test configuration $\chi \in \mathcal{T}_{\mathbb{R}}$ as the Monge–Ampère measure of the associated Fubini– Study function $FS(\chi) \in \mathcal{H}_{\mathbb{R}}$ (see Proposition 2.15), i.e.

$$MA(\chi) := MA(FS(\chi)).$$

The invariance/equivariance properties of the operators MA: $\mathcal{H}_{\mathbb{R}} \to \mathcal{M}^1$ and FS: $\mathcal{T}_{\mathbb{R}} \to \mathcal{H}_{\mathbb{R}}$ imply that if $\chi \in \mathcal{T}_{\mathbb{R}}$, $c \in \mathbb{R}$ and $t \in \mathbb{R}_{>0}$, then

$$MA(\chi + c) = MA(\chi)$$
 and $MA(t\chi) = t_{\star} MA(\chi)$.

When $\chi \in \mathcal{T}_{\mathbb{Z}}$, the Monge–Ampère measure can be computed geometrically as follows. By the Rees correspondence (A.7), χ is associated to an ample test configuration. Let $(\mathcal{X}, \mathcal{L})$ be its integral closure, with central fiber $\mathcal{X}_0 = \sum_i b_i E_i$. By Lemma A.12, $\varphi := FS(\chi) \in \mathcal{H}_{\mathbb{Q}}$ satisfies (A.5), so by [18, Proposition 7.19(ii)] we have

(7.1)
$$\operatorname{MA}(\chi) = \sum_{i} b_i \left(\mathcal{L}|_{E_i}^n \right) \delta_{v_i},$$

where $v_i \in X^{\text{div}}$ is the divisorial valuation associated to E_i .

For general $\chi \in \mathcal{T}_{\mathbb{R}}$, the support of MA(χ) is a finite subset of X^{lin} , see Lemma 4.1.

ANNALES DE L'INSTITUT FOURIER

LEMMA 7.1. — The Monge–Ampère operator above defines an isometry (7.2) $MA: (\mathcal{T}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1) \longrightarrow (\mathcal{M}^1, d_1)$

with dense image.

Proof. — By (5.11), the Fubini–Study operator defines an isometric surjection FS: $(\mathcal{T}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1) \rightarrow (\mathcal{H}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1)$. Now $\mathcal{H}_{\mathbb{R}}/\mathbb{R}$ is a dense subspace of $(\mathcal{E}^1/\mathbb{R}, \underline{d}_1)$, and by definition, the metric d_1 on \mathcal{M}^1 has the property that MA: $(\mathcal{E}^1/\mathbb{R}, \underline{d}_1) \rightarrow (\mathcal{M}^1, d_1)$ is an injective isometry with dense image, see Theorem 5.11. It therefore follows that (7.2) is an isometry with dense image.

7.2. Monge–Ampère measures of general norms

We now define the Monge–Ampère operator on general norms.

THEOREM 7.2. — The Monge–Ampère operator above extends uniquely to an isometry

(7.3)
$$\operatorname{MA}: (\mathcal{N}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1) \longrightarrow (\mathcal{M}^1, d_1),$$

with dense image.

As $(\mathcal{M}^1, \mathbf{d}_1)$ is complete, the Monge–Ampère operator thus realizes $(\mathcal{M}^1, \mathbf{d}_1)$ as the Hausdorff completion of the pseudo-metric spaces $(\mathcal{T}_{\mathbb{R}}/\mathbb{R}, \underline{\mathbf{d}}_1)$ and $(\mathcal{N}_{\mathbb{R}}/\mathbb{R}, \underline{\mathbf{d}}_1)$.

Proof. — By Theorem 3.18, $\mathcal{T}_{\mathbb{R}}$ is dense in $(\mathcal{N}_{\mathbb{R}}, d_1)$. As a consequence, $\mathcal{T}_{\mathbb{R}}/\mathbb{R}$ sits as a dense subspace of $(\mathcal{N}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1)$, and we conclude using Lemma 7.1.

Combining Theorem 7.2 with Lemma 3.15, we get:

COROLLARY 7.3. — The induced map MA: $(\mathcal{N}_{\mathbb{R}}, d_1) \rightarrow (\mathcal{M}^1, d_1)$ is 1-Lipschitz, and its nonempty fibers consist precisely of asymptotic equivalence classes of norms modulo translation.

The induced map MA: $\mathcal{N}_{\mathbb{R}} \to \mathcal{M}^1$ satisfies the following properties, the first of which gives a more concrete description.

PROPOSITION 7.4. — For any norm $\chi \in \mathcal{N}_{\mathbb{R}}$ we have:

- (i) the canonical approximants (χ_d) satisfy $\lim_d MA(\chi_d) = MA(\chi)$ strongly in \mathcal{M}^1 ;
- (ii) for $c \in \mathbb{R}$ and $t \in \mathbb{R}_{>0}$, we have $MA(\chi + c) = MA(\chi)$ and $MA(t\chi) = t_{\star} MA(\chi)$;

- (iii) $MA(\chi) = MA(\chi^{hom})$, where χ^{hom} is the homogenization of χ ;
- (iv) If (X, L) has the weak envelope property, then $MA(\chi) = MA(\chi^{max})$.
- (v) if $FS^{*}(\chi)$ is L-psh, then $MA(\chi) = MA(FS^{*}(\chi))$.

Recall that (v) applies if (X, L) has the envelope property, or for any continuous norm $\chi \in \mathcal{N}_{\mathbb{R}}^{\text{cont}}$.

Proof. — Theorem 3.18 shows that $\chi_d \in \mathcal{T}_{\mathbb{R}}$ satisfy $\lim_d d_1(\chi_d, \chi) = 0$, which implies (i). The equalities in (ii) follow, since $(\chi + c)_d = \chi_d + c$ and $(t\chi)_d = t\chi_d$, whereas (iii) and (iv) follow since $\chi^{\text{hom}} \sim \chi$ and $\chi^{\text{max}} \sim \chi$, respectively, see Corollary 5.2 and Theorem 6.22. If $\text{FS}^*(\chi)$ is *L*-psh, then $\text{FS}(\chi_d) \to \text{FS}^*(\chi)$ strongly in \mathcal{E}^1 (see Lemma 6.17), and hence $\text{MA}(\chi_d) =$ $\text{MA}(\text{FS}(\chi_d)) \to \text{MA}(\text{FS}^*(\chi))$ in \mathcal{M}^1 . This proves (v), in view of (i).

Recall the space $\mathcal{N}_{\mathbb{R}}^{\max} = \mathcal{N}_{\mathbb{R}}^{X^{\operatorname{div}}}$ from Section 6.3. By Corollary 6.24, the pseudometric d_1 restricts to a metric on $\mathcal{N}_{\mathbb{R}}^{\max}$. Lemma 5.10 thus implies that \underline{d}_1 restricts to a metric on $\mathcal{N}_{\mathbb{R}}^{\max}/\mathbb{R}$.

COROLLARY 7.5. — The Monge–Ampère operator MA: $\mathcal{N}_{\mathbb{R}} \to \mathcal{M}^1$ induces an isometric embedding

(7.4)
$$MA: \left(\mathcal{N}_{\mathbb{R}}^{\max}/\mathbb{R}, \underline{d}_{1}\right) \hookrightarrow \left(\mathcal{M}^{1}, d_{1}\right)$$

with dense image. If (X, L) has the weak envelope property, then the image equals $MA(\mathcal{N}_{\mathbb{R}})$.

Recall that the weak envelope property holds when char k = 0 or dim $X \leq 2$.

Proof. — Everything except for the last statement is clear by what precedes, and that statement is an immediate consequence of Theorem 6.22.

Remark 7.6. — Even if X is smooth (and of positive dimension) and char k = 0, MA($\mathcal{N}_{\mathbb{R}}$) is a strict subspace of \mathcal{M}^1 , which is not so easy to describe. See [21] for related questions.

For later use, we also show the following version of Theorem 5.8.

LEMMA 7.7. — For all $\chi, \chi' \in \mathcal{N}_{\mathbb{R}}$ we have

$$d_1(\chi, \chi') \approx \int |FS(\chi) - FS(\chi')| (MA(\chi) + MA(\chi')).$$

Proof. — Set $\varphi_d \coloneqq FS(\chi_d), \, \varphi'_d \coloneqq FS(\chi'_d)$ and

$$\mu_d \coloneqq \operatorname{MA}(\varphi_d) = \operatorname{MA}(\chi_d), \quad \mu'_d \coloneqq \operatorname{MA}(\varphi'_d) = \operatorname{MA}(\chi'_d),$$

with $(\chi_d), (\chi'_d)$ the canonical approximants of χ, χ' . Since FS: $(\mathcal{T}_{\mathbb{R}}, d_1) \to$ $(\mathcal{H}_{\mathbb{R}}, d_1)$ is an isometry, Theorem 5.8 yields

$$\mathbf{d}_1(\chi_d, \chi_d') = \mathbf{d}_1(\varphi_d, \varphi_d') \approx \int g_d \left(\mu_d + \mu_d'\right),$$

with $g_d := |\varphi_d - \varphi'_d|$. Since (φ_d) , (φ'_d) are uniformly bounded and $\mu_d \to$ $\mu := \mathrm{MA}(\chi), \, \mu'_d \to \mu' := \mathrm{MA}(\chi') \text{ strongly in } \mathcal{M}^1 \text{ (see Proposition 7.4(i))},$ (5.18) yields $\int g_d (\mu_d + \mu'_d) = \int g_d (\mu + \mu') + o(1)$. Since (g_d) is uniformly bounded and converges pointwise to $g := |FS(\chi) - FS(\chi')|$, dominated convergence applied to the cofinal sequence $(g_{m!})_m$ further yields $\int g_{m!}(\mu +$ $\mu') \rightarrow \int g(\mu + \mu')$. Combining this with Theorem 3.18, we conclude

$$d_1(\chi,\chi') = \lim_m d_1(\chi_{m!},\chi'_{m!}) \approx \lim_m \int g_{m!} (\mu + \mu') = \int g (\mu + \mu'),$$

h proves the result.

which proves the result.

7.3. Variational principle

As we next show, the Monge–Ampère equation $MA(\chi) = \mu$ with $\chi \in \mathcal{N}_{\mathbb{R}}$ and $\mu \in \mathcal{M}^1$ admits a variational characterization, that will be deduced from its counterpart for L-psh functions.

PROPOSITION 7.8. — For any $\mu \in \mathcal{M}^1$, we have

$$\mathrm{E}^{\vee}(\mu) = \sup_{\chi \in \mathcal{N}_{\mathbb{R}}} \left(\mathrm{vol}(\chi) - \int \mathrm{FS}(\chi) \, \mu \right).$$

Further, the supremum is achieved by $\chi \in \mathcal{N}_{\mathbb{R}}$ iff $MA(\chi) = \mu$.

Proof. — We have $E^{\vee}(\mu) = \sup_{\varphi \in \mathcal{H}_{\mathbb{Q}}} (E(\varphi) - \int \varphi \mu)$, and any $\varphi \in \mathcal{H}_{\mathbb{Q}}$ can be written as $\varphi = FS(\chi)$ with $\chi := IN(\varphi) \in \mathcal{N}_{\mathbb{R}}$. Since $E(\varphi) = vol(\chi)$, this yields

$$\mathrm{E}^{\vee}(\mu) \leqslant \sup_{\chi \in \mathcal{N}_{\mathbb{R}}} \left(\mathrm{vol}(\chi) - \int \mathrm{FS}(\chi) \, \mu \right).$$

For the reverse inequality, pick any $\chi \in \mathcal{N}_{\mathbb{R}}$, and consider the increasing net (φ_d) in $\mathcal{H}_{\mathbb{R}}$ defined by $\varphi_d \coloneqq FS(\chi_d)$, with (χ_d) the canonical approximants of χ . Then $E(\varphi_d) - \int \varphi_d \mu \leq E^{\vee}(\mu)$. By Theorem 3.18 and Theorem 5.1, $E(\varphi_d) = vol(\chi_d) \to vol(\chi)$, while $\int \varphi_d \mu \to \int FS(\chi) \mu$, by monotone convergence (applied to the cofinal sequence $\varphi_{d!}$). Thus $\operatorname{vol}(\chi) - \int \operatorname{FS}(\chi) \mu \leq$ $E^{\vee}(\mu)$, and equality holds iff $E(\varphi_d) - \int \varphi_d \mu \to E^{\vee}(\mu)$, i.e. (φ_d) is a maximizing net for μ . By [18, Corollary 10.13], the latter is also equivalent to $MA(\varphi_d) \to \mu$ strongly in \mathcal{M}^1 , and hence to $MA(\chi) = \mu$, since $MA(\varphi_d) =$ $MA(\chi_d) \to MA(\chi)$ strongly in \mathcal{M}^1 , by Corollary 7.3.

TOME 75 (2025), FASCICULE 2

Remark 7.9. — With a little bit of extra effort, one can show as in [18, Corollary 10.13] that a net (χ_i) in $\mathcal{N}_{\mathbb{R}}$ computes the supremum, that is $\lim_i (\operatorname{vol}(\chi_i) - \int \operatorname{FS}(\chi_i) \mu) = \operatorname{E}^{\vee}(\mu)$, iff $\operatorname{MA}(\chi_i) \to \mu$ strongly in \mathcal{M}^1 .

7.4. Divisorial norms and divisorial measures

The image $MA(\mathcal{N}_{\mathbb{R}})$ of the Monge–Ampère operator is a strict subset of \mathcal{M}^1 and not so easy to describe, but we now exhibit an important class of measures contained in the image.

Given any compact subset $\Sigma \subset X^{\mathrm{an}}$, denote by \mathcal{M}^{Σ} the set of Radon probability measures μ on X^{an} with support in Σ .

Example 7.10. — When $\Sigma \subset X^{\mathrm{an}}$ is finite, each $\mu \in \mathcal{M}^{\Sigma}$ is of the form $\mu = \sum_{v \in \Sigma} m_v \delta_v$, where $m_v := \mu(\{v\})$, and it is easy to see that $\mu \mapsto m = (m_v)$ defines a homeomorphism of \mathcal{M}^{Σ} (equipped with the weak topology) with the simplex $\{m \in \mathbb{R}_{\geq 0}^{\Sigma} \mid \sum_v m_v = 1\}$.

Recall that the strong topology of X^{lin} is defined by the metric d_{∞} (see (1.19)); the weak topology refers to the subset topology from X^{an} . For all $v, w \in X^{\text{lin}}$ we have

(7.5)
$$d_{\infty}(v,w) = \sup_{\varphi \in \mathrm{PSH}} |\varphi(v) - \varphi(w)|,$$

which shows that d_{∞} is the smallest metric on X^{lin} such that the restriction to X^{lin} of any *L*-psh function is 1-Lipschitz. By (7.5), the weak and strong topologies coincide on a given subset $\Sigma \subset X^{\text{lin}}$ iff PSH $|_{\Sigma}$ is equicontinuous for the weak topology of Σ . This is in particular the case when Σ is strongly compact (as the identity map (Σ , strong) \rightarrow (Σ , weak) is then a homeomorphism, being continuous and bijective on a compact Hausdorff space).

Example 7.11. — Every finite subset $\Sigma \subset X^{\text{lin}}$ is of course strongly compact. If X is smooth and char k = 0, then the dual complex $\Delta_{\mathcal{X}}$ of any snc test configurations \mathcal{X} also forms a strongly compact subset of X^{lin} , cf. [18, Theorem A.4].

LEMMA 7.12. — For any strongly compact subset $\Sigma \subset X^{\text{lin}}$, we have $\mathcal{M}^{\Sigma} \subset \mathcal{M}^{1}$, and the induced weak and strong topologies on \mathcal{M}^{Σ} coincide.

Proof. — Since Σ is strongly compact, $C \coloneqq \sup_{v \in \Sigma} T(v)$ is finite, and satisfies $\sup \varphi - \varphi(v) \leq C$ for each $\varphi \in PSH$ and $v \in \Sigma$. For each $\mu \in \mathcal{M}^{\Sigma}$

we thus have

$$\mathbf{E}^{\vee}(\mu) = \sup_{\varphi \in \mathcal{E}^1} \left\{ \mathbf{E}(\varphi) - \int \varphi \, \mu \right\} \leqslant \sup_{\varphi \in \mathcal{E}^1} \left\{ \sup \varphi - \int \varphi \, \mu \right\} \leqslant C,$$

and hence $\mu \in \mathcal{M}^1$. Now pick a weakly convergent net $\mu_i \to \mu$ in \mathcal{M}^{Σ} . Since $\mathrm{PSH}|_{\Sigma}$ is equicontinuous, we have $\int \varphi \, \mu_i \to \int \varphi \, \mu$ uniformly for $\varphi \in \mathrm{PSH}$. Thus

$$\mathbf{E}^{\vee}(\mu_i) = \sup_{\varphi \in \mathcal{E}^1} \left\{ \mathbf{E}(\varphi) - \int \varphi \, \mu \right\} \longrightarrow \sup_{\varphi \in \mathcal{E}^1} \left\{ \mathbf{E}(\varphi) - \int \varphi \, \mu \right\} = \mathbf{E}^{\vee}(\mu)$$

and hence $\mu_i \to \mu$ strongly in \mathcal{M}^1 .

THEOREM 7.13. — For any strongly compact subset $\Sigma \subset X^{\text{lin}}$, the Monge–Ampère operator induces a surjective isometry

MA:
$$(\mathcal{N}^{\Sigma}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1) \longrightarrow (\mathcal{M}^{\Sigma}, d_1).$$

If $\Sigma \subset X^{\text{div}}$, or if the weak envelope property holds (e.g., if char k = 0), then this map is an isometric isomorphism.

Recall that $\mathcal{N}_{\mathbb{R}}^{\Sigma}$ denotes the set of norms of the form $\chi = IN_{\Sigma}(\varphi)$ for a bounded function $\varphi \colon \Sigma \to \mathbb{R}$, see Section 6.1. We emphasize that Theorem 7.13 is true for an *arbitrary* polarized variety, whether or not the envelope property holds. The following important special case illustrates this.

Example 7.14. — For each $v \in X^{\text{lin}}$ we have $\operatorname{MA}(\chi_v) = \delta_v$. If the envelope property holds for (X, L), then the function $\varphi_v = \operatorname{FS}(\chi_v)$ belongs to CPSH, $\varphi_v(v) = 0$ and $\operatorname{MA}(\varphi_v) = \delta_v$. However, in general the equation $\operatorname{MA}(\varphi) = \delta_v$ may not have any solution in \mathcal{E}^1 . This is the case, for example, when X is a nodal curve and v is a divisorial valuation with center at the node (compare Example 6.15).

Another important special case is when $\Sigma \subset X^{\text{div}}$ is finite. Recall that the set $\mathcal{N}_{\mathbb{R}}^{\text{div}}$ of divisorial norms is the union of $\mathcal{N}_{\mathbb{R}}^{\Sigma}$ over all nonempty finite subset $\Sigma \subset X^{\text{div}}$. We similarly introduce:

DEFINITION 7.15. — The set \mathcal{M}^{div} of divisorial measures on X^{an} is defined by

(7.6)
$$\mathcal{M}^{\operatorname{div}} = \bigcup_{\Sigma \subset X^{\operatorname{div}} \text{ finite}} \mathcal{M}^{\Sigma}.$$

The set \mathcal{M}^{div} is used in [20] to define the notion of divisorial stability.

 \square

COROLLARY 7.16. — The Monge–Ampère operator induces an isometric isomorphism

MA:
$$(\mathcal{N}^{\mathrm{div}}_{\mathbb{R}}/\mathbb{R}, \underline{\mathrm{d}}_1) \xrightarrow{\sim} (\mathcal{M}^{\mathrm{div}}, \mathrm{d}_1).$$

Further, for any $\chi \in \mathcal{N}_{\mathbb{R}}^{\text{div}}$, $\Sigma \coloneqq \text{supp MA}(\chi)$ is the smallest finite subset of X^{div} such that $\chi \in \mathcal{N}_{\mathbb{R}}^{\Sigma}$.

Example 7.17. — If $v \in X^{\text{lin}}$, then χ_v is divisorial iff v is divisorial. Indeed, $\chi_v \in \mathcal{N}_{\mathbb{R}}^{\text{div}} \Longrightarrow \text{MA}(\chi_v) = \delta_v \in \mathcal{M}^{\text{div}} \Longrightarrow v \in X^{\text{div}}$.

We now turn to the proof of Theorem 7.13. For any $\chi \in \mathcal{N}_{\mathbb{R}}$ and $\varphi \in C^0(X)$, we set

$$\chi[\varphi] \coloneqq \mathrm{IN}(\mathrm{FS}(\chi) + \varphi) \in \mathcal{N}_{\mathbb{R}}.$$

Thus $\chi[0] = \chi^{\text{hom}}$ (see Theorem 2.16). The main ingredient in the proof is now the following version of [18, Theorem 8.5] (itself a consequence of [16, Theorem A]).

LEMMA 7.18. — For any $\chi \in \mathcal{N}_{\mathbb{R}}$ and $\varphi \in C^{0}(X)$, we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\Big|_{t=0} \operatorname{vol}\left(\chi[t\varphi]\right) = \int \varphi \,\operatorname{MA}(\chi).$$

Proof. — For *d* sufficiently divisible, set $\psi_d := FS(\chi_d) \in \mathcal{H}_{\mathbb{R}}$. By Theorem 5.1 and Corollary 5.3, we have $vol(\chi) = \widetilde{E}(FS(\chi)) = vol(\chi[0])$, and

(7.7)
$$\operatorname{vol}(\chi_d[\varphi]) = \widetilde{\mathrm{E}}(\psi_d + \varphi) \longrightarrow \widetilde{\mathrm{E}}(\mathrm{FS}(\chi) + \varphi) = \operatorname{vol}(\chi[\varphi]).$$

Assume first $\varphi \in PL(X)$. By [18, Theorem 8.5], we then have

$$\widetilde{\mathbf{E}}(\psi_d + t\varphi) = \mathbf{E}(\psi_d) + t \int \varphi \, \mathrm{MA}(\psi_d) + O(t^2)$$

as $t \to 0$, where the implicit contant in O is uniform with respect to d (but does depend on φ). Now $MA(\psi_d) = MA(\chi_d) \to MA(\chi)$ strongly in \mathcal{M}^1 (see Proposition 7.4(i)); combined with (7.7), this yields

$$\operatorname{vol}(\chi[t\varphi]) = \operatorname{vol}(\chi) + t \int \varphi \operatorname{MA}(\chi) + O(t^2),$$

which proves the result for $\varphi \in PL(X)$. Consider now an arbitrary $\varphi \in C^0(X)$. Since PL(X) is dense in $C^0(X)$ with respect to uniform convergence, we can find a sequence (φ_i) in PL(X) such that $\delta_i := \sup_{X^{an}} |\varphi_i - \varphi| \to 0$. Then $\varphi_i - \delta_i \leq \varphi \leq \varphi_i + \delta_i$, and hence

$$\operatorname{vol}(\chi[t\varphi_i]) - t\delta_i \leq \operatorname{vol}(\chi[t\varphi]) \leq \operatorname{vol}(\chi[t\varphi_i]) + t\delta_i.$$

ANNALES DE L'INSTITUT FOURIER

By the first part of the proof, this yields

$$\int \varphi_i \operatorname{MA}(\chi) - \delta_i \leqslant \liminf_{t \to 0_+} t^{-1} \left(\operatorname{vol}\left(\chi[t\varphi]\right) - \operatorname{vol}(\chi) \right)$$
$$\leqslant \limsup_{t \to 0_+} t^{-1} \left(\operatorname{vol}\left(\chi[t\varphi]\right) - \operatorname{vol}(\chi) \right) \leqslant \int \varphi_i \operatorname{MA}(\chi),$$

and letting $i \to \infty$ yields, as desired, $\lim_{t\to 0} t^{-1} \left(\operatorname{vol}(\chi[t\varphi]) - \operatorname{vol}(\chi) \right) = \int \varphi \operatorname{MA}(\chi).$

Proof of Theorem 7.13. — By Theorem 7.2, MA: $(\mathcal{N}_{\mathbb{R}}/\mathbb{R}, \underline{d}_1) \to (\mathcal{M}^1, d_1)$ is an isometry. Let us first show that it maps $\mathcal{N}_{\mathbb{R}}^{\Sigma}/\mathbb{R}$ into \mathcal{M}^{Σ} . Pick $\chi \in \mathcal{N}_{\mathbb{R}}^{\Sigma}$. We need to show that $\int \varphi \operatorname{MA}(\chi) = 0$ for any $\varphi \in C^0$ such that $\varphi|_{\Sigma} = 0$. Now, for any $t \in \mathbb{R}$, we have

$$\chi[t\varphi] = \mathrm{IN}(\mathrm{FS}(\chi) + t\varphi) \leqslant \mathrm{IN}_{\Sigma}(\mathrm{FS}(\chi) + t\varphi) = \mathrm{IN}_{\Sigma}(\mathrm{FS}(\chi)) = \chi$$

This implies $\operatorname{vol}(\chi[t\varphi]) \leq \operatorname{vol}(\chi)$ for all $t \in \mathbb{R}$, and hence $\int \varphi \operatorname{MA}(\chi) = 0$, thanks to Lemma 7.18.

We next show that MA: $\mathcal{N}_{\mathbb{R}}^{\Sigma} \to \mathcal{M}^{\Sigma}$ is onto. Pick $\mu \in \mathcal{M}^{\Sigma}$, and choose a maximizing sequence (φ_i) in $\mathcal{H}_{\mathbb{R}}$ for μ , i.e. $E(\varphi_i) - \int \varphi_i \mu \to E^{\vee}(\mu)$, normalized by $\sup \varphi_i = 0$. Since Σ is strongly compact, the restriction of $PSH_{\sup} = \{\varphi \in PSH \mid \sup \varphi = 0\}$ to Σ is equicontinuous and bounded, since $0 \leq \sup_{v \in \Sigma} (-\varphi(v)) \leq \sup_{v \in \Sigma} T(v) < \infty$ for $\varphi \in PSH_{\sup}$. By the Arzelà–Ascoli theorem, we may thus assume, after passing to a subsequence, that $\varphi_i|_{\Sigma}$ converges uniformly to some $\varphi \in C^0(\Sigma)$. We claim that $\chi \coloneqq IN_{\Sigma}(\varphi) \in \mathcal{N}_{\mathbb{R}}^{\Sigma}$ satisfies $MA(\chi) = \mu$, which will conclude the proof. By Proposition 7.8, it suffices to show $vol(\chi) - \int FS(\chi) \mu \ge E^{\vee}(\mu)$.

Set $\chi_i := IN_{\Sigma}(\varphi_i)$. As $\varphi_i \to \varphi$ uniformly on Σ , (6.2) implies $d_1(\chi_i, \chi) \to 0$, and hence $vol(\chi_i) \to vol(\chi)$. Further, $\chi_i \ge IN(\varphi_i)$, and hence $vol(\chi_i) \ge vol(IN(\varphi_i)) = E(\varphi_i)$ (see Corollary 5.3). By Proposition 6.5(i), we also have $FS(\chi)|_{\Sigma} \le \varphi$. This yields, as desired,

$$\operatorname{vol}(\chi) - \int \operatorname{FS}(\chi) \mu \ge \operatorname{vol}(\chi) - \int \varphi \mu = \lim_{i} \left(\operatorname{vol}(\chi_{i}) - \int \varphi_{i} \mu \right)$$
$$\ge \lim_{i} \left(\operatorname{E}(\varphi_{i}) - \int \varphi_{i} \mu \right) = \operatorname{E}^{\vee}(\mu).$$

Finally, if $\Sigma \subset X^{\text{div}}$, or if the weak envelope property holds, then $\mathcal{N}_{\mathbb{R}}^{\Sigma}$ is contained in $\mathcal{N}_{\mathbb{R}}^{\text{max}}$ (see Corollary 6.21), and the last point thus follows from Corollary 7.5.

7.5. Dervan's minimum norm

In [36], Dervan introduced the notion of the *minimum norm* of a test configuration. Here we extend his notion to arbitrary norms.

DEFINITION 7.19. — We define the minimum norm $\|\chi\|$ of $\chi \in \mathcal{N}_{\mathbb{R}}$ by

$$\|\chi\| \coloneqq \mathbf{E}^{\vee}(\mathbf{MA}(\chi)) \in \mathbb{R}_{\geq 0}.$$

By Corollary 7.3, the minimum norm is a continuous function on $(\mathcal{N}_{\mathbb{R}}, d_1)$.

PROPOSITION 7.20. — For any $\chi \in \mathcal{N}_{\mathbb{R}}$, we have:

- (i) if χ ∈ T_Z is associated to an ample test configuration, then ||χ|| coincides, up to normalization, with the minimum norm defined in [36];
- (ii) the canonical approximants (χ_d) satisfy $\|\chi\| = \lim_d \|\chi_d\|$;
- (iii) for any $c \in \mathbb{R}$ and $t \in \mathbb{R}_{>0}$ we have $\|\chi + c\| = \|\chi\|$, and $\|t\chi\| = t\|\chi\|$;
- (iv) $\|\chi\| \approx \underline{d}_1(\chi, \chi_{triv}) \approx \|\chi\|_1$; in particular, $\|\chi\| = 0$ iff $\chi \sim \chi_{triv} + c$ for some $c \in \mathbb{R}$;
- (v) if $\chi \in \mathcal{T}_{\mathbb{R}}$, then $\|\chi\| = E(\varphi) \int \varphi \operatorname{MA}(\varphi) = I(\varphi) J(\varphi)$ with $\varphi := FS(\chi)$;
- (vi) if $\chi \in \mathcal{T}_{\mathbb{Q}}$, then $\|\chi\| \in \mathbb{Q}$;
- (vii) $\|\chi\| = \|\chi^{\text{hom}}\|;$
- (viii) if (X, L) has the weak envelope property, then $\|\chi\| = \|\chi^{\max}\|$.

Here $\|\chi\|_1 = d_1(\chi, \chi_{triv} + vol(\chi))$ is the L¹-norm of χ , see Definition 3.16.

Proof. — If $\chi \in \mathcal{T}_{\mathbb{R}}$, then $MA(\chi) = MA(\varphi)$ with $\varphi := FS(\chi) \in \mathcal{H}_{\mathbb{R}}$, so (4.10) yields $\|\chi\| = E^{\vee}(\mu) = I(\varphi) - J(\varphi)$, proving (v), and also (vi), since $\varphi \in \mathcal{H}_{\mathbb{Q}}$ when $\chi \in \mathcal{T}_{\mathbb{Q}}$. If, further, $\chi \in \mathcal{T}_{\mathbb{Z}}$ is a test configuration, then [17, Remark 7.12] shows that $I(\varphi)-J(\varphi)$ coincides (up to normalization by V_L) with the minimum norm of χ as defined in [36, Definition 2.5]. Thus (i) holds. Now (ii), (iii), (vii) and (viii) are immediate consequence of the corresponding properties in Proposition 7.4.

It remains to prove (iv). That $\|\chi\| \approx \underline{d}_1(\chi, \chi_{triv})$ follows from (5.15) and Theorem 7.2, whereas $\|\chi\|_1 \approx \|\chi\|$ follows from (iv) and [17, Theorem 7.9] when $\chi \in \mathcal{T}_{\mathbb{R}}$, and hence in general, by density.

By d₁-density of $\mathcal{T}_{\mathbb{Z}}$ in $\mathcal{N}_{\mathbb{R}}$ (see Corollary 3.19), we infer:

COROLLARY 7.21. — The minimum norm functional $\mathcal{N}_{\mathbb{R}} \to \mathbb{R}_{\geq 0}$ is the unique d₁-continuous extension of Dervan's minimum norm from $\mathcal{T}_{\mathbb{Z}}$ to $\mathcal{N}_{\mathbb{R}}$.

7.6. Valuations of linear growth

In this final section, we specialize the above results to prove:

THEOREM 7.22. — For all $v, w \in X^{\text{lin}}$ with associated norms $\chi_v, \chi_w \in \mathcal{N}_{\mathbb{R}}$ and measures $\delta_v, \delta_w \in \mathcal{M}^1$, we have:

- (i) $d_{\infty}(v,w) = d_{\infty}(\chi_v,\chi_w) \approx \underline{d}_1(\chi_v,\chi_w) = d_1(\delta_v,\delta_w);$
- (ii) $d_{\infty}(v, v_{\text{triv}}) = T(v) = \lambda_{\max}(\chi_v);$
- (iii) $S(v) = vol(\chi_v) = ||\chi_v|| = E^{\vee}(\delta_v).$

Since $\underline{\mathbf{d}}_1 \leq \mathbf{d}_1 \leq \mathbf{d}_p \leq \mathbf{d}_\infty$ on $\mathcal{N}_{\mathbb{R}}$ for $1 \leq p \leq \infty$, this implies:

COROLLARY 7.23. — For any $p \in [1, \infty]$, the embeddings

$$(X^{\mathrm{lin}}, \mathrm{d}_{\infty}) \hookrightarrow (\mathcal{M}^{1}, \mathrm{d}_{1}), \quad (X^{\mathrm{lin}}, \mathrm{d}_{\infty}) \hookrightarrow (\mathcal{N}_{\mathbb{R}}, \mathrm{d}_{p})$$

respectively defined by $v \mapsto \delta_v$ and $v \mapsto \chi_v$ are bi-Lipschitz.

Note that this implies Corollary E in the introduction.

Proof of Theorem 7.22. — By Theorem 7.13, we have $MA(\chi_v) = \delta_v$, $MA(\chi_w) = \delta_v$, and hence

$$d_1(\delta_v, \delta_v) = \underline{d}_1(\chi_v, \chi_w) \leqslant d_\infty(\chi_v, \chi_w) = d_\infty(v, w),$$

by Theorem 7.2 and Corollary 2.10. Next, note that

(7.8)
$$FS(\chi_v)(w) = \sup\{m^{-1}(v(s) - w(s))\},\$$

where s runs over nonzero sections of mL with m sufficiently divisible. In particular, $FS(\chi_v) \ge 0$, and $FS(\chi_v)(v) = 0$. Comparing (7.8) with (1.19) yields

(7.9)
$$d_{\infty}(v,w) = \max\{\mathrm{FS}(\chi_v)(w),\mathrm{FS}(\chi_w)(v)\}.$$

On the other hand, for each $c \in \mathbb{R}$, Lemma 7.7 yields

$$d_1(\chi_v + c, \chi_w) \approx \int |FS(\chi_v) + c - FS(\chi_w)| (\delta_v + \delta_w)$$

= |FS(\chi_v)(w) + c| + |c - FS(\chi_w)(v)|
\ge FS(\chi_v)(w) + FS(\chi_w)(v) \ge d_\pi(v, w).

Thus

$$\underline{\mathbf{d}}_1(\chi_v, \chi_w) = \inf_{c \in \mathbb{R}} \mathbf{d}_1(\chi_v, \chi_w + c) \gtrsim \mathbf{d}_\infty(v, w).$$

This proves (i), and (ii) follows. Finally, $MA(\chi_v) = \delta_v$ implies $||\chi_v|| = E^{\vee}(\delta_v)$, by definition of the minimum norm. Since $FS(\chi_v)$ vanishes at v, Proposition 7.8 further yields $E^{\vee}(\delta_v) = vol(\chi_v)$, which coincides with S(v) (see Example 3.5). This proves (iii).

TOME 75 (2025), FASCICULE 2

Appendix A. Test configurations, integral closure, and homogenization

In this appendix we revisit the correspondence between test configurations and integral norms [17, 68], and provide a description of homogenization in terms of integral closure. We also provide a geometric description of \mathbb{R} -test configurations, following [49, 50].

A.1. The norm associated to a test configuration

A test configuration $(\mathcal{X}, \mathcal{L})$ for (X, L) consists of: a flat projective morphism $\pi \colon \mathcal{X} \to \mathbb{A}^1$; a \mathbb{Q} -line bundle \mathcal{L} on \mathcal{X} ; a \mathbb{G}_m -action on $(\mathcal{X}, \mathcal{L})$ that makes π equivariant; and a \mathbb{G}_m -equivariant isomorphism

(A.1)
$$(\mathcal{X}, \mathcal{L})|_{\mathbb{G}_{\mathrm{m}}} \simeq (X, L) \times \mathbb{G}_{\mathrm{m}}.$$

We denote by z the coordinate on $\mathbb{A}^1 = \operatorname{Spec} k[z]$ and $\mathbb{G}_{\mathrm{m}} = \operatorname{Spec} k[z^{\pm}]$.

Example A.1. — The trivial test configuration $(\mathcal{X}_{triv}, \mathcal{L}_{triv})$ is defined by $\mathcal{X}_{triv} = X \times \mathbb{A}^1, \ \mathcal{L}_{triv} = p_1^* L.$

As originally pointed out in [68], to any test configuration $(\mathcal{X}, \mathcal{L})$ is associated an integral norm $\chi_{\mathcal{L}} \in \mathcal{N}_{\mathbb{Z}}$, defined on $R_m = \mathrm{H}^0(X, mL)$ for any $m \in \mathbb{N}$ such that $m\mathcal{L}$ is a line bundle, as follows. Consider the embedding $\mathrm{H}^0(\mathcal{X}, m\mathcal{L}) \hookrightarrow R_m \otimes k[z^{\pm}]$ induced by (A.1). This yields a decomposition

(A.2)
$$\mathrm{H}^{0}(\mathcal{X}, m\mathcal{L}) = \bigoplus_{\lambda \in \mathbb{Z}} z^{-\lambda} F^{\lambda} R_{m},$$

corresponding to the weight decomposition with respect to the $\mathbb{G}_{\mathrm{m}}\text{-}\mathrm{action},$ where

(A.3)
$$F^{\lambda}R_m = \{s \in R_m \mid z^{-\lambda}s \in \mathrm{H}^0(\mathcal{X}, m\mathcal{L})\}$$

is a \mathbb{Z} -filtration of R_m , and we define $\chi_{\mathcal{L}}$ as the associated norm. It is clear that $\chi_{\mathcal{L}+c\mathcal{X}_0} = \chi_{\mathcal{L}} + c$ for any $c \in \mathbb{Q}$, and using flat base change, one easily checks:

LEMMA A.2. — If $(\mathcal{X}_d, \mathcal{L}_d)$ denotes the base change of $(\mathcal{X}, \mathcal{L})$ with respect to $z \mapsto z^d$, $d \ge 1$, then $\chi_{\mathcal{L}_d} = d\chi_{\mathcal{L}}$.

In order to further analyze the norm $\chi_{\mathcal{L}}$, recall from [18, Section 1.4] that a test configuration \mathcal{X} is *integrally closed* if \mathcal{X} is integrally closed in the generic fiber of π ; when X is normal, this is equivalent to \mathcal{X} being normal. If \mathcal{X}_0 is reduced, then \mathcal{X} is integrally closed. If \mathcal{X} is integrally closed, the local ring of \mathcal{X} at the generic point of any irreducible component E of \mathcal{X}_0 is a DVR, which defines a divisorial valuation ord_E on \mathcal{X} ; we denote by

(A.4)
$$b_E \coloneqq \operatorname{ord}_E(\mathcal{X}_0) = \operatorname{ord}_E(z)$$

the multiplicity of \mathcal{X}_0 along E. By (A.1) we have a function field extension $k(X) \hookrightarrow k(\mathcal{X})$, and the restriction of $b_E^{-1} \operatorname{ord}_E$ to k(X) is a divisorial valuation $v_E \in X^{\operatorname{div}}$, with values in $b_E^{-1}\mathbb{Z}$. Conversely, any divisorial valuation can be geometrically realized in this way.

Recall also from [18, Section 2.7] that any test configuration $(\mathcal{X}, \mathcal{L})$ for (X, L) determines a PL function $\varphi_{\mathcal{L}} \in PL(X)$, whose restriction to the dense subset $X^{\text{div}} \subset X^{\text{an}}$ is given as follows. Pick $v \in X^{\text{div}}$, and choose an integrally closed test configuration \mathcal{X}' for X such that $v = v_E$ is associated to an irreducible component $E \subset \mathcal{X}'_0$ and such that the canonical \mathbb{G}_{m} -equivariant birational maps $\mu \colon \mathcal{X}' \to \mathcal{X}$ and $\rho \colon \mathcal{X}' \to \mathcal{X}_{\text{triv}}$ are morphisms. Then $\mu^* \mathcal{L} - \rho^* \mathcal{L}_{\text{triv}} = D$ for a Q-Cartier divisor D supported on \mathcal{X}'_0 , and

(A.5)
$$\varphi_{\mathcal{L}}(v_E) = b_E^{-1} \operatorname{ord}_E(D).$$

Conversely, any PL function on X^{an} can be realized in this way (see [18, Theorem 2.31]).

PROPOSITION A.3. — Pick an integrally closed test configuration $(\mathcal{X}, \mathcal{L})$, set $\varphi \coloneqq \varphi_{\mathcal{L}}$, and denote by $\Sigma \subset X^{\text{div}}$ the (finite) set of valuations attached to the irreducible components of \mathcal{X}_0 . Then:

- (i) $\chi_{\mathcal{L}} = \lfloor IN_{\Sigma}(\varphi) \rfloor = \lfloor IN(\varphi) \rfloor;$
- (ii) $\chi_{\mathcal{L}}^{\text{hom}} = \text{IN}_{\Sigma}(\varphi) = \text{IN}(\varphi);$
- (iii) if \mathcal{X}_0 is reduced, then $\chi_{\mathcal{L}}$ is homogeneous.

Here $IN_{\Sigma}(\varphi)$ is defined in Section 6.1.

LEMMA A.4. — Under the above assumptions we have

$$\chi_{\mathcal{L}} = \chi_{\mu^{\star}\mathcal{L}}, \quad \varphi_{\mathcal{L}} = \varphi_{\mu^{\star}\mathcal{L}}$$

for any morphism of test configurations $\mu \colon \mathcal{X}' \to \mathcal{X}$.

Proof. — By Zariski's main theorem, we have $\mu_{\star}\mathcal{O}_{\mathcal{X}'} = \mathcal{O}_{\mathcal{X}}$ (see [18, Lemma 1.12]). Combined with the projection formula, this shows $\mathrm{H}^{0}(\mathcal{X}, m\mathcal{L}) = \mathrm{H}^{0}(\mathcal{X}', m\mu^{\star}\mathcal{L})$ for *m* sufficiently divisible. The first point follows, while the second one holds by (A.5).

Proof of Proposition A.3. — Pick $s \in R_m$ with m sufficiently divisible and $\lambda \in \mathbb{Z}$. Then $z^{-\lambda}s$ determines a rational section σ of $m\mathcal{L}$ which is regular outside \mathcal{X}_0 , and hence is regular on \mathcal{X} iff $\operatorname{ord}_E(\sigma) \ge 0$ for each irreducible component E of \mathcal{X} (see [18, Lemma 1.23]). Now (A.5) implies

$$b_E^{-1} \operatorname{ord}_E(\sigma) = -\lambda + v_E(s) + \varphi(v_E),$$

and we infer

$$\chi_{\mathcal{L}}(s) = \max\left\{\lambda \in \mathbb{Z} \mid \lambda \leqslant \min_{E} \{v_{E}(s) + \varphi(v_{E})\}\right\} = \lfloor \mathrm{IN}_{\Sigma}(\varphi)(s) \rfloor.$$

Next, pick $v \in X^{\text{div}}$, and choose an integrally closed test configuration \mathcal{X}' that dominates \mathcal{X} via $\mu \colon \mathcal{X}' \to \mathcal{X}$ and such that v lies in the corresponding set $\Sigma' \subset X^{\text{div}}$. Lemma A.4 and the first step of the proof yield

 $\chi_{\mathcal{L}}(s) = \chi_{\mu^{\star}\mathcal{L}}(s) = \lfloor \mathrm{IN}_{\Sigma'}(\varphi)(s) \rfloor \leqslant v(s) + \varphi(v).$

By density of X^{div} and continuity of φ , we infer

$$\chi_{\mathcal{L}}(s) \leqslant \inf_{v \in X^{\mathrm{div}}} \{v(s) + \varphi(v)\} = \mathrm{IN}(\varphi)(s) \leqslant \mathrm{IN}_{\Sigma}(\varphi)(s)$$

Since $\chi_{\mathcal{L}} = \lfloor \mathrm{IN}_{\Sigma}(\varphi) \rfloor$, this proves (i), and (ii) follows, cf. Example 2.5. Finally, if \mathcal{X}_0 is reduced, then each $v \in \Sigma$ is integer valued on $k(X)^{\times}$. Since $\varphi(v)$ is rational (see (A.5)), we get

$$IN_{\Sigma}(\varphi)(s) = \min_{v \in \Sigma} \{v(s) + m\varphi(v)\} \in \mathbb{Z}$$

for $s \in R_m$ with *m* sufficiently divisible, and (iii) now follows from (i) and (ii).

Remark A.5. — Proposition A.3(ii) implies $\chi_{\mathcal{L}}^{\text{hom}} \in \mathcal{N}_{\mathbb{Q}}^{\Sigma}$, and hence $MA(\chi_{\mathcal{L}}) = MA(\chi_{\mathcal{L}}^{\text{hom}}) \in \mathcal{M}^{\Sigma}$ (see Theorem 7.13). The coefficients of this measure admit an explicit description in terms of positive intersection classes on the canonical compactification $\overline{\mathcal{X}} \to \mathbb{P}^1$, see [56, Theorem 1.1].

Consider now an arbitrary test configuration $(\mathcal{X}, \mathcal{L})$, and denote by $(\widetilde{\mathcal{X}}, \widetilde{\mathcal{L}})$ its integral closure, i.e. $\widetilde{\mathcal{X}} \to \mathcal{X}$ is the integral closure of \mathcal{X} in the generic fiber of $\mathcal{X} \to \mathbb{A}^1$, and $\widetilde{\mathcal{L}}$ is the pullback of \mathcal{L} .

THEOREM A.6. — For any test configuration $(\mathcal{X}, \mathcal{L})$ for (X, L), we have

$$\chi_{\mathcal{L}}^{\text{hom}} = \chi_{\widetilde{\mathcal{L}}}^{\text{hom}} = \text{IN}(\varphi_{\mathcal{L}}) \text{ and } \chi_{\widetilde{\mathcal{L}}} = \lfloor \chi_{\mathcal{L}}^{\text{hom}} \rfloor.$$

In other words, integral closure is the round-down of homogenization.

LEMMA A.7. — We have $\chi_{\mathcal{L}} \leq \chi_{\widetilde{\mathcal{L}}} \leq \chi_{\mathcal{L}}^{\text{hom}}$.

Proof. — Pick $s \in R_m \setminus \{0\}$ with m sufficiently divisible. Since $\widetilde{\mathcal{L}}$ is the pullback of \mathcal{L} to $\widetilde{\mathcal{X}}, \chi_{\mathcal{L}}(s) \leq \chi_{\widetilde{\mathcal{L}}}(s) =: \mu$ follows directly from (A.3). Since $\sigma := z^{-\mu}s \in \mathrm{H}^0(\widetilde{\mathcal{X}}, m\widetilde{\mathcal{L}})$ is integral over $\mathcal{O}_{\mathcal{X}}$, it satisfies $\sigma^d + \sum_{i=1}^d \sigma_i \sigma^{d-i} =$

0 for some $d \ge 1$ and $\sigma_i \in \mathrm{H}^0(\mathcal{X}, im\mathcal{L})$. By (A.2), we have a Laurent expansion $\sigma_i = \sum_{\lambda \in \mathbb{Z}} \sigma_{i,\lambda} z^{-\lambda}$ with $\sigma_{i,\lambda} \in \mathrm{H}^0(X, mL)$ such that $\chi_{\mathcal{L}}(\sigma_{i,\lambda}) \ge \lambda$, and tracing the coefficient of $z^{-d\mu}$ yields $s^d + \sum_{i=1}^d \sigma_{i,i\mu} s^{d-i} = 0$. Since $\chi_{\mathcal{L}}^{\mathrm{hom}}(\sigma_{i,i\mu}) \ge \chi_{\mathcal{L}}(\sigma_{i,i\mu}) \ge i\mu$, we infer

$$d\chi_{\mathcal{L}}^{\mathrm{hom}}(s) = \chi_{\mathcal{L}}^{\mathrm{hom}}(s^d) \ge \min_{1 \le i \le d} \left\{ i\mu + (d-i)\chi_{\mathcal{L}}^{\mathrm{hom}}(s) \right\};$$

hence $\chi_{\mathcal{L}}^{\text{hom}}(s) \ge \mu = \chi_{\widetilde{\mathcal{L}}}(s)$, and we are done.

Proof of Theorem A.6. — Lemma A.7 implies $\chi_{\mathcal{L}}^{\text{hom}} = \chi_{\widetilde{\mathcal{L}}}^{\text{hom}}$, which is equal to $\text{IN}(\varphi_{\widetilde{\mathcal{L}}}) = \text{IN}(\varphi_{\mathcal{L}})$, by Proposition A.3(ii) and pullback invariance of $\varphi_{\mathcal{L}}$. The final identity follows from Proposition A.3.

As a consequence, we get the following geometric description of homogenization:

COROLLARY A.8. — For any test configuration $(\mathcal{X}, \mathcal{L})$, $\chi^{\text{hom}}_{\mathcal{L}}$ lies in $\mathcal{N}^{\text{div}}_{\mathbb{Q}}$, and is equal to $d^{-1}\chi_{\widetilde{\mathcal{L}}_d}$ for any sufficiently divisible $d \in \mathbb{Z}_{\geq 1}$.

As above, $(\mathcal{X}_d, \mathcal{L}_d)$ denotes the base change of $(\mathcal{X}, \mathcal{L})$ with respect to $z \mapsto z^d$, and $(\widetilde{\mathcal{X}}_d, \widetilde{\mathcal{L}}_d)$ is its integral closure.

Proof. — By [18, Corollary 2.35], the central fiber of $(\widetilde{\mathcal{X}}_d, \widetilde{\mathcal{L}}_d)$ is reduced for d sufficiently divisible. Then $\chi^{\text{hom}}_{\mathcal{L}_d} = \chi^{\text{hom}}_{\widetilde{\mathcal{L}}_d} = \chi_{\widetilde{\mathcal{L}}_d}$, by Theorem A.6 and Proposition A.3(iii). By Lemma A.2, we have, on the other hand, $\chi_{\mathcal{L}_d} = d\chi_{\mathcal{L}}$, and hence $\chi^{\text{hom}}_{\mathcal{L}_d} = d\chi^{\text{hom}}_{\mathcal{L}}$. Thus $\chi^{\text{hom}}_{\mathcal{L}} = d^{-1}\chi^{\text{hom}}_{\widetilde{\mathcal{L}}_d}$, which lies in $\mathcal{N}^{\text{div}}_{\mathbb{U}}$, by Proposition A.3(ii).

Remark A.9. — Corollary A.8 can be used to provide a more elementary proof of Theorem 2.3 in the case of rational norms.

Finally, we relate (integrally closed) test configurations and (rational) divisorial norms, as follows:

THEOREM A.10. — For any $\chi \in \mathcal{N}_{\mathbb{Z}}$, the following are equivalent:

- (i) χ = χ_L is associated to some integrally closed test configuration (X, L) for (X, L);
- (ii) $\chi = \lfloor \chi' \rfloor$ for some $\chi' \in \mathcal{N}_{\mathbb{Q}}^{\text{div}}$, which is then uniquely determined as $\chi' = \chi^{\text{hom}}$.

Proof. — That (i) implies (ii) follows from Proposition A.3. Conversely, pick $\chi' \in \mathcal{N}_{\mathbb{Q}}^{\text{div}}$, and set $\chi := \lfloor \chi' \rfloor$, so that $\chi' = \chi^{\text{hom}}$ (see Example 2.5). By Theorem 6.12, we have $\chi' = \text{IN}(\varphi)$ for some $\varphi \in \text{PL}(X)$, which can in turn be written $\varphi = \varphi_{\mathcal{L}}$ for some integrally closed test configuration $(\mathcal{X}, \mathcal{L})$. By Proposition A.3, we then have $\chi_{\mathcal{L}} = \lfloor \chi' \rfloor = \chi$, which shows (ii) \Rightarrow (i). \Box

 \square

A.2. The Rees correspondence

A test configuration $(\mathcal{X}, \mathcal{L})$ is ample if \mathcal{L} is ample. For d sufficiently divisible, the graded k[z]-algebra $R(\mathcal{X}, d\mathcal{L}) = \bigoplus_{m \in \mathbb{N}} \mathrm{H}^0(\mathcal{X}, md\mathcal{L})$ is then generated in degree 1, which shows that $\chi = \chi_{\mathcal{L}} \in \mathcal{T}_{\mathbb{Z}}$ is of finite type. Note further that

(A.6)
$$R(\mathcal{X}_0, d\mathcal{L}) \simeq \operatorname{gr}_{\gamma} R^{(d)}$$

for d sufficiently divisible. Thus $(\mathcal{X}_0, \mathcal{L}_0)$ can be identified with the central fiber of χ (see (1.14)).

Denoting by \mathcal{T} the set of ample test configurations, $(\mathcal{X}, \mathcal{L}) \mapsto \chi_{\mathcal{L}}$ yields a map $\mathcal{T} \to \mathcal{T}_{\mathbb{Z}}$. A map in the reverse direction is provided by the *Rees* construction. Given $\chi \in \mathcal{T}_{\mathbb{Z}}$, pick $d \ge 1$ such that χ is represented by an integral norm on $R^{(d)} = R(X, dL)$ generated in degree 1, with associated filtration $(F^{\lambda}R^{(d)})_{\lambda \in \mathbb{Z}}$. The *Rees* algebra

$$\mathcal{R} \coloneqq \bigoplus_{\lambda \in \mathbb{Z}} z^{-\lambda} F^{\lambda} R^{(d)}$$

is a graded k[z]-algebra (with respect to the N-grading inherited from that of $R^{(d)}$), generated in degree 1, and we set $\mathcal{X} := \operatorname{Proj}_{k[z]} \mathcal{R}$ and $\mathcal{L} = d^{-1}\mathcal{O}_{\mathcal{X}}(1)$. This yields a map $\mathcal{T}_{\mathbb{Z}} \to \mathcal{T}$ which is an inverse of the previous one (see [17, Proposition 2.15]). We shall refer to the 1–1 map

(A.7)
$$\mathcal{T} \simeq \mathcal{T}_{\mathbb{Z}}$$

so defined as the Rees correspondence.

By (A.6), the central fiber \mathcal{X}_0 of an ample test configuration $(\mathcal{X}, \mathcal{L})$ is reduced iff $\operatorname{gr}_{\chi_{\mathcal{L}}} R^{(d)}$ is reduced for *d* sufficiently divisible, which holds iff $\chi_{\mathcal{L}}$ is homogeneous (i.e. the converse of Proposition A.3(iii) holds for ample test configurations). The Rees correspondence thus induces a bijection between $\mathcal{T}_{\mathbb{Z}}^{\text{hom}}$ and the set of ample test configurations with reduced central fiber.

Denote by $\mathcal{T}^{int} \subset \mathcal{T}$ the set of ample integrally closed test configurations, and by $\mathcal{T}_{\mathbb{Z}}^{int} \subset \mathcal{T}_{\mathbb{Z}}$ its image under the Rees correspondence. Any test configuration with reduced central fiber is integrally closed, and hence

$$\mathcal{T}^{\mathrm{hom}}_{\mathbb{Z}} \subset \mathcal{T}^{\mathrm{int}}_{\mathbb{Z}} \subset \mathcal{T}_{\mathbb{Z}}.$$

THEOREM A.11. — Homogenization induces a bijection $\mathcal{T}_{\mathbb{Z}}^{int} \xrightarrow{\sim} \mathcal{T}_{\mathbb{Q}}^{hom}$, with inverse provided by round-down.

Proof. — Pick $\chi \in \mathcal{T}_{\mathbb{Z}}^{\text{int.}}$. Then $\chi^{\text{hom}} \in \mathcal{T}_{\mathbb{Q}}^{\text{hom}}$ (see Lemma 2.11, or Corollary A.8), and $\chi = \lfloor \chi^{\text{hom}} \rfloor$ (see Proposition A.3). Conversely, pick $\chi' \in \mathcal{T}_{\mathbb{Q}}^{\text{hom}}$. By Corollary 2.18, $\chi' = \chi^{\text{hom}}$ for some $\chi \in \mathcal{T}_{\mathbb{Z}}$, i.e. $\chi = \chi_{\mathcal{L}}$ for some

ample test configuration $(\mathcal{X}, \mathcal{L})$. After passing to the integral closure $(\tilde{\mathcal{X}}, \tilde{\mathcal{L}})$ (which remains ample, since $\tilde{\mathcal{X}} \to \mathcal{X}$ is finite), we may further assume that $(\mathcal{X}, \mathcal{L})$ is integrally closed (see Theorem A.6), and hence $\chi \in \mathcal{T}_{\mathbb{Z}}^{\text{int}}$. Then $\chi = \lfloor \chi' \rfloor$, by Proposition A.3 again, and $\chi' = \chi^{\text{hom}}$, which completes the proof.

Finally, we note:

LEMMA A.12. — For any ample test configuration $(\mathcal{X}, \mathcal{L})$ for (X, L) we have

$$\varphi_{\mathcal{L}} = \mathrm{FS}(\chi_{\mathcal{L}}) = \mathrm{FS}(\chi_{\mathcal{L}}^{\mathrm{hom}}).$$

Proof. — The second equality follows from Proposition 2.14. Since $\chi_{\mathcal{L}}^{\text{hom}} = \text{IN}(\varphi_{\mathcal{L}})$ (see Theorem A.6) and $\varphi_{\mathcal{L}} \in \mathcal{H}_{\mathbb{Q}}$, Proposition 2.29 further yields $\text{FS}(\chi_{\mathcal{L}}^{\text{hom}}) = \mathbb{Q}(\varphi_{\mathcal{L}}) = \varphi_{\mathcal{L}}$, which completes the proof.

Combining Theorem A.11 with the bijection FS: $\mathcal{T}_{\mathbb{Q}}^{\text{hom}} \xrightarrow{\sim} \mathcal{H}_{\mathbb{Q}}$, we thus recover [18, Corollary 2.32]:

COROLLARY A.13. — The map $(\mathcal{X}, \mathcal{L}) \mapsto \varphi_{\mathcal{L}}$ restricts to a bijection $\mathcal{T}^{\text{int}} \xrightarrow{\sim} \mathcal{H}_{\mathbb{Q}}$.

A.3. The case of higher rank

Following [49, Section 2.2] and [50, Section 2.2], we briefly discuss a version of the Rees correspondence for \mathbb{R} -test configurations.

DEFINITION A.14. — For any $r \in \mathbb{N}$, we define a rank r test configuration $(\mathcal{X}, \mathcal{L}, \xi)$ for (X, L) as the following data:

- a flat projective scheme morphism $\pi: \mathcal{X} \to \mathbb{A}^r$;
- a \mathbb{Q} -line bundle \mathcal{L} on \mathcal{X} ;
- a $\mathbb{G}_{\mathrm{m}}^{r}$ -action on $(\mathcal{X}, \mathcal{L})$ that makes π equivariant (with respect to the standard action on \mathbb{A}^{r});
- a $\mathbb{G}_{\mathrm{m}}^{r}$ -equivariant isomorphism $(\mathcal{X}, \mathcal{L})|_{\mathbb{G}_{\mathrm{m}}^{r}} \simeq (X, L) \times \mathbb{G}_{\mathrm{m}}^{r};$
- a vector $\xi \in \mathbb{R}^r_+$ with \mathbb{Q} -linearly independent components.

A usual test configuration as in Section A.1 is thus a rank 1 test configuration, up to the scaling factor $\xi \in \mathbb{R}_{>0}$.

Denote by z_1, \ldots, z_r the coordinates on $\mathbb{A}^r = \operatorname{Spec} k[z_1, \ldots, z_r]$ and $\mathbb{G}_{\mathrm{m}}^r = \operatorname{Spec} k[z_1^{\pm}, \ldots, z_r^{\pm}]$. The above data yields an embedding

$$\mathrm{H}^{0}(\mathcal{X}, m\mathcal{L}) \hookrightarrow R_{m}[z_{1}^{\pm}, \dots, z_{r}^{\pm}]$$

for m sufficiently divisible, and we define a norm $\chi_{\mathcal{L},\mathcal{E}} \in \mathcal{N}_{\mathbb{R}}$ by setting

$$\chi_{\mathcal{L},\xi}(s) = \max\left\{ \langle \xi, \alpha \rangle \mid \alpha \in \mathbb{Z}^r, \, z^{-\alpha}s \in \mathrm{H}^0(\mathcal{X}, m\mathcal{L}) \right\}$$

for $s \in R_m$, where $z^{\alpha} \coloneqq \prod_i z_i^{\alpha_i}$. Note that

$$\chi_{\mathcal{L},\xi} \in \mathcal{N}_{\Lambda} \text{ with } \Lambda \coloneqq \sum_{i} \mathbb{Z}\xi_{i} \simeq \mathbb{Z}^{r}.$$

PROPOSITION A.15. — For any rank r test configuration $(\mathcal{X}, \mathcal{L}, \xi)$ with \mathcal{L} relatively ample, the associated norm $\chi_{\mathcal{L},\xi}$ is of finite type; this norm is further of rank r, and its central fiber can be identified with the fiber $(\mathcal{X}_0, \mathcal{L}_0)$ of π over $0 \in \mathbb{A}^r$. Conversely, any \mathbb{R} -test configuration $\chi \in \mathcal{T}_{\mathbb{R}}$ arises in this way.

Proof. — We sketch the argument, and refer to [50, Proposition 2.20] for details. Assume \mathcal{L} is relatively ample, and set $\chi = \chi_{\mathcal{L},\xi}$. The restriction map $R(\mathcal{X}, d\mathcal{L}) \to R(\mathcal{X}_0, d\mathcal{L}_0)$ is surjective for d sufficiently divisible, and one checks that it induces an isomorphism $\operatorname{gr}_{\chi} R^{(d)} \simeq R(\mathcal{X}_0, d\mathcal{L}_0)$ as $\mathbb{N} \times \mathbb{Z}^r$ graded algebras. The rest easily follows.

Conversely, pick $\chi \in \mathcal{T}_{\mathbb{R}}$, of rank r. As in Example 1.11, one can find an embedding $X \hookrightarrow \mathbb{P}^N$ such that $\mathcal{O}(1)|_X = dL$, an action of a torus $T \simeq \mathbb{G}^r_{\mathrm{m}}$ on $(\mathbb{P}^N, \mathcal{O}(1))$ and $\xi \in N_{\mathbb{R}} \simeq \mathbb{R}^r$, such that the induced norm on $R(\mathbb{P}^N, \mathcal{O}(1))$ restricts to χ . Acting on X defines a T-equivariant morphism $T \to$ Hilb to the Hilbert scheme of \mathbb{P}^N . Pick a regular top-dimensional cone $\sigma \subset N_{\mathbb{R}}$ that contains ξ in its interior, and denote by $B \simeq \mathbb{A}^r$ the corresponding toric affine variety. After passing to a finer cone, one may assume, by toric resolution of singularities, that the corresponding T-equivariant rational map $B \dashrightarrow$ Hilb is a morphism, and pulling back the universal family yields the desired polarized family $(\mathcal{X}, \mathcal{L})$.

Appendix B. The toric case

We give a brief account of how some of the main results in the paper specialize to the toric setting [24, 45]. See also Appendix B in [18].

Consider an algebraic torus $T \simeq \mathbb{G}_{\mathrm{m}}^{n}$, with associated dual lattices $M := \mathrm{Hom}(T, \mathbb{G}_{\mathrm{m}})$ and $N := \mathrm{Hom}(\mathbb{G}_{\mathrm{m}}, T)$. We have a canonical embedding $M \hookrightarrow k(T)^{\times}$ given by $\alpha \mapsto z^{\alpha}$ onto the set *T*-invariant functions, and a dual canonical embedding $N_{\mathbb{R}} \hookrightarrow T^{\mathrm{val}}$ given by $\xi \mapsto v_{\xi}$ onto the set of T(k)-invariant valuations, such that $v_{\xi}(z^{\alpha}) = \langle \xi, \alpha \rangle$ for all $\xi \in N_{\mathbb{R}}$ and $\alpha \in M \hookrightarrow k(T)^{\times}$.

A polarized toric variety (X, L) is determined by a rational polytope $P \subset M_{\mathbb{R}}$, such that, for each m sufficiently divisible, the set of weights $\alpha \in M$ of the T(k)-module $R_m = \mathrm{H}^0(X, mL)$ coincides with $mP \cap M$, each with multiplicity 1. Denoting by $\widehat{P} \coloneqq \mathbb{R}_{\geq 0}(\{1\} \times P) \subset \mathbb{R} \times M_{\mathbb{R}}$ the (rational polyhedral) cone over P, this yields, for d sufficiently divisible, a 1–1 correspondence between:

- (a) the set of toric (i.e. T(k)-invariant) norms χ on $R^{(d)} = R(X, dL)$ and superadditive functions $h: \Gamma^{(d)} \to \mathbb{R}$ on the semigroup $\Gamma^{(d)} :=$ $(d\mathbb{N} \times M) \cap \widehat{P}$ such that $h(m, \alpha) = O(m)$:
- (b) the subset of toric homogeneous norms χ and concave, bounded functions $q: P \to \mathbb{R}$, the corresponding superadditive function on $\Gamma^{(d)}$ being $h(m, \alpha) = mq(m^{-1}\alpha)$.

Example B.1. — Each $\xi \in N_{\mathbb{R}}$ determines a toric homogeneous norm $\chi_{v_{\xi}}$, with associated function $g(\alpha) = \langle \xi, \alpha \rangle + c$ for $c \in \mathbb{R}$ such that $\inf_P g = 0$, i.e. $c = -\inf_{\alpha \in P} \langle \xi, \alpha \rangle$.

A function g as in (b) above is automatically lsc on P (see [46]), but might be discontinuous at some boundary points. Denote by $g^{\vee} \colon N_{\mathbb{R}} \to \mathbb{R}$ its (convex) Legendre transform, defined by

$$g^{\vee}(\xi) = \sup_{\alpha \in P} \{ \langle \alpha, \xi \rangle + g(\alpha) \},$$

and let also λ_P be the Lebesgue measure of P, normalized to mass 1. Then:

- (i) $FS(\chi)|_{N_{\mathbb{R}}} = g^{\vee} 0^{\vee}$, where 0^{\vee} coincides with the support function of P;
- (ii) $\operatorname{vol}(\chi) = \int g \lambda_P;$
- (iii) $d_{\infty}(\chi, \chi') = \sup_{P} |g g'|$, and $d_{p}(\chi, \chi') = ||g g'||_{L^{p}(\lambda_{P})}$ for $p \in$ $[1,\infty);$
- (iv) $\chi \in \mathcal{T}_{\mathbb{R}}$ (resp. $\mathcal{T}_{\mathbb{Q}}$) iff $g^{\vee}(\xi) = \max_i \{ \langle \xi, \alpha_i \rangle + \lambda_i \}$ for a finite set $\alpha_i \in P \cap M_{\mathbb{Q}}$ and $\lambda_i \in \mathbb{R}$ (resp. \mathbb{Q});
- (v) $\chi \in \mathcal{N}_{\mathbb{R}}^{\text{cont}} \iff g \in \mathcal{C}^{0}(P) \iff g \text{ usc } \iff \chi \in \mathcal{N}_{\mathbb{R}}^{\max};$ (vi) $\chi \in \mathcal{N}_{\mathbb{R}}^{\text{div}}$ (resp. $\mathcal{N}_{\mathbb{Q}}^{\text{div}}$) iff $g(\alpha) = \min_{j} \{ \langle \xi_{j}, \alpha \rangle + c_{j} \}$ for a finite set $\xi_i \in N_{\mathbb{Q}}$ and $c_i \in \mathbb{R}$ (resp. \mathbb{Q});
- (vii) $MA(\chi) = MA_{\mathbb{R}}(g^{\vee}) = (\nabla g)_{\star}\lambda_P$, where $MA_{\mathbb{R}}$ is the real Monge-Ampère operator and ∇g is the (λ_P -a.e. defined) gradient of g.

By (iv), (vi) and basic convex geometry, it follows that any toric homogeneous norm χ satisfies

$$\chi \in \mathcal{N}_{\mathbb{Q}}^{\mathrm{div}} \Longleftrightarrow \chi \in \mathcal{T}_{\mathbb{Q}}.$$

However, both implications fail when \mathbb{Q} is replaced with \mathbb{R} .

Example B.2. — Assume $(X, L) = (\mathbb{P}^1, \mathcal{O}(1))$, and consider the toric divisorial norm

 $\chi \coloneqq \min\{\chi_v, \chi_{\text{triv}} + c\} \in \mathcal{N}_{\mathbb{R}}^{\text{div}},$

where $v = \operatorname{ord}_0$ with 0 is the origin in $\mathbb{A}^1 \subset \mathbb{P}^1$, and $c \in [0, 1]$. The corresponding concave function is $g(\alpha) = \min\{\alpha, c\}$ with $\alpha \in P = [0, 1] \subset M_{\mathbb{R}} = \mathbb{R}$, and a simple computation yields

$$g^{\vee}(\xi) = \max\{\xi + c, c\xi + c, 0\}$$

for $v \in N_{\mathbb{R}} = \mathbb{R}$. Using (iv), this shows $\chi \in \mathcal{T}_{\mathbb{R}} \iff c \in \mathbb{Q}$.

Example B.3. — For any $\xi \in N_{\mathbb{R}} \subset X^{\text{lin}}$, χ_{ξ} is of finite type (see Example 1.10). However, χ_{ξ} is divisorial iff $\xi \in N_{\mathbb{Q}}$ (see Example 7.17).

BIBLIOGRAPHY

- V. G. BERKOVICH, Spectral theory and analytic geometry over non-Archimedean fields, Mathematical Surveys and Monographs, vol. 33, American Mathematical Society, 1990, x+169 pages.
- [2] ——, "Étale cohomology for non-Archimedean analytic spaces", Publ. Math., Inst. Hautes Étud. Sci. (1993), no. 78, p. 5-161.
- [3] R. J. BERMAN, S. BOUCKSOM, V. GUEDJ & A. ZERIAHI, "A variational approach to complex Monge-Ampère equations", Publ. Math., Inst. Hautes Étud. Sci. 117 (2013), p. 179-245.
- [4] R. J. BERMAN, S. BOUCKSOM & M. JONSSON, "A variational approach to the Yau-Tian-Donaldson conjecture", J. Am. Math. Soc. 34 (2021), no. 3, p. 605-652.
- [5] H. BLUM & M. JONSSON, "Thresholds, valuations, and K-stability", Adv. Math. 365 (2020), article no. 107062 (57 pages).
- [6] H. BLUM, Y. LIU & L. QI, "Convexity of multiplicities of filtrations on local rings", https://arxiv.org/abs/2208.04902, 2022.
- [7] H. BLUM, Y. LIU & C. XU, "Openness of K-semistability for Fano varieties", Duke Math. J. 171 (2022), no. 13, p. 2753-2797.
- [8] H. BLUM, Y. LIU, C. XU & Z. ZHUANG, "The existence of the Kähler–Ricci soliton degeneration", Forum Math. Pi 11 (2023), article no. e9 (28 pages).
- [9] H. BLUM, Y. LIU & C. ZHOU, "Optimal destabilization of K-unstable Fano varieties via stability thresholds", Geom. Topol. 26 (2022), no. 6, p. 2507-2564.
- [10] H. BLUM & C. XU, "Uniqueness of K-polystable degenerations of Fano varieties", Ann. Math. 190 (2019), no. 2, p. 609-656.
- [11] S. BOSCH, U. GÜNTZER & R. REMMERT, Non-Archimedean analysis. A systematic approach to rigid analytic geometry, Grundlehren der Mathematischen Wissenschaften, vol. 261, Springer, 1984, xii+436 pages.
- [12] S. BOUCKSOM, "Corps d'Okounkov (d'après Okounkov, Lazarsfeld–Mustaţă et Kaveh–Khovanskii)", in Séminaire Bourbaki. Volume 2012/2013. Exposés 1059– 1073. Avec table par noms d'auteurs de 1948/49 à 2012/13, Astérisque, vol. 361, Société Mathématique de France, 2014, p. vii, 1-41.
- [13] S. BOUCKSOM & H. CHEN, "Okounkov bodies of filtered linear series", Compos. Math. 147 (2011), no. 4, p. 1205-1229.

- [14] S. BOUCKSOM & D. ERIKSSON, "Spaces of norms, determinant of cohomology and Fekete points in non-Archimedean geometry", Adv. Math. 378 (2021), article no. 107501 (124 pages).
- [15] S. BOUCKSOM, C. FAVRE & M. JONSSON, "Singular semipositive metrics in non-Archimedean geometry", J. Algebr. Geom. 25 (2016), no. 1, p. 77-139.
- [16] S. BOUCKSOM, W. GUBLER & F. MARTIN, "Differentiability of relative volumes over an arbitrary non-Archimedean field", Int. Math. Res. Not. (2022), no. 8, p. 6214-6242.
- [17] S. BOUCKSOM, T. HISAMOTO & M. JONSSON, "Uniform K-stability, Duistermaat– Heckman measures and singularities of pairs", Ann. Inst. Fourier 67 (2017), no. 2, p. 743-841.
- [18] S. BOUCKSOM & M. JONSSON, "Global pluripotential theory over a trivially valued field", Ann. Fac. Sci. Toulouse, Math. 31 (2022), no. 3, p. 647-836.
- [19] —, "Addendum to the article 'Global pluripotential theory over a trivially valued field", to appear in Ann. Fac. Sci. Toulouse Math., https://arxiv.org/ abs/2206.07183, 2022.
- [20] ——, "A non-Archimedean approach to K-stability, II: divisorial stability and openness", to appear in J. Reine Angew. Math., https://arxiv.org/abs/2206. 09492, 2023.
- [21] ——, "Non-Archimedean Green's functions and Zariski decompositions", https: //arxiv.org/abs/2304.00144, 2023.
- [22] S. BOUCKSOM, A. KÜRONYA, C. MACLEAN & T. SZEMBERG, "Vanishing sequences and Okounkov bodies", Math. Ann. 361 (2015), no. 3-4, p. 811-834.
- [23] F. BRUHAT & J. TITS, "Groupes réductifs sur un corps local", Publ. Math., Inst. Hautes Étud. Sci. (1972), no. 41, p. 5-251.
- [24] J. I. BURGOS GIL, P. PHILIPPON & M. SOMBRA, Arithmetic geometry of toric varieties. Metrics, measures and heights, Astérisque, no. 360, Société Mathématique de France, 2014, vi+222 pages.
- [25] A. CHAMBERT-LOIR, "Mesures et équidistribution sur les espaces de Berkovich", J. Reine Angew. Math. 595 (2006), p. 215-235.
- [26] A. CHAMBERT-LOIR & A. DUCROS, "Formes différentielles réelles et courants sur les espaces de Berkovich", https://arxiv.org/abs/1204.6277, 2012.
- [27] H. CHEN & C. MACLEAN, "Distribution of logarithmic spectra of the equilibrium energy", Manuscr. Math. 146 (2015), no. 3-4, p. 365-394.
- [28] H. CHEN & A. MORIWAKI, "Extension property of semipositive invertible sheaves over a non-archimedean field", Ann. Sc. Norm. Super. Pisa, Cl. Sci. 18 (2018), no. 1, p. 241-282.
- [29] X. CHEN, S. SUN & B. WANG, "Kähler–Ricci flow, Kähler–Einstein metric, and K-stability", Geom. Topol. 22 (2018), no. 6, p. 3145-3173.
- [30] G. CODOGNI, "Tits buildings and K-stability", Proc. Edinb. Math. Soc., II. Ser. 62 (2019), no. 3, p. 799-815.
- [31] T. DARVAS, "The Mabuchi geometry of finite energy classes", Adv. Math. 285 (2015), p. 182-219.
- [32] ——, "The Mabuchi completion of the space of Kähler potentials", Am. J. Math. 139 (2017), no. 5, p. 1275-1313.
- [33] T. DARVAS, E. DI NEZZA & C. H. LU, "L¹ metric geometry of big cohomology classes", Ann. Inst. Fourier 68 (2018), no. 7, p. 3053-3086.
- [34] T. DARVAS, C. H. LU & Y. A. RUBINSTEIN, "Quantization in geometric pluripotential theory", Commun. Pure Appl. Math. 73 (2020), no. 5, p. 1100-1138.
- [35] T. DARVAS & M. XIA, "The closures of test configurations and algebraic singularity types", Adv. Math. 397 (2022), article no. 108198 (56 pages).

- [36] R. DERVAN, "Uniform stability of twisted constant scalar curvature Kähler metrics", Int. Math. Res. Not. (2016), no. 15, p. 4728-4783.
- [37] R. DERVAN & E. LEGENDRE, "Valuative stability of polarised varieties", Math. Ann. 385 (2023), no. 1-2, p. 357-391.
- [38] R. DERVAN & G. SZÉKELYHIDI, "The Kähler-Ricci flow and optimal degenerations", J. Differ. Geom. 116 (2020), no. 1, p. 187-203.
- [39] S. K. DONALDSON, "Scalar curvature and stability of toric varieties", J. Differ. Geom. 62 (2002), no. 2, p. 289-349.
- [40] A. DUCROS, "Les espaces de Berkovich sont excellents", Ann. Inst. Fourier 59 (2009), no. 4, p. 1443-1552.
- [41] L. EIN, R. LAZARSFELD & K. E. SMITH, "Uniform approximation of Abhyankar valuation ideals in smooth function fields", Am. J. Math. 125 (2003), no. 2, p. 409-440.
- [42] Y. FANG, "Study of positively metrized line bundles over a non-Archimedean field via holomorphic convexity", PhD Thesis, University of Paris, 2019.
- [43] K. FUJITA, "Uniform K-stability and plt blowups of log Fano pairs", Kyoto J. Math. 59 (2019), no. 2, p. 399-418.
- [44] ——, "A valuative criterion for uniform K-stability of Q-Fano varieties", J. Reine Angew. Math. 751 (2019), p. 309-338.
- [45] W. FULTON, Introduction to toric varieties. The William H. Roever Lectures in Geometry, Annals of Mathematics Studies, vol. 131, Princeton University Press, 1993, The William H. Roever Lectures in Geometry, xii+157 pages.
- [46] D. GALE, V. KLEE & R. T. ROCKAFELLAR, "Convex functions on convex polytopes", Proc. Am. Math. Soc. 19 (1968), p. 867-873.
- [47] W. GUBLER, "Local heights of subvarieties over non-Archimedean fields", J. Reine Angew. Math. 498 (1998), p. 61-113.
- [48] W. GUBLER, P. JELL, K. KÜNNEMANN & F. MARTIN, "Continuity of plurisubharmonic envelopes in non-archimedean geometry and test ideals. With an appendix by José Ignacio Burgos Gil and Martín Sombra", Ann. Inst. Fourier 69 (2019), no. 5, p. 2331-2376.
- [49] J. HAN & C. LI, "Algebraic uniqueness of Kähler-Ricci flow limits and optimal degenerations of Fano varieties", https://arxiv.org/abs/2009.01010, 2020.
- [50] E. INOUE, "Entropies in μ-framework of canonical metrics and K-stability, II Nonarchimedean aspect: non-archimedean μ-entropy and μK-semistability", https:// arxiv.org/abs/2202.12168, 2022.
- [51] A. KÜRONYA, "A divisorial valuation with irrational volume", J. Algebra 262 (2003), no. 2, p. 413-423.
- [52] R. LAZARSFELD & M. MUSTAŢĂ, "Convex bodies associated to linear series", Ann. Sci. Éc. Norm. Supér. 42 (2009), no. 5, p. 783-835.
- [53] C. LI, "K-semistability is equivariant volume minimization", Duke Math. J. 166 (2017), no. 16, p. 3147-3218.
- [54] —, "G-uniform stability and Kähler-Einstein metrics on Fano varieties", Invent. Math. 227 (2022), no. 2, p. 661-744.
- [55] —, "Geodesic rays and stability in the cscK problem", Ann. Sci. Éc. Norm. Supér. 55 (2022), no. 6, p. 1529-1574.
- [56] ——, "K-stability and Fujita approximation", in Birational geometry, Kähler-Einstein metrics and degenerations, Springer Proceedings in Mathematics & Statistics, vol. 409, Springer, 2023, p. 545-566.
- [57] C. LI & C. XU, "Stability of valuations and Kollár components", J. Eur. Math. Soc. 22 (2020), no. 8, p. 2573-2627.

- [58] Y. LIU, "Openness of uniformly valuative stability on the Kähler cone of projective manifolds", Math. Z. 303 (2023), no. 2, article no. 52 (26 pages).
- [59] D. MCKINNON & M. ROTH, "Seshadri constants, diophantine approximation, and Roth's theorem for arbitrary varieties", *Invent. Math.* 200 (2015), no. 2, p. 513-583.
- [60] D. MUMFORD, J. FOGARTY & F. KIRWAN, Geometric invariant theory, third ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 2. Folge, vol. 34, Springer, 1994, xiv+292 pages.
- [61] Y. ODAKA, "On parametrization, optimization and triviality of test configurations", Proc. Am. Math. Soc. 143 (2015), no. 1, p. 25-33.
- [62] J. POINEAU, "Les espaces de Berkovich sont angéliques", Bull. Soc. Math. Fr. 141 (2013), no. 2, p. 267-297.
- [63] R. REBOULET, "The asymptotic Fubini–Study operator over general non-Archimedean fields", Math. Z. 299 (2021), no. 3-4, p. 2341-2378.
- [64] ——, "Plurisubharmonic geodesics in spaces of non-Archimedean metrics of finite energy", J. Reine Angew. Math. 793 (2022), p. 59-103.
- [65] G. SZÉKELYHIDI, "Filtrations and test-configurations. With an appendix by Sebastien Boucksom", Math. Ann. 362 (2015), no. 1-2, p. 451-484.
- [66] M. TEMKIN, "On local properties of non-Archimedean analytic spaces. II", Isr. J. Math. 140 (2004), p. 1-27.
- [67] ——, "Introduction to Berkovich analytic spaces", in Berkovich spaces and applications, Lecture Notes in Mathematics, vol. 2119, Springer, 2015, p. 3-66.
- [68] D. WITT NYSTRÖM, "Test configurations and Okounkov bodies", Compos. Math. 148 (2012), no. 6, p. 1736-1756.
- [69] K. ZHANG, "Valuative invariants with higher moments", J. Geom. Anal. 32 (2022), no. 1, article no. 10 (33 pages).
- [70] S. ZHANG, "Positive line bundles on arithmetic varieties", J. Am. Math. Soc. 8 (1995), no. 1, p. 187-221.

Manuscrit reçu le 23 juillet 2021, révisé le 16 juin 2022, accepté le 4 août 2023.

Sébastien BOUCKSOM Sorbonne Université and Université Paris Cité CNRS, IMJ-PRG F-75005 Paris (France) sebastien.boucksom@imj-prg.fr

Mattias JONSSON Dept of Mathematics University of Michigan Ann Arbor, MI 48109-1043 (USA) mattiasj@umich.edu