



ANNALES DE L'INSTITUT FOURIER

Simon ALLAIS

**Morse estimates for translated points on unit tangent
bundles**

Article à paraître, mis en ligne le 22 septembre 2025, 20 p.

Article mis à disposition par son auteur selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE



<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les *Annales de l'Institut Fourier* sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org e-ISSN : 1777-5310

MORSE ESTIMATES FOR TRANSLATED POINTS ON UNIT TANGENT BUNDLES

by Simon ALLAIS (*)

ABSTRACT. — In this article, we study conjectures of Sandon on the minimal number of translated points of contactomorphisms in the special case of the unit tangent bundle of a Riemannian manifold. We restrict ourselves to contactomorphisms of a unit tangent bundle that lift diffeomorphisms of the base homotopic to the identity. We prove that there exist sequences (p_n, t_n) where p_n is a translated point of time-shift t_n with $t_n \rightarrow +\infty$ for a large class of manifolds. In the case of Zoll–Riemannian manifolds, we also prove estimates relating the number of translated points to either the sum of the Betti numbers of the bundle under a generic assumption or its cuplength under a C^0 -closedness assumption.

RÉSUMÉ. — Dans cet article, nous étudions des conjectures de Sandon concernant le nombre minimal de points translatés dans le cas particulier du fibré tangent unitaire d’une variété riemannienne. Nous restreignons aux contactomorphismes du fibré tangent unitaire relevant les difféomorphismes de la base homotopes à l’identité. Nous montrons qu’il existe des suites (p_n, t_n) où p_n est un point translaté de temps de décalage t_n avec $t_n \rightarrow +\infty$ pour une grande classe de variétés. Dans le cas des variétés riemanniennes–Zoll, nous montrons aussi des inégalités entre le nombre de points translatés et la somme des nombres de Betti du fibré sous une hypothèse générique ou la *cuplength* sous une hypothèse de proximité C^0 .

1. Introduction

In this article, we study conjectures essentially due to Sandon on the minimal number of translated points in the special case of the unit tangent bundle of a Riemannian manifold. Let us recall the definition of translated points. Let (V^{2n-1}, α) be a cooriented contact manifold with a fixed

Keywords: Arnold conjecture, Chas–Sullivan product, geodesics, Lusternik–Schnirelmann theory, Morse theory, translated points of contactomorphisms, unit tangent bundles, Zoll metrics.

2020 *Mathematics Subject Classification:* 53C22, 53D25, 57R17, 58E10.

(*) The first author was partially funded by the postdoctoral fellowships of the *Fondation Sciences Mathématiques de Paris*.

contact form α (i.e. $\alpha \wedge (d\alpha)^{n-1}$ does not vanish). A contactomorphism $\varphi \in \text{Cont}(V, \alpha)$ is a diffeomorphism of V such that $\varphi^*\alpha = e^g\alpha$ for some $g : V \rightarrow \mathbb{R}$. A point $p \in V$ is a discriminant point of φ if and only if it is fixed by φ and $g(p) = 0$ (this definition does not depend on the choice of the contact form associated with $\ker \alpha$). Let (ϕ_t^α) be the Reeb flow of α . A point $p \in V$ is a translated point of φ if and only if it is a discriminant point of $\phi_{-t}^\alpha \circ \varphi$ for some $t \in \mathbb{R}$ called a time-shift of p . Similarly to the Hamiltonian case, using a Weinstein neighborhood of the graph of the identity, one can prove that for every contactomorphism of a closed contact manifold $\varphi \in \text{Cont}(V, \alpha)$ which is C^1 -close to the identity,

$$(1.1) \quad \#\{p \in V \mid p \text{ is a translated point of } \varphi\} \geq \min_{f \in C^\infty(V, \mathbb{R})} \# \text{Crit}(f),$$

where $\text{Crit}(f)$ denotes the set of critical points of f [24]. Moreover, if the Reeb flow is periodic, this inequality is sharp (see e.g. the introduction of [3]). In [24], Sandon proved that this inequality still holds without the “ C^1 -close” assumption, as long as φ is isotopic to the identity in the case of the real projective spaces $\mathbb{RP}^{2n-1} = \mathbb{S}^{2n-1}/(z \sim -z)$ endowed with the contact form induced by $\alpha_0 = \frac{1}{2}(xdy - ydx) \in \Omega^1(\mathbb{S}^{2n-1})$. Therefore, Sandon asked the following question, where $\text{Cont}_0(V, \alpha)$ denotes the set of contactomorphisms isotopic to the identity.

QUESTION 1.1. — *Given a closed contact manifold endowed with a contact form (V, α) , does every $\varphi \in \text{Cont}_0(V, \alpha)$ satisfy (1.1)?*

Similarly to the Arnol’d conjecture on fixed points of Hamiltonian symplectomorphisms, one can ask weaker estimates on the number of translated points, using the cup-length estimate or the category estimate of Lusternik–Schnirelmann theory. Another variation of Question 1.1 can be asked for the non-degenerate contactomorphisms (see below), replacing the estimate on the minimal number of critical points of any $f : M \rightarrow \mathbb{R}$ by an estimate on the minimal number of critical points a Morse map can have. As mentioned earlier, Sandon proved that Question 1.1 is true for $(\mathbb{RP}^{2n-1}, \alpha_0)$ [24]. We proved that this is also the case for every quotient $L_k^{2n-1}(w)$ of $(\mathbb{S}^{2n-1}, \alpha_0)$ by free $\mathbb{Z}/k\mathbb{Z}$ -actions of the form $(z_j) \mapsto (e^{2i\pi w_j/k} z_j)$ (with $k \geq 2$) in [3], improving an estimate of Granja–Karshon–Pabiniak–Sandon [12]. A Morse estimate for generic contactomorphisms was also given by Albers–Fuchs–Merry [2] completed by Meiwes–Naef [19] in the case of hypertight contact manifolds (i.e. such that all Reeb orbits are non-contractible for some contact form supporting the contact structure).

This first question seems better suited for contact manifolds all of whose Reeb orbits are closed: the so-called Besse contact manifolds. A result of

Granja–Karshon–Pabiniak–Sandon suggests a second question that could cover a greater class of contact manifolds.

QUESTION 1.2. — *Given a closed contact manifold endowed with a contact form (V, α) , does every $\varphi \in \text{Cont}_0(V, \alpha)$ possess a sequence of couples $(p_n, t_n) \in V \times \mathbb{R}$ with $t_n \rightarrow +\infty$ such that p_n is a translated point of φ with time-shift t_n ?*

Let us remark that the Weinstein conjecture is satisfied by the (V, α) 's satisfying the conclusion of Question 1.2 (by taking $\varphi = \text{id}$). Let us further remark that a Besse contact manifold answering Question 1.1 positively also answers Question 1.2 positively. Granja–Karshon–Pabiniak–Sandon answered positively Question 1.2 for $(L_k^{2n-1}(w), \alpha)$, α being any contact form supporting the same contact structure as the quotient of α_0 .

In this paper, we want to give motivations that Question 1.1 should be answered positively for unit tangent bundles of Riemannian manifolds all of whose geodesics are closed and of same prime length whereas Question 1.2 should be answered positively for every unit tangent bundle. In order to do so, we will prove that this is indeed the case in a weak sense for a subclass of contactomorphisms of SM: the contactomorphic lift of the diffeomorphisms of M .

Let M be a Riemannian manifold, its unit tangent bundle SM is a contact manifold for the contact form α :

$$\alpha_{(x,v)} \cdot \xi = \langle v, d\pi_x \cdot \xi \rangle, \quad \forall (x, v) \in \text{SM}, \quad \forall \xi \in T_{(x,v)} \text{SM},$$

where $\pi : \text{SM} \rightarrow M$ is the bundle map and $\langle \cdot, \cdot \rangle$ is the Riemannian metric. Given a diffeomorphism $f : M \rightarrow M$, we denote $\tilde{f} : \text{SM} \rightarrow \text{SM}$ the associated contactomorphism

$$\tilde{f}(x, v) := \left(f(x), \frac{df_x^{-T} \cdot v}{\|df_x^{-T} \cdot v\|} \right), \quad \forall (x, v) \in \text{SM},$$

where df^{-T} denotes the inverse of the adjoint df^T . We will study the minimal number of translated points of \tilde{f} for f homotopic to the identity. The notion of translated points of \tilde{f} with time-shift t can be naturally generalized to smooth maps $f : M \rightarrow M$ (and not only diffeomorphisms) in such a way that, in particular, a translated point of a diffeomorphism $f : M \rightarrow M$ with time-shift t is exactly a translated point of the contactomorphism \tilde{f} with time-shift t .

DEFINITION 1.3. — A point $(x, v) \in \text{SM}$ is a translated point of $f : M \rightarrow M$ with time-shift $t \neq 0$ if there exists a geodesic $\gamma : [0, 1] \rightarrow \mathbb{R}$ of length $|t|$ such that

$$\begin{cases} \dot{\gamma}(0) = v \|\dot{\gamma}(0)\| & \text{and} & df_x^T \cdot \dot{\gamma}(1) = \dot{\gamma}(0) & \text{when } t > 0, \\ \dot{\gamma}(1) = v \|\dot{\gamma}(1)\| & \text{and} & df_x^T \cdot \dot{\gamma}(0) = \dot{\gamma}(1) & \text{when } t < 0; \end{cases}$$

it is a translated point with time-shift $t = 0$ if $df_x^T \cdot v = v$ (in particular, $f(x) = x$).

The correspondence between translated points of \widetilde{f} and translated points of f is due to the fact that the Reeb flow of SM is the geodesic flow (see e.g. [22, Section 1.3.3]). We give the following partial answer to Question 1.2.

THEOREM 1.4. — Let M be a closed Riemannian manifold that has a finite cover \widetilde{M} , the singular homology group $H_*(\Lambda\widetilde{M})$ of the free loop space of which is not finitely generated. Every smooth map $f : M \rightarrow M$ homotopic to the identity admits a sequence of couples $(p_n, t_n) \in \text{SM} \times (0, +\infty)$ with $t_n \rightarrow +\infty$ such that p_n is a translated point of f with time-shift t_n .

The assumption on M that $H_*(\Lambda\widetilde{M})$ is not finitely generated is satisfied by a large class of closed Riemannian manifolds (in fact, the author does not know if there are counter-examples). On the one hand, when the number of conjugacy classes of $\pi_1(M)$ is infinite, the group $H_0(\Lambda M)$ is already not finitely generated. On the other hand, when $\pi_1(M)$ is finite, Vigué-Poirrier and Sullivan proved that $H_*(\Lambda\widetilde{M})$ is not finitely generated for the universal cover \widetilde{M} [25]. Therefore, the only possible counter-examples are among the closed manifolds that have an infinite fundamental group with a finite number of conjugacy classes.

Our most satisfying answer to Question 1.1 concerns the non-degenerate case. Let us first discuss the notion of non-degenerate translated points. Let (G_t) be the geodesic flow of TM and, for a diffeomorphism $f : M \rightarrow M$, let $\widehat{f} : \text{TM} \rightarrow \text{TM}$ be its symplectic lift $\widehat{f}(x, v) = (f(x), df_x^{-T} \cdot v)$. In our particular case, a translated point $p \in \text{SM}$ of a contactomorphic lift \widetilde{f} of a diffeomorphism $f : M \rightarrow M$ is non-degenerate for its time-shift $t \in \mathbb{R}$ if and only if $d(G_{-t} \circ \widehat{f})(p)$ does not have 1 as an eigenvalue. One can check that this definition coincides with the definition given by Sandon in the general setting [24]. Similarly to the definition of translated points, one can extend this notion to any smooth map $M \rightarrow M$ (for simplicity, we only give the definition for positive time-shifts).

DEFINITION 1.5. — A translated point (x, v) of $f : M \rightarrow M$ is non-degenerate for the time-shift $t > 0$ if the subspace of the Jacobi fields J along the associated geodesic γ satisfying

$$\begin{cases} J(1) = df \cdot J(0), \\ (d^2f \cdot J(0))^T \cdot \dot{\gamma}(1) = \dot{J}(0) - df_x^T \cdot \dot{J}(1), \end{cases}$$

is reduced to 0 (the linear morphism $d^2f \cdot J(0)$ denotes $u \mapsto d^2f[J(0), u]$, see Section 2.4).

The equivalence of both definitions in the case of diffeomorphisms is proven in Proposition 2.8. We give the following partial answer to Question 1.1 in the non-degenerate case. Let us recall that a Riemannian manifold M is called Zoll (of length ℓ) if all its geodesics are closed and of the same prime length (equal to ℓ).

THEOREM 1.6. — Let M be a closed Zoll–Riemannian manifold and let $R := \mathbb{Z}$ if M is orientable and $R := \mathbb{Z}/2\mathbb{Z}$ otherwise. For every smooth map $f : M \rightarrow M$ homotopic to the identity with finitely many translated points in SM all of which are non-degenerate, the number of translated points is not less than $\sum_j \beta_j(\text{SM}; R)$, where $\beta_j(\text{SM}; R) = \text{rank } H_j(\text{SM}; R)$ denotes the j^{th} Betti number of SM.

In the degenerate case, our result is less satisfying as it requires a C^0 -closeness assumption. Let us recall that the cup-length $\text{CL}(X; R) \in \mathbb{N}$ of a space X is the maximal k such that $u_1 \smile \cdots \smile u_k \neq 0$ for some non-zero $u_j \in H^*(X; R)$ of positive degree. By a H^1 -homotopy (f_s) of maps $M \rightarrow M$, we will mean that $t \mapsto f_t(x)$ is in the Sobolev space $H^1([0, 1], M)$ for all $x \in M$. The set of time-shifts of $f : M \rightarrow M$ will denote the set of $t \in \mathbb{R}$ that are time-shifts of a translated point; when M is Zoll of length ℓ , this set is $\ell\mathbb{Z}$ -invariant so it can be seen as a subset of $\mathbb{R}/\ell\mathbb{Z}$.

THEOREM 1.7. — Let M be a closed Zoll–Riemannian manifold of length ℓ and let $R := \mathbb{Z}$ if M is orientable and $R := \mathbb{Z}/2\mathbb{Z}$ otherwise. Let (f_s) be a H^1 -homotopy of maps $M \rightarrow M$ such that $f_0 = \text{id}$ and

$$\int_0^1 \|\partial_t f_t(x)\|^2 dt < \left(\frac{\ell}{2}\right)^2, \quad \forall x \in M.$$

If the number of time-shifts of f_1 seen in $\mathbb{R}/\ell\mathbb{Z}$ is less than $1 + \text{CL}(\text{SM}; R)$, then f_1 has infinitely many translated points. In particular, the number of translated points of f_1 is not less than $1 + \text{CL}(\text{SM}; R)$.

The only closed Zoll–Riemannian manifolds known by the author are diffeomorphic to the compact rank-one symmetric spaces: \mathbb{S}^n , \mathbb{RP}^n , \mathbb{CP}^n , \mathbb{HP}^n

and CaP^2 (see however [5] for examples of manifolds with exotic structures all of whose geodesics starting from a special point go back to this point at the same length). According to a result of Bott and Samelson, every Zoll–Riemannian manifold has a cohomology ring isomorphic to the cohomology ring of one of these spaces [7, 23]. Let us refer to [6] for a comprehensive introduction to the theory of Zoll and Besse Riemannian manifolds. Let us also point out that the study of Besse Riemannian manifolds and more generally of Besse contact forms from the variational viewpoint has recently known significant advances [1, 10, 18].

The proofs of this article are based on a rather classical variational principle that goes back to Grove [13]. Indeed, we remark in Section 2.1 that geodesics corresponding to translated points of positive time-shift are exactly the critical points of the restriction of the energy functional to paths $\gamma : [0, 1] \rightarrow M$ such that $f(\gamma(0)) = \gamma(1)$. This variational principle was initially applied in the specific case where f is an isometry in order to study isometry-invariant geodesics [4, 13, 14, 15, 16, 17].

Theorem 1.4 is a direct application of Morse theory. Theorems 1.6 and 1.7 are more subtle consequences of Morse and Lusternik–Schnirelmann theories. In order to translate the symmetry $(p, t) \mapsto (p, t + \ell)$ of the set of couples (translated point, associated time-shift), we make use of the Chas–Sullivan product [9]. More precisely, we apply the results of Goresky–Hingston concerning the product-structure of the homology groups of the free loop space of Zoll–Riemannian manifolds [11].

Organization of the paper

In Section 2, we study the variational principle satisfied by translated points of positive time-shifts and prove Theorem 1.4. In Section 3, we study the special case of Zoll–Riemannian manifolds and prove Theorems 1.6 and 1.7.

Acknowledgment

I am grateful to Margherita Sandon and to my former advisor Marco Mazzucchelli for their supports and fruitful discussions and suggestions.

2. The variational principle

2.1. The variational principle

Let M be a closed Riemannian manifold. We will denote $\text{PM} := H^1([0, 1], M)$ the path space of M endowed with its usual structure of Hilbert manifold (here H^1 denotes the Sobolev space also denoted $W^{1,2}$). Given $f : M \rightarrow M$, let

$$\Lambda(f) := \{\gamma \in \text{PM} \mid \gamma(1) = f(\gamma(0))\};$$

in particular $\Lambda(\text{id}) = \Lambda M$ is the free loop space of M . We will also use the notation Λ_f for $\Lambda(f)$ and Λ for ΛM . This space was already introduced in [13] as $\Lambda_{G(f)}M$, we refer to this article for details about the properties recalled here. The space $\Lambda(f)$ is a Hilbert-submanifold of PM as $\gamma \mapsto (\gamma(0), \gamma(1))$ is a submersion. Given $\gamma \in \Lambda(f)$, a vector $U \in T_\gamma \Lambda(f)$ is a H^1 -vector field along γ satisfying $\text{df} \cdot U(0) = U(1)$. Let us denote $E : \text{PM} \rightarrow \mathbb{R}$ the energy functional,

$$E(\gamma) := \int_0^1 \|\dot{\gamma}\|^2 dt,$$

and $E_f : \Lambda(f) \rightarrow \mathbb{R}$ the restriction to $\Lambda(f)$ for $f : M \rightarrow M$. According to [13, Theorem 2.4], the functional E_f satisfies the Palais–Smale condition.

Let us denote ∇ the Levi-Civita covariant derivative of M . Taken along a curve $t \mapsto c(t)$, the covariant derivative of the vector field X will be either denoted $\nabla_{\dot{c}}X$, $\frac{DX}{dt}$ or \dot{X} (this last notation is reserved to the time variable t). By convention, a geodesic will always mean a geodesic with constant speed: $\gamma \in \text{PM}$ is a geodesic if and only if

$$\nabla_{\dot{\gamma}}\dot{\gamma} = \ddot{\gamma} = 0.$$

PROPOSITION 2.1. — *Given $f : M \rightarrow M$, $\gamma \in \Lambda(f)$ is a critical point of E_f if and only if γ is a geodesic and $\text{df}_{\gamma(0)}^T \cdot \dot{\gamma}(1) = \dot{\gamma}(0)$.*

Proof. — Let $\gamma \in \Lambda(f)$, let $U \in T_\gamma \Lambda(f)$ and let (γ_u) be a smooth family in $\Lambda(f)$ such that $U = \partial_u \gamma_u$ (the derivative being taken at $u = 0$). Then

$$\frac{1}{2} \text{d}E_f(\gamma) \cdot U = \int_0^1 \left\langle \frac{D}{\partial u} \dot{\gamma}_u, \dot{\gamma} \right\rangle dt = \int_0^1 \langle \dot{U}, \dot{\gamma} \rangle dt = [\langle U, \dot{\gamma} \rangle]_0^1 - \int_0^1 \langle U, \ddot{\gamma} \rangle dt.$$

As $\text{df} \cdot U(0) = U(1)$, one has

$$[\langle U, \dot{\gamma} \rangle]_0^1 = \langle U(0), \text{df}^T \cdot \dot{\gamma}(1) - \dot{\gamma}(0) \rangle.$$

The identity $\text{d}E_f(\gamma) \cdot U = 0$ for the vector fields along γ such that $U(0) = 0$ implies that $\ddot{\gamma} = 0$, i.e. γ is a geodesic. The identity $\text{d}E_f(\gamma) \cdot U = 0$ for

every $U \in T_\gamma \Lambda(f)$ then implies that $df^T \cdot \dot{\gamma}(1) = \dot{\gamma}(0)$ (although we will often write df^T without mention of the base point of M at which we take the adjoint, we point out that this notation can be ambiguous as df^T is defined on $f^* \text{TM}$ rather than TM). \square

COROLLARY 2.2. — *For every diffeomorphism $f : M \rightarrow M$, critical points of E_f with energy value $e > 0$ are in bijection with translated points (x, v) of \tilde{f} with time-shift \sqrt{e} .*

As the correspondence suggests, we will rather be interested in the values of the functional $P_f := \sqrt{E_f}$ (which is C^1 away from $\{E_f = 0\}$) as they correspond to time-shifts at critical points with positive critical values. For every $\lambda \geq 0$, let $\Lambda_f^{<\lambda}$ and $\Lambda_f^{\leq \lambda}$ be respectively the subsets $\{P_f < \lambda\}$ and $\{P_f \leq \lambda\}$ of $\Lambda(f)$; when λ^2 is not a critical value of E_f , these subsets are submanifolds (with boundary in the second case).

2.2. The topology of $\Lambda(f)$

Let us first extend the join of loops defined in [11, Section 2.3] to the spaces $\Lambda(f)$, for $f : M \rightarrow M$ smooth, in the obvious way. Let us define the submanifold

$$\text{PM} \times_M \text{PM} := \{(\alpha, \beta) \in \text{PM} \times \text{PM} \mid \alpha(1) = \beta(0)\}$$

as well as $\Lambda(f) \times_M \Lambda(g) := (\Lambda(f) \times \Lambda(g)) \cap (\text{PM} \times_M \text{PM})$. The concatenation $\phi : \text{PM} \times_M \text{PM} \rightarrow \text{PM}$ will denote the continuous map defined as follows. Given $(\alpha, \beta) \in \text{PM} \times_M \text{PM}$, if $E(\alpha) = E(\beta) = 0$, both paths α and β are constant $\equiv p$ and we set $\phi(\alpha, \beta) \equiv p$, otherwise

$$\phi(\alpha, \beta)(t) := \begin{cases} \alpha\left(\frac{t}{s}\right), & t \in [0, s] \\ \beta\left(\frac{t-s}{1-s}\right), & t \in [s, 1] \end{cases}, \quad \text{where } s = \frac{\sqrt{E(\alpha)}}{\sqrt{E(\alpha)} + \sqrt{E(\beta)}}.$$

This map satisfies

$$(2.1) \quad \sqrt{E(\phi(\alpha, \beta))} = \sqrt{E(\alpha)} + \sqrt{E(\beta)}.$$

The concatenation ϕ is associative: for every $\alpha, \beta, \gamma \in \text{PM}$ such that $\alpha(1) = \beta(0)$ and $\beta(1) = \gamma(0)$,

$$(2.2) \quad \phi(\phi(\alpha, \beta), \gamma) = \phi(\alpha, \phi(\beta, \gamma)).$$

By restriction, one gets a continuous map $\phi : \Lambda(f) \times_M \Lambda(g) \rightarrow \Lambda(g \circ f)$ satisfying $P_{g \circ f}(\phi(\alpha, \beta)) = P_f(\alpha) + P_g(\beta)$, for every $f, g : M \rightarrow M$.

Let $(f_s)_{s \in [0,1]}$ be a H^1 -homotopy of smooth maps $M \rightarrow M$, i.e. such that $s \mapsto f_s(x)$ is in PM for all $x \in M$. It induces the following map $\tau(f_s) : \Lambda(f_0) \rightarrow \Lambda(f_1)$,

$$\tau(f_s)(\alpha) := \phi(\alpha, t \mapsto f_t(\alpha(0))).$$

Let

$$\delta(f_s) := \sup_{x \in M} \sqrt{E(t \mapsto f_t(x))},$$

then $\tau(f_s) : \Lambda_{f_0}^{\leq \lambda} \rightarrow \Lambda_{f_1}^{\leq \lambda + \delta(f_s)}$ according to (2.1). Following the same lines as [13, Lemma 3.6] for the more usual concatenation of continuous paths, one proves the following counterpart for ϕ .

LEMMA 2.3 ([13, Lemma 3.6]). — *Let $(f_s)_{s \in [0,1]}$ be a homotopy of smooth maps $M \rightarrow M$ such that $s \mapsto f_s(x)$ is in PM for all $x \in M$. Then $\tau(f_s)$ and $\tau(f_{1-s})$ are homotopy inverses.*

In particular, when $f : M \rightarrow M$ is homotopic to identity, $\Lambda(f)$ is homotopy equivalent to the free loop space ΛM whereas when $f : M \rightarrow M$ is homotopic to a constant, $\Lambda(f)$ is homotopy equivalent to a point.

COROLLARY 2.4. — *Under the hypothesis of Lemma 2.3, the maps $\tau(f_s)$ and $\tau(f_{1-s})$ induce a $\delta(f_s)$ -interleaving between the persistence modules $t \mapsto H_*(\Lambda_{f_j}^{\leq t})$, $j \in \{0, 1\}$, i.e. denoting, for all $t \geq 0$,*

$$\begin{cases} \tau^t := \tau(f_s)_* : H_*(\Lambda_{f_0}^{\leq t}) \rightarrow H_*(\Lambda_{f_1}^{\leq t + \delta(f_s)}), \\ \bar{\tau}^t := \tau(f_{1-s})_* : H_*(\Lambda_{f_1}^{\leq t}) \rightarrow H_*(\Lambda_{f_0}^{\leq t + \delta(f_s)}), \end{cases}$$

the respective morphisms $\bar{\tau}^{t+\delta(f_s)} \circ \tau^t$ and $\tau^{t+\delta(f_s)} \circ \bar{\tau}^t$ coincide with the morphisms induced by the inclusions $\Lambda_{f_j}^{\leq t} \hookrightarrow \Lambda_{f_j}^{\leq t+2\delta(f_s)}$ for $j = 0$ and $j = 1$ respectively (the same is true for $t \mapsto H_(\Lambda_{f_j}^{\leq t})$).*

2.3. Proof of Theorem 1.4

The following fact is well-known to the experts when $f = \text{id}$ (and this case will be enough for us).

LEMMA 2.5. — *Given $f : M \rightarrow M$, for every $\lambda \geq 0$, the image of the inclusion morphism $H_*(\Lambda_f^{\leq \lambda}) \rightarrow H_*(\Lambda_f)$ is finitely generated.*

Proof. — Let $f : M \rightarrow M$. It is enough to prove that for every $\lambda \geq 0$, the inclusion morphism $H_*(\Lambda_f^{\leq \lambda}) \rightarrow H_*(\Lambda_f^{\leq K})$ is finitely generated for some $K \geq \lambda$. Let $K > 0$ be a regular value of E_f . Using a subspace of broken geodesics, one can retract $\Lambda_f^{\leq K}$ on a finite-dimensional submanifold N with a deformation retraction $r : \Lambda_f^{\leq K} \rightarrow N$ such that $E_f \circ r \leq E_f$ and the critical points of E_f are exactly the critical points of $g := E_f \circ r$ (see e.g. [20, Section 16]). Therefore, the desired result boils down to proving that the image of the homology morphism induced by $\{g \leq \lambda\} \hookrightarrow N$ has a finitely generated image for $g : N \rightarrow \mathbb{R}$ a map on a finite-dimensional manifold satisfying the Palais–Smale condition. As this property is C^0 -open, one can assume that g is a Morse map satisfying the Palais–Smale condition (one can adapt indeed the proof of [21, Theorem 2.7] using the fact that the critical set of g in $\{g \leq \lambda\}$ is compact and the norm of dg is uniformly bounded from below outside any neighborhood of this set). The Palais–Smale condition then implies that $\{g \leq \lambda\}$ contains only a finite number $m \in \mathbb{N}$ of critical points (which are non-degenerate), so $H_*(\{g \leq \lambda\})$ can be generated by m elements. \square

Proof of Theorem 1.4. — Let us first prove that one can assume $M = \widetilde{M}$. Let us assume that the theorem is true for a finite Riemannian cover $q : \widetilde{M} \rightarrow M$ satisfying the homological hypothesis. Let $f = f_1 : M \rightarrow M$ be homotopic to the id $= f_0$ through (f_t) . By applying the homotopy lifting property to $(f_t \circ q)$ one gets a homotopy (\widetilde{f}_t) in $\widetilde{M} \rightarrow \widetilde{M}$ from the identity to a map \widetilde{f}_1 commuting with f . If (\widetilde{p}_n, t_n) is a sequence of $\widetilde{M} \times \mathbb{R}$ satisfying the conclusion of the theorem for \widetilde{f}_1 , then $(q(\widetilde{p}_n), t_n)$ is the desired sequence for f .

Let us now assume that $M = \widetilde{M}$. Let $f : M \rightarrow M$ be a smooth map homotopic to the identity. According to Lemma 2.3, $H_*(\Lambda(f))$ is isomorphic to $H_*(\Lambda M)$ which is not finitely generated. Therefore, Lemma 2.5 implies that $H_*(\Lambda_f^{\leq \lambda}) \rightarrow H_*(\Lambda_f)$ is never onto for $\lambda \in [0, +\infty)$. According to the Morse deformation lemma, there exists a sequence (t_n) of positive critical values with $t_n \rightarrow +\infty$. The conclusion follows from Corollary 2.2. \square

2.4. Non-degeneracy of a translated point

In this section, we show the equivalence between the non-degeneracy of a translated point in the sense of contact geometry and the non-degeneracy of the associated critical point of E_f .

Let us denote R the Riemann tensor defined by

$$R(X, Y) := [\nabla_X, \nabla_Y] - \nabla_{[X, Y]},$$

so that a vector field J along γ is a Jacobi field if and only if

$$\ddot{J} = R(\dot{\gamma}, J)\dot{\gamma}.$$

Let us also recall that the second derivative d^2f of a map $f : M \rightarrow N$ is well-defined as $\nabla(df)$, i.e. by the tensorial expression: for all vector fields X, Y ,

$$d^2f[X, Y] := X \cdot (Y \cdot f) - df \cdot \nabla_X Y.$$

Let us recall that the Hessian $d^2E_f(\gamma)$ of the energy functional is well-defined at a critical point γ (see e.g. [20, Section 13]).

PROPOSITION 2.6. — *Let $\gamma \in \Lambda(f)$ be a critical point of E_f , for every $U, V \in T_\gamma \Lambda(f)$,*

$$\begin{aligned} \frac{1}{2}d^2E_f(\gamma)[U, V] \\ = \int_0^1 [R(V, \dot{\gamma})U, \dot{\gamma}] + \langle \dot{U}, \dot{V} \rangle dt + \langle d^2f_{\gamma(0)}[U(0), V(0)], \dot{\gamma}(1) \rangle. \end{aligned}$$

Proof. — Let $U, V \in T_\gamma \Lambda(f)$ and let $(\gamma_{u,v})$ be a smooth family of $\Lambda(f)$ such that $\gamma_{0,0} = \gamma$, $U = \partial_u \gamma_{u,0}$, $V = \partial_v \gamma_{0,v}$ (the partial derivatives being taken at 0). Then

$$\begin{aligned} \frac{1}{2}d^2E_f(\gamma)[U, V] &= \int_0^1 \frac{\partial}{\partial v} \left\langle \frac{D}{\partial u} \dot{\gamma}_{u,v}, \dot{\gamma}_{0,v} \right\rangle dt \\ &= \int_0^1 \left[\left\langle \frac{D^2}{\partial v \partial u} \dot{\gamma}_{u,v}, \dot{\gamma} \right\rangle + \left\langle \frac{D}{\partial u} \dot{\gamma}_{u,0}, \frac{D}{\partial v} \dot{\gamma}_{0,v} \right\rangle \right] dt \\ &= \int_0^1 [\langle R(V, \dot{\gamma})U, \dot{\gamma} \rangle + \langle \dot{U}, \dot{V} \rangle] dt + [\langle \nabla_V U, \dot{\gamma} \rangle]_0^1. \end{aligned}$$

By derivating the identity $f(\gamma_{u,v}(0)) = \gamma_{u,v}(1)$, one gets

$$d^2f_{\gamma(0)}[V(0), U(0)] + df_{\gamma(0)} \cdot \nabla_V U(0) = \nabla_V U(1),$$

so that, using $df^T \cdot \dot{\gamma}(1) = \dot{\gamma}(0)$,

$$\begin{aligned} [\langle \nabla_V U, \dot{\gamma} \rangle]_0^1 &= \langle \nabla_V U(1) - df \cdot \nabla_V U(0), \dot{\gamma}(1) \rangle \\ &= \langle d^2f[U(0), V(0)], \dot{\gamma}(1) \rangle. \end{aligned} \quad \square$$

COROLLARY 2.7. — *Let $\gamma \in \Lambda(f)$ be a critical point of E_f , a vector field $J \in T_\gamma \Lambda(f)$ belongs to the kernel of the quadratic form $d^2E_f(\gamma)$ if and only if J is a Jacobi field satisfying*

$$(d^2f \cdot J(0))^T \cdot \dot{\gamma}(1) = \dot{J}(0) - df^T \cdot \dot{J}(1),$$

where $d^2f \cdot J(0)$ denotes the linear morphism $T_{\gamma(0)}M \rightarrow T_{\gamma(1)}M$, $u \mapsto d^2f[J(0), u]$.

In particular, the kernel of $d^2E_f(\gamma)$ has a dimension bounded by $2 \dim M - 1$. Indeed, the vector space of Jacobi fields of γ has dimension $2 \dim M$, and $J \in T_\gamma \Lambda(f)$ implies that $J(1) = 0$ if $J(0) = 0$, so, for instance, the Jacobi field $J(t) = t\dot{\gamma}(t)$ does not belong to $T_\gamma \Lambda(f)$.

Proof. — Let $J \in \ker d^2E_f(\gamma)$. Applying Proposition 2.6 with $U := J$ and every vector field V with $V(0) = 0$, one proves that J is a Jacobi field in a classical manner. The conclusion now follows from the identity $\forall V \in T_\gamma \Lambda(f)$,

$$\begin{aligned} \int_0^1 \langle \dot{J}, \dot{V} \rangle dt &= [\langle \dot{J}, V \rangle]_0^1 - \int_0^1 \langle \ddot{J}, V \rangle dt \\ &= \langle df^T \cdot \dot{J}(1) - \dot{J}(0), V(0) \rangle - \int_0^1 \langle \ddot{J}, V \rangle dt. \quad \square \end{aligned}$$

PROPOSITION 2.8. — *Let $f : M \rightarrow M$ be a diffeomorphism, a translated point $(x, v) \in \text{SM}$ of \tilde{f} is non-degenerate for the time-shift $t > 0$ if and only if the associated geodesic $\gamma \in \Lambda(f)$ of length t is a non-degenerate critical point of E_f .*

Proof. — We will need the following differential identity: for every diffeomorphism $f : M \rightarrow M$ and every vector field U of M ,

$$(2.3) \quad \nabla_U(df^{-T}) = -df^{-T} \cdot (\nabla_U df)^T \cdot df^{-T}.$$

This identity can be derived by taking the covariant derivative of the identity $df^{-T} \cdot df^T = \text{id}$ (id meaning the section $x \mapsto \text{id}_{T_x M}$) and using $\nabla_U(df^T) = (\nabla_U df)^T$ (which can be obtained by derivating the definition of the adjoint operator).

Let $f : M \rightarrow M$ be a diffeomorphism and let $\hat{f} : \text{TM} \rightarrow \text{TM}$ be its symplectic lift,

$$\hat{f}(x, v) = (f(x), df_x^{-T} \cdot v), \quad \forall (x, v) \in \text{TM}.$$

The differential of \hat{f} at $(x, v) \in \text{TM}$ is

$$d\hat{f}_{(x,v)} \cdot (\xi_1, \xi_2) = (df_x \cdot \xi_1, df_x^{-T} \cdot \xi_2 + \nabla_{\xi_1}(df^{-T}) \cdot v), \quad \forall \xi_1, \xi_2 \in T_x M,$$

where the identification $T_{(x,v)} \text{TM} \simeq T_x M \times T_x M$ is given by the Levi-Civita connection (see e.g. [22, Section 1.3.1]). Let $(x, v) \in \text{SM}$ be a translated point of \tilde{f} for the time-shift $t > 0$ and $\gamma \in \Lambda(f)$ be the associated geodesic of length t . It is non-degenerate if and only if $d(\hat{f} \circ G_{-t})_{G_t(x,v)}$ does not have 1 as an eigenvalue. We recall that, for $t > 0$,

$$(dG_{-t})_{G_t(x,v)} \cdot \left(J(1), \frac{1}{t} \dot{J}(1) \right) = \left(J(0), \frac{1}{t} \dot{J}(0) \right),$$

for every Jacobi field J along γ (the t^{-1} factor being due to the reparametrization, see e.g. [22, Section 1.5]). Since $J \mapsto (J(1), \dot{J}(1))$ is an isomorphism and $tv = \dot{\gamma}(0)$, (x, v) is degenerate if and only if there exists a Jacobi field J along γ such that

$$(2.4) \quad \begin{cases} df \cdot J(0) = J(1), \\ df^{-T} \cdot \dot{J}(0) + \nabla_{J(0)}((df^{-T}) \cdot \dot{\gamma}(0)) = \dot{J}(1). \end{cases}$$

The first equation of (2.4) means that $J \in T_\gamma \Lambda(f)$. Using identity (2.3) and $df^{-T} \cdot \dot{\gamma}(0) = \dot{\gamma}(1)$, the second equation of (2.4) becomes

$$df^{-T} \cdot \dot{J}(0) - df^{-T} \cdot (\nabla_{J(0)} df)^T \cdot \dot{\gamma}(1) = \dot{J}(1).$$

Finally, applying df^T to this last equation, we find the equation of Corollary 2.7. \square

3. The Zoll case

3.1. The Chas–Sullivan product

Let us recall and extend in an obvious manner the filtered Chas–Sullivan product defined in [11]. We refer to [11] for technical details. For now, we do not need to assume that the closed Riemannian manifold M is Zoll. Let $f, g : M \rightarrow M$ be smooth maps, as the space $\Lambda(f) \times_M \Lambda(g)$ is a submanifold of $\Lambda(f) \times \Lambda(g)$ of codimension $n := \dim M$, there is a well-defined Gysin morphism

$$H_*(\Lambda(f) \times \Lambda(g)) \longrightarrow H_{*-n}(\Lambda(f) \times_M \Lambda(g))$$

(see e.g. [11, Proposition B.2]). The Chas–Sullivan product

$$* : H_*(\Lambda(f)) \otimes H_*(\Lambda(g)) \longrightarrow H_{*-n}(\Lambda(g \circ f))$$

is defined by composing this morphism with the morphism in homology induced by the concatenation map ϕ . Let $a, b > 0$ be regular values of P_f and P_g , then $\Lambda_f^{\leq a}$, $\Lambda_g^{\leq b}$ are submanifolds with boundary and one can thus define the Gysin morphism

$$H_*\left(\Lambda_f^{\leq a} \times \Lambda_g^{\leq b}\right) \longrightarrow H_{*-n}\left(\Lambda_f^{\leq a} \times_M \Lambda_g^{\leq b}\right).$$

According to (2.1), one can thus define a filtered Chas–Sullivan product

$$* : H_*\left(\Lambda_f^{\leq a}\right) \otimes H_*\left(\Lambda_g^{\leq b}\right) \longrightarrow H_{*-n}\left(\Lambda_{g \circ f}^{\leq a+b}\right)$$

that is compatible with the morphisms induced by inclusion of sets $\Lambda_h^{\leq c} \hookrightarrow \Lambda_h^{\leq d}$ for $h = f, g$ or $g \circ f$ and $c \leq d$. A relative version of this product is also available

$$\begin{aligned} H_*\left(\Lambda_f^{\leq a}, \Lambda_f^{\leq a'}\right) \otimes H_*\left(\Lambda_g^{\leq b}, \Lambda_g^{\leq b'}\right) &\longrightarrow H_{*-n}\left(\Lambda_{g \circ f}^{\leq a+b}, \Lambda_{g \circ f}^{\leq \max(a+b', a'+b)}\right), \\ H_*\left(\Lambda_f^{\leq a}\right) \otimes H_*\left(\Lambda_g^{\leq b}, \Lambda_g^{\leq b'}\right) &\longrightarrow H_{*-n}\left(\Lambda_{g \circ f}^{\leq a+b}, \Lambda_{g \circ f}^{\leq a+b'}\right), \\ H_*\left(\Lambda_f^{\leq a}, \Lambda_f^{\leq a}\right) \otimes H_*\left(\Lambda_g^{\leq b}, \Lambda_g^{\leq b}\right) &\longrightarrow H_{*-n}\left(\Lambda_{g \circ f}^{\leq a+b}, \Lambda_{g \circ f}^{\leq a+b}\right), \\ H_*\left(\Lambda_f^{\leq a}\right) \otimes H_*\left(\Lambda_g^{\leq b}, \Lambda_g^{\leq b}\right) &\longrightarrow H_{*-n}\left(\Lambda_{g \circ f}^{\leq a+b}, \Lambda_{g \circ f}^{\leq a+b}\right), \end{aligned}$$

that is compatible with morphisms induced by inclusion of relative sublevel sets.

LEMMA 3.1. — *Let $g : M \rightarrow M$ be a smooth map, let (f_s) be an H^1 -homotopy of smooth maps $M \rightarrow M$, let $\delta := \delta(f_s)$ (which is $\geq \delta(f_s \circ g)$). For every $\alpha \in H_*(\Lambda_g^{\leq a})$, $\beta \in H_*(\Lambda_{f_0}^{\leq b})$,*

$$\alpha * \tau(f_s)_* \beta = \tau(f_s \circ g)_* (\alpha * \beta) \in H_*\left(\Lambda_{f_1 \circ g}^{\leq a+b+\delta}\right),$$

as long as the product $*$ is well-defined.

Proof. — Let us consider the following diagram:

$$\begin{array}{ccccc} \Lambda_g^{\leq a} \times \Lambda_{f_0}^{\leq b} & \longleftarrow & \Lambda_g^{\leq a} \times_M \Lambda_{f_0}^{\leq b} & \xrightarrow{\phi} & \Lambda_{f_0 \circ g}^{\leq a+b} \\ \downarrow \text{id} \times \tau(f_s) & & \downarrow \text{id} \times \tau(f_s) & & \downarrow \tau(f_s \circ g) \\ \Lambda_g^{\leq a} \times \Lambda_{f_1}^{\leq b+\delta} & \longleftarrow & \Lambda_g^{\leq a} \times_M \Lambda_{f_1}^{\leq b+\delta} & \xrightarrow{\phi} & \Lambda_{f_1 \circ g}^{\leq a+b+\delta} \end{array},$$

where the unlabeled arrows are inclusion maps. By definition of $\tau(f_s)$ and $\tau(f_s \circ g)$ and by associativity of ϕ (2.2), this diagram commutes. By naturality of the Gysin morphisms and by definition of the product $*$, the conclusion follows. \square

In the same way, one proves that $*$ is associative.

The product structure of the homology group $H_*(\Lambda)$ was studied by Goresky-Hingston, especially in the case where all geodesics are closed and of the same prime length ℓ [11, Sections 13-15]. In the sequel, M will satisfy this assumption and the coefficient ring of the singular homology groups will be \mathbb{Z} if M is orientable and $\mathbb{Z}/2\mathbb{Z}$ otherwise. In this case, the energy functional E_{id} is a perfect Morse–Bott functional: this implies in particular

that the following sequences of inclusion morphisms are exact

$$(3.1) \quad \begin{aligned} 0 &\longrightarrow H_*(\Lambda^{\leq a}) \longrightarrow H_*(\Lambda^{\leq b}) \longrightarrow H_*(\Lambda^{\leq b}, \Lambda^{\leq a}) \longrightarrow 0, \\ 0 &\longrightarrow H^*(\Lambda^{\leq b}, \Lambda^{\leq a}) \longrightarrow H^*(\Lambda^{\leq b}) \longrightarrow H^*(\Lambda^{\leq a}) \longrightarrow 0, \end{aligned}$$

for every $0 \leq a \leq b \leq +\infty$, with $\Lambda^{\leq +\infty} := \Lambda$, the same being true replacing $\Lambda^{\leq \lambda}$ with $\Lambda^{< \lambda}$. Moreover, the filtered product $*$ can be defined for all values. The critical values of P_f correspond to the positive multiples of ℓ with associated critical submanifolds diffeomorphic to SM via $\gamma \mapsto \dot{\gamma}(0)/\|\dot{\gamma}(0)\|$. Let $\lambda_1 \geq 0$ be the Morse index of the critical submanifold Σ_1 associated with the critical value ℓ . The manifold Σ_1 can be oriented and one defines

$$\Theta \in H_{2n-1+\lambda_1}(\Lambda^{\leq \ell})$$

to be the image of the fundamental class of Σ_1 under the canonical isomorphism

$$(3.2) \quad H_*(\Lambda^{\leq \ell}) \simeq H_*(\Lambda^{\leq 0}) \oplus H_{*-\lambda_1}(\Sigma_1)$$

resulting from the splitting of the exact sequence (3.1) for the pair $(\Lambda^{\leq \ell}, \Lambda^{\leq 0})$ induced by $\text{ev} : \Lambda \rightarrow \Lambda^{\leq 0}$, $\text{ev}(\gamma) \equiv \gamma(0)$, and the Morse–Bott isomorphism $H_*(\Lambda^{\leq \ell}, \Lambda^{\leq 0}) \simeq H_{*-\lambda_1}(\Sigma_1)$. Let us set $b := \lambda_1 + n - 1$.

THEOREM 3.2 ([11, Theorem 13.4]). — *Let M be an n -dimensional closed Zoll–Riemannian manifold of length ℓ . For every positive integer r , the Chas–Sullivan product with the class Θ*

$$\Theta * \cdot : H_*(\Lambda^{\leq r\ell}, \Lambda^{< r\ell}) \longrightarrow H_{*+b}(\Lambda^{\leq (r+1)\ell}, \Lambda^{< (r+1)\ell})$$

induces an isomorphism.

3.2. Min-max critical values

Given $f : M \rightarrow M$, let us define min-max critical values of P_f associated with homology classes of $\Lambda(f)$. Given $\alpha \in H_*(\Lambda(f))$, let us define

$$c(\alpha, f) := \inf \left\{ \lambda \geq 0 \mid \alpha \in \text{im} \left(H_*(\Lambda_f^{\leq \lambda}) \longrightarrow H_*(\Lambda_f) \right) \right\}.$$

By the Morse deformation lemma, if $c(\alpha, f) > 0$ then it is a critical value of P_f . A consequence of Theorem 3.2 and the perfectness of E_{id} in the Zoll case is the following corollary.

COROLLARY 3.3. — *Let M be a Zoll–Riemannian manifold with prime length ℓ . For every $\alpha \in H_*(\Lambda)$, one has $c(\alpha, \text{id}) \in \mathbb{N}\ell$ and if $c(\alpha, \text{id}) > 0$, then*

$$c(\Theta^k * \alpha, \text{id}) = c(\alpha, \text{id}) + k\ell, \quad \forall k \in \mathbb{N}.$$

Proof. — Since the $c(\alpha, \text{id})$ are either 0 or critical values of P_{id} , one has $c(\alpha, \text{id}) \in \mathbb{N}\ell$ for all $\alpha \in H_*(\Lambda)$. Let $\alpha \in H_*(\Lambda)$ be such that $c(\alpha, \text{id}) > 0$, so $c(\alpha, \text{id}) = r\ell$ with $r \in \mathbb{N}^*$ and there exists $\beta \in H_*(\Lambda^{\leq r\ell})$ the image of which is α under the inclusion morphism. Since $\Theta^k * \cdot$ commutes with the inclusion morphisms, the image of $\Theta^k * \beta \in H_{*+kb}(\Lambda^{\leq (r+k)\ell})$ in $H_*(\Lambda)$ is $\Theta^k * \alpha$, so $c(\Theta^k * \alpha, \text{id}) \leq (r+k)\ell$.

The image γ of β under the inclusion morphism $H_*(\Lambda^{\leq r\ell}) \rightarrow H_*(\Lambda^{\leq r\ell}, \Lambda^{< r\ell})$ is non-zero since $c(\alpha, \text{id}) \geq r\ell$ (by looking at the long exact sequence of the couple $(\Lambda^{\leq r\ell}, \Lambda^{< r\ell})$). Let us consider the commuting diagram:

$$\begin{array}{ccc} H_*(\Lambda^{\leq r\ell}) & \xrightarrow{\Theta^k * \cdot} & H_{*+kb}(\Lambda^{\leq (r+k)\ell}) \\ \downarrow & & \downarrow \\ H_*(\Lambda^{\leq r\ell}, \Lambda^{< r\ell}) & \xrightarrow[\simeq]{\Theta^k * \cdot} & H_{*+kb}(\Lambda^{\leq (r+k)\ell}, \Lambda^{< (r+k)\ell}), \end{array}$$

where the vertical arrows are inclusion morphisms. Since the bottom arrow is an isomorphism according to Theorem 3.2, the image of γ is non-zero and so is the image of $\Theta^k * \beta$ under the right vertical arrow. This implies that $\Theta^k * \beta$ is not in the image of $H_{*+kb}(\Lambda^{< (r+k)\ell})$ under the inclusion morphism so $c(\Theta^k * \alpha, \text{id}) \geq (r+k)\ell$. \square

PROPOSITION 3.4. — *Let M be a Zoll-Riemannian manifold with prime length ℓ and let (f_s) be a H^1 -homotopy from id to f and let us set $\tau := \tau(f_s)$ and $\delta := \delta(f_s)$.*

- (1) *For every $\alpha \in H_*(\Lambda)$, $|c(\alpha, \text{id}) - c(\tau_*\alpha, f)| \leq \delta$.*
- (2) *For every $\alpha \in H_*(\Lambda)$,*

$$c(\Theta^{k+1} * \tau_*\alpha, f) = c(\Theta^k * \tau_*\alpha, f) + \ell - \varepsilon_k(\alpha), \quad \forall k \in \mathbb{N},$$

where $(\varepsilon_k(\alpha))$ is a sequence of non-negative reals that converges to 0 when $c(\alpha, \text{id}) > 0$.

Proof. — The first statement is a direct consequence of the fact that $\tau(f_s)$ and $\tau(f_{1-s})$ induce a δ -interleaving in the sense of Corollary 2.4. Let us consider the commuting diagram

$$\begin{array}{ccc} H_*(\Lambda_f^{\leq \lambda}) & \xrightarrow{\Theta * \cdot} & H_{*+b}(\Lambda_f^{\leq \lambda+\ell}) \\ \downarrow & & \downarrow \\ H_*(\Lambda_f) & \xrightarrow{\Theta * \cdot} & H_{*+b}(\Lambda_f), \end{array}$$

where the vertical arrows are inclusion morphisms and $\lambda \geq 0$. It implies that if $c(\alpha, f) < \lambda$ then $c(\Theta * \alpha, f) < \lambda + \ell$. In particular, for every $\alpha \in H_*(\Lambda)$,

$$c(\Theta^{k+1} * \tau_* \alpha, f) = c(\Theta^k * \tau_* \alpha, f) + \ell - \varepsilon_k(\alpha), \quad \forall k \in \mathbb{N},$$

for some non-negative sequence $(\varepsilon_k(\alpha))$. According to the first statement of this proposition, for every $k \in \mathbb{N}$,

$$\begin{aligned} \sum_{i=0}^{k-1} \varepsilon_i(\alpha) &= c(\tau_* \alpha, f) - c(\Theta^k * \tau_* \alpha, f) \\ &\quad + k\ell \leq 2\delta + c(\alpha, \text{id}) - c(\Theta^k * \alpha, \text{id}) + k\ell, \end{aligned}$$

where we have used that $\Theta^k * \tau_* \alpha = \tau_*(\Theta^k * \alpha)$ (Lemma 3.1). Therefore, Corollary 3.3 implies that the series $\sum_i \varepsilon_i(\alpha)$ is bounded by 2δ when $c(\alpha, \text{id}) > 0$, bringing the conclusion. \square

3.3. Proof of Theorems 1.6 and 1.7

Proof of Theorem 1.6. — Let $N := \sum_i \beta_i(\text{SM})$ and let $\alpha_1, \dots, \alpha_N \in H_*(\Lambda)$ be an independent family satisfying $c(\alpha_i, \text{id}) = \ell$ for all i . Such a family can be obtained as follows: one takes the image in $H_*(\Lambda^{\leq \ell})$ of an independent family of $0 \oplus H_*(\Sigma_1) \simeq H_*(\text{SM})$ under the isomorphism (3.2), then takes the image of this family under the inclusion morphism $H_*(\Lambda^{\leq \ell}) \rightarrow H_*(\Lambda)$ (this last morphism is injective by exactness of (3.1)). The exactness of (3.1) together with Theorem 3.2 implies that the family $(\Theta^k * \alpha_i)_{k,i}$, $k \in \mathbb{N}$, $1 \leq i \leq N$, is independent in $H_*(\Lambda)$.

Let (f_s) be a H^1 -homotopy from id to $f : M \rightarrow M$ and let $\tau := \tau(f_s)$. Applying the isomorphism $\tau_* : H_*(\Lambda) \rightarrow H_*(\Lambda(f))$, the family $(\Theta^k * \tau_* \alpha_i)_{k,i}$ is independent in $H_*(\Lambda(f))$ (we have used Lemma 3.1). Let us assume that f has finitely many translated points in SM. Since M is a Zoll–Riemannian manifold, the set of positive time-shifts of a translated point of f is $t + \mathbb{N}\ell$ for some $t \in (0, \ell]$. Therefore, the set of (positive) critical values of P_f is invariant by $t \mapsto t + \ell$, discrete and such that any interval $(a, a + \ell]$, $a \geq 0$, contains a fixed finite number of them. Let us study the sequences $(c_k^i) := (c(\Theta^k * \tau_* \alpha_i, f))$ for $1 \leq i \leq N$, the image of which is contained in this discrete set. Since $c_{k+1}^i = c_k^i + \ell - \varepsilon_k^i$ with $\varepsilon_k^i \rightarrow 0$ (Proposition 3.4), by discreteness of the $\mathbb{N}\ell$ -invariant set of critical values, $\varepsilon_k^i = 0$ for large k 's. We deduce that for $A > 0$ large, for each i , there exists a unique k_i such that $c_{k_i}^i \in (A, A + \ell]$. By non-degeneracy of the critical points of P_f , the Morse inequalities in the window of values $(A, A + \ell]$ imply that

the number of critical points in $P_f^{-1}(A, A + \ell]$ is not less than the cardinal of $(\Theta^{k_i} * \tau_* \alpha_i)_i$, which is N . For each translated point, only one time-shift belongs to $(A, A + \ell]$, which brings the conclusion. \square

Proof of Theorem 1.7. — Let $N := \text{CL}(\text{SM})$ and let $u_1, \dots, u_N \in H^*(\Lambda)$ be non-zero classes of positive degree such that

$$c(\Theta \smallfrown (u_1 \smallfrown \dots \smallfrown u_N), \text{id}) = \ell.$$

Such a family can be obtained as follows: let $v_1, \dots, v_N \in H^*(\text{SM})$ be non-zero classes of positive degree such that $v_1 \smallfrown \dots \smallfrown v_N \neq 0$ (which exist by definition of $\text{CL}(\text{SM})$), then $[\text{SM}] \smallfrown (v_1 \smallfrown \dots \smallfrown v_N) \neq 0$ and we apply (3.2) together with the exactness of (3.1) to send $[\text{SM}]$ to Θ (by definition of Θ) and the v_i 's to the u_i 's.

Let $\alpha_0, \dots, \alpha_N \in H_*(\Lambda)$ be the subordinated classes $\alpha_i := \Theta \smallfrown (u_1 \smallfrown \dots \smallfrown u_i)$ such that $c(\alpha_i, \text{id}) = \ell$ for all i (as by subordination, $c(\alpha_{i+1}, \text{id}) \leq c(\alpha_i, \text{id})$ and $c(\Theta, \text{id}) = \ell$). Let (f_s) be a H^1 -homotopy as in the statement of Theorem 1.7, so that $\delta(f_s) < \ell/2$, and let $\tau := \tau(f_s)$. By subordination, $c(\tau_* \alpha_i, f)$ is non-increasing with i and by the first point of Proposition 3.4,

$$\frac{\ell}{2} < c(\tau_* \alpha_N, f) \leq c(\tau_* \alpha_{N-1}, f) \leq \dots \leq c(\tau_* \alpha_0, f) < \frac{3\ell}{2}.$$

According to the Lusternik–Schnirelmann theorem (see e.g. [8, Section II.3.2]), if P_f has a finite number of critical points in the window $(\ell/2, 3\ell/2)$ then $c(\tau_* \alpha_i, f)$ is decreasing. Since the $c(\tau_* \alpha_i, f)$'s are critical values of P_f , the conclusion follows by Corollary 2.2. \square

BIBLIOGRAPHY

- [1] A. ABBONDANDOLO & G. BENEDETTI, “On the local systolic optimality of Zoll contact forms”, *Geom. Funct. Anal.* **33** (2023), no. 2, p. 299-363.
- [2] P. ALBERS, U. FUCHS & W. J. MERRY, “Orderability and the Weinstein conjecture”, *Compos. Math.* **151** (2015), no. 12, p. 2251-2272.
- [3] S. ALLAIS, “On the minimal number of translated points in contact lens spaces”, *Proc. Am. Math. Soc.* **150** (2022), no. 6, p. 2685-2693.
- [4] V. BANGERT, “Non-closed isometry-invariant geodesics”, *Arch. Math.* **106** (2016), no. 6, p. 573-580.
- [5] L. BÉRARD-BERGÉRY, “Quelques exemples de variétés Riemanniennes où toutes les géodésiques issues d’un point sont fermées et de même longueur, suivis de quelques résultats sur leur topologie”, *Ann. Inst. Fourier* **27** (1977), no. 1, p. 231-249.
- [6] A. L. BESSE, *Manifolds all of whose geodesics are closed*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, vol. 93, Springer, 1978, With appendices by D. B. A. Epstein, J.-P. Bourguignon, L. Bérard-Bergery, M. Berger and J. L. Kazdan.
- [7] R. BOTT, “On manifolds all of whose geodesics are closed”, *Ann. Math. (2)* **60** (1954), p. 375-382.

- [8] K.-C. CHANG, *Infinite-dimensional Morse theory and multiple solution problems*, Progress in Nonlinear Differential Equations and their Applications, vol. 6, Birkhäuser, 1993.
- [9] M. CHAS & D. SULLIVAN, “Closed string operators in topology leading to Lie bialgebras and higher string algebra”, in *The legacy of Niels Henrik Abel*, Springer, 2004, p. 771-784.
- [10] V. L. GINZBURG, B. Z. GÜREL & M. MAZZUCHELLI, “On the spectral characterization of Besse and Zoll Reeb flows”, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **38** (2021), no. 3, p. 549-576.
- [11] M. GORESKY & N. HINGSTON, “Loop products and closed geodesics”, *Duke Math. J.* **150** (2009), no. 1, p. 117-209.
- [12] G. GRANJA, Y. KARSHON, M. PABINIAK & S. SANDON, “Givental’s non-linear Maslov index on lens spaces”, *Int. Math. Res. Not.* **2021** (2021), no. 23, p. 18225-18299.
- [13] K. GROVE, “Condition (C) for the energy integral on certain path spaces and applications to the theory of geodesics”, *J. Differ. Geom.* **8** (1973), p. 207-223.
- [14] K. GROVE & M. TANAKA, “On the number of invariant closed geodesics”, *Acta Math.* **140** (1978), no. 1-2, p. 33-48.
- [15] L. MACARINI & M. MAZZUCHELLI, “Isometry-invariant geodesics and the fundamental group. II”, *Adv. Math.* **308** (2017), p. 671-698.
- [16] M. MAZZUCHELLI, “On the multiplicity of isometry-invariant geodesics on product manifolds”, *Algebr. Geom. Topol.* **14** (2014), no. 1, p. 135-156.
- [17] ———, “Isometry-invariant geodesics and the fundamental group”, *Math. Ann.* **362** (2015), no. 1-2, p. 265-280.
- [18] M. MAZZUCHELLI & S. SUHR, “A min-max characterization of Zoll Riemannian metrics”, *Math. Proc. Camb. Philos. Soc.* **172** (2022), no. 3, p. 591-615.
- [19] M. MEIWES & K. NAEF, “Translated points on hypertight contact manifolds”, *J. Topol. Anal.* **10** (2018), no. 2, p. 289-322.
- [20] J. MILNOR, *Morse theory*, Annals of Mathematics Studies, vol. 51, Princeton University Press, 1963, based on lecture notes by M. Spivak and R. Wells.
- [21] ———, *Lectures on the h-cobordism theorem. Notes by L. Siebenmann and J. Sondow*, Mathematical Notes (Princeton), Princeton University Press, 1965.
- [22] G. P. PATERNAIN, *Geodesic flows*, Progress in Mathematics, vol. 180, Birkhäuser, 1999.
- [23] H. SAMELSON, “On manifolds with many closed geodesics”, *Port. Math.* **22** (1963), p. 193-196.
- [24] S. SANDON, “A Morse estimate for translated points of contactomorphisms of spheres and projective spaces”, *Geom. Dedicata* **165** (2013), p. 95-110.
- [25] M. VIGUÉ-POIRRIER & D. SULLIVAN, “The homology theory of the closed geodesic problem”, *J. Differ. Geom.* **11** (1976), no. 4, p. 633-644.

Manuscrit reçu le 27 mai 2022,
révisé le 14 février 2024,
accepté le 21 mars 2024.

Simon ALLAIS
IRMA,
Université de Strasbourg,
7 rue Rene Descartes,
67084 Strasbourg (France)
simon.allais@math.unistra.fr