



ANNALES DE L'INSTITUT FOURIER

Emilien ZABETH

**Block decomposition via the geometric Satake
equivalence**

Article à paraître, mis en ligne le 26 janvier 2026, 73 p.

Article mis à disposition par son auteur selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION-NoDERIVS (CC-BY-ND) 3.0
 <http://creativecommons.org/licenses/by-nd/3.0/>



Les *Annales de l'Institut Fourier* sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org e-ISSN : 1777-5310

BLOCK DECOMPOSITION VIA THE GEOMETRIC SATAKE EQUIVALENCE

by Emilien ZABETH

ABSTRACT. — We give a new proof for the description of the blocks in the category of representations of a reductive algebraic group \mathbf{G} over a field of positive characteristic ℓ (originally due to Donkin), by working in the Satake category of the Langlands dual group and applying Smith–Treumann theory as developed by Riche and Williamson. On the representation theoretic side, our methods enable us to give a bound for the length of a minimum chain linking two weights in the same block, and to give a new proof for the block decomposition of a quantum group at an ℓ^{th} root of unity.

RÉSUMÉ. — Nous donnons une nouvelle preuve pour la description des blocs de la catégorie des représentations d'un groupe algébrique réductif sur un corps de caractéristique positive ℓ (originellement due à Donkin), en travaillant dans la catégorie de Satake du groupe dual de Langlands et en appliquant la théorie de Smith–Treumann telle que développée par Riche et Williamson. Du côté de la théorie des représentations, nos méthodes nous permettent de donner une borne pour la longueur minimale d'une chaîne reliant deux poids dans le même bloc, et de donner une nouvelle preuve de la décomposition en blocs d'un groupe quantique à une racine $\ell^{\text{ème}}$ de l'unité.

1. Introduction

In this paper, we give a proof for the description of the block decomposition of a reductive algebraic group over a field of prime characteristic ℓ using the geometry of the affine Grassmannian. We apply results from [30], where Riche and Williamson used and developed Smith–Treumann theory for sheaves to give a geometric proof of the Linkage principle, which is the first step towards the block decomposition. Most of our paper is dedicated to the study of equivalence classes on some subsets of the affine Weyl group. These equivalence classes will be defined using homomorphisms between

Keywords: Kazhdan–Lusztig polynomials, reductive algebraic groups, perverse sheaves.
2020 Mathematics Subject Classification: 20G05, 14F08, 22E57.

indecomposable parity complexes on some partial affine flag varieties, and we will show that their description implies the desired description for the block decomposition. Our methods will moreover allow us to give a bound for the length of a minimum chain linking two weights in the same block. Finally, using the fact that some of our techniques work in any characteristic (for the field of coefficients of our sheaves), we will give a proof for the description of the block decomposition of a quantum group at an ℓ^{th} root of unity.

In the rest of this introduction, we start by recalling the description of the blocks for a simply-connected simple algebraic group (due to Donkin), before giving a brief summary of some key results from [30] and an overview of our proof.

1.1. The block decomposition of a simply-connected simple algebraic group

Let \mathbf{G} be a simply-connected simple⁽¹⁾ algebraic group over a field \mathbf{k} of prime characteristic ℓ , with a split maximal torus and a Borel subgroup $\mathbf{T} \subset \mathbf{B}$. Let also $\mathfrak{R}_+ \subset \mathfrak{R} \subset \mathbf{X}$ denote the set of positive roots (with respect to the Borel subgroup B^+ satisfying $B^+ \cap B = T$) and roots inside the character lattice of \mathbf{T} , and W_0 be the Weyl group associated with (G, T) . The category $\text{Rep}(\mathbf{G})$ of finite dimensional algebraic \mathbf{G} -modules is a highest weight category, admitting the set of dominant characters \mathbf{X}_+ of \mathbf{T} as a weight poset. For each $\lambda \in \mathbf{X}_+$, denote by L_λ the associated simple \mathbf{G} -module (which is the simple socle of the induced \mathbf{G} -module of highest weight λ), and consider the equivalence relation \sim on \mathbf{X}_+ generated by the relation \mathcal{R}_1 :

$$\lambda \mathcal{R}_1 \mu \iff \text{Ext}_{\text{Rep}(\mathbf{G})}^1(L_\lambda, L_\mu) \neq 0.$$

For any $\lambda \in \mathbf{X}_+$, let $\bar{\lambda} \in \mathbf{X}_+/\sim$ be the associated equivalence class and define $\text{Rep}_{\bar{\lambda}}(\mathbf{G})$ as the Serre subcategory generated by the family $(L_\mu, \mu \sim \lambda)$ (this coincides with the full subcategory of $\text{Rep}(\mathbf{G})$ consisting of \mathbf{G} -modules whose composition factors are of the form L_μ , with $\mu \sim \lambda$). The so-called block decomposition (cf. [20, Lemma 7.1, Part II]) of $\text{Rep}(\mathbf{G})$ is then:

$$\text{Rep}(\mathbf{G}) = \bigoplus_{\bar{\lambda} \in \mathbf{X}_+/\sim} \text{Rep}_{\bar{\lambda}}(\mathbf{G}).$$

⁽¹⁾ By simple we mean that the root system of \mathbf{G} is irreducible.

Remark 1.1. — This formalism also makes sense when $\text{char}(\mathbf{k}) = 0$, in which case the semi-simplicity of the category $\text{Rep}(\mathbf{G})$ (cf. [25, Theorem 22.42]) tells us that $\bar{\lambda} = \{\lambda\}$ for every $\lambda \in \mathbf{X}_+$, so that the block $\text{Rep}_{\bar{\lambda}}(\mathbf{G})$ is just the additive subcategory generated by L_λ .

Fix $\lambda \in \mathbf{X}_+$ and denote by $\rho \in \mathbf{X}$ the half-sum of the positive roots (this belongs to \mathbf{X} thanks to our assumption that \mathbf{G} is simply connected). The linkage principle (cf. [20, Corollary 6.17]) tells us that

$$(1.1) \quad \bar{\lambda} \subset W \bullet_\ell \lambda \cap \mathbf{X}_+,$$

where $W := W_0 \ltimes \mathbb{Z}\mathfrak{R}$ is the affine Weyl group associated with G and \bullet_ℓ is the usual ℓ -dilated dot action on \mathbf{X} , i.e. we have

$$wt_\mu \bullet_\ell \lambda := w(\lambda + \ell\mu + \rho) - \rho$$

for any $\lambda \in \mathbf{X}$, $w \in W_0$ and $\mu \in \mathbb{Z}\mathfrak{R}$. A non-obvious result is that the inclusion (1.1) turns out to be an equality when λ is not contained in a special facet of $\mathbf{X} \otimes_{\mathbb{Z}} \mathbb{R}$ (which concerns the majority of weights), i.e. when $\lambda + \rho \in \mathbf{X} \setminus \ell \cdot \mathbf{X}$. An easy case, which has been treated in [19], is when λ is contained inside of an alcove, which means that $\langle \lambda + \rho, \alpha \rangle \notin \ell\mathbb{Z}$ for all α in the dual root system \mathfrak{R}^\vee (where $\langle \cdot, \cdot \rangle$ denotes the usual perfect pairing between \mathbf{X} and the cocharacter lattice \mathbf{X}^\vee). The case where $\langle \lambda + \rho, \alpha \rangle \in \ell\mathbb{Z}$ for some $\alpha \in \mathfrak{R}^\vee$ (but still with $\lambda + \rho \in \mathbf{X} \setminus \ell \cdot \mathbf{X}$) is however much more involved, and was treated by Donkin in [15].

The inclusion (1.1) is not an equality in general, and the exact description of the blocks is still due to Donkin (cf. [15]; as we will recall in Remark 1.4 below, the proof uses the case where λ is not contained in a special facet): let $r(\lambda + \rho)$ be the smallest integer satisfying $\lambda + \rho \in \ell^{r(\lambda + \rho)} \cdot \mathbf{X} \setminus \ell^{r(\lambda + \rho) + 1} \cdot \mathbf{X}$ and $W^{(r(\lambda + \rho))}$ be the affine Weyl group with translation part dilated by $\ell^{r(\lambda + \rho)}$ (cf. Subsection 6.7 for the precise definition), then we have:

$$\bar{\lambda} = W^{(r(\lambda + \rho))} \bullet_\ell \lambda \cap \mathbf{X}_+.$$

Remark 1.2. — More recently, M. De Visscher gave a shorter proof for the block decomposition [36]. Several of her ideas play a key role in our proof. Her proof requires some restrictions on ℓ and \mathfrak{R} ; we circumvent these restrictions in this paper.

The main goal of this article is to give a proof of this result by working in the setting of constructible sheaves via the geometric Satake equivalence. As we will see in the end (cf. Subsection 7.2), one can deduce from this case the block decomposition for a general reductive group (this description was known, but not explicitly written down in [15]).

1.2. A geometric proof of the linkage principle

Let \mathbb{F} be an algebraically closed field of prime characteristic $p \neq \ell$, and G be the Langlands dual group of \mathbf{G} over \mathbb{F} (in the main body of the text, we will change the notation and replace \mathbf{G} with G^\vee). The affine grassmannian Gr is an ind-scheme over \mathbb{F} , which can be defined as the fppf-quotient of the loop group ind-scheme LG (representing the functor $R \mapsto G(R((z)))$, where z is an indeterminate) by the positive loop group scheme L^+G (representing the functor $R \mapsto G(R[[z]])$). The geometric Satake equivalence (cf. [26]), asserts that there is an equivalence of monoidal categories

$$(\mathrm{Perv}_{L^+G}(\mathrm{Gr}, \mathbf{k}), \star) \xrightarrow{\sim} (\mathrm{Rep}(\mathbf{G}), \otimes_{\mathbf{k}}),$$

where the left-hand side denotes the *Satake category* (equipped with a convolution product), consisting of perverse (étale) sheaves on the affine Grassmannian Gr , with coefficients in \mathbf{k} and which respect an equivariance condition for the left action of the positive loop group L^+G on Gr .

In [30], Riche and Williamson managed to give a new proof of the linkage principle by working in $\mathrm{Perv}_{L^+G}(\mathrm{Gr}, \mathbf{k})$. Moreover, their methods allowed them to give a new character formula for tilting objects (valid in all characteristics), which involves certain ℓ -Kazhdan–Lusztig polynomials. This seems to be the first instance of the geometric Satake equivalence being able to bring us some knowledge on the combinatorics of the category $\mathrm{Rep}(\mathbf{G})$ in positive characteristic. Their proof applies Treumann’s “Smith theory for sheaves” to the Iwahori–Whittaker incarnation of the Satake category, which is a highest weight category $\mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})$ with⁽²⁾ weight poset $\mathbf{X}_{++} := \rho + \mathbf{X}_+$ admitting an equivalence of highest weight categories

$$(1.2) \quad \mathrm{Perv}_{L^+G}(\mathrm{Gr}, \mathbf{k}) \xrightarrow{\sim} \mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k}).$$

This equivalence, which comes from [12], sends an indecomposable tilting object associated with $\lambda \in \mathbf{X}_+$ to the indecomposable tilting object associated with $\lambda + \rho$, which we denote by $\mathcal{T}_{\lambda+\rho}^{\mathcal{IW}}$.

Remark 1.3. — Smith–Treumann theory originates in the work of Smith in the 1930’s concerning the cohomology of topological spaces with coefficients in $\mathbb{Z}/\ell\mathbb{Z}$, and was more recently revisited by Treumann in the setting of constructible sheaves ([35]). The fact that the latter results can be applied to the theory of parity sheaves was first pointed out by Leslie–Lonergan in [23]. See [30, Section 1.5] for more comments.

⁽²⁾One needs to assume that there exists a primitive p^{th} roots of 1 in \mathbf{k} to define $\mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})$.

Let μ_ℓ denote the \mathbb{F} -group scheme of ℓ^{th} root of unity, $(\text{Gr})^{\mu_\ell}$ be the fixed points for the action of $\mu_\ell \subset \mathbb{G}_m$ on Gr by rescaling the indeterminate, and put

$$\mathbf{a}_\ell := \{\mu \in \mathbf{X} \otimes_{\mathbb{Z}} \mathbb{R} : 0 < \langle \mu, \alpha^\vee \rangle < \ell, \forall \alpha \in \mathfrak{R}_+\}.$$

The set \mathbf{a}_ℓ is called the fundamental alcove, and its closure is a fundamental domain for the ℓ -dilated “box” action \square_ℓ of W on $\mathbf{X} \otimes_{\mathbb{Z}} \mathbb{R}$, defined by

$$wt_\mu \square_\ell \lambda := w(\lambda + \ell\mu)$$

for any $\lambda \in \mathbf{X} \otimes_{\mathbb{Z}} \mathbb{R}$, $w \in W_0$ and $\mu \in \mathbb{Z}\mathfrak{R}$. Notice that, for any $\lambda \in \mathbf{X} \otimes_{\mathbb{Z}} \mathbb{R}$ and $w \in W$, we have

$$(1.3) \quad w \bullet_\ell (\lambda - \rho) = w \square_\ell \lambda - \rho.$$

One of the main ingredients used in [30] is the decomposition into connected components

$$(\text{Gr})^{\mu_\ell} = \bigsqcup_{\lambda \in \overline{\mathbf{a}_\ell} \cap \mathbf{X}} \text{Fl}_{\mathbf{G}_\lambda}^{\ell, \circ},$$

where $\text{Fl}_{\mathbf{G}_\lambda}^{\ell, \circ}$ denotes the identity component in the partial affine flag variety associated with the facet $\mathbf{g}_\lambda \subset \overline{\mathbf{a}_\ell}$ containing λ (cf. [30, Proposition 4.7]). This partial affine flag variety is an ind-scheme defined as the fppf-quotient of the loop group ind-scheme $L_\ell G$ representing the functor $R \mapsto G(R((z^\ell)))$ by a positive loop group scheme $L_\ell^+ P_{\mathbf{g}_\lambda}$, representing $R \mapsto P_{\mathbf{G}_\lambda}(R[[z^\ell]])$ and associated with the “parahoric” group scheme $P_{\mathbf{g}_\lambda}$ arising from Bruhat–Tits theory. When $\lambda \in \mathbf{a}_\ell$, one recovers for instance the full affine flag variety for G and if $\lambda = 0$, then we get a copy of the affine Grassmannian.

The other main result is the construction of a fully-faithful functor

$$\Phi : \text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k}) \longrightarrow \text{Sm}_{\mathcal{IW}}((\text{Gr})^{\mu_\ell}, \mathbf{k}),$$

from the subcategory of tilting objects of $\text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$ to the so-called *Smith category* on $(\text{Gr})^{\mu_\ell}$, which involves a pull-back along the immersion $(\text{Gr})^{\mu_\ell} \hookrightarrow \text{Gr}$ followed by passing to a certain Verdier quotient.

As a consequence of the full-faithfulness (and of the fact that both categories are Krull–Schmidt), Φ sends indecomposable objects to indecomposable objects. Thus, one can see that for every $\lambda, \mu \in \mathbf{X}_{++}$, the space $\text{Hom}_{\text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})}(\mathcal{T}_\lambda^{\mathcal{IW}}, \mathcal{T}_\mu^{\mathcal{IW}})$ is non-zero only if the supports of $\Phi(\mathcal{T}_\lambda^{\mathcal{IW}})$ and $\Phi(\mathcal{T}_\mu^{\mathcal{IW}})$ lie in the same connected component $\text{Fl}_{\mathbf{G}_\gamma}^{\ell, \circ}$ of $(\text{Gr})^{\mu_\ell}$ for some $\gamma \in \overline{\mathbf{a}_\ell} \cap \mathbf{X}$, and observe that this happens only if $\lambda, \mu \in W \square_\ell \gamma$. This means that we must have $W \square_\ell \lambda = W \square_\ell \mu$, which is equivalent to $W \bullet_\ell (\lambda - \rho) = W \bullet_\ell (\mu - \rho)$ thanks to (1.3). In view of the equivalence (1.2) and of the geometric Satake equivalence, this means that we have

$$\text{Hom}_{\text{Rep}(\mathbf{G})}(T_{\lambda - \rho}, T_{\mu - \rho}) \neq 0 \implies W \bullet_\ell (\lambda - \rho) = W \bullet_\ell (\mu - \rho),$$

where $T_{\lambda-\rho}$ (resp. $T_{\mu-\rho}$) denotes the indecomposable tilting \mathbf{G} -module associated with $\lambda-\rho$ (resp. $\mu-\rho$). Standard arguments on highest weight categories (see Theorem 2.3 below) then show that this statement is equivalent to the linkage principle (1.1). Moreover, those same standard arguments and equivalences of categories show that, if we denote by $\bar{\lambda} \subset \mathbf{X}_{++}$ the equivalence class of λ for the equivalence relation on \mathbf{X}_{++} induced by the relation \mathcal{R}_2 :

$$\gamma \mathcal{R}_2 \gamma' \iff \text{Hom}_{\text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})}(\mathcal{T}_\gamma^{\mathcal{IW}}, \mathcal{T}_{\gamma'}^{\mathcal{IW}}) \neq 0,$$

then Donkin's theorem on blocks is equivalent to

$$\bar{\lambda} = W^{(r(\lambda))} \square_\ell \lambda \cap \mathbf{X}_{++}.$$

1.3. Summary of the proof

In the sequel, we will push this study further to get the full description of the blocks. Fix $\lambda, \mu \in \mathbf{X}_{++}$. A first step in this direction was actually made in an earlier unpublished version of [30]; namely, we have an isomorphism (cf. Proposition 6.9)

$$\text{Hom}_{\text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})}(\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}}, \mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}}) \simeq \text{Hom}_{\text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})}(\mathcal{T}_\lambda^{\mathcal{IW}}, \mathcal{T}_\mu^{\mathcal{IW}}).$$

This isomorphism enables us to only focus on the case where $\lambda \in \mathbf{X} \setminus \ell \cdot \mathbf{X}$, for which we want to show that the inclusion $\bar{\lambda} \subset W \square_\ell \lambda \cap \mathbf{X}_{++}$, provided by the linkage principle, is an equality. So from now on, let us assume that $r(\lambda) = r(\mu) = 0$, with $W \square_\ell \lambda = W \square_\ell \mu$; we are thus reduced to showing that $\lambda \sim \mu$. We let γ denote the unique element of $W \square_\ell \lambda \cap \bar{\mathbf{a}_\ell}$.

Remark 1.4. — For any $r \in \mathbb{Z}_{\geq 1}$ and $M \in \text{Rep}(\mathbf{G})$, let $M^{[r]}$ denote the twist of M by the r^{th} power of the Frobenius endomorphism. This first step is the analogue of the second step in [36], which uses the fact that the functor $M \mapsto M^{[r]} \otimes L_{(\ell^r-1) \cdot \rho}$ induces an equivalence of categories from $\text{Rep}_{\bar{\lambda}}(\mathbf{G})$ to $\text{Rep}_{\bar{\lambda}'}(\mathbf{G})$, where $\lambda' := (\ell^r-1) \cdot \rho + \ell^r \cdot \lambda$.

Let $\mathbf{g} \subset \bar{\mathbf{a}_\ell}$ be a facet, $W_{\mathbf{g}} \subset W$ denote the stabilizer of \mathbf{g} for \square_ℓ , and $\text{Par}_{\mathcal{IW}_\ell}(\text{Fl}_{\mathbf{g}}^{\ell, \circ}, \mathbf{k})$ denote the additive category of Iwahori–Whittaker-equivariant parity complexes⁽³⁾ on $\text{Fl}_{\mathbf{G}}^{\ell, \circ}$ (cf. Subsection 6.3). The isomorphism classes of indecomposable objects of this category are labelled

⁽³⁾In the body of the paper, we will mostly work in the equivalent context where \mathbf{G} is a facet for the box action \square_1 included in the closure of \mathbf{a}_1 and $\text{Fl}_{\mathbf{G}}^{1, \circ}$ replaces the isomorphic ind-variety $\text{Fl}_{\ell \cdot \mathbf{G}}^{\ell, \circ}$.

(up to a shift) by the set

$${}_f W^{\mathbf{g}} = \{w \in W : w \text{ is minimal in } W_0 w \text{ and maximal in } w W_{\mathbf{g}}\},$$

and we will denote by $\mathcal{E}_{\ell,w}^{\mathbf{g}}$ the indecomposable parity complex associated with w . Moreover, the assignment $w \mapsto w \square_{\ell} \lambda$ induces a bijection ${}_f W^{\mathbf{g}_{\gamma}} \xrightarrow{\sim} W^{\square_{\ell} \gamma} \cap \mathbf{X}_{++}$. In this article, we will extensively study the equivalence relation $\sim_{\mathbf{G}}$ on ${}_f W^{\mathbf{G}}$ generated by the relation $\mathcal{R}_{\mathbf{G}}$:

$$(1.4) \quad w \mathcal{R}_{\mathbf{G}} w' \iff \text{Hom}_{\text{Par}_{\mathcal{IW}_{\ell}}(\text{Fl}_{\mathbf{g}}^{\ell, \circ}, \mathbf{k})}^{\bullet} \left(\mathcal{E}_{\ell,w}^{\mathbf{g}}, \mathcal{E}_{\ell,w'}^{\mathbf{g}} \right) \neq 0, \quad \forall w, w' \in {}_f W^{\mathbf{g}}.$$

The reason for this interest is that we have an isomorphism

$$(1.5) \quad \text{Hom}_{\text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})} \left(\mathcal{T}_{u \square_{\ell} \gamma}^{\mathcal{IW}}, \mathcal{T}_{u' \square_{\ell} \gamma}^{\mathcal{IW}} \right) \simeq \text{Hom}_{\text{Par}_{\mathcal{IW}_{\ell}}(\text{Fl}_{\mathbf{g}_{\gamma}}^{\ell, \circ}, \mathbf{k})}^{\bullet} \left(\mathcal{E}_{\ell,u}^{\mathbf{g}_{\gamma}}, \mathcal{E}_{\ell,u'}^{\mathbf{g}_{\gamma}} \right)$$

for all $u, u' \in {}_f W^{\mathbf{g}_{\gamma}}$, proved in [30] and which arises once again from Smith–Treumann theory (cf. Proposition 6.8). Thus, if we let v and v' be the elements of ${}_f W^{\mathbf{g}_{\gamma}}$ such that $v \square_{\ell} \gamma = \lambda$, $v' \square_{\ell} \gamma = \mu$, we have that

$$\lambda \sim \mu \iff v \sim_{\mathbf{g}_{\gamma}} v'.$$

So we are reduced to proving that the set ${}_f W^{\mathbf{g}_{\gamma}}$ consists of a single equivalence class for $\sim_{\mathbf{g}_{\gamma}}$. The following statement (which is Theorem 5.31) is the main result of this article.

THEOREM 1.5. — *If $\mathbf{G} \subset \overline{\mathbf{a}_{\ell}}$ is a non-special facet, then ${}_f W^{\mathbf{g}}$ consists of a single equivalence class for $\sim_{\mathbf{g}}$.*

A facet \mathbf{g} is called *special* if it is of the form $\mathbf{g} = \{\ell \cdot \delta\}$ for some $\delta \in \mathbf{X}$. Since $r(\lambda) = 0$, the facet \mathbf{g}_{γ} is non-special, so this theorem implies the desired result. There might also be some non-special facets $\mathbf{g} \subset \overline{\mathbf{a}_{\ell}}$ which do not contain an element of \mathbf{X} ; this is for instance the case of \mathbf{a}_{ℓ} when ℓ is less than or equal to the Coxeter number of \mathbf{G} . So this theorem gives a more general result than the block decomposition. We will also describe completely the equivalence classes in ${}_f W^{\mathbf{g}}$ in the case where \mathbf{g} is a special facet and \mathfrak{R}^{\vee} is not irreducible (see Theorem 6.14).

Fix a non-special facet $\mathbf{g} \subset \overline{\mathbf{a}_{\ell}}$. By standard considerations on parity complexes, computing the dimension of the Hom-space on the right-hand side of (1.5) boils down to computing the dimension of the stalks of $\mathcal{E}_{\ell,w}^{\mathbf{g}}$ and $\mathcal{E}_{\ell,w'}^{\mathbf{g}}$, which are given by evaluating some anti-spherical ℓ -Kazhdan–Lusztig polynomials at 1 (cf. [29, Part III]). As the dimension of the Hom-space can only increase when passing from $\text{char}(\mathbf{k}) = 0$ to $\text{char}(\mathbf{k}) = \ell > 0$, we will get the following crucial implication (which is our Corollary 4.8):

$$\forall w, w' \in {}_f W^{\mathbf{g}}, \quad n_{w',w}(1) \neq 0 \implies w \mathcal{R}_{\mathbf{g}} w',$$




Figure 1.1. Hyperplanes arrangement when \mathfrak{R}^\vee is of type C_2 for (a) and A_2 for (b). The walls of the dominant cone \mathcal{C}_0^+ (resp. of $\ell \cdot \rho + \mathcal{C}_0^+$) are represented by thick black lines (resp. dashed lines), each alcove A_i for ($i \in \llbracket 0, r \rrbracket$) is gray, each alcove $w_i \square_\ell \mathbf{a}_\ell$ is labelled with w_i , and the facets $\{w_i \square_\ell \mathbf{G}, i \in \llbracket 0, r \rrbracket\}$ are represented by red lines. Notice that on both of these examples, it sometimes happens that there are two alcoves A_i and A_{i+1} containing a same red faced in their closure for some i , so that we have $w_i = w_{i+1}$.

where $(n_{x,y}, x, y \in {}_f W)$ denotes the usual anti-spherical Kazhdan–Lusztig polynomials studied in [31, Theorem 3.1]. This implication allows us to view our problem as a question of combinatorics for the anti-spherical Kazhdan–Lusztig polynomials. The proof of Theorem 1.5 then roughly goes into two steps:

(Step 1) *Moving away from the walls of the dominant cone.* We show that, for any $u \in {}_f W^g$, there exists some $\hat{u} \in {}_f W^g$ satisfying

$$n_{u,\hat{u}}(1) \neq 0 \quad \text{and} \quad \hat{u} \square_\ell g \subset \ell \cdot \rho + \overline{\mathcal{C}^+},$$

where $\overline{\mathcal{C}^+}$ denotes the closure of the dominant cone (see Proposition 5.16). This will allow us to only focus on polynomials $n_{w',w}$ with $w, w' \in {}_f W^g$ satisfying

$$w' \square_\ell g, w \square_\ell g \subset \ell \cdot \rho + \overline{\mathcal{C}^+},$$

over which we have a much better control.

(Step 2) *Linking close elements.* Let $v \in {}_f W^g$ be such that $v \square_\ell g \subset \ell \cdot \rho + \overline{a_\ell}$ and pick $w \in {}_f W^g$ such that $w \square_\ell g \subset \ell \cdot \rho + \overline{\mathcal{C}^+}$. We will prove that we have $w \sim_g v$ (in view of the first step, this will conclude the proof, since any element of ${}_f W^g$ will then be in relation with v). In order to do that, we will let A_0 be an alcove such that $w \square_\ell G \subset \overline{A_0}$,

$$A_r := \ell \cdot \rho + \mathbf{a}_\ell, A_{r-1}, \dots, A_0$$

be a sequence of alcoves included in $\ell \cdot \rho + \mathcal{C}_0^+$, where A_{i+1} is obtained by reflecting A_i along one of its walls for every $i \in \llbracket 0, r-1 \rrbracket$, and denote by w_i the element of ${}_f W^g$ satisfying $w_i \square_\ell g \subset \overline{A_i}$ for all i (so $w_0 = w$ and $w_r = v$)⁽⁴⁾. See Figure 1.1 for a representation of the situation in types C_2 and A_2 , where g is a wall. We will show that we have

$$w_i \sim_g w_{i-1}, \quad \forall i \in \llbracket 0, r-1 \rrbracket.$$

More specifically, we will find an element $u_i \in {}_f W^g$ such that $n_{w_i, u_i}(1) \neq 0$ and $n_{w_{i-1}, u_i}(1) \neq 0$ (occasionally we might have $u_i = w_{i-1}$, but not always).

Remarks 1.6.

(1) The need for Step 1 is due to the “cancellation effect” occurring for the anti-spherical Kazhdan–Lusztig polynomials when we are close to the walls of the dominant chamber, as it was observed

⁽⁴⁾Notice that we don’t necessarily have $w_i \square_\ell \mathbf{a}_i = A_i$.

in the similar context of [7] (cf. the introduction and section 9 of this reference).

- (2) The Step 2 will actually be split into two parts: the case where \mathbf{g} is not a point, and the case where \mathbf{g} is a point (and still a non-special facet). The latter uses the theory of “periodic polynomials” (originally due to Lusztig) as displayed in [31], and is the most involved part of the paper (cf. Subsection 5.5).
- (3) These steps were inspired to us by the steps used in [36, Section 2].
- (4) We will actually use “geometric” arguments (namely, parity complexes on partial affine flag varieties) to prove some “combinatorial” facts concerning the anti-spherical ℓ -Kazhdan–Lusztig polynomials (see Proposition 5.4), which hold in particular for the ordinary anti-spherical Kazhdan–Lusztig polynomials. However, we do not yet have an analogous geometric incarnation for the combinatorics we use concerning the periodic polynomials (cf. Proposition 5.24), but hope to come back to this question in the future.

1.4. Consequences

Our new proof of Donkin’s Theorem enables us to give a bound on the length of a minimum chain linking two weights in the same block for $\text{Rep}(\mathbf{G})$, in the same fashion as [36, Corollary 3.1], but without any restriction on the root system nor on the characteristic of \mathbf{k} , see Proposition 7.3. Such a result was not available via Donkin’s original proof, as his arguments required going arbitrarily far inside of the dominant cone.

Moreover, our determination of the equivalence classes in ${}_f W^{\mathbf{g}}$ for $\sim_{\mathbf{g}}$ also applies to the case where $\text{char}(\mathbf{k}) = 0$, from which we are able to deduce the block decomposition for a quantum group at an ℓ^{th} root of unity in Subsection 7.3 (this result was originally found in [34]).

1.5. Structure of the paper

After some preliminaries on highest weight categories in Section 2, we recall in Section 3 the construction of Iwahori–Whittaker equivariant derived categories on partial affine flag varieties of the form $\text{Fl}_{\mathbf{g}}^\circ$, for a facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$. In particular, we introduce indecomposable parity complexes, and study the effect of pushforward and pullback of these objects under the canonical

proper morphism $\pi_g : \mathrm{Fl}_{a_1}^\circ \rightarrow \mathrm{Fl}_g^\circ$ (Subsection 4.1). This allows us to introduce and study the equivalence relation \sim_g on the set ${}_f W^g$. Section 5 is the heart of the paper and is dedicated to the study of the equivalence classes for \sim_g . In particular, we apply the plan described above: Step 1 is dealt with at the end of Subsection 5.3, while Step 2 is taken care of in Subsection 5.4 (when g is not a point) and Subsection 5.5 (when g is a non-special point). The case where g is a special facet is done at the end of Section 6, as we need Smith–Treumann theory to deduce it from the previous cases. Finally, we harvest the consequences for representation theory in the last section: block decomposition for reductive groups in Subsection 7.2 and for quantum groups in Subsection 7.3.

Acknowledgements

I am deeply grateful to Simon Riche for his guidance through many suggestions and countless careful re-readings, and to Geordie Williamson for helpful discussions and comments. I also want to thank Joel Gibson for putting together the interactive visualization tool *Lievis* and making it accessible on his web page (cf. [16]): a lot (if not all) of the intuition behind the results of this paper concerning the anti-spherical Kazhdan–Lusztig polynomials came from there. This work was conducted during PhD studies at the Université Clermont Auvergne, co-supervised by Simon Riche and Geordie Williamson.

2. Blocks for a highest weight category

In this section, \mathbf{k} is any field. The goal of these formal preliminaries is to introduce two equivalence relations on the weight poset of a highest weight category. Both relations allow us to give block decompositions of the category, and we will show that these relations and decompositions are the same. These results will later be applied in Sections 6 and 7 to categories of perverse sheaves and representations. We start by briefly recalling some properties of a highest weight category.

2.1. Recollections on highest weight categories

Let \mathcal{A} be a highest weight category over \mathbf{k} , with weight poset (Λ, \leqslant) (cf. [28, Section 3.7] for a detailed treatment of highest weight categories).

In particular, we assume that each object of \mathcal{A} has finite length and that the vector space $\text{Hom}_{\mathcal{A}}(M, N)$ is finite dimensional for every $M, N \in \mathcal{A}$. To each $s \in \Lambda$ corresponds a simple object L_s , a *standard object* Δ_s and a *costandard object* ∇_s . By definition, the association $s \mapsto L_s$ induces a bijection between Λ and the isomorphism classes of simple objects of \mathcal{A} . For any $M \in \mathcal{A}$ and $s \in \Lambda$, we will denote by $[M : L_s]$ the number of times L_s appears in a composition series of M .

We denote by $\text{Tilt}(\mathcal{A})$ the full additive subcategory of \mathcal{A} consisting of tilting objects, i.e. those admitting both a filtration with sub-quotients being standard objects and a filtration with sub-quotients being costandard objects. For any $s \in \Lambda$ and $M \in \text{Tilt}(\mathcal{A})$, we will denote by $(M : \nabla_s)$, resp. $(M : \Delta_s)$, the number of times ∇_s appears in a filtration of M with costandard, resp. standard, sub-quotients. This number does not depend on the choice of such a filtration, as one can show that it is equal to $\dim_{\mathbf{k}} \text{Hom}_{\mathcal{A}}(\Delta_s, M)$, resp. $\dim_{\mathbf{k}} \text{Hom}_{\mathcal{A}}(M, \nabla_s)$. Indecomposable tilting objects are also parameterized (up to isomorphism) by Λ ; we denote by T_s the unique indecomposable tilting object such that

$$[T_s : L_s] = 1 \quad \text{and} \quad \forall t \in \Lambda, \quad [T_s : L_t] \neq 0 \implies t \leq s.$$

We have morphisms $L_s \xleftarrow{\quad} \Delta_s \hookrightarrow T_s$ for all $s \in \Lambda$. Finally recall that the canonical functor

$$(2.1) \quad \text{K}^b \text{Tilt}(\mathcal{A}) \longrightarrow D^b(\mathcal{A})$$

is an equivalence of categories (cf. [28, Proposition 7.17]).

2.2. Equivalence relations on Λ

We are going to consider two equivalence relations on the set Λ . The first one, denoted by \sim_1 , is generated by the relation \mathcal{R}_1 , defined by

$$s\mathcal{R}_1 t \iff \text{Ext}_{\mathcal{A}}^1(L_s, L_t) \neq 0.$$

Here, $\text{Ext}_{\mathcal{A}}^1(B, A)$ denotes the Ext-group of isomorphism classes of extensions of B by A , for two objects A, B . Recall (cf. [33, Lemma 13.27.6]) that this coincides with the \mathbf{k} -vector space $\text{Hom}_{D^b(\mathcal{A})}(B, A[1])$. For all $s \in \Lambda$, we denote by \bar{s} the associated equivalence class and by $\mathcal{A}_{\bar{s}}$ the Serre subcategory generated by the L_t 's, for $t \in \bar{s}$.

PROPOSITION 2.1. — *The canonical functor*

$$\bigoplus_{\bar{s} \in \Lambda / \sim_1} \mathcal{A}_{\bar{s}} \longrightarrow \mathcal{A}$$

is an equivalence of categories.

Proof. — The full faithfulness comes from the fact that, for each $(M, N) \in \mathcal{A}_{\bar{s}} \times \mathcal{A}_{\bar{t}}$ with $\bar{s} \neq \bar{t}$, we have $\text{Hom}_{\mathcal{A}}(M, N) = 0$. This is obvious when M and N are simple, and can be shown by induction on the length of these objects in the general case. With this fact and another induction on the length, one can also show using long exact sequences of cohomology that

$$(2.2) \quad \text{Ext}_{\mathcal{A}}^1(M, N) = 0.$$

It remains to show that the functor is essentially surjective. We still proceed by induction on the length of M , the case of simple objects being obvious. Let $N \subset M$ and $t \in \Lambda$ be such that we have a short exact sequence

$$(2.3) \quad 0 \longrightarrow N \xrightarrow{\varphi} M \longrightarrow L_t \longrightarrow 0.$$

By the induction hypothesis, we have a canonical isomorphism

$$N \simeq \bigoplus_{\bar{s} \in \Lambda / \sim_1} N_{\bar{s}},$$

with $N_{\bar{s}} \in \mathcal{A}_{\bar{s}}$ for all \bar{s} . Now, let $p : N \rightarrow N_{\bar{t}}$ be the canonical projection and $M' \in \mathcal{A}$ be such that we have the following commutative diagram, with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & N & \xrightarrow{\varphi} & M & \longrightarrow & L_t \longrightarrow 0 \\ & & \downarrow p & & \downarrow \psi & & \downarrow \text{id} \\ 0 & \longrightarrow & N_{\bar{t}} & \longrightarrow & M' & \longrightarrow & L_t \longrightarrow 0. \end{array}$$

The morphism ψ is clearly surjective, and restricting φ to $M'' := \bigoplus_{\bar{s} \neq \bar{t}} N_{\bar{s}}$ yields an exact sequence

$$0 \longrightarrow M'' \longrightarrow M \longrightarrow M' \longrightarrow 0.$$

Finally, observe that M' , resp. M'' , belongs to $\mathcal{A}_{\bar{t}}$, resp. $\bigoplus_{\bar{s} \neq \bar{t}} \mathcal{A}_{\bar{s}}$, so that $\text{Ext}_{\mathcal{A}}^1(M', M'') = 0$ thanks to (2.2). Thus, we have $M \simeq M' \oplus M''$, and this concludes the proof. \square

The existence of a quasi-inverse functor allows us to define projection functors $Q_{\bar{s}} : \mathcal{A} \rightarrow \mathcal{A}_{\bar{s}}$ for every \bar{s} . We thus have an isomorphism of functors

$$(2.4) \quad \text{id}_{\mathcal{A}} \simeq \bigoplus_{\bar{s} \in \Lambda / \sim_1} Q_{\bar{s}}$$

where there exists no non-zero morphism between the essential images of any two distinct functors $Q_{\bar{s}}$ and $Q_{\bar{t}}$. In particular, each $Q_{\bar{s}}$ is exact.

The second equivalence relation is denoted \sim_2 . It is generated by the relation \mathcal{R}_2 :

$$s\mathcal{R}_2 t \iff \text{Hom}_{\mathcal{A}}(T_s, T_t) \neq 0.$$

For every weight s , we denote by (s) the associated equivalence class and by $\text{Tilt}_{(s)}(\mathcal{A})$ the additive sub-category generated by the family $(T_t)_{t \in (s)}$. Since any tilting object is a sum of indecomposable tilting objects, it is easy to see that the canonical functor $\bigoplus_{(s) \in \Lambda / \sim_2} \text{Tilt}_{(s)}(\mathcal{A}) \rightarrow \text{Tilt}(\mathcal{A})$ is an equivalence, so that we can define projection functors $\pi_{(s)} : \text{Tilt}(\mathcal{A}) \rightarrow \text{Tilt}_{(s)}(\mathcal{A})$, inducing an isomorphism of functors

$$\text{id}_{\text{Tilt}(\mathcal{A})} \simeq \bigoplus_{(s) \in \Lambda / \sim_2} \pi_{(s)}.$$

The $\pi_{(s)}$'s induce endofunctors of the category $\text{K}^b \text{Tilt}(\mathcal{A})$ (that we still denote by $\pi_{(s)}$), and those still induce a decomposition of the identity functor. Conjugating the $\pi_{(s)}$'s with the equivalence (2.1), the obtained functors (which we still denote by $\pi_{(s)}$) provide a decomposition

$$(2.5) \quad \text{id}_{D^b(\mathcal{A})} \simeq \bigoplus_{(s) \in \Lambda / \sim_2} \pi_{(s)}.$$

This isomorphism implies that each functor $\pi_{(s)}$ preserves the sub-category \mathcal{A} , so we get an isomorphism $\text{id}_{\mathcal{A}} \simeq \bigoplus_{(s)} \pi_{(s)}$. It is also clear that each $\pi_{(s)}$ is an exact functor of the category \mathcal{A} and that there is no non-zero morphism between the essential images of two distinct such functors.

LEMMA 2.2. — *Let $s \in \Lambda$. For all $t \in \Lambda$, we have isomorphisms*

$$\pi_{(t)}(L_s) \simeq \begin{cases} L_s & \text{if } s \in (t) \\ 0 & \text{otherwise} \end{cases} \quad Q_{\bar{t}}(T_s) \simeq \begin{cases} T_s & \text{if } s \in \bar{t} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. — By the decompositions (2.4) and (2.5) and the fact that the objects L_s and T_s are indecomposable, we know that there exists a unique (t) , resp. a unique \bar{t} , such that $\pi_{(t)}(L_s) \neq 0$, resp. $Q_{\bar{t}}(T_s) \neq 0$, and we then have $\pi_{(t)}(L_s) \simeq L_s$, resp. $Q_{\bar{t}}(T_s) \simeq T_s$. By exactness of the projection functors, we have morphisms

$$Q_{\bar{u}}(L_s) \xleftarrow{\quad} Q_{\bar{u}}(\Delta_s) \xrightarrow{\quad} Q_{\bar{u}}(T_s) \quad \text{and} \quad \pi_{(u)}(L_s) \xleftarrow{\quad} \pi_{(u)}(\Delta_s) \xrightarrow{\quad} \pi_{(u)}(T_s)$$

for every $u \in \Lambda$. If $Q_{\bar{u}}(T_s) = 0$, then $Q_{\bar{u}}(L_s) = 0$, so $\bar{u} \neq \bar{s}$. This implies that $\bar{r} = \bar{s}$. Likewise, if $\pi_{(u)}(L_s) \neq 0$, then $\pi_{(u)}(T_s) \neq 0$, so $(u) = (s)$, whence $(t) = (s)$. \square

We have now all the necessary ingredients to prove the following theorem.

THEOREM 2.3. — *The relations \sim_1 and \sim_2 coincide.*

Proof. — Let $s, t \in \Lambda$ be such that $(s) \neq (t)$. By Lemma 2.2, we have

$$\mathrm{Ext}_{\mathcal{A}}^1(L_t, L_s) = \mathrm{Hom}_{D^b(\mathcal{A})}(L_t, L_s[1]) \simeq \mathrm{Hom}_{D^b(\mathcal{A})}(\pi_{(t)}(L_t), \pi_{(s)}(L_s[1])).$$

As there is no non-zero morphism between the essential images of $\pi_{(s)}$ and $\pi_{(t)}$, the right-hand side is zero, so $t \not\sim_1 s$. We have thus shown the implication $\bar{s} = \bar{t} \Rightarrow (s) = (t)$.

Conversely, if $s, t \in \Lambda$ are such that $\bar{s} \neq \bar{t}$, then it follows that

$$\mathrm{Hom}_{\mathcal{A}}(T_t, T_s) \simeq \mathrm{Hom}_{\mathcal{A}}(Q_{\bar{t}}(T_t), Q_{\bar{s}}(T_s)) = 0$$

so $t \not\sim_2 s$, and we have the implication $(s) = (t) \Rightarrow \bar{s} = \bar{t}$. \square

In the sequel, we will denote by \sim the equivalence relation on Λ considered in the previous theorem.

3. Recollections on (Iwahori–Whittaker) equivariant derived categories

In this section, we recall the construction of Iwahori–Whittaker-equivariant derived categories on some partial affine flag varieties arising from Bruhat–Tits theory. We fix a prime number $\ell \neq p$. Since our sheaves will be étale, we let \mathbf{k} be either a finite field of prime characteristic ℓ or a finite extension of \mathbb{Q}_ℓ .

3.1. Notations

From now on, G will denote a semi-simple algebraic group of adjoint type defined over an algebraically closed field \mathbb{F} of characteristic $p > 0$. Choose a Borel subgroup $B \subset G$ and a maximal torus $T \subset B$, and let $\mathfrak{R} \subset \mathbb{X}$ (resp. \mathfrak{R}_+) denote the subset of roots (resp. positive roots with respect to the opposite Borel subgroup of B with respect to T) inside the group of characters of T . Each root α defines a subgroup $U_\alpha \subset G$ isomorphic to the additive group \mathbb{G}_a , and the subgroup generated by the U_α 's, for $\alpha \in \mathfrak{R}_+$ (resp. for $\alpha \in -\mathfrak{R}_+$), is a unipotent group which we will denote by U^+ .

(resp. U). We will also denote by $\mathfrak{R}^\vee \subset \mathbb{X}^\vee$ the set of coroots inside the group of cocharacters of T and by \mathbb{X}_+^\vee (resp. \mathbb{X}_{++}^\vee) the set of dominant (resp. strictly dominant) cocharacters, i.e. those cocharacters which satisfy $\langle \lambda, \alpha \rangle \geq 0$ (resp. $\langle \lambda, \alpha \rangle > 0$) for any $\alpha \in \mathfrak{R}_+$.

Fix an integer $n \geq 1$. We will write $\mathcal{O}_n := \mathbb{F}[[z^n]]$, $\mathcal{K}_n := \mathbb{F}((z^n))$, $\mathcal{O} := \mathcal{O}_1$, $\mathcal{K} := \mathcal{K}_1$ where z is an indeterminate. For any affine \mathbb{F} -group scheme H , we define the functors $L_n^+ H := R \mapsto H(R[[z^n]])$ and $L_n H : R \mapsto H(R((z^n)))$ from \mathbb{F} -algebras to groups, which are representable by a \mathbb{F} -group scheme and a \mathbb{F} -group ind-scheme respectively. Note that these definitions also make sense if H is only defined over \mathcal{O}_n . We will write $L^+ H$, resp. $L H$, instead of $L_n^+ H$, resp. $L_n H$. We will denote by Iw_u^+ the inverse image of U^+ under the evaluation map $L^+ G \rightarrow G, z \mapsto 0$.

We assume that there exists a primitive p^{th} root of unity $\zeta \in \mathbf{k}$, and consider the Artin–Schreier map $\text{AS} : \mathbb{G}_a \rightarrow \mathbb{G}_a$ determined by the map of rings $x \mapsto x^p - x$. This morphism is a Galois cover of group $\mathbb{Z}/p\mathbb{Z}$, so determines a continuous group morphism $\pi_1(\mathbb{G}_a, 0) \rightarrow \mathbb{Z}/p\mathbb{Z}$, where $\pi_1(\mathbb{G}_a, 0)$ is the étale fundamental group of \mathbb{G}_a with geometric base point 0. The composition of this map with the morphism $\mathbb{Z}/p\mathbb{Z} \rightarrow \mathbf{k}^\times$ (induced by ζ) yields a continuous representation of the fundamental group, and thus a local system on \mathbb{G}_a of rank one. We denote this local system by \mathcal{L}_{AS} .

3.2. The affine Weyl group and some Bruhat–Tits theory

We let $N_G(T)$ be the normalizer of T in G . The finite Weyl group associated with (G, T) will be denoted by $W_0 := N_G(T)/T$, and we consider the affine Weyl group

$$W := W_0 \ltimes \mathbb{Z}\mathfrak{R}^\vee,$$

which acts naturally on $E := \mathbb{X}^\vee \otimes_{\mathbb{Z}} \mathbb{R}$ via the n -dilated “box” action, defined by

$$wt_\mu \square_n \lambda := w(\lambda + n\mu)$$

for any $\lambda \in \mathbb{X}^\vee \otimes_{\mathbb{Z}} \mathbb{R}$, $w \in W_0$ and $\mu \in \mathbb{Z}\mathfrak{R}^\vee$, where we have denoted by wt_μ the element of the affine Weyl group associated with the couple (w, μ) . The closure $\overline{\mathbf{a}_n}$ of the set

$$\mathbf{a}_n := \{\lambda \in E \mid 0 < \langle \lambda, \alpha \rangle < n, \quad \forall \alpha \in \mathfrak{R}_+\}$$

is a fundamental domain for this action, which stabilizes \mathbb{X}^\vee . Thus, $\overline{\mathbf{a}_n} \cap \mathbb{X}^\vee$ is a fundamental domain for the action of W on the set of cocharacters. Each root $\alpha \in \mathfrak{R}$ defines a reflection $s_\alpha \in W$, and we will denote by S_0 the set of simple reflections (i.e. associated with a simple root) of the finite

Weyl group, which is known to generate W_0 . It is also well known (cf. [20, Section II.6.3]) that the set of simple reflections

$$S := \{(s, 0), s \in S_0\} \cup \{(s_\beta, \beta^\vee)\},$$

where β^\vee runs through the set of largest short roots of the irreducible components of \mathfrak{R}^\vee , generates W . Moreover, W is a Coxeter group, with Coxeter generating system S , for which we will denote by $l : W \rightarrow \mathbb{Z}_{\geq 0}$ the associated length function.

It will also be useful for us to note that one can extend the translation action of $\mathbb{Z}\mathfrak{R}^\vee$ to the whole group of cocharacters \mathbb{X}^\vee , by putting $t_\mu \square_n \lambda := \lambda + n\mu$ for every $\mu \in E, \lambda \in \mathbb{X}^\vee$, and that this extends the action of W on E to an action of the extended affine Weyl group $\widetilde{W} := W_0 \ltimes \mathbb{X}^\vee$. The subgroup $W \subset \widetilde{W}$ is normal and one can extend the length function l to the whole group \widetilde{W} (cf. [5, Section 2.1] for more details), so that we can define the subgroup

$$\Omega := \left\{ w \in \widetilde{W} \mid l(w) = 0 \right\},$$

which acts on W by Coxeter group isomorphisms and induces an isomorphism

$$(3.1) \quad \Omega \ltimes W \simeq \widetilde{W}$$

such that $l(\omega w) = l(ww) = l(w)$ for all $(\omega, w) \in \Omega \times W$.

The action of \widetilde{W} on E defines a hyperplane arrangement in E , and hence a collection of facets (cf. [13, Chapter 5, Section 1.2]). To any facet $\mathbf{g} \subset \overline{\mathbf{a}_n}$, Bruhat–Tits theory associates a “parahoric group scheme” $P_{\mathbf{G}}$ defined over \mathcal{O}_n , whose generic fiber is isomorphic to $G \times_{\text{Spec}(\mathbb{F})} \text{Spec}(\mathcal{K}_n)$ and whose group of \mathcal{O}_n -points coincides with a subgroup of finite index of the pointwise stabilizer of $-\mathbf{g}^{(5)}$ for the action of $G(\mathcal{K}_n)$ on the Bruhat–Tits building associated with $G \times_{\text{Spec}(\mathbb{F})} \text{Spec}(\mathcal{K}_n)$. We fix such a facet \mathbf{g} . The partial affine flag variety associated with \mathbf{g} will be denoted by $\text{Fl}_{\mathbf{g}}^n$ and defined as the fppf-quotient $L_n G / L_n^+ P_{\mathbf{g}}$, which is an ind-projective ind-scheme over \mathbb{F} (cf. [27] for a detailed exposition on these partial affine flag varieties). The connected components of $\text{Fl}_{\mathbf{G}}^n$ are in bijection with the group $\mathbb{X}^\vee / \mathbb{Z}\mathfrak{R}^\vee$ (see [27, Theorem 0.1]), so we will denote by $\text{Fl}_{\mathbf{g}}^{n, \circ}$ the connected component associated with the neutral element. In the sequel, we will apply these results to the cases where $n = \ell$ or $n = 1$; when $n = 1$ (which will

(5) We follow the conventions of [30, Section 4] for the facet which $P_{\mathbf{g}}(\mathcal{O}_n)$ must stabilize. Note that the authors there defined \mathbf{a}_n to be the opposite of our current fundamental alcove.

be the case until Section 6), we will write $\mathrm{Fl}_{\mathbf{G}}^\circ$ (resp. $\mathrm{Fl}_{\mathbf{g}}$) instead of $\mathrm{Fl}_{\mathbf{g}}^{1,\circ}$ (resp. $\mathrm{Fl}_{\mathbf{g}}^1$).

3.3. Parity sheaves on partial affine flag varieties

Let $\chi_0 : U^+ \rightarrow \mathbb{G}_a$ be a morphism of \mathbb{F} -algebraic groups which restricts to a non-zero morphism on each subgroup U_α , for α a simple root, and let $\chi : \mathrm{Iw}_u^+ \rightarrow \mathbb{G}_a$ denote the composition of χ_0 with the evaluation map. For any $Y \subset \mathrm{Fl}_{\mathbf{g}}^\circ$ which is a locally closed finite union of Iw_u^+ -orbits, one can show (cf. [14, Lemma 3.2]) that the Iw_u^+ -action on Y factors through a quotient group of finite type J such that χ factors through $\chi_J : J \rightarrow \mathbb{G}_a$, where χ_J is induced by χ ; we can then consider the $(J, \chi_J^* \mathcal{L}_{\mathrm{AS}})$ -equivariant derived category of étale \mathbf{k} -sheaves $D_{J, \chi_J^* \mathcal{L}_{\mathrm{AS}}}^b(Y, \mathbf{k})$. This category is by definition the subcategory of $D_{\mathrm{c}}^b(Y, \mathbf{k})$ consisting of constructible complexes of étale \mathbf{k} -sheaves \mathcal{F} such that there exists an isomorphism $a^* \mathcal{F} \simeq \chi_J^* \mathcal{L}_{\mathrm{AS}} \boxtimes \mathcal{F}$, where $a : J \times Y \rightarrow Y$ is the action map; this definition does not depend on the choice of J . We then define the category $D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}^\circ, \mathbf{k})$ as a direct limit of the categories $D_{J, \chi_J^* \mathcal{L}_{\mathrm{AS}}}^b(Y, \mathbf{k})$, indexed by finite and closed unions of Iw_u^+ -orbits Y which are ordered by inclusion. Notice that, since the transition maps are push-forwards of closed immersions, they are fully faithful functors, and one can see this limit as an increasing union of categories.

We now recall the definition and some of the properties of parity complexes on partial affine flag varieties, since those will play a major role in the sequel. Let $Y \subset \mathrm{Fl}_{\mathbf{g}}^\circ$ be a finite locally closed union of Iw_u^+ -orbits. A complex $\mathcal{F} \in D_{\mathcal{IW}}^b(Y, \mathbf{k})$ is called $*$ -even, resp. $!$ -even, if for any inclusion $j : Y' \hookrightarrow Y$ of an Iw_u^+ -orbit Y' , the complex $j^* \mathcal{F}$, resp. $j^! \mathcal{F}$, is concentrated in even degrees. One defines similarly $*$ -odd and $!$ -odd complexes and say that a complex is even, resp. odd, if it is both $*$ -even and $!$ -even, resp. $*$ -odd and $!$ -odd. A complex is called parity if it is a direct sum of even and odd complexes.

Since the category $D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}^\circ, \mathbf{k})$ is defined as a direct limit, it makes sense to talk about even and odd objects in this category, and we will denote by $\mathrm{Par}_{\mathcal{IW}}(\mathrm{Fl}_{\mathbf{g}}^\circ, \mathbf{k})$ the additive full subcategory consisting of parity objects. The general theory of parity complexes (from [21]) allows us to state the following result.

PROPOSITION 3.1. — *Let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a facet. For each Iw_u^+ -orbit Y in $\mathrm{Fl}_{\mathbf{g}}^\circ$ which supports a non-zero Iwahori–Whittaker rank one local system \mathcal{L} ,*

there exists a unique indecomposable parity complex in $\text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})$ supported on \overline{Y} whose restriction to Y is isomorphic to $\mathcal{L}[\dim(Y)]$, and each indecomposable parity complex is isomorphic to such an object up to a cohomological shift. Moreover, each object in $\text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})$ is isomorphic to a sum of indecomposable parity complexes.

Remark 3.2. — See Proposition 4.3 below for a characterization of Iw_u^+ -orbits which supports a non-zero Iwahori–Whittaker rank one local system.

4. Parity sheaves on partial affine flag varieties and equivalence relations

4.1. Pushforward and pullback of indecomposable parity complexes

Let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a facet and $\pi_{\mathbf{g}} : \text{Fl}_{\mathbf{a}_1}^{\circ} \rightarrow \text{Fl}_{\mathbf{g}}^{\circ}$ be the canonical projection. In this section, we will explain what is the effect of applying the functors $\pi_{\mathbf{g}}^*$ and $\pi_{\mathbf{g}*}$ to the Iwahori–Whittaker-equivariant indecomposable parity complexes that we described in Proposition 3.1.

Let us first state a lemma which will be needed below. Let (Λ, \leqslant) be a finite set endowed with a partial order \leqslant , admitting a unique minimal element λ_0 and equipped with a function $l : \Lambda \rightarrow \mathbb{Z}_{\geqslant 0}$ compatible with the order \leqslant (i.e. such that $\lambda \leqslant \mu \Rightarrow l(\lambda) \leqslant l(\mu)$). Let also $(X, (X_{\lambda})_{\lambda \in \Lambda})$ be a stratified variety over \mathbb{F} . We make the following assumptions on the stratification:

- for every $\lambda \in \Lambda$, X_{λ} is an affine space of dimension $l(\lambda)$ over \mathbb{F} , and we have $\overline{X_{\lambda}} = \bigsqcup_{\mu \leqslant \lambda} X_{\mu}$;
- there exist isomorphisms of \mathbb{F} -schemes (which we fix) $X_{\lambda} \simeq X_{\lambda_0} \times_{\mathbb{F}} \mathbb{A}_{\mathbb{F}}^{l(\lambda)-l(\lambda_0)}$ for every $\lambda \in \Lambda$;
- there exists a morphism of \mathbb{F} -schemes $q : X \rightarrow X_{\lambda_0}$ such that $q|_{X_{\lambda}}$ is induced by the canonical projection on the first component $X_{\lambda_0} \times_{\mathbb{F}} \mathbb{A}_{\mathbb{F}}^{l(\lambda)-l(\lambda_0)} \rightarrow X_{\lambda_0}$ for every $\lambda \in \Lambda$.

LEMMA 4.1. — We have an isomorphism in $D_c^b(X_{\lambda_0}, \mathbf{k})$:

$$q_! \underline{\mathbf{k}}_X \simeq \bigoplus_{\lambda \in \Lambda} \underline{\mathbf{k}}_{X_{\lambda_0}} [-2(l(\lambda) - l(\lambda_0))].$$

Proof. — The proof is done by induction on the cardinality of Λ . First note that the result is trivial when $\#\Lambda = 1$, so we assume from now on that $\#\Lambda \geqslant 2$. Let $\lambda \in \Lambda$ be a maximal element for \leqslant , so that we have

an open immersion $i : X_\lambda \hookrightarrow X$. Denoting by j the closed immersion $X' := X \setminus X_\lambda \hookrightarrow X$, we get a distinguished triangle

$$i_! i^* \underline{\mathbf{k}}_X \longrightarrow \underline{\mathbf{k}}_X \longrightarrow j_! j^* \underline{\mathbf{k}}_X \xrightarrow{+1}$$

to which we can apply the triangulated functor $q_!$:

$$(4.1) \quad q_! i_! i^* \underline{\mathbf{k}}_X \longrightarrow q_! \underline{\mathbf{k}}_X \longrightarrow q_! j_! j^* \underline{\mathbf{k}}_X \xrightarrow{+1}.$$

Since $q \circ i$ is the projection induced by $X_{\lambda_0} \times_{\mathbb{F}} \mathbb{A}_{\mathbb{F}}^{l(\lambda)-l(\lambda_0)} \rightarrow X_{\lambda_0}$ and $i^* \underline{\mathbf{k}}_X \simeq \underline{\mathbf{k}}_{X_{\lambda_0}}$, we have an isomorphism $q_! i_! i^* \underline{\mathbf{k}}_X \simeq \underline{\mathbf{k}}_{X_{\lambda_0}}[-2(l(\lambda) - l(\lambda_0))]$. On the other hand, we have $j^* \underline{\mathbf{k}}_X \simeq \underline{\mathbf{k}}_{X'}$, and the morphism $q' := q \circ j : X' \rightarrow X_{\lambda_0}$ satisfies $q'|_{X_\mu} = q|_{X_\mu}$ for every $\mu \neq \lambda$. Thus, we can apply our induction hypothesis to the stratified space $(X', (X_\mu)_{\mu \neq \lambda})$ to find that

$$q_! j_! j^* \underline{\mathbf{k}}_X \simeq q'_! \underline{\mathbf{k}}_{X'} \simeq \bigoplus_{\mu \neq \lambda} \underline{\mathbf{k}}_{X_{\lambda_0}}[-2(l(\mu) - l(\lambda_0))].$$

In particular, we see that $q_! i_! i^* \underline{\mathbf{k}}_X$ and $q_! j_! j^* \underline{\mathbf{k}}_X$ only have cohomologies in even degrees, so that the distinguished triangle (4.1) is split (recall that X_{λ_0} is an affine space, so it only admits cohomology in even degrees). Therefore we have an isomorphism

$$q_! \underline{\mathbf{k}}_X \simeq q_! i_! i^* \underline{\mathbf{k}}_X \oplus q_! j_! j^* \underline{\mathbf{k}}_X.$$

This concludes the proof of Lemma 4.1. \square

Fix a facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$. We denote by $W_{\mathbf{g}} \subset W$ the stabilizer of \mathbf{g} for the box action \square_1 and put $S_{\mathbf{g}} := S \cap W_{\mathbf{g}}$. It is well known that $(W_{\mathbf{g}}, S_{\mathbf{g}})$ is a finite Coxeter system.⁽⁶⁾ We will denote by $w_{\mathbf{g}}$ the element of maximal length in $W_{\mathbf{g}}$, and by $W^{\mathbf{g}}$ the set of elements $w \in W$ for which w is maximal in the left coset $wW_{\mathbf{g}}$. We put $w_0 := w_{\{0\}}$ (notice that $W_0 = W_{\{0\}}$). The set of elements $w \in W$ which are minimal in $W_0 w$ will be denoted by ${}_f W$, and we put

$${}_f W^{\mathbf{g}} = W^{\mathbf{g}} \cap {}_f W.$$

The following well-known result (which is a direct consequence of [5, Lemma 2.2]) will be useful throughout all the rest of this paper.

PROPOSITION 4.2. — *Let $w \in W^{\mathbf{g}}$. We have the following equivalences*

$$w \in {}_f W^{\mathbf{g}} \iff \forall r \in W_{\mathbf{g}}, wr \in {}_f W.$$

⁽⁶⁾ Note that $W_{\mathbf{g}}$ coincides with the stabilizer of the facet $\ell \cdot \mathbf{g} \subset \overline{\mathbf{a}_\ell}$ for the dilated box action \square_ℓ of W .

For any $w \in W^{\mathbf{g}}$, we choose a lift \dot{w} of w in $N_G(T)(\mathbb{K})$ (see [30, Section 4.2] for the construction of such a lift), and denote by $\mathcal{X}_w^{\mathbf{g}}$ the Iw_u^+ -orbit in $\mathrm{Fl}_{\mathbf{g}}^{\circ}$ of the image of \dot{w} under the canonical projection $LG(\mathbb{F}) \rightarrow \mathrm{Fl}_{\mathbf{g}}^{\circ}(\mathbb{F})$. We then have a stratification

$$(4.2) \quad (\mathrm{Fl}_{\mathbf{g}}^{\circ})_{\mathrm{red}} = \bigsqcup_{w \in W^{\mathbf{g}}} \mathcal{X}_w^{\mathbf{g}},$$

of the reduced ind-scheme associated with $\mathrm{Fl}_{\mathbf{g}}^{\circ}$, where each orbit $\mathcal{X}_w^{\mathbf{g}}$ is isomorphic to an affine space over \mathbb{F} whose dimension is equal to the length of the minimal element in $w_0 w W_{\mathbf{g}}$. All the properties concerning this stratification that we use in the sequel follow from the analogous standard facts concerning the opposite Iwahori subgroup $\mathrm{Iw}_u := \dot{w}_0 \mathrm{Iw}_u^+ \dot{w}_0$, cf. [2, Appendix A].

PROPOSITION 4.3. — *For any $w \in W^{\mathbf{g}}$, the Iw_u^+ -orbit $\mathcal{X}_w^{\mathbf{g}}$ supports a non-zero Iwahori–Whittaker equivariant local system if and only if $w \in {}_f W$, which means that $w \in W^{\mathbf{g}}$.*

For any $w \in {}_f W^{\mathbf{g}}$, we will denote by $\mathcal{L}_w^{\mathbf{g}}$, resp. $\mathcal{E}_w^{\mathbf{g}} \in \mathrm{Par}_{\mathcal{IW}}(\mathrm{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})$, resp. $\nabla_w^{\mathbf{G}} \in D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})$, the rank one Iwahori–Whittaker equivariant local system, resp. the indecomposable parity complex from Proposition 3.1, resp. the costandard perverse sheaf, associated with $\mathcal{X}_w^{\mathbf{g}}$.

Recall that $\pi_{\mathbf{g}} : \mathrm{Fl}_{\mathbf{a}_1}^{\circ} \rightarrow \mathrm{Fl}_{\mathbf{g}}^{\circ}$ is the canonical projection, which is a proper morphism of ind-schemes, and denote by $N_{\mathbf{g}}$ the length of $w_{\mathbf{g}}$. For any $v \in {}_f W$, we write \mathcal{X}_v (resp. \mathcal{L}_v , resp. \mathcal{E}_v , resp. ∇_v) instead of $\mathcal{X}_v^{\mathbf{a}_1}$ (resp. $\mathcal{L}_v^{\mathbf{a}_1}$, resp. $\mathcal{E}_v^{\mathbf{a}_1}$, resp. $\nabla_v^{\mathbf{a}_1}$). For every $w \in {}_f W^{\mathbf{g}}$ and $v \in {}_f W$, we fix isomorphisms of \mathbb{F} -schemes

$$(4.3) \quad \mathcal{X}_v \simeq \mathbb{A}_{\mathbb{F}}^{l(w_0 v)}, \quad \mathcal{X}_w^{\mathbf{G}} \simeq \mathbb{A}_{\mathbb{F}}^{l(w_0 w w_{\mathbf{g}})}.$$

Let $w \in {}_f W^{\mathbf{g}}$. It is important to note that $\pi_{\mathbf{g}}$ is a locally trivial fibration, with $\pi_{\mathbf{g}}^{-1}(\mathcal{X}_w^{\mathbf{G}}) = \bigsqcup_{x \in W_{\mathbf{g}}} \mathcal{X}_{wx}$, and that for any $h \in W_{\mathbf{g}}$ the morphism $\pi_{\mathbf{g}}|_{\mathcal{X}_{w w_{\mathbf{g}} h}} : \mathcal{X}_{w w_{\mathbf{g}} h} \rightarrow \mathcal{X}_w^{\mathbf{g}}$ identifies with the canonical projection on the first component

$$\mathcal{X}_{w w_{\mathbf{g}} h} \simeq \mathcal{X}_{w w_{\mathbf{g}}} \times \mathcal{X}_{w_0 h} \longrightarrow \mathcal{X}_w^{\mathbf{g}}.$$

The projection on the first component comes from the identification of $\mathcal{X}_{w w_{\mathbf{g}}}$ and $\mathcal{X}_w^{\mathbf{g}}$ with an affine space of dimension $l(w_0 w w_{\mathbf{g}})$ thanks to (4.3); fix $h \in W_{\mathbf{g}}$, and let us explain the first isomorphism above. For any $v \in W$, denote by \bar{v} the image of \dot{v} under the canonical projection $LG(\mathbb{F}) \rightarrow \mathrm{Fl}_{\mathbf{a}_1}^{\circ}(\mathbb{F})$. By construction we have

$$\mathcal{X}_{w w_{\mathbf{g}} h} = \dot{w}_0 \mathrm{Iw}_u \dot{w}_0 \cdot \overline{w w_{\mathbf{g}} h} = \dot{w}_0 \mathrm{Iw}_u \cdot \overline{w_0 w w_{\mathbf{g}} h}.$$

But thanks to our hypothesis that $w \in W^{\mathbf{g}}$, we know that $ww_{\mathbf{g}}$ is minimal in $ww_{\mathbf{g}}W_{\mathbf{g}}$, and our hypothesis that $w \in {}_f W$ implies that $wr \in {}_f W$ for all $r \in W_{\mathbf{g}}$ thanks to Proposition 4.2 (in particular this holds when $r \in \{w_{\mathbf{g}}, w_{\mathbf{g}}h\}$); these facts imply that we have

$$l(w_0ww_{\mathbf{g}}h) = l(w_0) + l(ww_{\mathbf{g}}h) = l(w_0) + l(ww_{\mathbf{g}}) + l(h) = l(w_0ww_{\mathbf{g}}) + l(h).$$

From these equalities of lengths and the fact that $\mathrm{Iw}_u \cdot \bar{x} \simeq \mathbb{A}_{\mathbb{F}}^{l(x)}$ for any $x \in W$, we deduce the first isomorphism below

$$\mathrm{Iw}_u \cdot \overline{w_0ww_{\mathbf{g}}h} \simeq \mathrm{Iw}_u \cdot \overline{w_0ww_{\mathbf{g}}} \times \mathrm{Iw}_u \cdot \overline{h} = \mathrm{Iw}_u \dot{w}_0 \cdot \overline{ww_{\mathbf{g}}} \times \mathrm{Iw}_u \dot{w}_0 \cdot \overline{w_0h},$$

and hence finally

$$\dot{w}_0 \mathrm{Iw}_u \cdot \overline{w_0ww_{\mathbf{g}}h} \simeq \dot{w}_0 \mathrm{Iw}_u \dot{w}_0 \cdot \overline{ww_{\mathbf{g}}} \times \dot{w}_0 \mathrm{Iw}_u \dot{w}_0 \cdot \overline{w_0h}.$$

The right-hand side corresponds to $\mathcal{X}_{ww_{\mathbf{g}}} \times \mathcal{X}_{w_0h}$.

LEMMA 4.4.

- (1) *The functors $\pi_{\mathbf{g}}^*$ and $\pi_{\mathbf{g}*}$ send parity complexes to parity complexes.*
- (2) *For any $v \in {}_f W^{\mathbf{g}}$, we have isomorphisms*

$$\pi_{\mathbf{g}}^*[N_{\mathbf{g}}](\mathcal{E}_v^{\mathbf{g}}) \simeq \mathcal{E}_v \quad \text{and} \quad \pi_{\mathbf{g}*}\mathcal{E}_v \simeq \bigoplus_{w \in W_{\mathbf{g}}} \mathcal{E}_v^{\mathbf{G}}[-2(l(w_0w) - l(w_0v))].$$

- (3) *Let $w \in {}_f W$, and write $w = w^{\mathbf{g}}h$ for some $w^{\mathbf{g}} \in W^{\mathbf{g}}, h \in W_{\mathbf{g}}$. Then we have*

$$\begin{aligned} \pi_{\mathbf{g}*}\nabla_w &\simeq \nabla_{w^{\mathbf{g}}}^{\mathbf{g}}[l(w_{\mathbf{g}}h)] \quad \text{if } w^{\mathbf{g}} \in {}_f W, \\ \pi_{\mathbf{g}*}\nabla_w &= 0 \quad \text{otherwise.} \end{aligned}$$

Proof. — The first (resp. third) point can be proven just as in [2, Proposition A.2] (resp. [2, Lemma A.1]).

Let us prove the second point, for which we will take back most of the arguments of [2, Lemma A.5]. The second isomorphism will arise while proving the first one.

We first make the following observation: let us write $X := \bigsqcup_{w \in W_{\mathbf{g}}} \mathcal{X}_{vw}$ and put $q := \pi_{\mathbf{g}}|_X$, the object $q^*\mathcal{L}_v^{\mathbf{g}}$ is a rank one Iwahori–Whittaker local system on X , whose restriction to each orbit \mathcal{X}_{vw} coincides with \mathcal{L}_{vw} (recall vw does belong to ${}_f W$ thanks to Proposition 4.2). In particular, $q^*\mathcal{L}_v^{\mathbf{g}}$ is indecomposable.

The object $\pi_{\mathbf{g}}^*[N_{\mathbf{g}}](\mathcal{E}_v^{\mathbf{g}})$ is parity by (1), with \mathcal{X}_v open in its support and its restriction to this stratum coinciding with $\mathcal{L}_v[l(w_0v)]$ (recall that \mathcal{X}_v is of dimension $l(w_0v)$ thanks to (4.3)). Therefore we can write

$$(4.4) \quad \pi_{\mathbf{g}}^*[N_{\mathbf{g}}](\mathcal{E}_v^{\mathbf{g}}) \simeq \mathcal{E}_v \oplus \mathcal{G},$$

for some object $\mathcal{G} \in \text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})$. Since the restriction of $\pi_{\mathbf{g}}^*[N_{\mathbf{g}}](\mathcal{E}_v^{\mathbf{g}})$ to X is the indecomposable object $q^*\mathcal{L}_v^{\mathbf{g}}[l(w_0v)]$, it must be isomorphic to $\mathcal{E}_v|_X$, because the latter is non-zero on \mathcal{X}_v .

Thus, the base change theorem implies the first isomorphism below

$$(4.5) \quad \begin{aligned} (\pi_{\mathbf{g}*}\mathcal{E}_v)|_{\mathcal{X}_v^{\mathbf{g}}} &\simeq q_*q^*\mathcal{L}_v^{\mathbf{g}}[l(w_0v)] \simeq q_!\mathbf{k}_X \otimes^L \mathcal{L}_v^{\mathbf{g}}[l(w_0v)] \\ &\simeq \bigoplus_{w \in W_{\mathbf{g}}} \mathcal{L}_v^{\mathbf{G}}[-2(l(w_0w) - l(w_0v))], \end{aligned}$$

the second isomorphism is implied by the projection formula (and the fact that $q_! = q_*$ because q is proper), and the third isomorphism is a direct application of Lemma 4.1. Since $\mathcal{X}_v^{\mathbf{g}}$ is open in the support of the parity complex $\pi_{\mathbf{g}*}\mathcal{E}_v$, we get an isomorphism

$$(4.6) \quad \pi_{\mathbf{g}*}\mathcal{E}_v \simeq \bigoplus_{w \in W_{\mathbf{g}}} \mathcal{E}_v^{\mathbf{g}}[-2(l(w_0w) - l(w_0v))] \oplus \mathcal{G}',$$

for some $\mathcal{G}' \in \text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})$. Applying $\pi_{\mathbf{G}*}$ to (4.4) and using (4.6), we deduce the following isomorphism of parity complexes:

$$\pi_{\mathbf{g}*}\pi_{\mathbf{g}}^*[N_{\mathbf{g}}]\mathcal{E}_v^{\mathbf{g}} \simeq \bigoplus_{w \in W_{\mathbf{g}}} \mathcal{E}_v^{\mathbf{g}}[-2(l(w_0w) - l(w_0v))] \oplus \pi_{\mathbf{g}*}\mathcal{G} \oplus \mathcal{G}'.$$

Now, for any parity complex $\mathcal{E} \in \text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})$ and $w \in {}_f W^{\mathbf{g}}$, one may write (following [37]):

$$\mathcal{E}|_{\mathcal{X}_w^{\mathbf{g}}} \simeq V(\mathcal{E})_w \otimes_{\mathbf{k}} \mathcal{L}_w^{\mathbf{g}},$$

where $V(\mathcal{E})_w$ is a finite-dimensional graded \mathbf{k} -vector space (and one can do likewise for any $\mathcal{E} \in \text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})$ and $w \in {}_f W$). From the description of $\pi_{\mathbf{g}}$ as a locally trivial fibration and from the first observation that was made at the beginning of the proof, we see that

$$\dim_{\mathbf{k}} \left(V(\pi_{\mathbf{g}}^*[N_{\mathbf{g}}]\mathcal{E}_v^{\mathbf{g}})_{uw} \right) = \dim_{\mathbf{k}} (V(\mathcal{E}_v^{\mathbf{g}})_u), \quad \forall (u, w) \in {}_f W^{\mathbf{g}} \times W_{\mathbf{g}},$$

where we forgot the grading of our vector spaces when taking their dimension. Moreover, the same arguments that were used to prove (4.5) imply that

$$\dim_{\mathbf{k}} \left(V(\pi_{\mathbf{g}*}\pi_{\mathbf{g}}^*[N_{\mathbf{g}}]\mathcal{E}_v^{\mathbf{g}})_u \right) = |W_{\mathbf{g}}| \cdot \dim_{\mathbf{k}} \left(V(\pi_{\mathbf{g}}^*[N_{\mathbf{g}}]\mathcal{E}_v^{\mathbf{g}})_u \right), \quad \forall u \in {}_f W^{\mathbf{g}}.$$

Thus we get

$$\dim_{\mathbf{k}} \left(V(\pi_{\mathbf{g}*}\pi_{\mathbf{g}}^*[N_{\mathbf{g}}]\mathcal{E}_v^{\mathbf{g}})_u \right) = \dim_{\mathbf{k}} (V(\mathcal{F})_u), \quad \forall u \in {}_f W^{\mathbf{g}},$$

where $\mathcal{F} := \bigoplus_{w \in W_{\mathbf{g}}} \mathcal{E}_v^{\mathbf{g}}[-2(l(w_0w) - l(w_0v))]$. We deduce that $V(\pi_{\mathbf{g}*}\mathcal{G})_u = V(\mathcal{G}')_u = 0$ for all $u \in {}_f W^{\mathbf{g}}$, so that $\pi_{\mathbf{g}*}\mathcal{G} \simeq \mathcal{G}' \simeq 0$ and $\pi_{\mathbf{g}*}\pi_{\mathbf{g}}^*[N_{\mathbf{g}}]\mathcal{E}_v^{\mathbf{g}} \simeq \mathcal{F}$.

This also implies that $\mathcal{G} \simeq 0$, and therefore we get the desired isomorphism

$$\pi_{\mathbf{g}}^*[N_{\mathbf{g}}](\mathcal{E}_v^{\mathbf{g}}) \simeq \mathcal{E}_v. \quad \square$$

4.2. The antispherical module

Let \mathcal{H} be the Hecke algebra associated with (W, S) , with standard basis $(H_w, w \in W)$, and denote by \mathcal{N} its antispherical module (we follow the notation of [31, Section 3], the antispherical module is denoted by $\mathcal{M}^{\text{asph}}$ in [30]), with standard basis $(N_w, w \in {}_f W)$ and ℓ -canonical basis $({}^\ell N_w, w \in {}_f W)$. We denote by $[\text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})]$ the split Grothendieck group of the additive category $\text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})$. We have a canonical isomorphism of groups

$$\text{ch} : [\text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})] \xrightarrow{\sim} \mathcal{N}$$

defined by

$$\text{ch}([\mathcal{F}]) := \sum_{w \in {}_f W} \left(\sum_{n \in \mathbb{Z}} \dim_{\mathbf{k}} \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})}(\mathcal{F}, \nabla_w[n]) v^n \right) N_w.$$

The morphism ch sends the indecomposable parity complex \mathcal{E}_w onto ${}^\ell N_w$. The ℓ -Kazhdan–Lusztig polynomials $({}^\ell n_{x,y}, x, y \in {}_f W)$ are defined by the equality

$$(4.7) \quad {}^\ell N_y = \sum_{x \in {}_f W} {}^\ell n_{x,y} N_x.$$

It makes sense to extend the definition of these polynomials to the whole group W , simply by putting ${}^\ell n_{x,y} = 0$ whenever x or y does not belong to ${}_f W$ (this consideration will slightly simplify the statement of the first point of Proposition 5.4). For any objects $\mathcal{E}, \mathcal{F} \in \text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})$, we set

$$\text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})}^{\bullet}(\mathcal{E}, \mathcal{F}) := \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})}(\mathcal{E}, \mathcal{F}[n]).$$

It follows that we have

$$(4.8) \quad {}^\ell n_{x,y}(1) = \dim_{\mathbf{k}} \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})}^{\bullet}(\mathcal{E}_y, \nabla_x).$$

When $\ell = 0$, the ℓ -canonical basis coincides with the usual Kazhdan–Lusztig basis from [31, Theorem 3.1], which is denoted by $(n_{x,y}, x, y \in {}_f W)$ there, so we have ${}^0 n_{x,y} = n_{x,y}$ (this is a consequence of the fact that the perversely shifted indecomposable parity complexes coincide with the intersection cohomology complexes when $\ell = 0$). We have the following easy and useful observation.

PROPOSITION 4.5. — Let $x, y \in {}_f W$. For any prime number ℓ , we have ${}^\ell n_{x,y}(1) \geq n_{x,y}(1)$.

Proof. — The arguments are the same as the one used in [6, Lemma 3.4], replacing tilting objects by indecomposable parity sheaves. \square

4.3. Equivalence relations on ${}_f W^g$

Let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a facet. In the sequel, we will consider a relation $\mathcal{R}_{\mathbf{g}}$ on the set ${}_f W^g$, defined by

$$w \mathcal{R}_{\mathbf{g}} w' \iff \text{Hom}_{\text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{g}}, \mathbf{k})}^{\bullet}(\mathcal{E}_w^g, \mathcal{E}_{w'}^g) \neq 0$$

for any $w, w' \in {}_f W^g$. The equivalence relation on ${}_f W^g$ generated by $\mathcal{R}_{\mathbf{g}}$ will be denoted by $\sim_{\mathbf{g}}$. The following results will be constantly used.

PROPOSITION 4.6. — Let $u, v \in {}_f W$ be such that ${}^\ell n_{u,v}(1) \neq 0$. Then $v \mathcal{R}_{\mathbf{a}_1} u$. Moreover, we get the same conclusion if $n_{u,v}(1) \neq 0$.

Proof. — Recall that, thanks to Proposition 4.5, we have

$$n_{u,v}(1) \leq {}^\ell n_{u,v}(1) = \dim_{\mathbf{k}} \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{a}_1}, \mathbf{k})}^{\bullet}(\mathcal{E}_v, \nabla_u),$$

so that our claim is a direct consequence of [21, Proposition 2.6]. \square

The relations $\mathcal{R}_{\mathbf{a}_1}$ and $\mathcal{R}_{\mathbf{g}}$ are actually equivalent on ${}_f W^g$.

PROPOSITION 4.7. — Let $w, w' \in {}_f W^g$. We have $w \mathcal{R}_{\mathbf{g}} w'$ iff $w \mathcal{R}_{\mathbf{a}_1} w'$.

Proof. — Using Lemma 2.2, we have

$$\begin{aligned} \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{a}_1}, \mathbf{k})}^{\bullet}(\mathcal{E}_w, \mathcal{E}_{w'}) &\simeq \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{a}_1}, \mathbf{k})}^{\bullet}(\pi_{\mathbf{g}}^*[N_{\mathbf{g}}]\mathcal{E}_w^g, \pi_{\mathbf{g}}^*[N_{\mathbf{g}}]\mathcal{E}_{w'}^g) \\ &\stackrel{\text{adjunction}}{\simeq} \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}, \mathbf{k})}^{\bullet}(\mathcal{E}_w^g, \pi_{\mathbf{g}*}\pi_{\mathbf{g}}^*\mathcal{E}_{w'}^g). \end{aligned}$$

From Lemma 2.2, we know that $\pi_{\mathbf{g}*}\pi_{\mathbf{g}}^*\mathcal{E}_{w'}^g$ is isomorphic to a finite direct sum of shifts of $\mathcal{E}_{w'}^g$, so the hom-spaces above are non-zero if and only if $\text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}, \mathbf{k})}^{\bullet}(\mathcal{E}_w^g, \mathcal{E}_{w'}^g)$ is non-zero. \square

Putting together the two previous propositions yields:

COROLLARY 4.8. — Let $w, w' \in {}_f W^g$. We have

$$n_{w',w}(1) \neq 0 \implies w \mathcal{R}_{\mathbf{g}} w'.$$

Remark 4.9. — Define a relation $\mathcal{R}'_{\mathbf{g}}$ on ${}_{\mathbf{f}}W^{\mathbf{g}}$ by

$$w\mathcal{R}'_{\mathbf{g}}w' \iff {}^{\ell}n_{w',w}(1) \neq 0.$$

Then, using Proposition 4.7 together with [21, Proposition 2.6], it is not difficult to show that the equivalence relation generated by $\mathcal{R}'_{\mathbf{g}}$ is equal to $\sim_{\mathbf{g}}$.

5. Determination of the equivalence classes

The main goal of this section is to show that, when the root system \mathfrak{R}^{\vee} (or, equivalently, \mathfrak{R}) is indecomposable and $\mathbf{g} \subset \overline{\mathbf{a}_1}$ is a non-special facet, the set ${}_{\mathbf{f}}W^{\mathbf{g}}$ consists of a single equivalence class for $\sim_{\mathbf{g}}$ (we will eventually see that the case where the root system is not indecomposable follows from this first case, see Proposition 5.18). As we will see in the end (cf. Subsection 6.7), the case where \mathbf{g} is a special facet can be deduced from the non-special case with the help of Smith–Treumann theory, see Proposition 6.9.

5.1. The regular case

In this short paragraph, we show that the set ${}_{\mathbf{f}}W$ is a single equivalence class for $\sim_{\mathbf{a}_1}$, i.e. that all the elements of ${}_{\mathbf{f}}W$ are in relation for $\sim_{\mathbf{a}_1}$. Although this case will be included in the more general statement of Theorem 5.21 (where \mathbf{a}_1 is replaced with a facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$ which is not a point), we give an independent proof here, which is much simpler since many difficulties do not appear yet.

Remark 5.1. — The result we get in Proposition 5.3 below implies (via the discussion at the end of Subsection 1.2) that the block associated with a “regular” dominant weight $\lambda \in \mathbb{X}_+^{\vee}$ (regular means inside of an alcove) is $W \bullet_{\ell} \lambda \cap \mathbb{X}_+^{\vee}$ (here we see \mathbb{X}_+^{\vee} as the weight poset of $\text{Rep}_{\mathbf{k}}(G^{\vee})$). This last result was already much simpler to get than the general description of the block associated with an arbitrary dominant weight, see [19, Section 2.4].

LEMMA 5.2. — *Let $w \in {}_{\mathbf{f}}W$. If $w \neq e$, then there exists $w' < w$ such that $w' \in {}_{\mathbf{f}}W$ and $w' \mathcal{R}_{\mathbf{a}_1} w$.*

Proof. — Writing a reduced expression for w , it is easy to see that there exists an $s \in S$ such that $ws < w$. Let \mathbf{g} be the wall fixed by s . Since $w \in {}_f W^{\mathbf{g}}$, Proposition 4.2 implies that $ws \in {}_f W$. By Lemma 2.2, we get

$$\begin{aligned} \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{a}_1}, \mathbf{k})}^\bullet(\mathcal{E}_w, \nabla_{ws}) &\simeq \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{a}_1}^\circ, \mathbf{k})}^\bullet(\pi_{\mathbf{g}}^*[1]\mathcal{E}_w^{\mathbf{g}}, \nabla_{ws}) \\ &\simeq \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}, \mathbf{k})}^\bullet(\mathcal{E}_w^{\mathbf{g}}, \pi_{\mathbf{g}*}[-1]\nabla_{ws}) \\ &\simeq \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}, \mathbf{k})}^\bullet(\mathcal{E}_w^{\mathbf{g}}, \nabla_w^{\mathbf{g}}) \neq 0 \end{aligned}$$

where the second line is obtained by adjunction. So if we put $w' := ws$, we get that ${}^\ell n_{w',w}(1) \neq 0$, and Proposition 4.6 allows us to conclude. \square

It is now straightforward to conclude.

PROPOSITION 5.3. — *The set ${}_f W$ consists of a single class for the equivalence relation $\sim_{\mathbf{a}_1}$.*

Proof. — Let $w \in {}_f W$. By the previous lemma and an induction on the length of w , we can see that $w \sim_{\mathbf{a}_1} e$. This concludes the proof. \square

The proof of the fact that all of the elements of ${}_f W^{\mathbf{g}}$ are in relation when $\mathbf{g} \subset \overline{\mathbf{a}_1}$ is an arbitrary non-special facet will be much more involved. This is due to the fact that, for an arbitrary $w \in {}_f W^{\mathbf{g}}$, there is no obvious choice of an element $w' < w$ in ${}_f W^{\mathbf{g}}$ such that $w' \sim_{\mathbf{g}} w$. The goal of the next subsection is to find such relations.

5.2. Some invariance properties of the ℓ -anti-spherical Kazhdan–Lusztig polynomials

The following result is a generalization of Lemma 5.2.

PROPOSITION 5.4. — *Let \mathbf{q} be a facet inside $\overline{\mathbf{a}_1}$ and $w \in {}_f W^{\mathbf{q}}$.*

- (1) *For all $w' \in {}_f W$, we have ${}^\ell n_{w',r,w}(1) = {}^\ell n_{w',w}(1)$ for all $r \in W_{\mathbf{q}}$.*
- (2) *Assume moreover that $\mathbf{g} \subset \overline{\mathbf{a}_1}$ is a facet such that $\mathbf{q} \subset \overline{\mathbf{g}}$. For all $w_1, w_2 \in wW_{\mathbf{q}} \cap W^{\mathbf{g}}$, we have $w_1 \sim_{\mathbf{g}} w_2$.*

Proof.

(1). — Let $w' \in {}_f W$ and write $w' = uh$, where u is the maximal element in $w'W_{\mathbf{q}}$, and $h \in W_{\mathbf{q}}$. We have the following sequence of isomorphisms⁽⁷⁾

⁽⁷⁾ Notice that we forget the grading of the vector spaces in those isomorphisms.

of \mathbf{k} -vector spaces for all $r \in W_{\mathbf{q}}$:

$$\begin{aligned} \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})}^{\bullet}(\mathcal{E}_w, \nabla_{w'r}) &\simeq \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{a}_1}^{\circ}, \mathbf{k})}^{\bullet}(\pi_{\mathbf{q}}^*[N_{\mathbf{q}}]\mathcal{E}_w^{\mathbf{q}}, \nabla_{w'r}) \\ &\stackrel{\text{adjunction}}{\simeq} \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{q}}, \mathbf{k})}^{\bullet}(\mathcal{E}_w^{\mathbf{q}}, (\pi_{\mathbf{q}})_*\nabla_{uhr}) \\ &\simeq \begin{cases} \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{q}}, \mathbf{k})}^{\bullet}(\mathcal{E}_w^{\mathbf{q}}, \nabla_u^{\mathbf{q}}) & \text{if } u \in {}_f W, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

where the first and last isomorphisms are consequences of the first and third point of Lemma 2.2 respectively. Thanks to the definition of the ℓ -anti-spherical polynomials (see (4.7)), this means that ${}^{\ell}n_{w'r, w}(1) = 0$ for all $r \in W_{\mathbf{q}}$ when $u \notin {}_f W$ (which trivially implies that ${}^{\ell}n_{w'r, w}(1) = {}^{\ell}n_{w', w}(1)$), and ${}^{\ell}n_{w'r, w}(1) = {}^{\ell}n_{u, w}(1) = {}^{\ell}n_{w', w}(1)$ otherwise.

(2). — First notice that, since $w \in {}_f W^{\mathbf{q}}$, we have that $wW_{\mathbf{q}} \subset {}_f W$ thanks to Proposition 4.2, so that

$$(5.1) \quad wW_{\mathbf{q}} \cap W^{\mathbf{g}} \subset {}_f W \cap W^{\mathbf{g}} = {}_f W^{\mathbf{g}}.$$

Applying the previous point to $w' = w$, we get that $n_{wr, w}(1) = n_{w, w}(1) = 1$ for every $r \in W_{\mathbf{q}}$, and therefore that $w\mathcal{R}_{\mathbf{g}}wr$ if $wr \in W^{\mathbf{g}}$ (which implies that $wr \in {}_f W^{\mathbf{g}}$ by (5.1)) thanks to Corollary 4.8. Thus, we have $w\mathcal{R}_{\mathbf{g}}w_i$ for $i \in \{1, 2\}$, and we conclude by transitivity that $w_1 \sim_{\mathbf{g}} w_2$. \square

The goal of the next subsection is to get a precise picture of the geometry of the affine space $\mathbb{X}^{\vee} \otimes_{\mathbb{Z}} \mathbb{R}$. By doing so, we will be able to use the previous relations together with the bijection between ${}_f W^{\mathbf{g}}$ and $\mathcal{A}_{\mathbf{g}}^+$ (the set of dominant facets which are in $W \cap \mathbf{g}$) to show that ${}_f W^{\mathbf{g}}$ consists of a single class for $\sim_{\mathbf{g}}$ (when \mathbf{g} is not a point).

5.3. Geometry of $\mathbb{X}^{\vee} \otimes_{\mathbb{Z}} \mathbb{R}$

In this subsection, we follow the terminology and some of the notations⁽⁸⁾ of both [24, Section 1] and [31, Section 4]. The box action \square_1 of W on the affine space $E := \mathbb{X}^{\vee} \otimes_{\mathbb{Z}} \mathbb{R}$ defines a set of hyperplanes \mathcal{H} (one could also work with the dilated box action \square_n for some $n \geq 1$ or with \bullet_n , cf. the setting of [24, Section 1]). For any facet \mathbf{p} (not necessarily included in $\overline{\mathbf{a}_1}$), we will denote by $W_{\mathbf{p}}$ its stabilizer in W for the box action. We define the set of strictly dominant elements by

$$\mathcal{C}_0^+ := \{\lambda \in E \mid \langle \lambda, \alpha \rangle > 0, \quad \forall \alpha \in \mathfrak{R}_+\}.$$

(8) The general reference, which is used in [24, Section 1], is [13].

Recall that the connected components of $E \setminus \bigcup_{H \in \mathcal{H}} H$ are products of open simplices called alcoves.⁽⁹⁾ An example is

$$\mathbf{a}_1 = \{\mu \in E \mid 0 < \langle \mu, \alpha \rangle < 1, \forall \alpha \in \mathfrak{R}_+\},$$

which we call the fundamental alcove. The box action on the set of alcoves \mathcal{A} can be extended to an action of \widetilde{W} , in which case the stabilizer of \mathbf{a}_1 is equal to $\Omega = \{w \in \widetilde{W} \mid l(w) = 0\}$. Thus, the assignment $w \mapsto w \square_1 \mathbf{a}_1$ yields a bijection $W \simeq \widetilde{W}/\Omega \xrightarrow{\sim} \mathcal{A}$, so that any alcove A can be written as $A = w \square_1 \mathbf{a}_1$ for a unique $w \in W$. This bijection $W \xrightarrow{\sim} \mathcal{A}$ allows us to define a right action of W on \mathcal{A} , induced by right multiplication on itself. We may also define an order \leqslant on \mathcal{A} , induced by the Bruhat order on W . Thus, for any $s \in S$ and $A \in \mathcal{A}$, we will have that $A < As$ if and only if $l(w) < l(ws)$, where $w \in W$ is such that $w \square_1 \mathbf{a}_1 = A$. Since As is obtained by reflecting A along its wall which is the W -conjugate of the wall of \mathbf{a}_1 fixed by s , this means that $A < As$ if and only if the number of hyperplanes separating A from \mathbf{a}_1 (i.e. the number of hyperplanes $H \in \mathcal{H}$ such that A and \mathbf{a}_1 are included in different connected components of $E \setminus H$) is smaller than the number of hyperplanes separating As from \mathbf{a}_1 (see [24, Section 1.4]). In the sequel, we will denote by $d(A)$ the number of hyperplanes separating A from \mathbf{a}_1 , so that we have $d(A) = l(w)$.

The set of dominant alcoves (i.e. those included in \mathcal{C}_0^+) will be denoted by \mathcal{A}^+ ; we have a bijection ${}_f W \xrightarrow{\sim} \mathcal{A}^+, w \mapsto w \square_1 \mathbf{a}_1$.

For any facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$, we define $\mathcal{A}_\mathbf{g} := \{w \square_1 \mathbf{g}, w \in W\}$ and $\mathcal{A}_\mathbf{g}^+ := \{w \square_1 \mathbf{g}, w \in {}_f W^\mathbf{g}\}$. One can show that $\mathcal{A}_\mathbf{g}^+$ is exactly the subset of facets of $\mathcal{A}_\mathbf{g}$ which are included in \mathcal{C}_0^+ . Note that the map $w \mapsto w \square_1 \mathbf{g}$ induces a bijection between $W^\mathbf{g}$, resp. ${}_f W^\mathbf{g}$, and $\mathcal{A}_\mathbf{g}$, resp. $\mathcal{A}_\mathbf{g}^+$. We will consider the left action of W on $\mathcal{A}_\mathbf{g}$ given by $w\mathbf{h} := w \square_1 \mathbf{h}$ for any $\mathbf{h} \in \mathcal{A}_\mathbf{G}$. Thus, if $\mathbf{h} = u\mathbf{G}$ for some $u \in W^\mathbf{G}$, then $w\mathbf{h} = wu\mathbf{G}$, and the element of $W^\mathbf{g}$ corresponding to the facet $w\mathbf{h}$ is the maximal element of $wuW_\mathbf{g}$. Also note that, by continuity of the action of W on E , if A is an alcove containing \mathbf{h} in its closure, then $w\mathbf{h}$ is the unique element of $\mathcal{A}_\mathbf{g}$ which is contained in the closure of wA . Let us state and prove the following easy result, which will be used a lot in the sequel.

PROPOSITION 5.5. — *Let \mathbf{h} be a facet containing a facet \mathbf{p} in its closure such that $\mathbf{p} \subset \mathcal{C}_0^+$. Then we have $\mathbf{h} \subset \mathcal{C}_0^+$. In particular if $\mathbf{g} \subset \overline{\mathbf{a}_1}$ is the*

(9) When the root system \mathfrak{R}^\vee is irreducible, the alcoves are open simplices. In general, \mathfrak{R}^\vee decomposes into a product of irreducible root systems, from which we deduce the decomposition of any alcove into a product of open simplices (cf. Subsection 5.4).

facet such that $\mathbf{h} \in \mathcal{A}_{\mathbf{g}}$, then there exists a unique element $w \in {}_f W^{\mathbf{g}}$ such that $w\mathbf{g} = \mathbf{h}$.

Proof. — By definition of \mathcal{C}_0^+ , any facet is either included in \mathcal{C}_0^+ or in $E \setminus \mathcal{C}_0^+$, the latter being a closed subset of E . If $\mathbf{h} \subset E \setminus \mathcal{C}_0^+$, then $\bar{\mathbf{h}} \subset E \setminus \mathcal{C}_0^+$, contradicting our assumption on \mathbf{p} . Therefore $\mathbf{h} \subset \mathcal{C}_0^+$. The rest of the proposition follows. \square

A *special facet* will be a zero dimensional facet by which a maximal number of hyperplanes passes through. An example is the point $\{0\}$, and all the other special facets are translates of this point by elements of \mathbb{X}^\vee , i.e. are of the form $\{\lambda\}$ for $\lambda \in \mathbb{X}^\vee$ (cf.[13, Chapter VI, Section 2, Proposition 3]). For any special facet \mathbf{v} , we will denote by $W_{\mathbf{v}}$ the stabilizer of \mathbf{v} , which is the subgroup of W generated by the reflections with respect to the hyperplanes passing through \mathbf{v} . If $\mathbf{v} = \{\lambda\}$, then $W_{\mathbf{v}} = t_\lambda W_0 t_{-\lambda}$. We will write $w_{\mathbf{v}} := t_\lambda w_0 t_{-\lambda}$, which is an element of $W_{\mathbf{v}}$.

LEMMA 5.6. — *Let \mathbf{u}, \mathbf{v} be special facets. Then $w_{\mathbf{u}} w_{\mathbf{v}}$ is a translation.*

Proof. — Write $\mathbf{v} = \{\lambda\}$, $\mathbf{u} = \{\mu\}$ for some $\lambda, \mu \in \mathbb{X}^\vee$, then we have

$$\begin{aligned} w_{\mathbf{u}} w_{\mathbf{v}} &= t_\mu w_0 t_{-\mu} t_\lambda w_0 t_{-\lambda} = w_0 t_{w_0 \mu} t_{-\mu} t_\lambda w_0 t_{-\lambda} \\ &= w_0^2 t_\mu t_{-w_0 \mu} t_{w_0 \lambda} t_{-\lambda} = t_{\mu - \lambda + w_0(\lambda - \mu)}. \end{aligned}$$

\square

Let \mathbf{v} be a special facet, a quarter with vertex \mathbf{v} is a connected component of

$$E \setminus \bigcup_{\substack{H \in \mathcal{H} \\ \mathbf{v} \subset H}} H.$$

When $\mathbf{v} = 0$, one such quarter is the dominant cone \mathcal{C}_0^+ . For a general \mathbf{v} , we denote by $\mathcal{C}_{\mathbf{v}}^+$ the quarter with vertex \mathbf{v} which is a translate of \mathcal{C}_0^+ , and we put $\mathcal{C}_{\mathbf{v}}^- := w_{\mathbf{v}} \mathcal{C}_{\mathbf{v}}^+$. We will also denote by $A_{\mathbf{v}}^+$, resp. $A_{\mathbf{v}}^-$, the unique alcove contained in $\mathcal{C}_{\mathbf{v}}^+$, resp. in $\mathcal{C}_{\mathbf{v}}^-$, containing \mathbf{v} in its closure. We clearly have $A_{\mathbf{v}}^- = w_{\mathbf{v}} A_{\mathbf{v}}^+$. Furthermore, let \mathcal{H}^* be the set of hyperplanes H of \mathcal{H} such that H is a wall of some $\mathcal{C}_{\mathbf{v}}^+$ for a special facet \mathbf{v} , we define *boxes* to be the connected components of $E \setminus \bigcup_{H \in \mathcal{H}^*} H$. Each alcove is contained in a unique box, and we denote by $\Pi_{\mathbf{v}}$ the box containing $A_{\mathbf{v}}^+$. In our context, if $\mathbf{v} = \{\lambda\}$ for some $\lambda \in \mathbb{X}^\vee$, we put $\Pi_\lambda := \Pi_{\mathbf{v}}$, and we get

$$(5.2) \quad \Pi_\lambda = \{\mu \in E \mid 0 < \langle \mu - \lambda, \alpha \rangle < 1, \quad \forall \alpha \in S_0\}.$$

Remark 5.7. — In [24], Lusztig defines Ω to be the group generated by the orthogonal reflections through the hyperplanes of \mathcal{H} , which is seen as acting on the right on E . In our context, this right action is the left action

of $W = \Omega$ on E , and the left action of W on the set of alcoves from [24] is our right action on \mathcal{A} .

PROPOSITION 5.8. — *Let $H \in \mathcal{H}$. Then, there exists a unique connected component E_1 of $E \setminus H$ that has a non-empty intersection with \mathcal{C}_v^+ for any special facet v . Moreover, there exists at least one special facet u such that $\mathcal{C}_u^- \cap E_1 = \emptyset$.*

Proof. — Let $f : E \rightarrow \mathbb{R}$ be a linear form (where E is seen as a real vector space with origin 0) and $x \in \mathbb{X}^\vee$ such that $H = x + f^{-1}(\{0\})$. Then the connected components of $E \setminus H$ are $E_1 := x + f^{-1}(\mathbb{R}_+^*)$ and $E_2 := x + f^{-1}(\mathbb{R}_-^*)$. Since $f^{-1}(\{0\}) \in \mathcal{H}$, \mathcal{C}_0^+ must be inside $f^{-1}(\mathbb{R}_+^*)$ or $f^{-1}(\mathbb{R}_-^*)$. We may assume that $\mathcal{C}_0^+ \subset f^{-1}(\mathbb{R}_+^*)$. Let $y \in \mathbb{X}$ and put $v := \{y\}$, so that $\mathcal{C}_v^+ = y + \mathcal{C}_0^+$. For any $u \in \mathcal{C}_0^+$, we can find $r > 0$ big enough so that $f(y - x + ru) > 0$ (because $f(u) > 0$), which implies that

$$y + ru \in (y + \mathcal{C}_0^+) \cap (x + f^{-1}(\mathbb{R}_+^*)) = \mathcal{C}_v^+ \cap E_1.$$

So E_1 has a non-empty intersection with any \mathcal{C}_v^+ . On the other hand, we have seen that $f^{-1}(\mathbb{R}_-^*) \cap \mathcal{C}_0^+ = \emptyset$, which means that $E_2 \cap \mathcal{C}_x^+ = \emptyset$. This shows that E_1 is the only connected component of $E \setminus H$ having the desired property.

Finally, we want to show that there exists a special facet u such that $\mathcal{C}_u^- \cap E_1 = \emptyset$. Recall that we have

$$\mathcal{C}_0^- = w_0 \mathcal{C}_0^+ = -\mathcal{C}_0^+.$$

Since $\mathcal{C}_0^+ \subset f^{-1}(\mathbb{R}_+^*)$, we must have $\mathcal{C}_0^- \cap f^{-1}(\mathbb{R}_+^*) = \emptyset$, and therefore $\mathcal{C}_x^- \cap E_1 = \emptyset$. \square

Let $H \in \mathcal{H}$, then $E \setminus H$ consists of two connected components E_H^- and E_H^+ . Thanks to Proposition 5.8, we can set E_H^+ to be the one connected component that has a non-empty intersection with \mathcal{C}_v^+ for any special facet v . Following [31, Section 4] (see also [24, Section 1.5]), we define a partial order \preceq on \mathcal{A} generated by the relations

$$A \preceq s_H A \quad \text{if} \quad A \in \mathcal{A}, H \in \mathcal{H}, s_H A \subset E_H^+,$$

where $s_H \in W$ denotes the affine reflection associated with the hyperplane H . One can show (cf. the proof of [31, Claim 4.14]) that this partial order \preceq coincides with the usual Bruhat order \leqslant on \mathcal{A}^+ . The following result will be crucial in the sequel. We let A be an alcove, $s \in S$ and H be the hyperplane containing the wall separating A and As .

PROPOSITION 5.9. — Assume that $A, As \in \mathcal{A}^+$ and that there exists a special point \mathbf{v} such that $w_{\mathbf{v}}A$ and $w_{\mathbf{v}}As$ belong to \mathcal{A}^+ . Then $A < As$ if and only if $w_{\mathbf{v}}As < w_{\mathbf{v}}A$.

Proof. — Since the two orders \leqslant and \preceq coincide on \mathcal{A}^+ and $As = s_H A$, the fact that $A < As$ is equivalent to $As \subset E_H^+$.

Because the actions of W on the right and on the left on \mathcal{A} commute, the hyperplane $H' := w_{\mathbf{v}} \square_1 H$ is the one separating $w_{\mathbf{v}}As$ from $w_{\mathbf{v}}A$. Using the fact that $w_{\mathbf{v}}A, w_{\mathbf{v}}As \in \mathcal{A}^+$, we see that $w_{\mathbf{v}}As < w_{\mathbf{v}}A$ if and only if $w_{\mathbf{v}}A$ is included in $E_{H'}^+$. Thus, to conclude it is enough to show that $E_{H'}^+ = w_{\mathbf{v}}E_H^-$, or equivalently that $E_{H'}^- = w_{\mathbf{v}}E_H^+$. Notice that since $w_{\mathbf{v}}E_H^+$ is connected, it equals either $E_{H'}^+$ or $E_{H'}^-$. By Proposition 5.8, we can find a special facet \mathbf{u} such that $\mathcal{C}_{\mathbf{u}}^- \cap E_H^+ = \emptyset$. Then we have $w_{\mathbf{v}}\mathcal{C}_{\mathbf{u}}^- \cap w_{\mathbf{v}}E_H^+ = \emptyset$. But

$$w_{\mathbf{v}}\mathcal{C}_{\mathbf{u}}^- = w_{\mathbf{v}}w_{\mathbf{u}}\mathcal{C}_{\mathbf{u}}^+,$$

and since $w_{\mathbf{v}}w_{\mathbf{u}}$ is a translation (thanks to Lemma 5.6), $w_{\mathbf{v}}\mathcal{C}_{\mathbf{u}}^-$ is of the form $\mathcal{C}_{\mathbf{w}}^+$ for some special facet \mathbf{w} . So $\mathcal{C}_{\mathbf{w}}^+ \cap w_{\mathbf{v}}E_H^+ = \emptyset$, which implies that $E_{H'}^- = w_{\mathbf{v}}E_H^+$. \square

Following [31, Section 4, Section 5], we consider the bijection $\mathcal{A} \rightarrow \mathcal{A}$, $A \mapsto \check{A}$. This bijection sends an alcove A inside of $\Pi_{\mathbf{v}}$ to the alcove $w_{\mathbf{v}}A$. We will denote by $A \mapsto \hat{A}$ the inverse of this bijection. For any alcove $B \subset \Pi := \Pi_0$, we see that $\check{B} = w_0B$. For a general alcove A , write it uniquely as $A = \lambda + B$ for some $\lambda \in \mathbb{X}^{\vee}$, $B \subset \Pi$, and it follows that $\check{A} = \lambda + w_0B$.

LEMMA 5.10. — Let $A \in \mathcal{A}$ and write $A = \lambda + B$, with $\lambda \in \mathbb{X}^{\vee}$, $B \subset \Pi$. We have $\hat{A} = \lambda + t_{2\rho^{\vee}}w_0B$.

Proof. — For all $\mu \in \Pi$, we have that $w_0(\mu - \rho^{\vee}) \in \Pi$. Thus, $w_0(-\rho^{\vee} + B) \subset \Pi$ for any alcove $B \subset \Pi$, so if we set $C := \rho^{\vee} + w_0(-\rho^{\vee} + B)$, we get $\check{C} = B$ and so

$$\hat{B} = C = 2\rho^{\vee} + w_0B = t_{2\rho^{\vee}}w_0B.$$

The general result follows easily, since any alcove A can be written uniquely as $A = \lambda + B$, with $\lambda \in \mathbb{X}^{\vee}$, $B \subset \Pi$, and $\hat{A} = \lambda + \hat{B}$. \square

PROPOSITION 5.11. — Let $A \in \mathcal{A}^+$. Then $\hat{A} \subset \rho^{\vee} + \mathcal{C}_0^+$.

Proof. — Let $A \in \mathcal{A}^+$, and write $A = \lambda + B$, for $\lambda \in \mathbb{X}^{\vee}$, $B \subset \Pi$. Since A is in \mathcal{A}^+ , we must have $\lambda \in \overline{\mathcal{C}^+} \cap \mathbb{X}^{\vee}$. Indeed, if $\mu \in \Pi$ and $\lambda \in \mathbb{X}^{\vee} \setminus \overline{\mathcal{C}^+}$, then one sees using (5.2) that $\lambda + \mu \notin \mathcal{C}_0^+$.

Thus, it is enough to prove the proposition for $A \subset \Pi$. But in this case, we have by Lemma 5.10

$$\hat{A} = t_{2\rho^\vee} w_0 A = \rho^\vee + w_0(-\rho^\vee + A).$$

Since $w_0(-\rho^\vee + A) \subset \Pi$, this concludes the proof. \square

We also need to define a bijection $\mathcal{A}_g \rightarrow \mathcal{A}_g, \mathbf{h} \mapsto \hat{\mathbf{h}}$ for an arbitrary facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$. One can do it as follows: let $\mathbf{h} \in \mathcal{A}_g$ and $A \in \mathcal{A}$ be such that $\mathbf{h} \subset \overline{A}$ and $A \preceq Aw$ for all $w \in W_g$ (such an alcove exists and is unique, thanks to the existence and unicity of a minimal element in a coset of a parabolic subgroup in a Coxeter group), then we define $\hat{\mathbf{h}}$ as the element of \mathcal{A}_g which sits inside the closure of \hat{A} .

COROLLARY 5.12. — For any $\mathbf{h} \in \mathcal{A}_g^+$, we have $\hat{\mathbf{h}} \subset \rho^\vee + \overline{\mathcal{C}^+}$.

Proof. — By definition, \mathbf{h} sits inside the closure of some alcove A . Notice that, since A contains \mathbf{h} in its closure, we must have $A \in \mathcal{A}^+$ thanks to Proposition 5.5. Thus $\hat{A} \subset \rho^\vee + \mathcal{C}_0^+$ by Proposition 5.11, so $\hat{\mathbf{h}} \subset \rho^\vee + \overline{\mathcal{C}^+}$. \square

Corollary 5.12 implies in particular that the operation $\mathbf{h} \mapsto \hat{\mathbf{h}}$ preserves \mathcal{A}_g^+ .

PROPOSITION 5.13. — Let $A \in \mathcal{A}^+$ be such that $Ar \geq A$ and $Ar \in \mathcal{A}^+$ for all $r \in W_g$. Then, for all $r \in W_g$, we have $\hat{A}r \leq \hat{A}$ and $\hat{A}r \in \mathcal{A}^+$.

Proof. — Let \mathbf{v} be the special point such that $\hat{A} \in \Pi_{\mathbf{v}}$. First notice that \hat{A} is included in $\rho^\vee + \mathcal{C}_0^+$ thanks to Proposition 5.11. Thus, if we denote by \mathbf{h} the facet of \mathcal{A}_g included in the closure of \hat{A} , we see that $\mathbf{h} \subset \rho^\vee + \overline{\mathcal{C}^+}$ and that \mathbf{h} is included in the closure of all the alcoves of $\hat{A}W_g$, so that they all belong to \mathcal{A}^+ thanks to Proposition 5.5. Since $A = w_{\mathbf{v}}\hat{A}$, we can apply Proposition 5.9 to get that $\hat{A}s < \hat{A}$ for all $s \in S_g$. From the general theory of maximal elements in cosets of parabolic subgroups in Coxeter groups, this implies that $\hat{A}w \leq \hat{A}$ for all $w \in W_g$. \square

The anti-spherical ℓ -Kazhdan–Lusztig polynomial $\ell n_{x,y}$ will also be denoted by $\ell n_{B,A}$ for the alcoves B, A corresponding to $x, y \in {}_{\mathbf{f}} W$. The following result will be of great importance for us.

LEMMA 5.14. — For any dominant alcove $A \in \mathcal{A}^+$, we have $n_{A,\hat{A}}(1) = 1$.

Proof. — This is an immediate consequence of [31, Theorem 5.1] (see also [31, Section 7]). \square

LEMMA 5.15. — Let $A \in \mathcal{A}^+$ and $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a facet such that $Ar \geq A$ and $Ar \in \mathcal{A}^+$ for all $r \in W_g$. Then we have $n_{Ar,\hat{A}}(1) = 1$ for all $r \in W_g$.

Proof. — By Lemma 5.14, we have that $n_{A, \hat{A}}(1) = 1$ and, by Proposition 5.4 (applied to the elements $w \in {}_f W^{\mathbf{g}}$, $w' \in {}_f W$ such that $w' \mathbf{a}_{\ell} = A$ and $w \mathbf{a}_{\ell} = \hat{A}$) combined with Proposition 5.13, we deduce that $n_{Ar, \hat{A}}(1) = 1$ for all $r \in W_{\mathbf{g}}$. \square

The following proposition will allow us to only consider facets which are far inside of the dominant cone.

PROPOSITION 5.16. — *Let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a facet and $\mathbf{h} \in \mathcal{A}_{\mathbf{g}}^+$. Denote by u , resp. \hat{u} , the elements of ${}_f W^{\mathbf{g}}$ corresponding to \mathbf{h} , resp. $\hat{\mathbf{h}}$ (i.e. such that $u\mathbf{g} = \mathbf{h}$, resp. $\hat{u}\mathbf{g} = \hat{\mathbf{h}}$). Then we have $n_{u, \hat{u}}(1) \neq 0$, so that*

$$\mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}, \mathbf{k})}^{\bullet}(\mathcal{E}_u^{\mathbf{g}}, \mathcal{E}_{\hat{u}}^{\mathbf{g}}) \neq 0.$$

Proof. — Recall that $w_{\mathbf{g}}$ is the element of maximal length in $W_{\mathbf{g}}$, and set $A := uw_{\mathbf{g}}\mathbf{a}_{\ell}$. Notice that $A \in \mathcal{A}^+$ thanks to Proposition 4.2 and, by construction, A is the alcove such that $\mathbf{h} \subset \overline{A}$, $Ar \in \mathcal{A}^+$ and $A \leqslant Ar$ for every $r \in W_{\mathbf{g}}$ (in particular $Ar \preceq A$ for all $r \in W_{\mathbf{g}}$), so that $\hat{\mathbf{h}} \subset \overline{\hat{A}}$. By Lemma 5.15, we have that $n_{Aw_{\mathbf{g}}, \hat{A}}(1) = 1 \neq 0$, and $\hat{A}w \leqslant \hat{A}$ for every $w \in W_{\mathbf{g}}$ thanks to Proposition 5.13. In particular we deduce that $\hat{u}\mathbf{a}_{\ell} = \hat{A}$. Thanks to Proposition 4.6, this implies that

$$\mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{a}_1}, \mathbf{k})}^{\bullet}(\mathcal{E}_u, \mathcal{E}_{\hat{u}}) \neq 0.$$

By Proposition 4.7, we get the desired result. \square

5.4. Facets which are not points

The root system \mathfrak{R}^{\vee} decomposes uniquely into a disjoint union $\mathfrak{R}^{\vee} = \mathfrak{R}_1^{\vee} \sqcup \cdots \sqcup \mathfrak{R}_t^{\vee}$ of irreducible root systems. This decomposition comes with a decomposition of the affine Weyl group $W = W_1 \times \cdots \times W_t$ (where each W_i is the affine Weyl group associated with \mathfrak{R}_i^{\vee}) and of the affine space $E = E_1 \times \cdots \times E_t$ (where each E_i is associated with \mathfrak{R}_i^{\vee}). Note that one also has dually a decomposition $\mathfrak{R} = \mathfrak{R}_1 \sqcup \cdots \sqcup \mathfrak{R}_t$ into irreducible root systems, and an isomorphism $G_1 \times \cdots \times G_r \xrightarrow{\sim} G$ induced by multiplication from the product of the minimal closed connected normal subgroups of G of positive dimension over \mathbb{F} , each G_i admitting \mathfrak{R}_i as a root system (cf. [18, Theorem and Corollary 27.5]). The point for us is that, for every i , the alcoves determined by the box action of W_i on E_i are open simplices, and that each alcove $A \in \mathcal{A}$ is a product of such alcoves $A_1 \times \cdots \times A_t$. More generally, each facet $\mathbf{h} \subset \overline{A}$ decomposes into a product of facets $\mathbf{h}_1 \times \cdots \times \mathbf{h}_t$, with $\mathbf{h}_i \subset \overline{A_i}$ for every i .

In this subsection we prove that, for any facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$ such that each \mathbf{g}_i is not a point (where $\mathbf{g} = \mathbf{g}_1 \times \cdots \times \mathbf{g}_t$), the set ${}_f W^{\mathbf{g}}$ is single equivalence class. We will prove in the next subsection that ${}_f W^{\mathbf{g}}$ is a single equivalence class exactly when each \mathbf{g}_i is a non-special facet. Let us first explain how we can reduce ourselves to the case where the root system is irreducible; we start with a technical lemma, where \mathbf{k} could be taken to be any field.

LEMMA 5.17. — *Let A be a local \mathbf{k} -algebra and B be a connected \mathbf{k} -algebra (we do not assume A and B to be commutative). Assume that there exists a \mathbf{k} -algebra morphism $A \rightarrow \mathbf{k}$ and that B is a finite dimensional \mathbf{k} -vector space. Then the ring $A \otimes_{\mathbf{k}} B$ is connected.*

Proof. — Denote by \mathfrak{m} the maximal ideal of A . Since we have a morphism $A \rightarrow \mathbf{k}$, we get an injection $A/\mathfrak{m} \hookrightarrow \mathbf{k}$, which is in fact an isomorphism $A/\mathfrak{m} \simeq \mathbf{k}$ as A/\mathfrak{m} is a \mathbf{k} -vector space. Therefore we get an isomorphism

$$A/\mathfrak{m} \otimes_{\mathbf{k}} B \simeq B,$$

from which we deduce that $A/\mathfrak{m} \otimes_{\mathbf{k}} B$ is a connected ring.

Thanks to Nakayama's lemma (cf. [22, Section 2, Proposition 4.2.3]), a morphism of finitely generated A -modules $f : M \rightarrow N$ is surjective if and only if

$$f \otimes \text{id}_{A/\mathfrak{m}} : M/\mathfrak{m}M \longrightarrow N/\mathfrak{m}N$$

is surjective. Since $A \otimes_{\mathbf{k}} B$ is finitely generated over A (because $\dim_{\mathbf{k}} B < \infty$), we can apply this last result to any endomorphism of the finitely generated A -module $A \otimes_{\mathbf{k}} B$, and deduce that an element $e \in A \otimes_{\mathbf{k}} B$ is invertible if and only if its image \bar{e} under the canonical projection $A \otimes_{\mathbf{k}} B \rightarrow A/\mathfrak{m} \otimes_{\mathbf{k}} B$ is invertible.

Let $e \in A \otimes_{\mathbf{k}} B$ be an idempotent, i.e. an element satisfying $e(e - 1) = 0$. Then $\bar{e} \in A/\mathfrak{m} \otimes_{\mathbf{k}} B$ is an idempotent, from which we deduce that $\bar{e} \in \{1, 0\}$ by connectedness of $A/\mathfrak{m} \otimes_{\mathbf{k}} B$. If $\bar{e} = 1$, then $e \in (A \otimes_{\mathbf{k}} B)^\times$ thanks to the previous paragraph, so $e(e - 1) = 0 \Rightarrow e = 1$. If $\bar{e} = 0$, then

$$\overline{e - 1} = \bar{e} - 1 = -1 \in (A/\mathfrak{m} \otimes_{\mathbf{k}} B)^\times,$$

from which we deduce once again that $e - 1 \in (A \otimes_{\mathbf{k}} B)^\times$ and thus $e = 0$. Therefore any idempotent of $A \otimes_{\mathbf{k}} B$ is trivial, which means that this ring is connected. \square

For the next result, notice that the decomposition $W = W_1 \times \cdots \times W_t$ carries on at the level of parabolic subgroups, so that we have equalities $W^{\mathbf{g}} = W_1^{\mathbf{g}_1} \times \cdots \times W_t^{\mathbf{g}_t}$ and ${}_f W^{\mathbf{g}} = {}_f W_1^{\mathbf{g}_1} \times \cdots \times {}_f W_t^{\mathbf{g}_t}$.

PROPOSITION 5.18. — Let $\mathbf{g} = \mathbf{g}_1 \times \cdots \times \mathbf{g}_t$ be a facet included in $\overline{\mathbf{a}_1}$, and $w = (w_1, \dots, w_t)$, $w' = (w'_1, \dots, w'_t)$ be elements of ${}_{\mathbf{f}} W^{\mathbf{g}}$. Then we have

$$(5.3) \quad w\mathcal{R}_{\mathbf{g}} w' \iff w_i \mathcal{R}_{\mathbf{G}_i} w'_i, \quad \forall i.$$

Moreover there is an equality

$$(5.4) \quad {}^{\ell}n_{w,w'}(1) = {}^{\ell}n_{w_1,w'_1}(1) \times \cdots \times {}^{\ell}n_{w_r,w'_r}(1).$$

Proof. — The isomorphism $G_1 \times \cdots \times G_r \xrightarrow{\sim} G$ induces an isomorphism from the product of the partial affine flag varieties $\mathrm{Fl}_{\mathbf{g}_i}^{\circ}$, each being relative to G_i , to the partial affine flag variety $\mathrm{Fl}_{\mathbf{g}}^{\circ}$ relative to G . This morphism is equivariant for the action of the product of the Iwahori subgroups $\mathrm{Iw}_1^+ \times \cdots \times \mathrm{Iw}_t^+$ on the left (where $\mathrm{Iw}_i^+ = L^+ G_i \cap \mathrm{Iw}^+$ for each i) and of Iw^+ on the right. Thus, external tensor product yields a functor

$$F : \prod_{i=1}^t D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}_i}^{\circ}, \mathbf{k}) \longrightarrow D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k}),$$

and one can easily check with the Künneth formula that F sends tuples of parity complexes to parity complexes.

Next, we claim that we have an isomorphism

$$F((\mathcal{E}_{w_1}^{\mathbf{g}_1}, \dots, \mathcal{E}_{w_t}^{\mathbf{g}_t})) \simeq \mathcal{E}_w^{\mathbf{g}}.$$

The two objects above are parity complexes and coincide when restricted to the stratum $\mathcal{X}_w^{\mathbf{g}}$, so we only need to check that the object $F((\mathcal{E}_{w_1}^{\mathbf{g}_1}, \dots, \mathcal{E}_{w_t}^{\mathbf{g}_t}))$ is indecomposable, which we will do by showing that its endomorphism ring is connected.

We claim that we have the following isomorphism of graded \mathbf{k} -vector spaces

$$(5.5) \quad \begin{aligned} \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})}^{\bullet} \left(\mathcal{E}_{w_1}^{\mathbf{g}_1} \boxtimes \cdots \boxtimes \mathcal{E}_{w_t}^{\mathbf{g}_t}, \mathcal{E}_{w'_1}^{\mathbf{g}_1} \boxtimes \cdots \boxtimes \mathcal{E}_{w'_t}^{\mathbf{g}_t} \right) \\ \simeq \bigotimes_{i=1}^t \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}_i}^{\circ}, \mathbf{k})}^{\bullet} \left(\mathcal{E}_{w_i}^{\mathbf{g}_i}, \mathcal{E}_{w'_i}^{\mathbf{g}_i} \right). \end{aligned}$$

Let us briefly explain how to obtain (5.5). Thanks to [1, Proposition 1.4.6] (the result is stated with sheaves for the analytic topology there, but is easily translated for the étale topology), we know that for every \mathcal{F} , $\mathcal{G} \in D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})$ we have a natural isomorphism of graded vector spaces

$$(5.6) \quad \begin{aligned} \mathrm{H}^{\bullet}(R\Gamma(R\mathrm{Hom}(\mathcal{F}, \mathcal{G}))) &\simeq \mathrm{H}^{\bullet}(R\mathrm{Hom}(\mathcal{F}, \mathcal{G})) \\ &\simeq \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k})}^{\bullet}(\mathcal{F}, \mathcal{G}), \end{aligned}$$

where $\underline{\text{Hom}}$ is the internal hom functor, and $H^\bullet(-)$ means the direct sum of cohomology groups. Thus, we only need to show that there exists an isomorphism

$$R\underline{\text{Hom}}(\mathcal{F}_1 \boxtimes \cdots \boxtimes \mathcal{F}_t, \mathcal{G}_1 \boxtimes \cdots \boxtimes \mathcal{G}_t) \simeq \bigotimes_{i=1}^t R\underline{\text{Hom}}(\mathcal{F}_i, \mathcal{G}_i)$$

for every $\mathcal{F}_i, \mathcal{G}_i \in D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}^\circ, \mathbf{k})$. This fact is proved in [11, Section 4.2.7(b)] for constructible complexes.

Next, notice that the \mathbf{k} -algebra $\text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}_i}^\circ, \mathbf{k})}(\mathcal{E}_{w_i}^{\mathbf{g}_i}, \mathcal{E}_{w'_i}^{\mathbf{g}_i})$ is local for each i (because $\mathcal{E}_{w_i}^{\mathbf{g}_i}$ is indecomposable), so that

$$A_i := \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}_i}^\circ, \mathbf{k})}^\bullet(\mathcal{E}_{w_i}^{\mathbf{g}_i}, \mathcal{E}_{w'_i}^{\mathbf{g}_i})$$

is also a local \mathbf{k} -algebra as its degree zero part is local (here we apply [17, Theorem 3.1]); moreover, restriction to the stratum $\mathcal{X}_{w_i}^{\mathbf{g}_i}$ yields a \mathbf{k} -algebra morphism from A_i to \mathbf{k} . Therefore we can apply Lemma 5.17 (recall that a local ring is connected) to deduce that the ring on the right-hand side of (5.5) is connected. Hence we deduce that the degree zero part of the left-hand side of (5.5) is connected (applying once again [17, Theorem 3.1]), proving the claim.

Thus, (5.5) can be rewritten as

$$\text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}^\circ, \mathbf{k})}^\bullet(\mathcal{E}_w^{\mathbf{g}}, \mathcal{E}_{w'}^{\mathbf{g}}) \simeq \bigotimes_{i=1}^t \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}_i}^\circ, \mathbf{k})}^\bullet(\mathcal{E}_{w_i}^{\mathbf{g}_i}, \mathcal{E}_{w'_i}^{\mathbf{g}_i}).$$

Since a finite tensor product of vector spaces is non-zero if and only if each vector space appearing in the product is non-zero, this isomorphism finishes implies the equivalence (5.3).

Now we prove (5.4). First notice that, by compatibility of the $*$ -pushforward with the external tensor product (cf. [11, Section 4.2.7(a)]), we get an isomorphism $\nabla_{w'}^{\mathbf{g}} \simeq \nabla_{w'_1}^{\mathbf{g}_1} \boxtimes \cdots \boxtimes \nabla_{w'_t}^{\mathbf{g}_t}$. Next, another use of (5.6) yields the second isomorphism below:

$$\begin{aligned} & \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}^\circ, \mathbf{k})}^\bullet(\mathcal{E}_w^{\mathbf{g}}, \nabla_{w'}^{\mathbf{g}}) \\ & \simeq \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}^\circ, \mathbf{k})}^\bullet(\mathcal{E}_{w_1}^{\mathbf{g}_1} \boxtimes \cdots \boxtimes \mathcal{E}_{w_t}^{\mathbf{g}_t}, \nabla_{w'_1}^{\mathbf{g}_1} \boxtimes \cdots \boxtimes \nabla_{w'_t}^{\mathbf{g}_t}) \\ & \simeq \bigotimes_{i=1}^t \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}_i}^\circ, \mathbf{k})}^\bullet(\mathcal{E}_{w_i}^{\mathbf{g}_i}, \nabla_{w'_i}^{\mathbf{g}_i}). \end{aligned}$$

Taking the dimensions in this isomorphism yields the desired equality. \square

Now that we understand how to reduce the study to the case of an irreducible root system thanks to (5.3), we treat this case in detail. We

start with the following easy lemma (which does not require \mathfrak{R}^\vee to be irreducible).

LEMMA 5.19. — *Let A be an alcove contained in $\rho^\vee + \mathcal{C}_0^+$. Then there exist alcoves*

$$A_r, A_{r-1}, \dots, A_1, A_0$$

contained in $\rho^\vee + \mathcal{C}_0^+$ such that

$$\rho^\vee + \mathbf{a}_1 = A_r \leqslant A_{r-1} \leqslant \dots \leqslant A_1 \leqslant A_0 = A$$

and A_i is obtained by reflecting A_{i-1} along one of its walls for each $i \in \llbracket 1, r-1 \rrbracket$.

Proof. — Since the order \leqslant is invariant under translation by ρ^\vee in \mathcal{A}^+ (because it coincides with the periodic order \preceq), it is equivalent to prove that for any alcove A contained in \mathcal{C}_0^+ , there exist alcoves $(A_i)_{0 \leqslant i \leqslant r}$ contained in \mathcal{C}_0^+ such that $(A_r, A_0) = (\mathbf{a}_1, A)$, $A_i \leqslant A_{i-1}$ and A_i is obtained by reflecting A_{i-1} along one of its walls. This last condition is equivalent to requiring that $A_i = A_{i-1}s$ for some simple reflection s . Now if we let w be the element of ${}_f W$ such that $w\mathbf{a}_1 = A$, we know by the proof of Lemma 5.2 that there exists some simple reflection s such that $w_1 := ws < w$, with $w_1 \in {}_f W$, so that we can put $A_1 := w_1\mathbf{a}_1$. We then conclude by induction. \square

See Figure 1.1 for an illustration of Lemma 5.19 (but notice that one needs to dilate the affine plane on the picture by ℓ^{-1} to find back the same setting).

LEMMA 5.20. — *Let $\mathbf{h} \in \mathcal{A}_g$. Pick a facet $\mathbf{p} \subset \bar{\mathbf{h}}$ and consider $\mathbf{h}' := v\mathbf{h}$ with $v \in W_p$. If $\mathbf{p} \subset \mathcal{C}_0^+$, then \mathbf{h} and \mathbf{h}' belong to \mathcal{A}_g^+ , and we have $w \sim_g w'$, where $w, w' \in {}_f W^g$ are such that $\mathbf{h} = w\mathbf{g}$ and $\mathbf{h}' = w'\mathbf{g}$.*

Proof. — Since \mathbf{p} is inside \mathcal{C}_0^+ , the same is true for all the facets $v\mathbf{h}$, with $v \in W_p$, since those contain \mathbf{p} in their closure. In particular \mathbf{h} and \mathbf{h}' belong to \mathcal{A}_g^+ thanks to Proposition 5.5.

Let $w \in {}_f W^g$ be such that $w\mathbf{g} = \mathbf{h}$. By construction, there exists a facet $\mathbf{q} \subset \bar{\mathbf{g}}$ such that $w\mathbf{q} = \mathbf{p}$. In particular, we get that $W_p = wW_q w^{-1}$ and, for any $v \in W_p$, we have

$$v\mathbf{h} = vw\mathbf{g} = wr\mathbf{g},$$

where $r \in W_q$ is such that $wrw^{-1} = v$. Let $u \in W_g$ be such that wru is maximal in wW_g . We still have $wrug = \mathbf{h}'$, and since $\mathbf{h}' \in \mathcal{A}_g^+$, we must have $wru \in {}_f W^g$, so $w' = wru$, with $ru \in W_q$. Thanks to the second point of Proposition 5.4, we conclude that $w \sim_g w'$ (more precisely, if we let

$w'' \in {}_f W^g$ be the maximal element in wW_q (see Figure 5.1 below), then we have $n_{w,w''}(1) \neq 0$ and $n_{w',w''}(1) \neq 0$. \square

We have now enough ingredients to conclude.

THEOREM 5.21. — *Assume that \mathfrak{R}^\vee is irreducible and let $g \subset \overline{\mathbf{a}_1}$ be a facet which is not a point. Then ${}_f W^g$ consists of a single class for the equivalence relation \sim_g .*

Proof. — We introduce a new notation for this proof: if $\mathbf{h}, \mathbf{h}' \in \mathcal{A}_g^+$ are such that

$$\mathbf{h} = w\mathbf{g}, \quad \mathbf{h}' = w'\mathbf{g}$$

for some $w, w' \in {}_f W^g$, then we write $\mathbf{h} \sim_g \mathbf{h}'$ if $w \sim_g w'$. We want to show that \mathcal{A}_g^+ consists of a single equivalence class for \sim_g .




Figure 5.1. Lemma 5.20 in type C_2 . The black dot is \mathbf{p} , while the $W_{\mathbf{p}}$ -conjugates of \mathbf{h} are represented by blue lines, and the red lines are the other walls fixed by $W_{\mathbf{p}}$.

By Proposition 5.16, we have $\mathbf{h} \sim_g \hat{\mathbf{h}}$ for any $\mathbf{h} \in \mathcal{A}_g^+$ and, thanks to Corollary 5.12, we know that $\hat{\mathbf{h}} \subset \rho^\vee + \overline{\mathcal{C}^+}$. Thus, it is enough to show that all the elements of \mathcal{A}_g^+ lying inside of $\rho^\vee + \overline{\mathcal{C}^+}$ are in relation. For this, we will show that any facet satisfying this condition is in relation with the unique representative of its W -orbit in $\rho^\vee + \overline{\mathbf{a}_\ell}$.

Pick \mathbf{h} inside of $\rho^\vee + \overline{\mathcal{C}^+}$ and let $A \subset \rho^\vee + \mathcal{C}_0^+$ be an alcove containing \mathbf{h} in its closure. Let also $A_r, A_{r-1}, \dots, A_1, A_0 := A$ be alcoves as in Lemma 5.19, and denote by $s_i \in S$ the reflection such that $A_i s_i = A_{i-1}$.

Since g is not a point and $\overline{\mathbf{a}_1}$ is a simplex (because \mathfrak{R}^\vee is indecomposable), the face of the simplex fixed by any $s \in S$ (which is the closure of

the facet fixed by s) must have a nonzero intersection⁽¹⁰⁾ with $\bar{\mathbf{g}}$ inside $\bar{\mathbf{a}_1}$. This means that for any $s \in S$, there exists a facet $\mathbf{q} \subset \bar{\mathbf{g}}$ such that $s \in W_{\mathbf{q}}$.

For $i \in \llbracket 1, r \rrbracket$, denote by \mathbf{q}_i a facet such that $\mathbf{q}_i \subset \bar{\mathbf{g}}$ and $s_i \in W_{\mathbf{q}_i}$; for $i \in \{0, \dots, r\}$, denote by w_i the element of ${}_f W$ such that $w_i \mathbf{a}_1 = A_i$ and put $\mathbf{h}_i := w_i \mathbf{g}$, $\mathbf{p}_i := w_i \mathbf{q}_i$. Notice that $A_i s_i = w_i s_i w_i^{-1} A_i$ and that $\mathbf{h}_i \subset \bar{A}_i$. By construction, we have the following data

$$\begin{aligned} \mathbf{p}_i &\subset \bar{\mathbf{h}}_i, \quad \text{with } \mathbf{h}_0 = \mathbf{h} \quad \text{and} \quad \mathbf{h}_r \subset \rho^\vee + \bar{\mathbf{a}_1}, \\ \mathbf{h}_{i-1} &= w_i s_i w_i^{-1} \mathbf{h}_i, \quad \forall i \in \llbracket 1, r \rrbracket \quad \text{with } w_i s_i w_i^{-1} \in W_{\mathbf{p}_i}. \end{aligned}$$

Since the alcoves A_i are inside of $\rho^\vee + \mathcal{C}_0^+$, the facets \mathbf{p}_i are inside of \mathcal{C}_0^+ for all i . Thus we can apply Lemma 5.20 to get that $\mathbf{h}_i \sim_{\mathbf{g}} \mathbf{h}_{i-1}$ for every i , and by transitivity $\mathbf{h}_r \sim_{\mathbf{g}} \mathbf{h}$. This concludes the proof of Theorem 5.21. \square

5.5. Non-special facets which are points

In order to get the statement of Theorem 5.21 for non-special facets which are points, we need to be able to say more about the anti-spherical Kazhdan–Lusztig polynomials $n_{B,A}$. For that, we will make use of the combinatorial data linking the “periodic” polynomial $p_{B,A}$ of [31, Remark 4.4] (which is very closely related to Lusztig’s polynomial $Q_{B,A}$ from [24]) and the anti-spherical polynomials.

Recall the partial order \preceq introduced in Subsection 5.3. The following lemma will be helpful throughout all of this subsection.

LEMMA 5.22. — *Let \mathbf{v} be a special facet and C be an alcove contained in \mathcal{C}_0^+ . Then C is maximal in $W_{\mathbf{v}}C$ for \preceq .*

Proof. — If $D \in W_{\mathbf{v}}C$ is different from C , then there exists a hyperplane H passing through \mathbf{v} which separates D from \mathcal{C}_0^+ (one may take a well chosen wall of \mathcal{C}_0^+), so that $D \subset E_H^-$ and $D \prec s_H D$, with $s_H D \in W_{\mathbf{v}}D$. Assume that $s_H D \neq C$, then $s_H D$ is not included in \mathcal{C}_0^+ (because C is the only alcove of $W_{\mathbf{v}}D$ contained in \mathcal{C}_0^+), so that we can once again find some alcove $D_2 \in W_{\mathbf{v}}D$ satisfying $s_H D \prec D_2$. Since $W_{\mathbf{v}}D$ is a finite set, we can repeat this process a finite number of times until finding an alcove $D_n \in W_{\mathbf{v}}D$ which is contained in \mathcal{C}_0^+ and such that $D \prec D_n$, so that $D_n = C$. Therefore C is the maximal element in $W_{\mathbf{v}}C$ for \preceq . \square

(10) Recall that, for a positive integer n , an n -simplex Δ inside of an affine space E of dimension n is defined as the convex hull of $n+1$ vertices (those vertices being $n+1$ points not lying on a same hyperplane). For $0 \leq k \leq n$, a k -face of Δ is a subset consisting of the convex hull of $k+1$ vertices of Δ . From these definitions, it is straightforward to show that an $(n-1)$ -face of Δ has a non-empty intersection with any k -face, for $k \geq 1$.

We now recall the definition of the “periodic” module \mathcal{P} , which is the free left $\mathbb{Z}[v^{\pm 1}]$ -module with basis \mathcal{A} , equipped with a structure of right \mathcal{H} -module satisfying

$$(5.7) \quad \forall s \in S, \quad AC_s = \begin{cases} As + vA, & \text{if } A \prec As \\ As + v^{-1}A, & \text{if } As \prec A, \end{cases}$$

where $C_s := H_s + v$ (these data do define a right action of \mathcal{H} thanks to [31, Lemma 4.1]). One then defines the submodule $\mathcal{P}^\circ \subset \mathcal{P}$ as the right \mathcal{H} -submodule generated by the elements of the form

$$E_\lambda := \sum_{z \in W_0} v^{\ell(z)}(\lambda + z\mathbf{a}_1), \quad \lambda \in \mathbb{X}^\vee.$$

Notice that this definition parallels Lusztig’s definition of

$$e_{\{\lambda\}} := \sum_{A \in \mathcal{A}, \{\lambda\} \subset \bar{A}} A$$

from [24, Section 1.7], because the set of alcoves $\{\lambda + z\mathbf{a}_1, z \in W_0\}$ is equal to the set $\{A \in \mathcal{A}, \{\lambda\} \subset \bar{A}\}$.

Recall that a morphism $f : M \rightarrow N$ of right \mathcal{H} -modules is called *skew linear* if it satisfies $f(xH_w) = f(x)(H_{w^{-1}})^{-1}$, $f(xv) = f(x)v^{-1}$ for every $x \in M, w \in W$. It can be shown (cf. [31, Theorem 4.3]) that \mathcal{P}° admits a unique \mathcal{H} -skew linear involution $(\cdot) : \mathcal{P}^\circ \rightarrow \mathcal{P}^\circ$ such that $\overline{E_\lambda} = E_\lambda$ for all $\lambda \in \mathbb{X}^\vee$, and that for all $A \in \mathcal{A}$ there exists a unique $\underline{P}_A \in \mathcal{P}^\circ$ which is self dual with respect to this involution, with $\underline{P}_A \in A + \sum_B v\mathbb{Z}[v]B$. The \underline{P}_A form a $\mathbb{Z}[v^{\pm 1}]$ -basis of \mathcal{P}° , and the periodic polynomials are defined via the formula

$$\underline{P}_A = \sum_B p_{B,A} B.$$

The following result follows quite directly from the constructions, but will be of great importance for us.

LEMMA 5.23. — *Let $A \in \mathcal{A}$ and \mathbf{v} be the special facet such that $A \subset \Pi_{\mathbf{v}}$. Then we have*

$$p_{wC,A}(1) = p_{C,A}(1) \quad \text{for all } C \in \mathcal{A} \quad \text{and} \quad w \in W_{\mathbf{v}}.$$

Proof. — Denote by \mathcal{P}_1 the free left \mathbb{Z} -module $\mathbb{Z} \otimes_{\mathbb{Z}[v^{\pm 1}]} \mathcal{P}$, where \mathbb{Z} is seen as a $\mathbb{Z}[v^{\pm 1}]$ -module through the map $\mathbb{Z}[v^{\pm 1}] \rightarrow \mathbb{Z}, v \mapsto 1$, and let $\varphi : \mathcal{P} \rightarrow \mathcal{P}_1$ be the morphism of \mathbb{Z} -modules induced by sending v to 1. In particular we have

$$\varphi(\underline{P}_A) = \sum_B p_{B,A}(1)B.$$

The left and right actions of W on \mathcal{A} endow \mathcal{P}_1 with a structure of left and right $\mathbb{Z}[W]$ -module, and one can check with (5.7) that $\varphi(PH_w) = \varphi(P)w$ for every $w \in W$ and $P \in \mathcal{P}$. In particular, $\varphi(PH_w)$ belongs to the right $\mathbb{Z}[W]$ -submodule generated by $\varphi(P)$, which we denote by $\varphi(P)\mathbb{Z}[W]$. From [31, Remark 4.4], we know that $p_{B,A}(1) = Q_{B,A}(1)$ for every alcoves A, B , where $Q_{B,A}$ is Lusztig's polynomial from [24]. But $\underline{D}_A := \sum_B Q_{B,A}B$ belongs to the right \mathcal{H} -submodule generated by e_v thanks to [24, Theorem 2.15], from which we deduce that $\varphi(\underline{P}_A)$ belongs to $e_v\mathbb{Z}[W]$. Now e_v is invariant under the left action of W_v by construction, but since the left and right actions of W on \mathcal{P}_1 commute, we get that $wP = P$ for every $P \in e_v\mathbb{Z}[W]$, concluding the proof. \square

For any $x \in W$ and $A \in \mathcal{A}$, write $A = \lambda + B$ for a unique $\lambda \in \mathbb{X}^\vee$ and $B \subset \Pi$, and put $x * A := x\lambda + B$. We will denote by N_A the element that we denoted by N_x in Subsection 3.3, where $x\mathbf{a}_1 = A$, and we define (following [31, Proposition 5.2]) the $\mathbb{Z}[v^{\pm 1}]$ -linear application

$$\text{res} : \mathcal{P} \longrightarrow \mathcal{N},$$

which sends an alcove A to N_A if $A \in \mathcal{A}^+$, and to 0 otherwise. Finally, for any $A \in \mathcal{A}$, set

$$\text{alt } \underline{P}_A := \sum_{x \in W_0} (-1)^{l(x)} \underline{P}_{x * A}.$$

The link between periodic and antispherical polynomials is made explicit by the following result, which is [31, Theorem 5.3(1)].

PROPOSITION 5.24. — *For any alcove $A \subset \rho^\vee + \mathcal{C}_0^+$, we have*

$$\underline{N}_A = \text{res alt } \underline{P}_A.$$

For any special facet v and alcove $A \in \mathcal{A}$ such that $A \subset \Pi_v$, we define the set

$$S_A = \{C \mid wC \preceq A, \forall w \in W_v\}.$$

The next few results will help us utilize Proposition 5.24. More precisely, our main goal in Proposition 5.28 will be to determine for which alcoves C one has $n_{C,A} = p_{C,A}$, where the alcove $A \subset \rho^\vee + \mathcal{C}_0^+$ is fixed. As recalled in the lemma below, our interest for the set S_A comes from the fact that it contains the support of \underline{P}_A .

LEMMA 5.25. — *Let $A \in \mathcal{A}$ and v be the special facet such that $A \subset \Pi_v$.*

- (1) *We have the implication $p_{B,A} \neq 0 \Rightarrow B \in S_A$.*
- (2) *We have*

$$S_A = \{wC \mid w \in W_v, C \preceq A \text{ and } C \subset \mathcal{C}_v^+\}.$$

(3) Assume that there exists some facet $\mathbf{G} \subset \overline{\mathbf{a}_1}$ such that $A \in A_{\mathbf{v}}^+ W_{\mathbf{g}}$, and that $\mathbf{v} \subset x\mathcal{C}_0^+$ for some $x \in W_0 \setminus \{\text{id}\}$ (this simply means that \mathbf{v} is inside another quarter with vertex $\{0\}$ than \mathcal{C}_0^+). Then we have

$$D \in S_A \implies D \not\subset \rho^\vee + \mathcal{C}_0^+.$$

Proof.

(1). — This is [31, Proposition 4.22].

(2). — Let us write

$$S'_A = \{wC \mid w \in W_{\mathbf{v}}, C \preceq A \text{ and } C \subset \mathcal{C}_{\mathbf{v}}^+\}.$$

The inclusion $S_A \subset S'_A$ just follows from the fact that any alcove $C \in \mathcal{A}$ has a $W_{\mathbf{v}}$ conjugate wC inside of $\mathcal{C}_{\mathbf{v}}^+$. Conversely, if C is an alcove such that $C \subset \mathcal{C}_{\mathbf{v}}^+$ and $C \preceq A$, then we have $wC \preceq C$ for every $w \in W_{\mathbf{v}}$ thanks to Lemma 5.22, so $wC \preceq A$ for every $w \in W_{\mathbf{v}}$. This proves the inclusion $S'_A \subset S_A$, and concludes the proof of the first point.

(3). — Let $C \subset \mathcal{C}_{\mathbf{v}}^+$ be an alcove such that $C \preceq A$.

We claim that C belongs to $A_{\mathbf{v}}^+ W_{\mathbf{g}}$. Since \preceq is invariant under translations, we can translate everything by $-\mu$, where $\mathbf{v} = \{\mu\}$, so that we are reduced to the case where $\mathbf{v} = \{0\}$. More precisely, if we denote by $w_\mu \in W$ the element such that $w_\mu \mathbf{a}_1 = A_{\mathbf{v}}^+$, by $w \in W_{\mathbf{g}}$ the element such that $A = A_{\mathbf{v}}^+ w$ and set $\omega_\mu := t_{-\mu} w_\mu$, then we get that $\omega_\mu \mathbf{a}_1 = \mathbf{a}_1$ and

$$\begin{aligned} C \preceq A &\iff t_{-\mu} C \preceq t_{-\mu} A = t_{-\mu}(w_\mu \mathbf{a}_1 w) = t_{-\mu}(w_\mu w \mathbf{a}_1) = \omega_\mu w \mathbf{a}_1 \\ &= (\omega_\mu w \omega_\mu^{-1}) \omega_\mu \mathbf{a}_1 = (\omega_\mu w \omega_\mu^{-1}) \mathbf{a}_1. \end{aligned}$$

So now, if we set $\mathbf{g}' := \omega_\mu \mathbf{g}$, we get the inequality $t_{-\mu} C \preceq w' \mathbf{a}_1$, with $w' = \omega_\mu w \omega_\mu^{-1} \in W_{\mathbf{g}'}$. Since the alcoves $t_{-\mu} C$ and $w' \mathbf{a}_1$ belong to \mathcal{A}^+ (because they are translates by $-\mu$ of alcoves in $\mathcal{C}_{\mathbf{v}}^+$), we get that $t_{-\mu} C \leq w' \mathbf{a}_1$, so that $t_{-\mu} C = w'' \mathbf{a}_1$ for some $w'' \leq w'$, from which we easily deduce that $w'' \in W_{\mathbf{g}'}$; this means that $w'' = \omega_\mu u \omega_\mu^{-1}$ for some $u \in W_{\mathbf{g}}$, so

$$\begin{aligned} C &= t_\mu(\omega_\mu u \omega_\mu^{-1}) \mathbf{a}_1 = t_\mu(\omega_\mu u \omega_\mu^{-1}) t_{-\mu} t_\mu \mathbf{a}_1 \\ &= (w_\mu u)(w_\mu^{-1} t_\mu) \mathbf{a}_1 = (w_\mu u) \omega_\mu^{-1} \mathbf{a}_1 = w_\mu u \mathbf{a}_1 = A_{\mathbf{v}}^+ u. \end{aligned}$$

Denote by \mathbf{h} the W -conjugate of \mathbf{g} contained in $\overline{A_{\mathbf{v}}^+}$. In particular, we get that \mathbf{h} belongs to both $A_{\mathbf{v}}^+$ and \overline{C} (because $C \in A_{\mathbf{v}}^+ W_{\mathbf{g}}$ thanks to the previous paragraph), so that $\overline{C} \cap \overline{A_{\mathbf{v}}^+} \neq \emptyset$. From this and the description of S_A given in (2), we deduce that for any alcove $D \in S_A$, there exists some $w \in W_{\mathbf{v}}$ such that

$$\overline{D} \cap w \overline{A_{\mathbf{v}}^+} \neq \emptyset$$

(just write $D = wC$ for some alcove C as above). But $wA_{\mathbf{v}}^+ \subset x\mathcal{C}_0^+$ for any $w \in W_{\mathbf{v}}$ (because $x^{-1}wA_{\mathbf{v}}^+$ is an alcove containing $x^{-1}\mathbf{v} \subset \mathcal{C}_0^+$ in its closure, so that $x^{-1}wA_{\mathbf{v}}^+ \subset \mathcal{C}_0^+$ thanks to Proposition 5.5), so we get ultimately that if an alcove $D \in S_A$ is contained in \mathcal{C}_0^+ , then \bar{D} must have a non-empty intersection with a hyperplane separating \mathcal{C}_0^+ from $x\mathcal{C}_0^+$, so that D is not contained in $\rho^{\vee} + \mathcal{C}_0^+$. \square

LEMMA 5.26. — Assume that the root system \mathfrak{R}^{\vee} is irreducible. Let λ be a nonzero weight contained in $\overline{\mathcal{C}^+} \cap \mathbb{X}^{\vee}$, \mathbf{v} be a special facet, C be an alcove contained in $\prod_{\mathbf{v}}$ and put $A := \lambda + C$. If $\bar{A} \cap \overline{A_{\mathbf{v}}^+} \neq \emptyset$, we must have $C = A_{\mathbf{v}}^+$.

Proof. — In order to simplify the notations, we may and will reduce ourselves to the case where $\mathbf{v} = \{0\}$. Let us write

$$C = \{\mu \in E \mid n_{\alpha} - 1 < \langle \mu, \alpha \rangle < n_{\alpha}, \quad \forall \alpha \in \mathfrak{R}_+\}$$

for some integers n_{α} . First notice that if $\alpha', \alpha \in \mathfrak{R}_+$ are such that $\alpha \leq \alpha'$ (i.e. $\alpha' - \alpha$ is a sum of positive roots), then we must have $n_{\alpha} \leq n_{\alpha'}$. Indeed, since $C \subset \mathcal{C}_0^+$, we know that $\langle \mu, \alpha \rangle \leq \langle \mu, \alpha' \rangle$ for every $\mu \in C$, so in particular we get that

$$n_{\alpha} - 1 < \langle \mu, \alpha \rangle \leq \langle \mu, \alpha' \rangle < n_{\alpha'},$$

proving the claim.

Since $C \subset \prod_{\mathbf{v}}$, we know that $n_{\alpha} = 1$ for every $\alpha \in S_0$, and thus $n_{\alpha} \geq 1$ for every $\alpha \in \mathfrak{R}_+$. The assumption that $\bar{A} \cap \overline{\mathbf{a}_1} \neq \emptyset$ implies that we have

$$n_{\alpha} + \langle \lambda, \alpha \rangle \in \{1, 2\} \quad \text{for every } \alpha \in \mathfrak{R}_+.$$

Therefore we must have $\langle \lambda, \alpha \rangle \in \{0, 1\}$ for every $\alpha \in \mathfrak{R}_+$. Since $\lambda \neq 0$, there exists some $\alpha_0 \in S_0$ such that $\langle \lambda, \alpha_0 \rangle > 0$. So let $\beta = \sum_{S_0} c_{\alpha} \alpha \in \mathfrak{R}_+$ be the longest root (cf. [13, Chapter 6, Proposition 25], this is where we need the assumption that \mathfrak{R} is irreducible). In particular we have $c_{\alpha} \geq 1$ for every $\alpha \in S_0$, so that $\langle \lambda, \beta \rangle > 0$, and thus $n_{\beta} = 1$. But for every $\alpha \in \mathfrak{R}_+$, we have $\alpha \leq \beta$, so we get that $n_{\alpha} \leq n_{\beta}$, and finally $n_{\alpha} = 1$. This means that $C = \mathbf{a}_1$. \square

The following corollary can be visualized on Figure 5.2 below.

COROLLARY 5.27. — Let \mathbf{v} be a special facet, $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a non-special facet and $A \in A_{\mathbf{v}}^+ W_{\mathbf{g}}$ be the alcove which is maximal in $A_{\mathbf{v}}^+ W_{\mathbf{g}}$ for \preceq . Then A belongs to $\prod_{\mathbf{v}}$.

Proof. — Let us first explain why $A \subset \mathcal{C}_{\mathbf{v}}^+$. Denote by \mathbf{h}' the element of $\mathcal{A}_{\mathbf{g}}$ which belongs to the closure of $A_{\mathbf{v}}^+$. We have the inclusion $A_{\mathbf{v}}^+ \subset \mathcal{C}_{\mathbf{v}}^+$,

so we know that \mathbf{h}' is included in $\overline{\mathcal{C}_v^+}$. If $C \in A_v^+ W_{\mathbf{g}}$ is an alcove which is not included in \mathcal{C}_v^+ , then there exists a hyperplane H of the boundary of \mathcal{C}_v^+ which contains \mathbf{h}' and separates C from \mathcal{C}_v^+ , so we clearly have the inclusion $C \subset E_H^-$, and therefore $C \prec s_H C$. Since $s_H \in W_{\mathbf{h}'}$, the alcove $s_H C$ must belong⁽¹¹⁾ to $A_v^+ W_{\mathbf{g}}$. This shows that the maximal element in $A_v^+ W_{\mathbf{g}}$ for \preceq must be contained in \mathcal{C}_v^+ , whence $A \subset \mathcal{C}_v^+$.

Now let us show that $A \subset \Pi_v$. For this we may and will assume until the end of this proof that \mathfrak{R} is irreducible. Assume that this inclusion does not hold, then because $A \subset \mathcal{C}_v^+$, we can write $A = \lambda + C$ for some nonzero weight $\lambda \in \overline{\mathcal{C}_v^+} \cap \mathbb{X}^\vee$ and some alcove $C \subset \Pi_v$. But we also have $\overline{A} \cap \overline{A_v^+} \neq \emptyset$ (both of those sets contain \mathbf{h}'), therefore we can apply Lemma 5.26 (here we use the assumption that \mathfrak{R} is irreducible) to see that $C = A_v^+$. So we are in the following situation:

$$(5.8) \quad \lambda + A_v^+ \in A_v^+ W_{\mathbf{g}}.$$

Let $\mu \in \mathbb{X}^\vee$ be such that $\{\mu\} = \mathbf{v}$, and denote by $w_\mu \in W$ the element such that $w_\mu \mathbf{a}_1 = A_v^+$. Putting $\omega_\mu := t_{-\mu} w_\mu$ and applying $t_{-\mu}$ to (5.8), we get that

$$\lambda + \omega_\mu \mathbf{a}_1 \in \omega_\mu \mathbf{a}_1 W_{\mathbf{g}} = (\omega_\mu W_{\mathbf{g}} \omega_\mu^{-1}) \omega_\mu \mathbf{a}_1,$$

and, using the fact that ω_μ fixes the fundamental alcove \mathbf{a}_1 , we obtain

$$(5.9) \quad \lambda + \mathbf{a}_1 \in W_{\mathbf{g}'} \mathbf{a}_1, \quad \text{with} \quad \mathbf{g}' := \omega_\mu \mathbf{g}.$$

Since $\omega_\mu \in \Omega$ (because ω_μ fixes \mathbf{a}_1) and \mathbf{g} is a non-special facet, \mathbf{g}' is also a non-special facet⁽¹²⁾ included in the closure of the fundamental alcove. From (5.9) we deduce that $\mathbf{g}' \subset \lambda + \overline{\mathbf{a}_1}$. But $\lambda \in \overline{\mathcal{C}_v^+} \cap \mathbb{X}^\vee$ is nonzero, so $\langle \lambda, \beta \rangle \geq 1$, where $\beta \in \mathfrak{R}_+$ is the longest root. Since $\mathbf{g}' \subset \overline{\mathbf{a}_1}$, we have that $\langle \mu, \beta \rangle \leq 0$ for every $\mu \in -\lambda + \mathbf{g}'$ (because $\langle \mu', \beta \rangle \leq 1$ for all $\mu' \in \overline{\mathbf{a}_1}$); therefore the inclusion $-\lambda + \mathbf{g}' \subset \overline{\mathbf{a}_1}$ is possible only when $\langle \mu, \beta \rangle = 0$ for all $\mu \in -\lambda + \mathbf{g}'$. Since $\alpha \leq \beta$ for all $\alpha \in \mathfrak{R}_+$ we get that

$$0 \leq \langle \mu, \alpha \rangle \leq \langle \mu, \beta \rangle = 0, \quad \forall \alpha \in \mathfrak{R}_+, \mu \in -\lambda + \mathbf{g}'$$

which means that $\mu = 0$ and so $\mathbf{g}' = \{\lambda\}$, contradicting our assumption that \mathbf{g}' is not a special facet. This concludes the proof of the Corollary 5.27. \square

We are now ready to prove our central result.

(11) Indeed, write $C = w \mathbf{a}_1$ for some $w \in W$, then $\mathbf{h}' = w \mathbf{g}$ and $w^{-1} s_H w \in W_{\mathbf{g}}$, so that $s_H C = C(w^{-1} s_H w) \in C W_{\mathbf{g}} = A_v^+ W_{\mathbf{g}}$.

(12) Indeed, the action of W (and hence of Ω) on E obviously permutes the set of hyperplanes \mathcal{H} , so it also permutes the set of special facets by construction.

PROPOSITION 5.28. — Let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a non-special facet, $\mathbf{h} \in \mathcal{A}_{\mathbf{g}}^+$ be such that $\mathbf{h} \subset \rho^\vee + \overline{\mathcal{C}^+}$ and $B \subset \rho^\vee + \mathcal{C}_0^+$ be an alcove containing \mathbf{h} in its closure. Let also \mathbf{v} be a special facet such that $\mathbf{v} \subset \overline{B}$, and A be the alcove which is maximal in $A_{\mathbf{v}}^+ W_{\mathbf{g}}$ for \leqslant . Then we have $A \subset \Pi_{\mathbf{v}}$ and

$$n_{Br,A}(1) = 1, \quad \forall r \in W_{\mathbf{g}}.$$

Proof. — Let \mathbf{h}' be the element of $\mathcal{A}_{\mathbf{g}}$ which is included in $\overline{A_{\mathbf{v}}^+}$. By construction (i.e. because $\mathbf{v} \subset \rho^\vee + \overline{\mathcal{C}^+}$), we have that $A_{\mathbf{v}}^+ \subset \rho^\vee + \mathcal{C}_0^+$, so that in particular $\mathbf{h}' \subset \mathcal{C}_0^+$. Thus, applying Proposition 5.5, we know that all the alcoves of $A_{\mathbf{v}}^+ W_{\mathbf{g}}$ are contained in \mathcal{A}^+ (because they contain \mathbf{h}' in their closure), so that the orders \preceq and \leqslant coincide in $A_{\mathbf{v}}^+ W_{\mathbf{g}}$. Therefore A is maximal in $A_{\mathbf{v}}^+ W_{\mathbf{g}}$ for \preceq , so is included in $\Pi_{\mathbf{v}}$ thanks to Corollary 5.27.

By Lemma 5.24 and because $A \subset \rho^\vee + \mathcal{C}_0^+$, we have that

$$(5.10) \quad \sum_C n_{C,A} N_C = \sum_{x \in W_0} (-1)^{l(x)} \sum_C p_{C,x*A} \operatorname{res} C.$$

Our next goal is to use this formula to show that

$$n_{C,A} = p_{C,A} \quad \text{for all } C \subset \rho^\vee + \mathcal{C}_0^+.$$

For that we will prove that when the alcove $C \subset \rho^\vee + \mathcal{C}_0^+$ is fixed and $x \in W_0 \setminus \{\text{id}\}$, we have $p_{C,x*A} = 0$; this will follow from a careful study of the periodic polynomials.

Fix $x \in W_0 \setminus \{\text{id}\}$, and notice that $x*A \subset \Pi_{x\mathbf{v}}$ since $A \subset \Pi_{\mathbf{v}}$. We can then apply the first point of Lemma 5.25:

$$p_{D,x*A} \neq 0 \implies D \in S_{x*A}.$$

But we have that $x*A \in A_{x\mathbf{v}}^+ W_{\mathbf{g}}$ (because $x*A$ and $A_{x\mathbf{v}}^+$ are just translates of A and $A_{\mathbf{v}}^+$ respectively, by the same translation of $\mathbb{Z}\mathfrak{R}^\vee$) and $x\mathbf{v} \subset x\mathcal{C}_0^+$, so Lemma 5.25(3) tells us that

$$D \in S_{x*A} \implies D \not\subset \rho^\vee + \mathcal{C}_0^+.$$

This fact, combined with (5.10), implies that we have

$$(5.11) \quad n_{C,A} = p_{C,A}, \quad \forall C \subset \rho^\vee + \mathcal{C}_0^+.$$

On the other hand, recall the result of Lemma 5.23:

$$(5.12) \quad p_{wC,A}(1) = p_{C,A}(1), \quad \forall w \in W_{\mathbf{v}}.$$

Thus, for any alcove C included in $\rho^\vee + \mathcal{C}_0^+$, (5.11) and (5.12) yield

$$(5.13) \quad n_{wC,A}(1) = n_{C,A}(1), \quad \forall w \in W_{\mathbf{v}} \text{ such that } wC \subset \rho^\vee + \mathcal{C}_0^+.$$

By construction, there exist elements $w \in W_{\mathbf{v}}$, $r_0 \in W_{\mathbf{g}}$ such that $B = wA_{\mathbf{v}}^+$ and $A_{\mathbf{v}}^+ = Ar_0$, so that $B = wAr_0$. Since $B \subset \rho^\vee + \mathcal{C}_0^+$, we can apply (5.13) to get that

$$n_{B,A}(1) = n_{Ar_0,A}(1).$$

Finally, since A is maximal in $AW_{\mathbf{g}}$ for \leqslant , we can apply successively the first point of Proposition 5.4 (where the facet \mathbf{q} from this proposition is our current \mathbf{g} , and where the elements $w \in {}_f W^{\mathbf{g}}$, $w' \in {}_f W$ are such that $w\mathbf{a}_1 = A$ and $w'\mathbf{a}_1 = B$) to obtain

$$n_{Br,A}(1) = n_{B,A}(1) = n_{Ar_0,A}(1) = n_{A,A}(1) = 1, \quad \forall r \in W_{\mathbf{g}}. \quad \square$$

See Figure 5.2 for an illustration of the setting of the previous Proposition.




Figure 5.2. Proposition 5.28 in type C_2 . The black dot is \mathbf{v} and the thick black lines represent the affine hyperplanes fixed by the reflections in $W_{\mathbf{v}}$. The red dots are the $W_{\mathbf{v}}$ -conjugates of \mathbf{h} , and the set of gray alcoves represents $\{Br, r \in W_{\mathbf{g}}\}$.

Remark 5.29. — Keep the notations of the previous proposition. By positivity of the coefficients of the anti-spherical Kazhdan–Lusztig polynomials, the fact that $n_{Br,A}(1) = 1$ for all $r \in W_{\mathbf{g}}$ means that the polynomial $n_{Br,A}$ is always a non-zero monomial.

COROLLARY 5.30. — Let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a non-special facet, $\mathbf{h} \in \mathcal{A}_{\mathbf{g}}^+$ be such that $\mathbf{h} \subset \rho^\vee + \overline{\mathcal{C}^+}$, $B \subset \rho^\vee + \mathcal{C}_0^+$ be an alcove such that $\mathbf{h} \subset \overline{B}$, and let $\mathbf{v} \subset \rho^\vee + \overline{\mathcal{C}^+}$ be a special facet such that $\mathbf{v} \subset \overline{B}$. If we let $u \in W_{\mathbf{v}}$ be such that $B' := uB$ is included in $\rho^\vee + \overline{\mathcal{C}^+}$ and denote by \mathbf{h}' the element of $\mathcal{A}_{\mathbf{g}}^+$ included in \overline{B}' , then we have $w' \sim_{\mathbf{g}} w$, where $w', w \in {}_f W^{\mathbf{g}}$ are such that $\mathbf{h} = wg$ and $\mathbf{h}' = w'\mathbf{g}$.

Proof. — Denote by A the alcove which is maximal in $A_{\mathbf{v}}^+ W_{\mathbf{g}}$ for \leqslant , and by \mathbf{h}'' the element of $\mathcal{A}_{\mathbf{g}}$ included in \bar{A} . Recall that $A \subset \Pi_{\mathbf{v}}$ thanks to Proposition 5.28, so that in particular $\mathbf{h}'' \in \mathcal{A}_{\mathbf{g}}^+$. So if we let w'' be the element of ${}_f W^{\mathbf{g}}$ which satisfies $w'' \mathbf{g} = \mathbf{h}''$ and $r \in W_{\mathbf{g}}$ be the element such that Br is maximal in $BW_{\mathbf{g}}$ for \leqslant (so that $w\mathbf{a}_1 = Br$ and $w''\mathbf{a}_1 = A$), then Proposition 5.28 tells us that

$$n_{Br, A}(1) = 1,$$

from which we deduce that $w \sim_{\mathbf{g}} w''$ thanks to Corollary 4.8. The same reasoning (considering this time the alcove B' containing both \mathbf{h}' and \mathbf{v} in its closure) shows that $w' \sim_{\mathbf{g}} w''$, so finally $w \sim_{\mathbf{g}} w'$ by transitivity. \square

We can finally complete the proof of Theorem 5.21.

THEOREM 5.31. — *Assume that \mathfrak{R}^\vee is irreducible and let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a non-special facet. Then ${}_f W^{\mathbf{g}}$ consists of a single class for the equivalence relation $\sim_{\mathbf{g}}$.*

Proof. — Thanks to Theorem 5.21, we may and will assume that \mathbf{g} is a point. With the notation of the proof of Theorem 5.21, recall that we want to show that $\mathcal{A}_{\mathbf{g}}^+$ consists of a single equivalence class for $\sim_{\mathbf{g}}$. Also recall from the proof of Theorem 5.21 that it is enough to show that all the elements of $\mathcal{A}_{\mathbf{g}}^+$ lying inside of $\rho^\vee + \overline{\mathcal{C}^+}$ are in relation.

Pick \mathbf{h} inside of $\rho^\vee + \overline{\mathcal{C}^+}$ and let $A \subset \rho^\vee + \mathcal{C}_0^+$ be an alcove containing \mathbf{h} in its closure. Let also $A_r, A_{r-1}, \dots, A_1, A_0 := A$ be alcoves as in Lemma 5.19, and denote by $s_i \in S$ the reflection such that $A_i s_i = A_{i-1}$.

Since \mathfrak{R} is irreducible, there exists a unique simple reflection $\sigma \in S$ which does not belong to the finite Weyl group W_0 . Moreover, because $\overline{\mathbf{a}_1}$ is a simplex, there is only one facet inside $\overline{\mathbf{a}_1}$ which is a point and which is not included in the hyperplane fixed by σ : this is the special facet $\{0\}$. Thus, since \mathbf{g} is not special, it must be included in the wall fixed by σ , or in other words we have $\sigma \in W_{\mathbf{g}}$.

For $i \in \llbracket 0, \dots, r \rrbracket$, denote by w_i the element of ${}_f W$ such that $w_i \mathbf{a}_1 = A_i$ and put $\mathbf{h}_i := w_i \mathbf{g}$. Notice that $A_i s_i = w_i s_i w_i^{-1} A_i$ and that $\mathbf{h}_i \subset \overline{A_i}$. By construction, we also have

$$\mathbf{h}_{i-1} = w_i s_i w_i^{-1} \mathbf{h}_i, \quad \forall i \in \llbracket 1, \dots, r \rrbracket \quad \text{and} \quad \mathbf{h}_0 = \mathbf{h}.$$

Fix $i \in \llbracket 1, \dots, r \rrbracket$. Now, if $s_i = \sigma$, then $w_i s_i w_i^{-1}$ fixes \mathbf{h}_i , so that $\mathbf{h}_i = \mathbf{h}_{i-1}$. So assume that $s_i \neq \sigma$, which implies that $s_i \in W_0$, and put $\mathbf{v}_i := w_i \{0\}$. Then we see that \mathbf{v}_i is a special facet included in $\overline{A_i}$, which thus satisfies the inclusion $\mathbf{v}_i \subset \rho^\vee + \overline{\mathcal{C}^+}$, with $w_i s_i w_i^{-1} \in W_{\mathbf{v}_i}$. Therefore

we may apply Corollary 5.30 to see that $\mathbf{h}_i \sim_{\mathbf{g}} \mathbf{h}_{i-1}$. By transitivity of the relation we get $\mathbf{h} \sim_{\mathbf{g}} \mathbf{h}_r$, concluding the proof. \square

Remark 5.32. — Take back the notations of the previous proof. Corollary 5.30 (when \mathbf{g} is a point) and Lemma 5.20 (when \mathbf{g} is not a point) actually allow us to be a bit more precise: for all $i \in \llbracket 1, \dots, r \rrbracket$, there exists an element $u_i \in {}_f W^{\mathbf{g}}$ (which is the maximal element in $W_{\mathbf{v}_i} w_i$ in Corollary 5.30, and the maximal element in $w_i W_{\mathbf{q}_i}$ in Lemma 5.20) such that $n_{w_i, u_i}(1) \neq 0$ and $n_{w_{i-1}, u_i}(1) \neq 0$.

One of the advantages of our proof is that it gives an explicit way of linking an element $w \in {}_f W^{\mathbf{g}}$ to the unique element v of ${}_f W^{\mathbf{g}}$ satisfying $v\mathbf{g} \subset \rho^\vee + \overline{\mathbf{a}_1}$. This allows us to give the following result.

COROLLARY 5.33. — *Assume that \mathfrak{R}^\vee is irreducible. Let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a non-special facet and w, w' be elements of ${}_f W^{\mathbf{g}}$ which are in the same equivalence class for $\sim_{\mathbf{g}}$. Denote by A_w (resp. $A_{w'}$) the alcove which contains $w\mathbf{g}$ (resp. $w'\mathbf{g}$) in its closure and which is minimal in $A_w W_{\mathbf{g}}$ (resp. $A_{w'} W_{\mathbf{g}}$) for \preceq . Then there exists a positive integer s and a chain of elements of ${}_f W^{\mathbf{g}}$*

$$w_s = w, w_{s-1}, \dots, w_0 = w'$$

such that, for all $i \in \llbracket 0, s-1 \rrbracket$, there exists an element $u_i \in {}_f W^{\mathbf{g}}$ satisfying

$$n_{w_i, u_i}(1) \neq 0 \quad \text{and} \quad n_{w_{i+1}, u_i}(1) \neq 0,$$

and such that

$$s \leq 2 + d(\widehat{A}_w - \rho^\vee) + d(\widehat{A}_{w'} - \rho^\vee).$$

Proof. — Denote by v the unique element of ${}_f W^{\mathbf{g}}$ such that $v\mathbf{g} \subset \rho^\vee + \overline{\mathbf{a}_1}$. Let us also denote by \widehat{w} the element of ${}_f W^{\mathbf{g}}$ such that $\widehat{w}\mathbf{g} \subset \widehat{A}_w$ and $A_0 := \widehat{A}_w, A_{r-1}, \dots, A_r := \rho^\vee + \mathbf{a}_1$ be the alcoves of Lemma 5.19. By Remark 5.32, we know that there exist elements $w_i, u_i \in {}_f W^{\mathbf{g}}$ such that $(w_r, w_0) = (v, \widehat{w})$ and

$$w_i \mathbf{g} \subset \overline{A_i}, n_{w_i, u_i}(1) \neq 0 \quad \text{and} \quad n_{w_{i+1}, u_i}(1) \neq 0, \quad \forall i \in \llbracket 0, r-1 \rrbracket.$$

Therefore, the chain $w, \widehat{w}, w_1, \dots, w_r = v$ linking w to v is of length equal to 1 plus the number of hyperplanes separating \widehat{A}_w from $\mathbf{a}_1 + \rho^\vee$, and this last number is equal to the number of hyperplanes separating $\widehat{A}_w - \rho^\vee$ from \mathbf{a}_1 , i.e. to $d(\widehat{A}_w - \rho^\vee)$. Repeating the same process for w' , we find a chain linking w' to v of length equal to $1 + d(\widehat{A}_{w'} - \rho^\vee)$, so that the concatenation of these two chains yields the desired result. \square

6. Smith–Treumann theory and consequences

6.1. Geometric Satake equivalence

The affine Grassmannian Gr associated with G can be defined as the fppf-quotient LG/L^+G , which can be shown to be an ind-projective ind-scheme over \mathbb{F} . In particular, Gr coincides with the partial affine flag variety $\mathrm{Fl}_{\{0\}}$ (cf. Subsection 3.2). For any $\lambda \in \mathbb{X}_+^\vee$, we denote by $z^\lambda \in T(\mathcal{O})$ the image of z by $\lambda : \mathbb{G}_m(\mathcal{K}) \rightarrow T(\mathcal{K})$, and define $[\lambda] := z^\lambda L^+G$. The orbit $L^+G \cdot [\lambda]$ will be denoted X_λ . Using the Cartan decomposition, one can show that the action of L^+G on Gr induces the following equality

$$(\mathrm{Gr})_{\mathrm{red}} = \bigsqcup_{\lambda \in \mathbb{X}_+^\vee} X_\lambda,$$

where $(\mathrm{Gr})_{\mathrm{red}}$ is the reduced ind-scheme associated with Gr . Moreover, each orbit X_λ is a smooth \mathbb{F} -scheme of finite type and

$$\overline{X_\lambda} = \bigsqcup_{\substack{\mu \leqslant \lambda \\ \mu \in \mathbb{X}_+^\vee}} X_\mu$$

is a projective \mathbb{F} -scheme, where $\mu \leqslant \lambda$ means as usual that $\lambda - \mu$ is a sum of positive coroots.

For any X which is a locally closed finite union of L^+G -orbits, one can show that the L^+G -action on X factors through a quotient group of finite type J ; this quotient may and will be chosen so that the kernel of the map $L^+G \rightarrow J$ is contained in $\ker(L^+G \rightarrow G)$. More concretely, one can take J to be the group representing the functor $R \mapsto G(R[z]/z^n)$, for a large enough integer n . We can then consider the J -equivariant derived category of constructible étale \mathbf{k} -sheaves $D_J^b(X, \mathbf{k})$, which does not depend on our choice of J (see [14, Section 3.3]). We are now allowed to define the category $D_{L^+G}^b(\mathrm{Gr}, \mathbf{k})$ as the direct limit of the categories $D_J^b(X, \mathbf{k})$, indexed by finite and closed unions of orbits X which are ordered by inclusion and where the transition maps are given by direct images. This category admits a canonical perverse t -structure, and we denote by $\mathrm{Perv}_{L^+G}(\mathrm{Gr}, \mathbf{k})$ its heart.

For $\lambda \in \mathbb{X}_+^\vee$, let $i^\lambda : X_\lambda \hookrightarrow \mathrm{Gr}$ denote the inclusion and define

$$\Delta_\lambda^{\mathrm{sph}} := {}^p\mathrm{H}^0(i_!^\lambda \underline{\mathbf{k}}_{X_\lambda}[\dim(X_\lambda)]), \quad \nabla_\lambda^{\mathrm{sph}} := {}^p\mathrm{H}^0(i_*^\lambda \underline{\mathbf{k}}_{X_\lambda}[\dim(X_\lambda)]).$$

The complex $\Delta_\lambda^{\mathrm{sph}}$, resp. $\nabla_\lambda^{\mathrm{sph}}$, is called the standard perverse sheaf, resp. the costandard perverse sheaf attached to λ . By construction, there is a canonical map $\Delta_\lambda^{\mathrm{sph}} \rightarrow \nabla_\lambda^{\mathrm{sph}}$, and we define the intersection cohomology complex $\mathrm{IC}_\lambda^{\mathrm{sph}}$ associated with λ as the image of this map. Intersection

cohomology complexes cover (up to isomorphism) all of the simple objects of the category $\mathrm{Perv}_{L+G}(\mathrm{Gr}, \mathbf{k})$ when λ runs through \mathbb{X}_+^\vee .

PROPOSITION 6.1 ([10, Proposition 12.4]). — *The category $\mathrm{Perv}_{L+G}(\mathrm{Gr}, \mathbf{k})$ is a highest weight category, with weight poset \mathbb{X}_+^\vee , standard objects $\{\Delta_\lambda^{\mathrm{sph}}, \lambda \in \mathbb{X}_+^\vee\}$ and costandard objects $\{\nabla_\lambda^{\mathrm{sph}}, \lambda \in \mathbb{X}_+^\vee\}$.*

The category $D_{L+G}^b(\mathrm{Gr}, \mathbf{k})$ admits a monoidal product \star , which restricts to a monoidal product on $\mathrm{Perv}_{L+G}(\mathrm{Gr}, \mathbf{k})$. In the following crucial result, we denote by G^\vee the Langlands dual group of G over \mathbf{k} , and by $\mathrm{Rep}_{\mathbf{k}}(G^\vee)$ the category of algebraic finite dimensional \mathbf{k} -representations of G^\vee .

THEOREM 6.2 ([26, Theorem 14.1]). — *There is an equivalence of monoidal categories*

$$(\mathrm{Perv}_{L+G}(\mathrm{Gr}, \mathbf{k}), \star) \xrightarrow{\sim} (\mathrm{Rep}_{\mathbf{k}}(G^\vee), \otimes_{\mathbf{k}}).$$

Remark 6.3. — Let $\lambda \in \mathbb{X}_+^\vee$. It is well known that the category $\mathrm{Rep}_{\mathbf{k}}(G^\vee)$ also admits a highest weight structure, and we denote by Δ_λ , resp. ∇_λ , resp. L_λ , the standard object, resp. the costandard object, resp. the simple module, associated with λ . Then, the previous equivalence of categories sends $\nabla_\lambda^{\mathrm{sph}}$, resp. $\Delta_\lambda^{\mathrm{sph}}$, resp. $\mathrm{IC}_\lambda^{\mathrm{sph}}$, on ∇_λ , resp. Δ_λ , resp. L_λ (cf. [26, Proposition 13.1]). We will say that this equivalence is an *equivalence of highest weight categories*.

6.2. Iwahori–Whittaker variant

Recall the construction of $D_{\mathcal{IW}}^b(\mathrm{Gr}, \mathbf{k})$ from Subsection 3.3. If we denote by Y_λ the orbit of $[\lambda]$ under the action of Iw_u^+ , then we have a decomposition

$$(\mathrm{Gr})_{\mathrm{red}} = \bigsqcup_{\lambda \in \mathbb{X}^\vee} Y_\lambda,$$

where each Y_λ is a finite dimensional affine space over \mathbb{F} . One can show that an orbit Y_λ supports a non-zero Iwahori–Whittaker local system iff λ is strictly dominant, and that in this case there exists exactly one (up to isomorphism) such local system of rank one on Y_λ , which we will denote by $\mathcal{L}_{\mathrm{AS}}^\lambda$ (here AS stands for Artin–Schreier).

Once again, the category $D_{\mathcal{IW}}^b(\mathrm{Gr}, \mathbf{k})$ admits a canonical perverse t -structure, and we will denote by $\mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})$ its heart. Thanks to the fact that the Iw_u^+ -orbits are affine spaces over \mathbb{F} , this category of perverse sheaves admits a transparent highest weight structure (cf. [12, Corollary 3.6]). Namely, the weight poset is given by \mathbb{X}_{++}^\vee , and the standard,

costandard and simple objects associated with some $\lambda \in \mathbb{X}_{++}^\vee$ are respectively given by

$$\Delta_\lambda^{\mathcal{IW}} := j_!^\lambda \mathcal{L}_{\text{AS}}^\lambda[\dim(Y_\lambda)], \quad \nabla_\lambda^{\mathcal{IW}} := j_*^\lambda \mathcal{L}_{\text{AS}}^\lambda[\dim(Y_\lambda)], \quad \text{IC}_\lambda^{\mathcal{IW}},$$

where $j^\lambda : Y_\lambda \hookrightarrow \text{Gr}$ is the inclusion, and $\text{IC}_\lambda^{\mathcal{IW}}$ is obtained as the image of the canonical morphism $\Delta_\lambda^{\mathcal{IW}} \rightarrow \nabla_\lambda^{\mathcal{IW}}$. In the sequel, we will denote by $\text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$ the category of tilting objects associated with this highest weight category, and by $\mathcal{T}_\lambda^{\mathcal{IW}}$ the indecomposable tilting object of highest weight $\lambda \in \mathbb{X}_{++}^\vee$.

Although the category $D_{\mathcal{IW}}^b(\text{Gr}, \mathbf{k})$ is not endowed with a canonical convolution product making it a monoidal category, it admits a right action of the monoidal category $D_{L+G}^b(\text{Gr}, \mathbf{k})$. The following result was found by the authors of [12], and gives another incarnation of the category $\text{Rep}_{\mathbf{k}}(G^\vee)$. Note that by our assumptions on G , the element ρ^\vee , defined as the half-sum of positive coroots, belongs to \mathbb{X}^\vee .

THEOREM 6.4. — *The functor $\mathcal{F} \mapsto \nabla_{\rho^\vee}^{\mathcal{IW}} \star \mathcal{F}$ induces an equivalence of highest weight categories*

$$\text{Perv}_{L+G}(\text{Gr}, \mathbf{k}) \xrightarrow{\sim} \text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k}).$$

Remark 6.5. — At this stage, we can note that the action of \mathbb{G}_m on Gr by rescaling z stabilizes each Iw_u^+ -orbit. This allows us to consider the loop rotation equivariant Iwahori–Whittaker derived category of \mathbf{k} -sheaves

$$D_{\mathcal{IW}, \mathbb{G}_m}^b(\text{Gr}, \mathbf{k}),$$

which comes with a natural t -exact forgetful functor to $D_{\mathcal{IW}}^b(\text{Gr}, \mathbf{k})$ (cf. [30, Section 5.2]). It is then not difficult to show (cf. [30, Lemma 5.2]) that the forgetful functor

$$(6.1) \quad \text{Perv}_{\mathcal{IW}, \mathbb{G}_m}(\text{Gr}, \mathbf{k}) \longrightarrow \text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$$

is an equivalence of categories. These considerations will become useful in Section 6.

6.3. Fixed points of the affine Grassmannian and connected components

As in Subsection 3.1, we fix an integer $n \geq 1$. We denote by μ_n the finite group scheme of n^{th} roots of unity, which acts on L^+G and LG by rescaling the indeterminate z ; in particular, μ_n acts on Gr . The following fact, which will only be used with $n = \ell$, is one of the fundamental tools

used by the authors of [30] (cf. [30, Proposition 4.7]). For any $\lambda \in \overline{\mathbf{a}_n} \cap \mathbb{X}^\vee$, we will denote by $\mathbf{g}_\lambda \subset \overline{\mathbf{a}_n}$ the facet which contains λ .

PROPOSITION 6.6. — *For any $\lambda \in \overline{\mathbf{a}_n} \cap \mathbb{X}^\vee$, the map $g \mapsto g \cdot [\lambda]$ factors through an open and closed embedding*

$$\mathrm{Fl}_{\mathbf{g}_\lambda}^{n, \circ} \hookrightarrow (\mathrm{Gr})^{\mu_n}$$

and the induced map

$$\bigsqcup_{\lambda \in \overline{\mathbf{a}_n} \cap \mathbb{X}^\vee} \mathrm{Fl}_{\mathbf{g}_\lambda}^{n, \circ} \longrightarrow (\mathrm{Gr})^{\mu_n}$$

is an isomorphism of ind-schemes.

If we denote by $\mathrm{Iw}_{u, \ell}^+$ the inverse image of U^+ under the evaluation map $L_\ell^+ G \rightarrow G, z^\ell \mapsto 0$, then we get $(\mathrm{Iw}_u^+)^{\mu_\ell} = \mathrm{Iw}_{u, \ell}^+$, and the orbits of $(\mathrm{Gr})^{\mu_\ell}$ under the action of $\mathrm{Iw}_{u, \ell}^+$ are still parametrized by \mathbb{X}^\vee (cf. [30, Lemma 4.8]). Thus, using the morphism $\mathrm{Iw}_{u, \ell}^+ \rightarrow \mathbb{G}_a$ induced by χ , we can take back the constructions of Subsection 6.2 to define the category $D_{\mathcal{IW}_\ell}^b(Y, \mathbf{k})$, where $Y \subset (\mathrm{Gr})^{\mu_\ell}$ is a finite locally closed union of $\mathrm{Iw}_{u, \ell}^+$ -orbits. The theory of parity complexes also adapts to the present context: a complex $\mathcal{F} \in D_{\mathcal{IW}_\ell}^b(Y, \mathbf{k})$ is called $*$ -even, resp. $!$ -even, if for any⁽¹³⁾ $\lambda \in \mathbb{X}_{++}^\vee$ such that $(Y_\lambda)^{\mu_\ell} \subset Y$ the complex $(j_{\mu_\ell}^\lambda)^* \mathcal{F}$, resp. $(j_{\mu_\ell}^\lambda)! \mathcal{F}$, is concentrated in even degrees, where $j_{\mu_\ell}^\lambda : (Y_\lambda)^{\mu_\ell} \hookrightarrow Y$ is the inclusion. The definition for $*$ -odd and $!$ -odd complexes is similar and one says that a complex is even, resp. odd, if it is both $*$ -even and $!$ -even, resp. $*$ -odd and $!$ -odd. As usual, a complex is called parity if it is a direct sum of even and odd complexes.

Defining the category $D_{\mathcal{IW}_\ell}^b((\mathrm{Gr})^{\mu_\ell}, \mathbf{k})$ (resp. $D_{\mathcal{IW}_\ell}^b(\mathrm{Fl}_{\mathbf{g}}^{\ell, \circ}, \mathbf{k})$ for a facet $\mathbf{G} \subset \overline{\mathbf{a}_\ell}$) as a direct limit, it then makes sense to talk about even and odd objects in this category, and we will denote by $\mathrm{Par}_{\mathcal{IW}_\ell}((\mathrm{Gr})^{\mu_\ell}, \mathbf{k})$ (resp. $\mathrm{Par}_{\mathcal{IW}_\ell}(\mathrm{Fl}_{\mathbf{g}}^{\ell, \circ}, \mathbf{k})$) the additive full subcategory consisting of parity objects. By Proposition 6.6, it is clear that the additive category $\mathrm{Par}_{\mathcal{IW}_\ell}((\mathrm{Gr})^{\mu_\ell}, \mathbf{k})$ splits into a direct sum of subcategories of the form $\mathrm{Par}_{\mathcal{IW}_\ell}(\mathrm{Fl}_{\mathbf{g}}^{\ell, \circ}, \mathbf{k})$, where \mathbf{g} runs through the facets inside $\overline{\mathbf{a}_\ell}$.

It is important to note that we have a bijection $\mathbf{g} \mapsto \ell \cdot \mathbf{G}$ between facets inside $\overline{\mathbf{a}_1}$ associated with the action \square_1 of W on E and facets inside $\overline{\mathbf{a}_\ell}$ associated with the action \square_ℓ of W on E , and that a simple change of variable (namely, replacing z with z^ℓ) induces canonical isomorphisms of

(13) Notice that, if $\lambda \notin \mathbb{X}_{++}^\vee$, the restriction and co-restriction of \mathcal{F} to $(Y_\lambda)^{\mu_\ell}$ is zero (because this orbit supports an Iwahori–Whittaker local system only when $\lambda \in \mathbb{X}_{++}^\vee$), so that we can restrict ourselves to the case where $\lambda \in \mathbb{X}_{++}^\vee$.

ind-schemes $\mathrm{Fl}_{\mathbf{g}}^{\circ} := \mathrm{Fl}_{\mathbf{g}}^{1,\circ} \simeq \mathrm{Fl}_{\ell \cdot \mathbf{g}}^{\ell,\circ}$, $\mathrm{Iw}_u^+ \simeq \mathrm{Iw}_{u,\ell}^+$, sending Iw_u^+ -orbits in $\mathrm{Fl}_{\mathbf{g}}^{\circ}$ to $\mathrm{Iw}_{u,\ell}^+$ -orbits in $\mathrm{Fl}_{\ell \cdot \mathbf{g}}^{\ell,\circ}$. For any facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$, we thus have a canonical equivalence of categories

$$D_{\mathcal{IW}}^b(\mathrm{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k}) \simeq D_{\mathcal{IW}_{\ell}}^b(\mathrm{Fl}_{\ell \cdot \mathbf{g}}^{\ell,\circ}, \mathbf{k}),$$

restricting to an equivalence between the corresponding categories of parity complexes:

$$(6.2) \quad \mathrm{Par}_{\mathcal{IW}}(\mathrm{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k}) \simeq \mathrm{Par}_{\mathcal{IW}_{\ell}}(\mathrm{Fl}_{\ell \cdot \mathbf{g}}^{\ell,\circ}, \mathbf{k}).$$

In the sequel, we will denote by $\mathcal{E}_{\ell,w}^{\ell \cdot \mathbf{g}}$ the indecomposable parity complex in $\mathrm{Par}_{\mathcal{IW}_{\ell}}(\mathrm{Fl}_{\ell \cdot \mathbf{g}}^{\ell,\circ}, \mathbf{k})$ corresponding to $\mathcal{E}_w^{\mathbf{g}}$ via the equivalence (6.2). It is clear that, up to a shift, all of the indecomposable parity complexes in $\mathrm{Par}_{\mathcal{IW}_{\ell}}(\mathrm{Fl}_{\ell \cdot \mathbf{g}}^{\ell,\circ}, \mathbf{k})$ (coming from Proposition 3.1) arise in this way.

6.4. Smith category and the linkage principle

Since the action of \mathbb{G}_m on Gr stabilizes the fixed points $(\mathrm{Gr})^{\mu_{\ell}}$, we can take back the construction recalled in Remark 6.5 to define the category $D_{\mathcal{IW}_{\ell}, \mathbb{G}_m}^b(Y, \mathbf{k})$, where $Y \subset (\mathrm{Gr})^{\mu_{\ell}}$ is a locally closed finite union of $\mathrm{Iw}_{u,\ell}^+$ -orbits. We will denote by $D_{\mathcal{IW}_{\ell}, \mathbb{G}_m}^b(Y, \mathbf{k})_{\mu_{\ell}-\mathrm{perf}}$ the full subcategory whose objects are the \mathcal{F} for which the object $\mathrm{Res}_{\mu_{\ell}}^{\mathbb{G}_m}(\mathcal{F})$ has perfect geometric stalks in the sense of [30, Section 3.3]. The *Smith category* $\mathrm{Sm}_{\mathcal{IW}}(Y, \mathbf{k})$ on Y is by definition the Verdier quotient

$$D_{\mathcal{IW}_{\ell}, \mathbb{G}_m}^b(Y, \mathbf{k}) / D_{\mathcal{IW}_{\ell}, \mathbb{G}_m}^b(Y, \mathbf{k})_{\mu_{\ell}-\mathrm{perf}}.$$

Let $X \subset \mathrm{Gr}$ be a locally closed finite union of Iw_u^+ -orbits, and $i_X : (X)^{\mu_{\ell}} \hookrightarrow X$ denote the inclusion. We define the functor

$$i_X^{!*} : \mathrm{Perv}_{\mathcal{IW}, \mathbb{G}_m}(X, \mathbf{k}) \longrightarrow \mathrm{Sm}_{\mathcal{IW}}((X)^{\mu_{\ell}}, \mathbf{k})$$

as the composition of the inverse image i_X^* with the canonical quotient map

$$Q : D_{\mathcal{IW}_{\ell}, \mathbb{G}_m}^b((X)^{\mu_{\ell}}, \mathbf{k}) \longrightarrow \mathrm{Sm}_{\mathcal{IW}}((X)^{\mu_{\ell}}, \mathbf{k}).$$

A crucial result says that taking $i_X^!$ instead of i_X^* in the previous construction gives isomorphic functors (cf. [30, Section 6.2]), whence the notation. It then takes a bit more work (cf. [30, Lemma 6.1]) to prove that, for two locally closed finite unions of $\mathrm{Iw}_{u,\ell}^+$ -orbits $Z \subset Y$, the canonical functor

$f_*^{\text{Sm}} : \text{Sm}_{\mathcal{IW}}(Z, \mathbf{k}) \rightarrow \text{Sm}_{\mathcal{IW}}(Y, \mathbf{k})$ induced by the direct image in the corresponding derived categories is fully faithful, and fits into the following commutative diagram

$$\begin{array}{ccc} D_{\mathcal{IW}_\ell, \mathbb{G}_m}^b(Z, \mathbf{k}) & \xrightarrow{f_*} & D_{\mathcal{IW}_\ell, \mathbb{G}_m}^b(Y, \mathbf{k}) \\ \downarrow & & \downarrow \\ \text{Sm}_{\mathcal{IW}}(Z, \mathbf{k}) & \xrightarrow{f_*^{\text{Sm}}} & \text{Sm}_{\mathcal{IW}}(Y, \mathbf{k}), \end{array}$$

where the vertical arrows are the quotient maps. One can thus define the category $\text{Sm}_{\mathcal{IW}}((\text{Gr})^{\mu_\ell}, \mathbf{k})$ as a direct limit indexed by finite closed unions of $\text{Iw}_{u, \ell}^+$ -orbits, and consider the functor

$$i_{\text{Gr}}^{!*} : \text{Perv}_{\mathcal{IW}, \mathbb{G}_m}(\text{Gr}, \mathbf{k}) \longrightarrow \text{Sm}_{\mathcal{IW}}((\text{Gr})^{\mu_\ell}, \mathbf{k}).$$

The following statement is [30, Theorem 7.4].

THEOREM 6.7. — *The composition of functors*

$$\text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k}) \xrightarrow{\text{Remark 6.5}} \text{Perv}_{\mathcal{IW}, \mathbb{G}_m}(\text{Gr}, \mathbf{k}) \xrightarrow{i_{\text{Gr}}^{!*}} \text{Sm}_{\mathcal{IW}}((\text{Gr})^{\mu_\ell}, \mathbf{k})$$

restricts to a fully faithful functor $\Phi : \text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k}) \rightarrow \text{Sm}_{\mathcal{IW}}((\text{Gr})^{\mu_\ell}, \mathbf{k})$.

We can now reformulate the proof of the linkage principle from [30, Theorem 8.5]. Thanks to the decomposition of $(\text{Gr})^{\mu_\ell}$ into its connected components from Proposition 6.6, we deduce from Theorem 6.7 above that two strictly dominant weights λ and μ which are not in the same orbit for the box action cannot be in relation for \mathcal{R}_2 (seen as a relation on the weight poset \mathbb{X}_{++}^\vee of the highest weight category $\text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$, cf. Section 2). Indeed, let δ , resp. γ , be the element of $W \square_\ell \lambda \cap \overline{\mathbf{a}_\ell}$, resp. of $W \square_\ell \mu \cap \overline{\mathbf{a}_\ell}$. We have

$$\begin{aligned} \text{Hom}_{\text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})}(\mathcal{T}_\lambda^{\mathcal{IW}}, \mathcal{T}_\mu^{\mathcal{IW}}) \\ \simeq \text{Hom}_{\text{Sm}_{\mathcal{IW}}((\text{Gr})^{\mu_\ell}, \mathbf{k})}(\Phi(\mathcal{T}_\lambda^{\mathcal{IW}}), \Phi(\mathcal{T}_\mu^{\mathcal{IW}})), \end{aligned}$$

and since the object $\Phi(\mathcal{T}_\lambda^{\mathcal{IW}})$, resp. $\Phi(\mathcal{T}_\mu^{\mathcal{IW}})$, is indecomposable (thanks to the full faithfulness of Φ), its support must be contained in a single connected component of $(\text{Gr})^{\mu_\ell}$, which is easily seen to be the parametrized by δ , resp. γ , in the isomorphism of Proposition 6.6 applied to the case $n = \ell$.

Thanks to Theorem 6.4, this implies that two dominant weights λ', μ' which are not in the same orbit for the dot action cannot be in relation for \mathcal{R}_2 (where we now consider the highest weight category $\text{Perv}_{L^+ G}(\text{Gr}, \mathbf{k})$).

Finally, the geometric Satake equivalence (Theorem 6.2) and Theorem 2.3 allow to prove the linkage principle for G^\vee (cf. [20, Corollary 6.17, Part II]):

$$(6.3) \quad \forall \lambda', \mu' \in \mathbb{X}_+^\vee \text{ such that } W \bullet_\ell \lambda' \neq W \bullet_\ell \mu',$$

we have $\text{Ext}_{G^\vee}^1(L_{\lambda'}, L_{\mu'}) = 0$.

6.5. Consequences on equivalence relations

Recall the definition of the “dot” action of W , which acts on \mathbb{X}^\vee by

$$w \bullet_\ell \mu := w \square_\ell (\mu + \rho^\vee) - \rho^\vee,$$

for any $w \in W$, $\mu \in \mathbb{X}^\vee$. Let $\mu \in \mathbb{X}^\vee \cap \overline{\mathbf{a}_\ell}$ and \mathbf{g}_μ be the facet (for \square_ℓ) containing μ . Therefore, $\ell^{-1} \cdot \mathbf{g}_\mu \subset \overline{\mathbf{a}_1}$ is a facet for \square_1 . The assignment $w \mapsto w \square_\ell \mu$, resp. $w \mapsto w \bullet_\ell (\mu - \rho^\vee)$, induces a bijection

$$(6.4) \quad {}_f W^{\ell^{-1} \cdot \mathbf{g}_\mu} \xrightarrow{\sim} W \square_\ell \mu \cap \mathbb{X}_{++}^\vee, \text{ resp. } {}_f W^{\ell^{-1} \cdot \mathbf{g}_\mu} \xrightarrow{\sim} W \bullet_\ell (\mu - \rho^\vee) \cap \mathbb{X}_+^\vee$$

(this follows from the fact that, if we denote by $W_{\mathbf{g}_\mu, \square_\ell} \subset W$ the stabilizer of \mathbf{g}_μ for \square_ℓ , then we have an equality $W_{\mathbf{g}_\mu, \square_\ell} = W_{\ell^{-1} \cdot \mathbf{g}_\mu}$).

In the sequel, we will want to determine an exact description of the blocks in \mathbb{X}_{++}^\vee (seen as the weight poset of $\text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$) for \mathcal{R}_2 (cf. Subsection 2.2). The following results will help us to do so (cf. the proof of [30, Theorem 8.9]).

PROPOSITION 6.8. — *Let $\mu \in \mathbb{X}^\vee \cap \overline{\mathbf{a}_\ell}$, \mathbf{g}_μ be the facet (for \square_ℓ) containing μ , and w, w' be elements of ${}_f W^{\ell^{-1} \cdot \mathbf{g}_\mu}$. We have an isomorphism*

$$\text{Hom}_{\text{Tilt}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})}(\mathcal{T}_{w \square_\ell \mu}^{\mathcal{IW}}, \mathcal{T}_{w' \square_\ell \mu}^{\mathcal{IW}}) \simeq \text{Hom}_{\text{Par}_{\mathcal{IW}_\ell}(\text{Fl}_{\mathbf{g}_\mu, \square_\ell}^{\ell, \circ}, \mathbf{k})}^{\bullet}(\mathcal{E}_{\ell, w}^{\mathbf{g}_\mu}, \mathcal{E}_{\ell, w'}^{\mathbf{g}_\mu}).$$

Proof. — This is a direct consequence of [30, Proposition 8.11] and of the equivalence of highest weight categories $\text{Rep}_{\mathbf{k}}(G^\vee) \simeq \text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$, which sends the indecomposable tilting module of highest weight $w \bullet_\ell (\mu - \rho^\vee)$, resp. $w' \bullet_\ell (\mu - \rho^\vee)$, to $\mathcal{T}_{w \square_\ell \mu}^{\mathcal{IW}}$, resp. $\mathcal{T}_{w' \square_\ell \mu}^{\mathcal{IW}}$. \square

Let $\lambda \in \overline{\mathbf{a}_\ell} \cap \mathbb{X}^\vee$. Note that, thanks to the equivalence (6.2), we have

$$\begin{aligned} \text{Hom}_{\text{Par}_{\mathcal{IW}}(\text{Fl}_{\ell^{-1} \cdot \mathbf{g}_\lambda}^{\circ}, \mathbf{k})}^{\bullet}(\mathcal{E}_w^{\ell^{-1} \cdot \mathbf{g}_\lambda}, \mathcal{E}_{w'}^{\ell^{-1} \cdot \mathbf{g}_\lambda}) \\ \simeq \text{Hom}_{\text{Par}_{\mathcal{IW}_\ell}(\text{Fl}_{\mathbf{g}_\lambda, \square_\lambda}^{\ell, \circ}, \mathbf{k})}^{\bullet}(\mathcal{E}_{\ell, w}^{\mathbf{g}_\lambda}, \mathcal{E}_{\ell, w'}^{\mathbf{g}_\lambda}). \end{aligned}$$

Therefore Proposition 6.8 yields

$$(6.5) \quad \forall w, w' \in {}_f W^{\ell^{-1} \cdot \mathbf{g}_\lambda}, \quad w \mathcal{R}_{\ell^{-1} \cdot \mathbf{g}_\lambda} w' \iff (w \square_\ell \lambda) \mathcal{R}_2 (w' \square_\ell \lambda).$$

6.6. Dilating weights by ℓ

Smith–Treumann theory allows us to understand the effect of dilating dominant weights by ℓ on the equivalence relation \sim .

PROPOSITION 6.9. — *Let $\lambda, \mu \in \mathbb{X}_{++}^\vee$. We have an isomorphism*

$$\mathrm{Hom}_{\mathrm{Tilt}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})}(\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}}, \mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}}) \simeq \mathrm{Hom}_{\mathrm{Tilt}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})}(\mathcal{T}_\lambda^{\mathcal{IW}}, \mathcal{T}_\mu^{\mathcal{IW}}).$$

Proof. — First recall that, by full-faithfulness of Φ , we have an isomorphism

$$(6.6) \quad \begin{aligned} \mathrm{Hom}_{\mathrm{Tilt}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})}(\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}}, \mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}}) \\ \simeq \mathrm{Hom}_{\mathrm{Sm}_{\mathcal{IW}}((\mathrm{Gr})^{\mu_\ell}, \mathbf{k})}(\Phi(\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}}), \Phi(\mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}})). \end{aligned}$$

Next, notice that we have an embedding

$$L_\ell G / L_\ell^+ G \hookrightarrow (\mathrm{Gr})^{\mu_\ell}$$

thanks to [30, Remark 4.8], and that the left-hand side identifies (via Proposition 6.6) with the union of the connected components $\mathrm{Fl}_{\mathbf{g}_\lambda}^{\ell, \circ}$, with λ running through $\overline{\mathbf{a}_\ell} \cap \ell \cdot \mathbb{X}^\vee$. Moreover, the $L_\ell^+ G$ -orbits in $L_\ell G / L_\ell^+ G$ are parametrized by $\ell \cdot \mathbb{X}_{++}^\vee$, and a simple change of variable $z \mapsto z^\ell$ together with (6.1) induce an equivalence of categories

$$R : \mathrm{Tilt}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k}) \simeq \mathrm{Tilt}_{\mathcal{IW}_\ell, \mathbb{G}_m}(L_\ell G / L_\ell^+ G, \mathbf{k}),$$

which sends $\mathcal{T}_\lambda^{\mathcal{IW}}$ (resp. $\mathcal{T}_\mu^{\mathcal{IW}}$) to the indecomposable tilting object associated with $\ell \cdot \lambda$ (resp. $\ell \cdot \mu$). Therefore, the indecomposable tilting objects in $\mathrm{Tilt}_{\mathcal{IW}_\ell, \mathbb{G}_m}(L_\ell G / L_\ell^+ G, \mathbf{k})$ are parity complexes (see [12, Proposition 4.12]), so that we can take back the same arguments as in the proof of [30, Theorem 7.4] to show that restricting the functor

$$Q : D_{\mathcal{IW}_\ell, \mathbb{G}_m}^b((\mathrm{Gr})^{\mu_\ell}, \mathbf{k}) \longrightarrow \mathrm{Sm}_{\mathcal{IW}}((\mathrm{Gr})^{\mu_\ell}, \mathbf{k})$$

to $L_\ell G / L_\ell^+ G$ and composing it with R yields a fully-faithful functor

$$Q \circ R : \mathrm{Tilt}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k}) \longrightarrow \mathrm{Sm}_{\mathcal{IW}}(L_\ell G / L_\ell^+ G, \mathbf{k}),$$

which satisfies the following isomorphisms:

$$Q \circ R(\mathcal{T}_\lambda^{\mathcal{IW}}) \simeq \Phi(\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}}) \quad \text{and} \quad Q \circ R(\mathcal{T}_\mu^{\mathcal{IW}}) \simeq \Phi(\mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}}).$$

The desired result thus follows from comparing (6.6) with the following isomorphism

$$\mathrm{Hom}(\mathcal{T}_\lambda^{\mathcal{IW}}, \mathcal{T}_\mu^{\mathcal{IW}}) \simeq \mathrm{Hom}(Q \circ R(\mathcal{T}_\lambda^{\mathcal{IW}}), Q \circ R(\mathcal{T}_\mu^{\mathcal{IW}})). \quad \square$$

In the next corollary, we consider the equivalence relation \sim on the set \mathbb{X}_{++}^\vee , seen as the weight poset of the highest weight category $\text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$ (cf. Subsection 2.2).

COROLLARY 6.10. — *For any $\gamma \in \mathbb{X}_{++}^\vee$, denote by $\bar{\gamma}$ the equivalence class of γ for the equivalence relation \sim . Then we have*

$$\forall \lambda \in \mathbb{X}_{++}^\vee, \quad \ell \cdot \bar{\lambda} \subset \overline{\ell \cdot \lambda}.$$

Proof. — This is a direct consequence of Proposition 6.9 and of the fact that \sim is generated by \mathcal{R}_2 (cf. Section 2). \square

COROLLARY 6.11. — *Let $\lambda, \mu \in \mathbb{X}_{++}^\vee$. We have an equality*

$$(\mathcal{T}_\lambda^{\mathcal{IW}} : \nabla_\mu^{\mathcal{IW}}) = (\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}} : \nabla_{\ell \cdot \mu}^{\mathcal{IW}}).$$

Proof. — For any $\gamma \in \mathbb{X}_{++}^\vee$, let us denote by $n(\gamma)$ the number of elements $\nu \in \mathbb{X}_{++}^\vee$ such that $\nu < \gamma$. We may and will assume that $\mu \leq \lambda$ to prove the desired equality, and proceed by induction on $n(\ell \cdot \mu)$. Standard arguments on highest weight categories (and the fact that Verdier duality sends $\Delta_\nu^{\mathcal{IW}}$, resp. $\mathcal{T}_\nu^{\mathcal{IW}}$, to $\nabla_\nu^{\mathcal{IW}}$, resp. $\mathcal{T}_\nu^{\mathcal{IW}}$, for all $\nu \in \mathbb{X}_{++}^\vee$; namely, we use the same arguments as the one used in [3, Section 6.2]) imply that we have

$$(6.7) \quad (\mathcal{T}_\lambda^{\mathcal{IW}} : \nabla_\mu^{\mathcal{IW}}) = \dim_{\mathbf{k}} \text{Hom}(\mathcal{T}_\lambda^{\mathcal{IW}}, \mathcal{T}_\mu^{\mathcal{IW}}) - \sum_{\nu < \mu, \nu \in \mathbb{X}_{++}^\vee} (\mathcal{T}_\lambda^{\mathcal{IW}} : \nabla_\nu^{\mathcal{IW}}) \cdot (\mathcal{T}_\mu^{\mathcal{IW}} : \nabla_\nu^{\mathcal{IW}}).$$

Now when $n(\ell \cdot \mu) = 0$, we get that $n(\mu) = 0$ and so

$$\begin{aligned} (\mathcal{T}_\lambda^{\mathcal{IW}} : \nabla_\mu^{\mathcal{IW}}) &= \dim_{\mathbf{k}} \text{Hom}(\mathcal{T}_\lambda^{\mathcal{IW}}, \mathcal{T}_\mu^{\mathcal{IW}}) \\ &= \dim_{\mathbf{k}} \text{Hom}(\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}}, \mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}}) \\ &= (\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}} : \nabla_{\ell \cdot \mu}^{\mathcal{IW}}), \end{aligned}$$

where the second equality is due to Proposition 6.9. This proves the first step of the induction. The induction step still follows from using (6.7) and Proposition 6.9, together with the fact that for any $\nu \in \mathbb{X}_{++}^\vee$ such that $(\mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}} : \nabla_\nu^{\mathcal{IW}}) \neq 0$, we have that $\nu \in \ell \cdot \mathbb{X}_{++}^\vee$ (this is because we must have $\nu \in W \square_\ell (\ell \cdot \mu)$ by the linkage principle). \square

6.7. The general case for equivalence relations on ${}_f W^g$

In this subsection, we apply the equivalence (6.5) and Proposition 6.9 to finish the study of Section 5 by treating the case of special facets. The arguments that we have used until now do not apply when g is a special facet

(notice that Corollary 5.27 is false in this case). With good reason: we are going to see that the set ${}_f W^{\mathbf{g}}$ splits into infinitely many classes when some component of \mathbf{g} is a special facet. As in Subsections 5.4 and 5.5, we are first going to deal with the case where the root system is irreducible before generalizing.

Let $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be a facet, $w \in {}_f W^{\mathbf{g}}$, and recall that if \mathbf{g} special, then $w\mathbf{g}$ can be written as $w\mathbf{g} = \{\lambda\}$ for a unique coweight λ . We will denote by \bar{w} the equivalence class of w for $\sim_{\mathbf{g}}$, and define $r_{\mathbf{g}}(w)$ to be equal to -1 if \mathbf{g} is not special, and to be the unique (non-negative) integer such that

$$\lambda \in \ell^{r_{\mathbf{g}}(w)} \mathbb{X}^{\vee} \setminus \ell^{r_{\mathbf{g}}(w)+1} \mathbb{X}^{\vee}, \quad \text{with } w\mathbf{g} = \{\lambda\}$$

if \mathbf{g} is special. For any positive integer r , we also define the group $W^{(r)} := W_0 \ltimes \ell^r \mathbb{Z} \mathfrak{R}^{\vee}$, which is a subgroup of W isomorphic to it, with $W^{(0)} = W$. Finally, notice that for any $\lambda \in \mathbb{X}^{\vee}$ and positive integer r , we have $\ell^r \cdot W \square_{\ell} \lambda = W^{(r)} \square_{\ell} (\ell^r \cdot \lambda)$. We start with an easy lemma.

LEMMA 6.12. — *Let $\nu \in \overline{\mathbf{a}_1} \cap \mathbb{X}^{\vee}$, $\mathbf{g} \subset \overline{\mathbf{a}_1}$ be the facet containing ν , $w \in {}_f W^{\mathbf{g}}$ and r be a positive integer. If we set $\lambda := w \square_1 \nu$, then the application $W \rightarrow \mathbb{X}^{\vee}$, $u \mapsto u \square_{\ell} (\ell \cdot \nu)$ induces a bijection*

$$\left(W^{(r)} w W_{\mathbf{g}} \right) \cap {}_f W^{\mathbf{g}} \simeq W^{(r)} \square_{\ell} (\ell \cdot \lambda) \cap \mathbb{X}_{++}^{\vee}.$$

Proof. — Notice that $\ell \cdot \lambda = w \square_{\ell} (\ell \cdot \nu)$, so that $W^{(r)} \square_{\ell} (\ell \cdot \lambda) = W^{(r)} w \square_{\ell} (\ell \cdot \nu)$. The desired isomorphism then follows from the fact that $W_{\mathbf{g}}$ is the stabilizer of $\ell \cdot \nu$ for \square_{ℓ} , and from the well-known fact that $u \mapsto u \square_{\ell} (\ell \cdot \nu)$ induces a bijection

$${}_f W^{\mathbf{g}} \simeq W \square_{\ell} (\ell \cdot \nu) \cap \mathbb{X}_{++}^{\vee}.$$

□

PROPOSITION 6.13. — *Assume that \mathfrak{R}^{\vee} is irreducible. For any facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$ and $w \in {}_f W^{\mathbf{g}}$, we have*

$$\bar{w} = \begin{cases} \{w\} & \text{if } \mathbf{g} \text{ is special and } \text{char}(\mathbf{k}) = 0 \\ (W^{(r_{\mathbf{g}}(w)+1)} w W_{\mathbf{G}}) \cap {}_f W^{\mathbf{g}} & \text{otherwise.} \end{cases}$$

Proof. — If \mathbf{g} is not special, then we have $r_{\mathbf{g}}(w) = -1$ by definition and

$$\left(W^{(r_{\mathbf{g}}(w)+1)} w W_{\mathbf{g}} \right) \cap {}_f W^{\mathbf{g}} = W \cap {}_f W^{\mathbf{g}} = {}_f W^{\mathbf{g}},$$

which does coincide with \bar{w} thanks to Theorem 5.31. So we assume from now on that \mathbf{g} is a special facet. We will first treat the case where $\text{char}(\mathbf{k}) = \ell$.

Let $w, v \in {}_f W^{\mathbf{g}}$, $\nu \in \mathbb{X}^{\vee} \cap \overline{\mathbf{a}_1}$ be such that $\mathbf{g} = \{\nu\}$, and put $w\mathbf{G} = \{\lambda\}$, $v\mathbf{g} = \{\mu\}$. Notice that $w \square_{\ell} (\ell \cdot \nu) = \ell \cdot \lambda$, $v \square_{\ell} (\ell \cdot \nu) = \ell \cdot \mu$, and recall that,

thanks to Proposition 6.8 coupled with (6.2), we have

$$\mathrm{Hom}_{\mathrm{Tilt}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})}(\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}}, \mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}}) \simeq \mathrm{Hom}_{\mathrm{Par}_{\mathcal{IW}}(\mathrm{Fl}_{\mathbf{g}}, \mathbf{k})}^{\bullet}(\mathcal{E}_w^{\mathbf{g}}, \mathcal{E}_v^{\mathbf{g}}),$$

which implies that $w\mathcal{R}_{\mathbf{g}}v \Leftrightarrow (\ell \cdot \lambda)\mathcal{R}_2(\ell \cdot \mu)$. Moreover, recall that by the geometric linkage principle ([30, Theorem 8.5]) we have

$$\forall \gamma \in \mathbb{X}_{++}^{\vee}, \quad (\ell \cdot \lambda)\mathcal{R}_2\gamma \implies \gamma \in W \square_{\ell}(\ell \cdot \lambda),$$

so that in particular

$$(6.8) \quad (\ell \cdot \lambda)\mathcal{R}_2\gamma \implies \gamma \in \ell \cdot \mathbb{X}_{++}^{\vee}.$$

Therefore, Lemma 6.12 (applied to $r = r_{\mathbf{g}}(w) + 1$) implies that our claim on \bar{w} is equivalent to showing that

$$\overline{\ell \cdot \lambda} = W^{(r_{\mathbf{g}}(w)+1)} \square_{\ell}(\ell \cdot \lambda) \cap \mathbb{X}_{++}^{\vee},$$

where $\overline{\ell \cdot \lambda}$ denotes the equivalence class of $\ell \cdot \lambda$ for the equivalence relation \sim on \mathbb{X}_{++}^{\vee} .

By definition of $r_{\mathbf{g}}(w)$ and $r_{\mathbf{g}}(v)$, we know that there exist $\lambda', \mu' \in \mathbb{X}_{++}^{\vee}$ such that $\ell^{r_{\mathbf{g}}(w)} \cdot \lambda' = \lambda$ and $\ell^{r_{\mathbf{g}}(v)} \cdot \mu' = \mu$. Assume (without any loss of generality, up to switching the roles of λ and μ) that $r_{\mathbf{g}}(v) \geq r_{\mathbf{g}}(w)$, so that Proposition 6.9 yields

$$(6.9) \quad \begin{aligned} \mathrm{Hom}_{\mathrm{Tilt}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})}(\mathcal{T}_{\ell \cdot \lambda}^{\mathcal{IW}}, \mathcal{T}_{\ell \cdot \mu}^{\mathcal{IW}}) \\ \simeq \mathrm{Hom}_{\mathrm{Tilt}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k})}(\mathcal{T}_{\lambda'}^{\mathcal{IW}}, \mathcal{T}_{\ell^{r_{\mathbf{g}}(v)-r_{\mathbf{g}}(w)} \cdot \mu'}^{\mathcal{IW}}). \end{aligned}$$

If we denote by $\mathbf{g}' \subset \overline{\mathbf{a}_{\ell}}$ the facet (for \square_{ℓ}) containing the element of $W \square_{\ell} \lambda' \cap \overline{\mathbf{a}_{\ell}}$, then $\ell^{-1} \cdot \mathbf{g}'$ is a facet (for \square_1) contained in $\overline{\mathbf{a}_1}$ which is *not* special (because otherwise we could write $\mathbf{g}' = \{\ell \cdot \lambda''\}$ for some weight λ'' , and $\lambda' = w'' \square_{\ell}(\ell \cdot \lambda'') = \ell \cdot (w'' \square_1 \lambda'')$ for some $w'' \in W$, contradicting the definition of $r_{\mathbf{g}}(w)$). So thanks to (6.5) and Theorem 5.31 we get

$$\overline{\lambda'} = W \square_{\ell} \lambda' \cap \mathbb{X}_{++}^{\vee}.$$

In particular, if $r_{\mathbf{g}}(v) > r_{\mathbf{g}}(w)$, then we easily see that $\ell^{r_{\mathbf{g}}(v)-r_{\mathbf{g}}(w)} \cdot \mu' \notin W \square_{\ell} \lambda'$, so the right-hand side in (6.9) is non-zero only if $r_{\mathbf{g}}(v) = r_{\mathbf{g}}(w)$. This observation, together with (6.9) and (6.8), allows us to deduce that $\overline{\ell^{r_{\mathbf{g}}(w)+1} \cdot \lambda'} \subset \ell^{r_{\mathbf{g}}(w)+1} \cdot \overline{\lambda'}$. The reversed inclusion $\ell^{r_{\mathbf{g}}(w)+1} \cdot \overline{\lambda'} \subset \overline{\ell^{r_{\mathbf{g}}(w)+1} \cdot \lambda'}$ is obtained by applying Corollary 6.10. We finally get the desired equality:

$$\begin{aligned} \overline{\ell \cdot \lambda} &= \overline{\ell^{r_{\mathbf{g}}(w)+1} \cdot \lambda'} = \ell^{r_{\mathbf{g}}(w)+1} \cdot \overline{\lambda'} \\ &= \ell^{r_{\mathbf{g}}(w)+1} \cdot W \square_{\ell} \lambda' \cap \mathbb{X}_{++}^{\vee} \\ &= W^{(r_{\mathbf{g}}(w)+1)} \square_{\ell}(\ell \cdot \lambda) \cap \mathbb{X}_{++}^{\vee}. \end{aligned}$$

Now we pass to the case $\text{char}(\mathbf{k}) = 0$. We want to show that for any $w' \in {}_f W^{\mathbf{g}}$ such that $w' \neq w$, we have

$$(6.10) \quad \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{G}}, \mathbf{k})}^{\bullet}(\mathcal{E}_{w'}^{\mathbf{g}}, \mathcal{E}_w^{\mathbf{g}}) = 0.$$

We claim that there exists $\omega \in \Omega$ such that $\omega\{0\} = \mathbf{g}$. Indeed, the map $u \mapsto u\{0\}$ is actually a bijection from Ω to $\overline{\mathbf{a}_1} \cap \mathbb{X}^{\vee}$ thanks to the first point of the remark in [13, Chapter VI, Section 2.3]. We then have an isomorphism $L^+ P_{\mathbf{g}} \simeq \dot{\omega} L^+ G \dot{\omega}^{-1}$, and conjugation by $\dot{\omega}$ yields an isomorphism

$$\text{Fl}_{\{0\}}^{\circ} \simeq \text{Fl}_{\mathbf{g}}^{\circ}.$$

Since $\dot{\omega}$ belongs to Ω , conjugation by $\dot{\omega}$ preserves Iw_u^+ , so that we get an equivalence of categories

$$(6.11) \quad D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}^{\circ}, \mathbf{k}) \simeq D_{\mathcal{IW}}^b(\text{Fl}_{\{0\}}^{\circ}, \mathbf{k}).$$

This equivalence implies that we only need to prove (6.10) with \mathbf{g} replaced by $\{0\}$. Now, recall that $\text{Fl}_{\{0\}}^{\circ} \simeq \text{Gr}$ and let $\lambda, \lambda' \in \mathbb{X}_{++}^{\vee}$ be such that $w\mathbf{g} = \{\lambda\}$, $w'\mathbf{g} = \{\lambda'\}$. By [12, Remark 3.5], the complex \mathcal{E}_w is isomorphic to $\text{IC}_{\lambda}^{\mathcal{IW}}$ when $\text{char}(\mathbf{k}) = 0$ (which denotes the intersection cohomology complex on Gr associated with λ , cf. Subsection 6.2). Therefore in characteristic zero we have an isomorphism

$$\text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\{0\}}^{\circ}, \mathbf{k})}^{\bullet}(\mathcal{E}_{w'}, \mathcal{E}_w) \simeq \text{Hom}_{D_{\mathcal{IW}}^b(\text{Gr}, \mathbf{k})}^{\bullet}(\text{IC}_{\lambda'}^{\mathcal{IW}}, \text{IC}_{\lambda}^{\mathcal{IW}}).$$

But we know from [12, Section 3.2] that we have an equivalence of triangulated categories $D_{\mathcal{IW}}^b(\text{Gr}, \mathbf{k}) \simeq D^b \text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$, and that $\text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$ is semi-simple when $\text{char}(\mathbf{k}) = 0$ thanks to [12, Corollary 3.6]. Therefore the right-hand side above is zero when $\lambda \neq \lambda'$ and $\text{char}(\mathbf{k}) = 0$, which is equivalent to $w \neq w'$.⁽¹⁴⁾ \square

This last result coupled with Proposition 5.18 now enables us to deal with the case where \mathfrak{R}^{\vee} splits into irreducible root systems, inducing decompositions $W = W_1 \times \cdots \times W_r$ and $\mathbf{g} = \mathbf{g}_1 \times \cdots \times \mathbf{g}_r$ (see the beginning of Subsection 5.4).

THEOREM 6.14. — *Assume that $\text{char}(\mathbf{k}) = \ell$. For any facet $\mathbf{g} \subset \overline{\mathbf{a}_1}$ and $w = (w_1, \dots, w_r) \in {}_f W^{\mathbf{g}}$, we have*

$$\overline{w} = \prod_{i=1}^r W_i^{(r(w_i)+1)} w_i W_{\mathbf{g}_i} \cap {}_f W_i^{\mathbf{g}_i}.$$

⁽¹⁴⁾ Instead of using the equivalence (6.11), we could have argued by saying that $\text{Fl}_{\mathbf{g}}^{\circ}$ is a connected component of Gr .

When $\text{char}(\mathbf{k}) = 0$ and $r(w_i) \geq 0$ for some i , one replaces the i 'th component in the above product by $\{w_i\}$.

7. Applications to representation theory

7.1. Preliminaries on alcoves

Just as in Subsection 5.3, the action \bullet_ℓ of W on E defines a hyperplane arrangement \mathcal{H}' , where the shift by ρ^\vee induces a bijection between \mathcal{H}' and \mathcal{H} . We will call a connected component of $E \setminus \mathcal{H}'$ an alcove for \bullet_ℓ , and a facet contained in the closure of an alcove for \bullet_ℓ will be called a facet for \bullet_ℓ ; once again shifting by ρ^\vee induces a bijection between facets for \bullet_ℓ and facets for \square_ℓ , and composing this bijection with the dilation by ℓ^{-1} gives a bijection with facets for \square_1 . We put $C_\ell := \mathbf{a}_\ell - \rho^\vee$. The bijection between sets of alcoves allows us to define right and left actions of W on these sets, together with the Bruhat order \leq and the periodic order \preceq , by transport of structure from the set of alcoves for \square_1 (or equivalently, by seeing every alcove for \bullet_ℓ , resp. for \square_ℓ , as a W -translate of C_ℓ , resp. of \mathbf{a}_ℓ). Note that if A, B are two alcoves for \bullet_ℓ , then we have $A \preceq B \Leftrightarrow A \uparrow B$, where the order \uparrow is defined in [20, Section 6]: this is an immediate consequence of [20, Chapter II, Section 6.6, (4)] (which says that \uparrow is invariant under translation) together with [4, Section II, Lemma 10.1] (which implies that \uparrow coincides with the Bruhat order inside of $\mathcal{C}_0^+ - \rho^\vee$, and thus with \preceq). For any alcove C for \bullet_ℓ , we define the alcove \hat{C} for \bullet_ℓ , resp. the integer $d(C)$, by transport of structure using once again the bijection between alcoves for \bullet_ℓ and \square_1 (notice that $d(C)$ is then the number of hyperplanes of \mathcal{H}' separating C from C_ℓ).

For any facet \mathbf{h} for \bullet_ℓ , we will denote by $W_{\mathbf{h}, \bullet_\ell} \subset W$ the stabilizer of \mathbf{h} for \bullet_ℓ . One can easily check that we have

$$W_{\mathbf{h}, \bullet_\ell} = W_{\mathbf{h} + \rho^\vee, \square_\ell} = W_{\ell^{-1} \cdot (\mathbf{h} + \rho^\vee)},$$

where the second and third sets denote stabilizers for \square_ℓ and \square_1 respectively.

Let $\lambda \in \mathbb{X}_+^\vee$ and $\mathbf{h} \subset E$ be the facet for \bullet_ℓ containing λ , which is of the form

$$\mathbf{h} = \left\{ \mu \in E \left| \begin{array}{l} \langle \mu + \rho^\vee, \alpha \rangle = \ell \cdot n_\alpha, \forall \alpha \in \mathfrak{R}_+^0(\mathbf{h}), \\ \ell \cdot (n_\alpha - 1) < \langle \mu + \rho^\vee, \alpha \rangle < \ell \cdot n_\alpha, \forall \alpha \in \mathfrak{R}_+^1(\mathbf{h}) \end{array} \right. \right\}$$

for suitable integers n_α and a partition $\mathfrak{R}_+ = \mathfrak{R}_+^0(\mathbf{h}) \sqcup \mathfrak{R}_+^1(\mathbf{h})$. If we let C_λ be the alcove for \bullet_ℓ defined by the integers $(n_\alpha)_{\alpha \in \mathfrak{R}_+}$, then C_λ is the only

alcove satisfying

$$\lambda \in \tilde{C}_\lambda := \{\mu \in E \mid \ell \cdot (n_\alpha - 1) < \langle \mu + \rho^\vee, \alpha \rangle \leq \ell \cdot n_\alpha, \forall \alpha \in \mathfrak{R}_+\}.$$

The set \tilde{C}_λ (which is denoted by \hat{C}_λ in [20]) is called the upper closure of C_λ . The alcove C_λ can be characterized as the only alcove containing \mathbf{h} in its closure which is minimal in $W_{\mathbf{h}, \bullet_\ell} \bullet_\ell C_\lambda$ for the order \preceq (cf. [20, Section 6.11]). Now, let us denote by μ the element of $W \bullet_\ell \lambda$ contained in $\overline{C_\ell}$, by $\mathbf{g}' \subset \overline{\mathbf{a}_\ell}$ the facet containing $\mu + \rho^\vee$ and by w the element of ${}_f W^{\ell^{-1} \cdot \mathbf{g}'}$ such that $w \square_\ell (\mu + \rho^\vee) = \lambda + \rho^\vee$ (cf. the first isomorphism of (6.4)). Then the alcove $A := \rho^\vee + C_\lambda$ is the only alcove for \square_ℓ containing $\lambda + \rho^\vee$ in its closure which is minimal in

$$W_{\mathbf{h} + \rho^\vee, \square_\ell} A = W_{w \mathbf{g}', \square_\ell} A = A W_{\mathbf{g}', \square_\ell}$$

for \preceq . Likewise, the alcove $A_w := \ell^{-1} \cdot A$ is the only alcove (for \square_1) containing $\ell^{-1} \cdot (\mathbf{h} + \rho^\vee)$ in its closure which is minimal in $A_w W_{\mathbf{g}}$ for \preceq , where $\mathbf{g} := \ell^{-1} \cdot \mathbf{g}'$. By definition of the operation $\hat{\cdot}$ on the set of alcoves for \bullet_ℓ , we have $\hat{C}_\lambda = \ell \cdot \hat{A}_w - \rho^\vee$, and so $\hat{C}_\lambda - \ell \cdot \rho^\vee = \ell \cdot (\hat{A}_w - \rho^\vee) - \rho^\vee$. In particular (once again by definition of the application $d(\cdot)$ on the set of alcoves for \bullet_ℓ), we have

$$(7.1) \quad d(\hat{C}_\lambda - \ell \cdot \rho^\vee) = d(\hat{A}_w - \rho^\vee).$$

7.2. A new proof of Donkin's Theorem

As an application of the study that was made in the previous sections, we can give a new proof of the description of blocks of $\text{Rep}_k(G^\vee)$. Unless specified otherwise, G is assumed to be a semi-simple algebraic group of adjoint type over \mathbb{F} (so G^\vee is simply connected). For any $\lambda \in \mathbb{X}^\vee$, we define $r(\lambda)$ to be the unique non-negative integer such that

$$\lambda \in \ell^{r(\lambda)} \cdot \mathbb{X}^\vee \setminus \ell^{r(\lambda)+1} \cdot \mathbb{X}^\vee.$$

If we let λ' be the W -conjugate (for the action \square_ℓ) of λ contained in $\overline{\mathbf{a}_\ell}$, $\mathbf{g}_{\lambda'} \subset \overline{\mathbf{a}_\ell}$ be the facet containing λ' , put $\mathbf{g} := \ell^{-1} \cdot \mathbf{g}_{\lambda'}$ and let w be the unique element of ${}_f W^{\mathbf{g}}$ such that $w \square_\ell \lambda' = \lambda$ (cf. (6.4)), then one can easily check that

$$r(\lambda) = r_{\mathbf{g}}(w) + 1,$$

where $r_{\mathbf{g}}(w)$ is as it was defined in Subsection 6.7.

We take back the setting of the beginning of Subsection 5.4 for the next statement: since G^\vee is semi-simple and simply connected, we get a decomposition $G^\vee = G_1^\vee \times \cdots \times G_t^\vee$ into simply connected simple algebraic groups,

each G_i^\vee admitting \mathfrak{R}_i^\vee as a root system; this decomposition induces a decomposition of the root system, affine Weyl group and dominant characters attached to G^\vee . Moreover, for every i , we will denote by $\rho_i^\vee \in \mathbb{X}_{i,+}^\vee$ the half sum of positive coroots relative to G_i^\vee and, for any positive integer r , we will denote by $W_i^{(r)}$ the subgroup of W_i whose translation part has been dilated by ℓ^r (cf. Subsection 6.7 for the precise definition).

THEOREM 7.1. — *Let $\mu = (\mu_1, \dots, \mu_r) \in \mathbb{X}_+^\vee = \prod_i \mathbb{X}_{i,+}^\vee$, and denote by $\bar{\mu}$ the equivalence class of μ for the equivalence relation \sim (cf. Section 2) on \mathbb{X}_+^\vee (seen as the weight poset of $\text{Rep}_\mathbf{k}(G^\vee)$). We have*

$$\bar{\mu} = \prod_{i=1}^r W_i^{(r(\mu_i + \rho_i^\vee))} \bullet_\ell \mu_i \cap \mathbb{X}_{i,+}^\vee.$$

Proof. — Recall that thanks to the geometric Satake equivalence (Theorem 6.2) and Theorem 6.4, we have an equivalence of highest weight categories

$$(7.2) \quad \text{Rep}_\mathbf{k}(G^\vee) \xrightarrow{\sim} \text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$$

sending a tilting module T_μ to the tilting Iwahori–Whittaker perverse sheaf $\mathcal{G}_\lambda^{\mathcal{IW}}$, where $\lambda := \mu + \rho^\vee$. Therefore, proving the claim is equivalent to showing that for any $\lambda \in \mathbb{X}_{++}^\vee = \mathbb{X}_+^\vee + \rho^\vee = \prod_i (\mathbb{X}_{i,+}^\vee + \rho_i^\vee)$, we have the equality

$$(7.3) \quad \bar{\lambda} = \prod_{i=1}^r W_i^{(r(\lambda_i))} \square_\ell \lambda_i \cap \mathbb{X}_{i,+}^\vee,$$

where $\bar{\lambda}$ now denotes the equivalence class of $\lambda = (\lambda_1, \dots, \lambda_r)$ for the equivalence relation \sim on \mathbb{X}_{++}^\vee , seen as the weight poset of $\text{Perv}_{\mathcal{IW}}(\text{Gr}, \mathbf{k})$.

Denote by λ' the unique W -conjugate (for the dilated box action \square_ℓ) of λ which is contained in $\overline{\mathbf{a}_\ell}$, by $\mathbf{g}_{\lambda'} \subset \overline{\mathbf{a}_\ell}$ the facet containing λ' , put $\mathbf{g} := \ell^{-1} \cdot \mathbf{g}_{\lambda'} \subset \overline{\mathbf{a}_1}$ and let w the element of ${}_f W^{\mathbf{g}}$ such that $w \square_\ell \lambda' = \lambda$. Recall that ${}_f W^{\mathbf{g}} \square_\ell \lambda' = W \square_\ell \lambda' \cap \mathbb{X}_{++}^\vee$. By (6.5), we have

$$(w \square_\ell \lambda') \sim (w' \square_\ell \lambda') \iff w \sim_{\mathbf{g}} w', \quad \forall w, w' \in {}_f W^{\mathbf{g}},$$

and by Proposition 5.18 we have

$$w \sim_{\mathbf{g}} w' \iff w_i \sim_{\mathbf{g}_i} w'_i, \quad \forall i,$$

where $w = (w_1, \dots, w_r)$, $w' = (w'_1, \dots, w'_r)$.

On the other hand, the linkage principle tells us that

$$(w \square_\ell \lambda') \sim \mu \implies \mu \in W \square_\ell \lambda',$$

from which we deduce (also using Lemma 6.12) that proving (7.3) is equivalent to proving that

$$\bar{w} = \prod_{i=1}^r W_i^{(r(\lambda_i))} w_i W_{\mathbf{g}_i} \cap {}_{\mathbf{f}} W_i^{\mathbf{g}_i}.$$

But this last equality was proved in Theorem 6.14 (because $r(\lambda_i) = r_{\mathbf{g}_i}(w_i) + 1$ for all i). \square

We can also apply our Corollary 5.33 to give a bound on the length of a minimum chain linking two weights in the same block. We start with a lemma.

LEMMA 7.2. — Let $\lambda, \mu \in \mathbb{X}_+^\vee$, $r \in \mathbb{Z}_{\geq 0}$, and put $\lambda' := \ell^r \cdot (\lambda + \rho^\vee) - \rho^\vee$, $\mu' := \ell^r \cdot (\mu + \rho^\vee) - \rho^\vee$. We have

$$(T_{\lambda'} : \nabla_{\mu'}) = (T_\lambda : \nabla_\mu).$$

Proof. — By the geometric Satake equivalence (Theorem 6.2) coupled with Theorem 6.4, we have an equivalence of highest weight categories

$$\mathrm{Rep}_{\mathbf{k}}(G^\vee) \xrightarrow{\sim} \mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr}, \mathbf{k}),$$

sending T_γ to $\mathcal{T}_{\gamma+\rho^\vee}^{\mathcal{IW}}$ for every $\gamma \in \mathbb{X}_+^\vee$. We can thus apply Corollary 6.11:

$$\begin{aligned} (T_{\lambda'} : \nabla_{\mu'}) &= \left(\mathcal{T}_{\ell^r \cdot (\lambda + \rho^\vee)}^{\mathcal{IW}} : \nabla_{\ell^r \cdot (\mu + \rho^\vee)}^{\mathcal{IW}} \right) \\ &\stackrel{\text{Corollary 6.11}}{=} (\mathcal{T}_{\lambda+\rho^\vee}^{\mathcal{IW}}, \nabla_{\mu+\rho^\vee}^{\mathcal{IW}}) = (T_\lambda : \nabla_\mu). \quad \square \end{aligned}$$

PROPOSITION 7.3. — Assume that G^\vee is a simple and simply connected algebraic group over \mathbf{k} . Let λ, λ' be two elements of \mathbb{X}_+^\vee in the same equivalence class for \sim , and denote by $\tilde{\lambda}$ (resp. $\tilde{\lambda}'$) the unique element of \mathbb{X}_+^\vee which satisfies $\lambda + \rho^\vee = \ell^r \cdot (\tilde{\lambda} + \rho^\vee)$ (resp. $\lambda' + \rho^\vee = \ell^r \cdot (\tilde{\lambda}' + \rho^\vee)$), where $r = r(\lambda + \rho^\vee)$ (notice that $\lambda = \tilde{\lambda}$ when $r = 0$). Also denote by $C_{\tilde{\lambda}}$ (resp. $C_{\tilde{\lambda}'}$) the alcove containing $\tilde{\lambda}$ (resp. $\tilde{\lambda}'$) in its upper closure. Then there exists a chain of dominant characters

$$\lambda_s = \lambda, \lambda_{s-1}, \dots, \lambda_0 = \lambda'$$

such that, for all $i \in \llbracket 0, s-1 \rrbracket$, there exists an indecomposable G^\vee -module M_i satisfying

$$[M_i : L_{\lambda_i}] \neq 0 \quad \text{and} \quad [M_i : L_{\lambda_{i+1}}] \neq 0$$

and such that

$$s \leq 2 + d\left(\hat{C}_{\tilde{\lambda}} - \ell \cdot \rho^\vee\right) + d\left(\hat{C}_{\tilde{\lambda}'} - \ell \cdot \rho^\vee\right).$$

Proof. — Denote by μ the W -conjugate (for \bullet_ℓ) of λ which is included in $\overline{C_\ell}$. We will first deal with the case where $r = 0$, which means that the facets (for \square_ℓ) containing $\lambda + \rho^\vee$ and $\lambda' + \rho^\vee$ are not special (notice that, since λ and λ' are in the same equivalence class, we must have $r(\lambda + \rho^\vee) = r(\lambda' + \rho^\vee)$ thanks to Theorem 7.1).

We denote by $\mathbf{g}_\mu \subset \overline{\mathbf{a}_\ell}$ the facet containing $\mu + \rho^\vee$, put $\mathbf{g} := \ell^{-1} \cdot \mathbf{g}_\mu$ and let w, w' be the elements of ${}_f W^{\mathbf{g}}$ satisfying $w \square_\ell (\mu + \rho^\vee) = \lambda + \rho^\vee$, $w' \square_\ell (\mu + \rho^\vee) = \lambda' + \rho^\vee$. The facet \mathbf{g} for \square_1 is a non-special facet included in the closure of $\overline{\mathbf{a}_1}$. Let us also denote by A_w , resp. $A_{w'}$, the alcove containing $w\mathbf{g}$, resp. $w'\mathbf{g}$, in its closure and which is minimal in $A_w W_{\mathbf{g}}$, resp. $A_{w'} W_{\mathbf{g}}$, for the order \preceq . Thus, we can apply Corollary 5.33, and pick a chain of elements of ${}_f W^{\mathbf{g}}$

$$w_s = w, w_{s-1}, \dots, w_0 = w'$$

as in this corollary, with $s \leq 2 + d(\widehat{A}_w - \rho^\vee) + d(\widehat{A}_{w'} - \rho^\vee)$. Thanks to the equation (7.1), we have that $d(\widehat{A}_w - \rho^\vee) = d(\widehat{C}_\lambda - \ell \cdot \rho^\vee)$ and $d(\widehat{A}_{w'} - \rho^\vee) = d(\widehat{C}_{\lambda'} - \ell \cdot \rho^\vee)$, so that s is bounded by the desired integer. This same corollary tells us that for all $i \in \llbracket 0, s-1 \rrbracket$, there exists an element $u_i \in {}_f W^{\mathbf{g}}$ satisfying

$$n_{w_i, u_i}(1) \neq 0 \quad \text{and} \quad n_{w_{i+1}, u_i}(1) \neq 0.$$

By Proposition 4.5, we can replace n with ℓn in the above, so that the character formula of tilting modules given in [30, Theorem 8.9] yields

$$(T_{u_i \bullet_\ell \mu} : \nabla_{w_i \bullet_\ell \mu}) \neq 0 \quad \text{and} \quad (T_{u_i \bullet_\ell \mu} : \nabla_{w_{i+1} \bullet_\ell \mu}) \neq 0,$$

which implies that

$$[T_{u_i \bullet_\ell \mu} : L_{w_i \bullet_\ell \mu}] \neq 0 \quad \text{and} \quad [T_{u_i \bullet_\ell \mu} : L_{w_{i+1} \bullet_\ell \mu}] \neq 0.$$

So we get the result by putting $\lambda_i = w_i \bullet_\ell \mu$ and $M_i = T_{u_i \bullet_\ell \mu}$.

Now we assume that $r > 0$. Since $r(\tilde{\lambda} + \rho) = r(\tilde{\lambda}' + \rho) = 0$, we can apply the previous step and find two sequences of dominant characters $(\tilde{\lambda}_i), (\tilde{\nu}_i)$ of the desired length such that $\tilde{\lambda}_s = \tilde{\lambda}, \tilde{\lambda}_0 = \tilde{\lambda}'$ and

$$(T_{\tilde{\nu}_i} : \nabla_{\tilde{\lambda}_i}) \neq 0 \quad \text{and} \quad (T_{\tilde{\nu}_i} : \nabla_{\tilde{\lambda}_{i+1}}) \neq 0.$$

By Lemma 7.2, we see that if we put $\nu_i := \ell^r \cdot (\tilde{\nu}_i + \rho) - \rho$ and $\lambda_i := \ell^r \cdot (\tilde{\lambda}_i + \rho) - \rho$, we get:

$$(T_{\nu_i} : \nabla_{\lambda_i}) = (T_{\tilde{\nu}_i} : \nabla_{\tilde{\lambda}_i}) \neq 0 \quad \text{and} \quad (T_{\nu_i} : \nabla_{\lambda_{i+1}}) = (T_{\tilde{\nu}_i} : \nabla_{\tilde{\lambda}_{i+1}}) \neq 0,$$

from which we deduce that

$$[\nu_i : L_{\lambda_i}] \neq 0 \quad \text{and} \quad [\nu_i : L_{\lambda_{i+1}}] \neq 0.$$

This concludes the proof of Proposition 7.3. \square

We will now use the process described in [20, Section II.7.3] to deduce from Theorem 7.1 the block decomposition of $\text{Rep}_{\mathbf{k}}(G^\vee)$ in the general case where G^\vee is a reductive group⁽¹⁵⁾, starting with a lemma.

LEMMA 7.4. — *Let H_1, H_2 be reductive algebraic groups over \mathbf{k} , with $T_1 \subset B_1$, resp. $T_2 \subset B_2$, a maximal torus and a Borel subgroup of H_1 , resp. of H_2 , and $\varphi : H_1 \rightarrow H_2$ a central isogeny such that $\varphi(T_1) = T_2$, $\varphi(B_1) = B_2$. Denote by $\mathbb{X}(T_1)_+ \subset \mathbb{X}(T_1)$ (resp. $\mathbb{X}(T_2)_+ \subset \mathbb{X}(T_2)$) the dominant characters and characters associated with $T_1 \subset B_1$ (resp. $T_2 \subset B_2$).*

Then φ induces an injective morphism $\mathbb{X}(T_2)_+ \hookrightarrow \mathbb{X}(T_1)_+$, and the blocks of H_2 are the blocks of H_1 contained in $\mathbb{X}(T_2)_+$.

Proof. — Since φ is a central isogeny, the morphism induced on root data identifies the root systems of H_1 and H_2 , (cf. [20, Section II.1.17]). Therefore, the injection $\mathbb{X}(T_2) \hookrightarrow \mathbb{X}(T_1)$ (induced by $\varphi|_{T_1}$) induces an injection $\mathbb{X}(T_2)_+ \hookrightarrow \mathbb{X}(T_1)_+$. On the other hand, pulling back by φ induces a fully-faithful functor $\varphi^* : \text{Rep}_{\mathbf{k}}(H_2) \rightarrow \text{Rep}_{\mathbf{k}}(H_1)$, which sends an indecomposable tilting module of highest weight λ to the indecomposable tilting module of same highest weight (cf. [20, Section E.7]), and such that there exists no non-zero morphism between an indecomposable tilting module in the essential image of φ^* and an indecomposable tilting module of $\text{Rep}_{\mathbf{k}}(H_1)$ which is not in this essential image. The conclusion follows easily, thanks to the description of blocks via tilting modules (cf. Theorem 2.3). \square

COROLLARY 7.5. — *Assume that G is a general reductive group. Denote by $\mathcal{D}G^\vee$ the derived subgroup of G^\vee , by T_2 the reduced part of the neutral connected component of the centre of G^\vee and let H_1, \dots, H_t be the simply connected covers of the minimal closed connected normal subgroups of positive dimension of $\mathcal{D}G^\vee$. For each $i \in \llbracket 1, t \rrbracket$, also denote by T_i^\vee the split maximal torus of H_i determined by the split maximal torus of G^\vee which is Langlands dual to T , by W_i (resp. by $\mathbb{X}_{i,+}^\vee$, resp. by ρ_i^\vee) the affine Weyl group (resp. the set of dominant characters of T_i^\vee determined by \mathfrak{R}_+^\vee , resp. the half-sum of positive roots) associated with H_i , seen as a subgroup of the affine Weyl group associated with $\mathcal{D}G^\vee$, and by $\mathbb{X}(T_2)$ the group of characters of T_2 . Then there is a central isogeny*

$$\varphi : \mathcal{D}G^\vee \times T_2 \longrightarrow G^\vee,$$

⁽¹⁵⁾We warn the reader that the statement (3) given in [20, Section II.7.3] is wrong, since one needs to assume the semi-simple group of loc. cit. to be simple for it to be correct.

which induces an injection $\mathbb{X}_+^\vee \hookrightarrow \prod_i \mathbb{X}_{i,+}^\vee \times \mathbb{X}(T_2)$ and such that, for every

$$\lambda = (\lambda_1, \dots, \lambda_t, \lambda_{t+1}) \in \mathbb{X}_+^\vee,$$

we have

$$\bar{\lambda} = \prod_{i=1}^t W_i^{(r(\lambda_i + \rho_i^\vee))} \bullet_\ell \lambda_i \cap \mathbb{X}_{i,+}^\vee \times \{\lambda_{t+1}\}.$$

Proof. — The fact that φ exists follows from [20, Section II.1.18]. Moreover, we know that $\mathcal{D}G^\vee$ is a semi-simple algebraic group, so that we have a central isogeny $H'_1 \times \dots \times H'_t \rightarrow \mathcal{D}G^\vee$, where H'_1, \dots, H'_t are the minimal closed normal subgroups of positive dimension of $\mathcal{D}G^\vee$, which are simple algebraic groups. We then get a central isogeny $H_1 \times \dots \times H_t \rightarrow \mathcal{D}G^\vee$ by replacing each H'_i with its simply connected cover. Therefore, Lemma 7.4 and Theorem 7.1 tell us that for any dominant character λ' of the maximal torus $T \cap \mathcal{D}G^\vee$ of $\mathcal{D}G^\vee$, the associated block of $\text{Rep}_\mathbf{k}(\mathcal{D}G^\vee)$ is equal to

$$\prod_{i=1}^t W_i^{(r(\lambda'_i + \rho_i^\vee))} \bullet_\ell \lambda'_i \cap \mathbb{X}_{i,+}^\vee,$$

where we see λ' as the element $(\lambda'_1, \dots, \lambda'_t)$ of $\prod_i \mathbb{X}_{i,+}^\vee$ (notice that the above set is included in the set of dominant characters of $T \cap \mathcal{D}G^\vee$, because each W_i is a subgroup of the affine Weyl group associated with $\mathcal{D}G^\vee$).

Finally, since T_2 is a torus (in particular, the category $\text{Rep}_\mathbf{k}(T_2)$ is semi-simple), one can easily check that the block of λ in $\mathcal{D}G^\vee \times T_2$ is equal to the product of blocks associated to $(\lambda_1, \dots, \lambda_t)$ and λ_{t+1} in $\text{Rep}_\mathbf{k}(\mathcal{D}G^\vee)$ and $\text{Rep}_\mathbf{k}(T_2)$ respectively, i.e. to

$$\prod_{i=1}^t W_i^{(r(\lambda_i + \rho_i^\vee))} \bullet_\ell \lambda_i \cap \mathbb{X}_{i,+}^\vee \times \{\lambda_{t+1}\}.$$

This concludes the proof by Lemma 7.4, since the above set is included in \mathbb{X}_+^\vee . \square

Remark 7.6. — We take back the context of Corollary 7.5, and let λ, λ' be dominant characters in the same equivalence class. The result of Proposition 7.3 can also be generalized to the case where G^\vee is a general reductive group. Indeed, using the central isogeny $\mathcal{D}G^\vee \times T_2 \rightarrow G^\vee$, one is reduced to proving it for $\mathcal{D}G^\vee$, and since $\mathcal{D}G^\vee$ is a central isogeny of the product of t simple subgroups (because $\mathcal{D}G^\vee$ is semi-simple), we can further reduce to the case where G^\vee is a product of simply connected simple groups $H_1 \times \dots \times H_t$. Finally, the equality (5.4) of Proposition 5.18 allows us to bound the length of a minimum chain linking two weights $(\lambda_1, \dots, \lambda_t)$ and $(\lambda'_1, \dots, \lambda'_t)$ in the same block by $\max\{s_i, i \in \llbracket 1, t \rrbracket\}$, where s_i is the bound

obtained in Proposition 7.3 for the length of a minimum chain linking λ_i and λ'_i .

7.3. Block decomposition for a quantum group

In this subsection, G is semi-simple of adjoint type. We assume that $\text{char}(\mathbf{k}) = 0$, and that there exists a primitive ℓ^{th} root of unity q in \mathbf{k} . Moreover, we assume that ℓ is odd, greater than the Coxeter number of \mathfrak{R}^\vee and not equal to 3 if \mathfrak{R}^\vee has a component of type G_2 . We then denote by $U_{q,\mathbf{k}}$ Lusztig's quantized enveloping algebra specialized at q and associated with G^\vee (we take the conventions of [20, Appendix H]). The category of finite dimensional representations of $U_{q,\mathbf{k}}$, which we will denote by $\text{Rep}(U_{q,\mathbf{k}})$, has many features in common with the category $\text{Rep}(G_{\mathbf{k}'}^\vee)$, where $G_{\mathbf{k}'}^\vee$ denotes the Langlands dual group of G over some field \mathbf{k}' of characteristic ℓ . In particular, $\text{Rep}(U_{q,\mathbf{k}})$ is a highest weight category with weight poset \mathbb{X}_+^\vee ; for any $\lambda \in \mathbb{X}_+^\vee$, we will denote by $T_q(\lambda)$, resp. $\nabla_q(\lambda)$, the indecomposable tilting module, resp. the costandard object, with highest weight λ .

We fix a weight $\lambda \in \bar{C}_\ell \cap \mathbb{X}^\vee$, let \mathbf{g}' be the facet for \square_ℓ containing $\lambda + \rho^\vee$ and w, w' be elements of ${}_f W^{\mathbf{g}}$, where $\mathbf{g} := \ell^{-1} \cdot \mathbf{g}'$ (recall the bijections (6.4)). Most importantly for us, the multiplicity of costandard objects in tilting objects is known.

PROPOSITION 7.7. — We have

$$(T_q(w \bullet_\ell \lambda) : \nabla_q(w' \bullet_\ell \lambda)) = n_{w',w}(1).$$

Moreover, we have that $(T_q(w \bullet_\ell \lambda) : \nabla_q(\lambda')) = 0$ whenever $\lambda' \notin W \bullet_\ell \lambda$.

Proof. — The second claim is due to the linkage principle for quantum groups, see [8, Section 8].

Let us denote by $W_{(0)}$ the subset of W consisting of elements w which are minimal in wW_0 , and recall that the action of Iw by left multiplication on $\text{Fl}_{\mathbf{a}_1}^\circ$, resp. on Gr° , yields stratifications

$$(\text{Fl}_{\mathbf{a}_1}^\circ)_{\text{red}} = \bigsqcup_{u \in W} X_u, \quad (\text{Gr}^\circ)_{\text{red}} = \bigsqcup_{v \in W_{(0)}} Y_v,$$

where $Y_u := \text{Iw} \cdot \dot{u}$, resp. $X_v := \text{Iw} \cdot \dot{v}$, is an affine \mathbb{F} -space of dimension $l(u)$, resp. $l(v)$. Moreover, the canonical projection $\pi : \text{Fl}_{\mathbf{a}_1}^\circ \rightarrow \text{Gr}^\circ$ is ind-proper, Iw -equivariant and satisfies $\pi^{-1}(Y_w) = \bigsqcup_{z \in W_0} X_{wz}$ for every $w \in W_{(0)}$. We denote by $\text{Perv}_{(\text{Iw})}(\text{Fl}_{\mathbf{a}_1}^\circ, \mathbf{k})$, resp. $\text{Perv}_{(\text{Iw})}(\text{Gr}^\circ, \mathbf{k})$, the category of perverse sheaves on $\text{Fl}_{\mathbf{a}_1}^\circ$, resp. Gr° , constant along the Iw -orbits, and by $\text{Rep}_{\bar{0}}(U_{q,\mathbf{k}})$

the Serre subcategory of $\text{Rep}(U_{q,\mathbf{k}})$ generated by the simple objects whose highest weight belongs to $W \bullet_\ell 0 \cap \mathbb{X}_+^\vee$ (the fact that $\text{Rep}_{\bar{0}}(U_{q,\mathbf{k}})$ is truly a block will follow from Proposition 7.10).

Thanks to [9], we have an equivalence of highest weight categories

$$\text{Rep}_{\bar{0}}(U_{q,\mathbf{k}}) \simeq \text{Perv}_{(\text{Iw})}(\text{Gr}^\circ, \mathbf{k}),$$

sending $T_q(x \bullet_\ell 0)$ (resp. $\nabla_q(y \bullet_\ell 0)$) to the indecomposable tilting object $T_{x^{-1}}^{(\text{Iw})}$ (resp. to the costandard object $\nabla_{y^{-1}}^{(\text{Iw})}$) associated with x , for any $x \in {}_f W$ (resp. $y \in {}_f W$).⁽¹⁶⁾

For any $u \in W$, let us denote by $\mathcal{T}_u \in \text{Perv}_{(\text{Iw})}(\text{Fl}_{\mathbf{a}_1}^\circ, \mathbf{k})$ the indecomposable tilting object associated with u ; by [38, Proposition 3.4.1], we have an isomorphism $\pi_! \mathcal{T}_w \simeq T_w^{(\text{Iw})}$ for any $w \in W_{(0)}$. Therefore, combining [38, (3.4.1)] with [38, Theorem 5.3.1], we get

$$\left(T_u^{(\text{Iw})} : \nabla_v^{(\text{Iw})} \right) = \sum_{z \in W_0} (-1)^{l(z)} h_{vz,u}(1), \quad \forall u, v \in W_{(0)},$$

where $(h_{x,y}, x, y \in W)$ denotes the ordinary Kazhdan–Lusztig polynomials associated with W , as described in [31]. Using [31, Proposition 3.4] together with the fact that $h_{x,y} = h_{x^{-1},y^{-1}}$ for all $x, y \in W$ (which is a consequence of the fact that the anti-automorphism i from the proof of [31, Theorem 2.7] commutes with the involution $d : \mathcal{H} \rightarrow \mathcal{H}$ defined in loc. cit.) show that the right-hand side in the above equation coincides with $n_{v^{-1},u^{-1}}(1)$. So we get

$$(T_q(w \bullet_\ell 0) : \nabla_q(w' \bullet_\ell 0)) = \left(T_{w^{-1}}^{(\text{Iw})} : \nabla_{w'^{-1}}^{(\text{Iw})} \right) = n_{w',w}(1),$$

which proves our claim in the case where $\lambda = 0$. Using the second point of [31, Remark 7.2], one can deduce from this the general case. \square

Remark 7.8. — The character formula of Proposition 7.7 was originally stated as a conjecture in [31, Conjecture 7.1], and proved in [32] for $\ell > 33$.

In particular, the second assertion of Proposition 7.7 tells us that $\bar{\mu} \subset W \bullet_\ell \mu$ for any $\mu \in \mathbb{X}_+^\vee$.

COROLLARY 7.9. — We have an equality

$$\begin{aligned} \dim_{\mathbf{k}} \text{Hom}_{\text{Rep}(U_{q,\mathbf{k}})}(T_q(w \bullet_\ell \lambda), T_q(w' \bullet_\ell \lambda)) \\ = \dim_{\mathbf{k}} \text{Hom}_{\text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{a}_1}^\circ, \mathbf{k})}^{\bullet}(\mathcal{E}_w^{\mathbf{g}}, \mathcal{E}_{w'}^{\mathbf{g}}). \end{aligned}$$

⁽¹⁶⁾Notice that $w \mapsto w^{-1}$ induces a bijection ${}_f W \simeq W_{(0)}$.

Proof. — Proposition 7.7 and standard arguments (see [3, Section 6.2]) show that we have

$$(7.4) \quad \dim_{\mathbf{k}} \text{Hom}_{\text{Rep}(U_{q,\mathbf{k}})}(T_q(w \bullet_{\ell} \lambda), T_q(w' \bullet_{\ell} \lambda)) = \sum_{y \in {}_t W^{\mathbf{g}}} n_{y,w}(1) \cdot n_{y,w'}(1).$$

But we also know that

$$\dim_{\mathbf{k}} \text{Hom}_{D_{\mathcal{IW}}^b(\text{Fl}_{\mathbf{g}}, \mathbf{k})}^{\bullet}(\mathcal{E}_x^{\mathbf{g}}, \nabla_y^{\mathbf{g}}) = n_{y,x}(1), \quad \forall x, y \in {}_t W^{\mathbf{g}}$$

thanks to (4.8) together with Lemma 2.2. So by [21, Proposition 2.6] the right-hand side in (7.4) coincides with

$$\dim_{\mathbf{k}} \text{Hom}_{\text{Par}_{\mathcal{IW}}(\text{Fl}_{\mathbf{g}}, \mathbf{k})}^{\bullet}(\mathcal{E}_w^{\mathbf{g}}, \mathcal{E}_{w'}^{\mathbf{g}}).$$

□

Therefore, we get a very similar situation as the one in Subsection 6.5:

$$w \bullet_{\ell} \lambda \sim w' \bullet_{\ell} \lambda \iff w \sim_{\mathbf{g}} w'.$$

For any $\mu \in \mathbb{X}_+^{\vee}$, recall the definition of $r(\mu)$ from Subsection 7.2, and put

$$\delta(\mu) := \begin{cases} 1 & \text{if } r(\mu) = 0 \\ 0 & \text{otherwise.} \end{cases}$$

We define W^{δ} to be equal to W when $\delta = 1$, and to $\{\text{id}\}$ otherwise. The next result is obtained from Theorem 6.14 in the same way that Theorem 7.1 was, from which we take back the notations.

PROPOSITION 7.10. — *Let $\mu = (\mu_1, \dots, \mu_r) \in \mathbb{X}_+^{\vee} = \prod_i \mathbb{X}_{i,+}^{\vee}$, and denote by $\bar{\mu}$ the equivalence class of μ in \mathbb{X}_+^{\vee} (seen as the weight poset of $\text{Rep}(U_{q,\mathbf{k}})$) for the equivalence relation \sim from Section 2. We have*

$$\bar{\mu} = \prod_{i=1}^r W_i^{\delta(\mu_i + \rho_i^{\vee})} \bullet_{\ell} \mu_i \cap \mathbb{X}_{i,+}^{\vee}.$$

Remarks 7.11.

- (1) The proof of the previous proposition does not require Smith–Treumann theory (because the proof of Theorem 6.14 does not require it when $\text{char}(\mathbf{k}) = 0$).
- (2) One can use Corollary 5.33 to give a bound for the length of a minimal chain linking two weights in the same block for $\text{Rep}(U_{q,\mathbf{k}})$, of the same kind as Proposition 7.3.
- (3) Another proof of Proposition 7.10 was found in [34] (under the same assumptions on ℓ). However, the results of loc. cit. use the proof of Donkin from [15], so it does not allow to give a bound as in the previous point.

BIBLIOGRAPHY

- [1] P. N. ACHAR, *Perverse Sheaves and Applications to Representation Theory*, Mathematical Surveys and Monographs, vol. 258, American Mathematical Society, 2021.
- [2] P. N. ACHAR, N. COONEY & S. RICHE, “The parabolic exotic t-structure”, *Épjournal de Géom. Algébr.*, *EPIGA* **2** (2018), article no. 8 (31 pages).
- [3] P. N. ACHAR & S. RICHE, “Modular perverse sheaves on flag varieties I: tilting and parity sheaves”, *Ann. Sci. Éc. Norm. Supér. (4)* **49** (2016), no. 2, p. 325–370, with a joint appendix by Geordie Williamson.
- [4] ———, “Reductive groups, the loop Grassmannian, and the Springer resolution”, *Invent. Math.* **214** (2018), no. 1, p. 289–436.
- [5] ———, “A geometric Steinberg formula”, 2021, <https://arxiv.org/abs/2109.11980v1>.
- [6] ———, “A geometric model for blocks of Frobenius kernels”, 2022, <https://arxiv.org/abs/2203.03530v1>.
- [7] H. H. ANDERSEN, “An inversion formula for the Kazhdan–Lusztig polynomials for affine Weyl groups”, *Adv. Math.* **60** (1986), no. 2, p. 125–153.
- [8] H. H. ANDERSEN, P. POLO & W. KEXIN, “Representations of quantum algebras”, *Invent. Math.* **104** (1991), no. 1, p. 1–59.
- [9] S. ARKHIPOV, R. BEZRUKAVNIKOV & V. GINZBURG, “Quantum groups, the loop Grassmannian, and the Springer resolution”, *J. Am. Math. Soc.* **17** (2004), no. 3, p. 595–678.
- [10] P. BAUMANN & S. RICHE, “Notes on the geometric Satake equivalence”, in *Relative aspects in Representation Theory, Langlands Functoriality and Automorphic Forms* (V. Heiermann & D. Prasad, eds.), Lecture Notes in Mathematics, vol. 2221, Springer, 2018, p. 1–134.
- [11] A. BEILINSON, J. BERNSTEIN, P. DELIGNE & O. GABBER, *Faisceaux pervers. Proceedings of the colloquium “Analysis and topology on singular spaces”*, Luminy, France, July 6–10, 1981. Part I, 2nd ed., Astérisque, vol. 100, Société Mathématique de France, 2018.
- [12] R. BEZRUKAVNIKOV, D. GAITSGORY, I. MIRKOVIĆ, S. RICHE & L. RIDER, “An Iwahori–Whittaker model for the Satake category”, *J. Éc. Polytech., Math.* **6** (2019), p. 707–735.
- [13] N. BOURBAKI, *Éléments de mathématique. Groupes et algèbres de Lie: Chapitres 4 à 6*, Springer, 2007.
- [14] J. CIAPPARA, “Hecke category actions via Smith–Treumann theory”, 2021, <https://arxiv.org/abs/2103.07091>.
- [15] S. DONKIN, “The blocks of a semisimple algebraic group”, *J. Algebra* **67** (1980), no. 1, p. 36–53.
- [16] J. GIBSON, “Lievis: Lie theory visualisations”, <https://www.jgibson.id.au/lievis/>, 2022.
- [17] R. GORDON & E. L. GREEN, “Graded artin algebras”, *J. Algebra* **76** (1982), no. 1, p. 111–137.
- [18] J. E. HUMPHREYS, *Linear Algebraic Groups*, Graduate Texts in Mathematics, vol. 21, Springer, 1975.
- [19] J. E. HUMPHREYS & J. C. JANTZEN, “Blocks and indecomposable modules for semisimple algebraic groups”, *J. Algebra* **54** (1978), no. 2, p. 494–503.
- [20] J. C. JANTZEN, *Representations of Algebraic Groups*, 2nd ed., Mathematical Surveys and Monographs, vol. 107, American Mathematical Society, 2003.
- [21] D. JUTEAU, C. MAUTNER & G. WILLIAMSON, “Parity sheaves”, *J. Am. Math. Soc.* **27** (2014), no. 4, p. 1169–1212.

- [22] M.-A. KNUS, *Quadratic and Hermitian forms over rings*, Grundlehren der Mathematischen Wissenschaften, vol. 294, Springer, 1991, with a foreword by I. Bertuccioni.
- [23] S. LESLIE & G. LONERGAN, “Parity sheaves and Smith theory”, *J. Reine Angew. Math.* **777** (2021), p. 49-87.
- [24] G. LUSZTIG, “Hecke algebras and Jantzen’s generic decomposition patterns”, *Adv. Math.* **37** (1980), no. 2, p. 121-164.
- [25] J. S. MILNE, *Algebraic groups. The theory of group schemes of finite type over a field*, Cambridge Studies in Advanced Mathematics, vol. 170, Cambridge University Press, 2017.
- [26] I. MIRKOVIĆ & K. VILONEN, “Geometric Langlands duality and representations of algebraic groups over commutative rings”, *Ann. Math.* **166** (2007), no. 1, p. 95-143.
- [27] G. PAPPAS & M. RAPOPORT, “Twisted loop groups and their affine flag varieties”, *Adv. Math.* **219** (2008), no. 1, p. 118-198.
- [28] S. RICHE, “Geometric Representation Theory in positive characteristic”, Habilitation à diriger des recherches, Université Blaise Pascal (Clermont Ferrand 2), 2016.
- [29] S. RICHE & G. WILLIAMSON, *Tilting modules and the p -canonical basis*, Astérisque, vol. 397, Société Mathématique de France, 2018.
- [30] ———, “Smith–Treumann theory and the linkage principle”, *Publ. Math., Inst. Hautes Étud. Sci.* **136** (2022), p. 225-292.
- [31] W. SOERGEL, “Kazhdan–Lusztig polynomials and a combinatoric for tilting modules”, *Represent. Theory* **1** (1997), p. 83-114.
- [32] ———, “Character formulas for tilting modules over Kac–Moody algebras”, *Represent. Theory* **2** (1998), no. 13, p. 432-448.
- [33] THE STACKS PROJECT AUTHORS, “The Stacks project”, <https://stacks.math.columbia.edu>, 2022.
- [34] L. THAMS, “The blocks of a quantum algebra”, *Commun. Algebra* **22** (1994), no. 5, p. 1617-1628.
- [35] D. TREUMANN, “Smith theory and geometric Hecke algebras”, *Math. Ann.* **375** (2019), no. 1-2, p. 595-628.
- [36] M. DE VISSCHER, “On the blocks of semisimple algebraic groups and associated generalized Schur algebras”, *J. Algebra* **319** (2008), no. 3, p. 952-965.
- [37] G. WILLIAMSON, “Modular intersection cohomology complexes on flag varieties”, *Math. Z.* **272** (2012), no. 3-4, p. 697-727, with an appendix by T. Braden.
- [38] Z. YUN, “Weights of mixed tilting sheaves and geometric Ringel duality”, *Sel. Math., New Ser.* **14** (2009), no. 2, p. 299-320.

Manuscrit reçu le 7 septembre 2022,
 révisé le 28 mars 2024,
 accepté le 17 juin 2024.

Emilien ZABETH
 Université Clermont Auvergne
 CNRS, LMBP
 F-63000 Clermont-Ferrand (France)
 emilien.zabeth@uca.fr