

# ANNALES DE L'INSTITUT FOURIER

CHARLES FEFFERMAN

## **Convergence on almost every line for functions with gradient in $L^p(\mathbf{R}^n)$**

*Annales de l'institut Fourier*, tome 24, n° 3 (1974), p. 159-164

<[http://www.numdam.org/item?id=AIF\\_1974\\_\\_24\\_3\\_159\\_0](http://www.numdam.org/item?id=AIF_1974__24_3_159_0)>

© Annales de l'institut Fourier, 1974, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>*

## CONVERGENCE ON ALMOST EVERY LINE FOR FUNCTIONS WITH GRADIENT IN $L^p(\mathbf{R}^n)$

by Charles FEFFERMAN

This note answers a question asked by L.D. Kudrjačev ([1], p. 264, problem 1). The question was suggested by the following result of Uspenskiĭ [2] : if  $u(x)$  is a smooth function  $\mathbf{R}^n \rightarrow \mathbf{R}$  and  $\int_{\mathbf{R}^n} |\operatorname{grad} u|^p dx < \infty$  ( $1 < p < n$ ), there exists a constant  $c$  such that  $\lim_{r \rightarrow \infty} u(rx') = c$  for almost every  $x' \in \mathbf{S}^{n-1}$ . Professor Kudrjačev kindly informed me that V. Portnov answered his question independently by another method [3].

We use now the notation

$$x = (x_1, x'), \quad \text{where } x_1 \in \mathbf{R} \quad \text{and} \quad x' \in \mathbf{R}^{n-1}.$$

**THEOREM.** — Let  $u(x_1, x')$  be a smooth function :  $\mathbf{R}^n \rightarrow \mathbf{R}$  and suppose  $\int_{\mathbf{R}^n} |\operatorname{grad} u|^p dx < \infty$  ( $1 < p < n$ ). Then for a constant  $c$ ,  $\lim_{x_1 \rightarrow \infty} u(x_1, x') = c$  for almost all  $x'$ .

*Prof.* — Set  $u_j = \frac{\partial u}{\partial x_j}$ , let  $R_j$  denote  $j^{\text{th}}$  Riesz transform, and let  $I^1$  denote fractional integration of first order in  $\mathbf{R}^n$ .

We begin with the standard formula

$$\sum_i R_i R_j v_j = v_i + \sum_j R_j I^1 \left( \frac{\partial v_j}{\partial x_i} - \frac{\partial v_i}{\partial x_j} \right),$$

valid for  $C^\infty$  functions of compact support. (To check this, just take Fourier transforms). Take a function  $\varphi_N$  on  $\mathbf{R}^n$  equal to one for  $|x| \leq 2^N$ , supported in  $|x| \leq 2^{N+1}$ , and satisfying  $|\nabla \varphi_N| \leq C 2^{-N}$ ; and apply the above formula to the functions  $v_j = \varphi_N u_j \in C_0^\infty$ . As

$N \rightarrow \infty$ , the left-hand side of the formula tends to  $\sum_j R_i R_j u_j$  in  $L^p$ , since the Riesz transforms are bounded on  $L^p$ . On the other hand,  $\frac{\partial v_j}{\partial x_i} - \frac{\partial v_i}{\partial x_j} = -\left(\frac{\partial u_i}{\partial x_j} - \frac{\partial u_j}{\partial x_i}\right) \varphi_N + u_j \frac{\partial \varphi_N}{\partial x_i} - u_i \frac{\partial \varphi_N}{\partial x_j}$ , and the first term is zero since  $(u_i)$  is a gradient. Thus,

$$\frac{\partial v_j}{\partial x_i} - \frac{\partial v_i}{\partial x_j} \underset{p}{\leq} \|\operatorname{grad} u\|_p \|\operatorname{grad} \varphi_N\|_\infty \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

Since  $R_j I^1$  is bounded from  $L^p$  to  $L^s$  with  $\frac{1}{s} = \frac{1}{p} - \frac{1}{n}$  ( $1 < s < \infty$

since  $1 < p < n$ ), we see that as  $N \rightarrow \infty$ , the final term in our formula tends to zero in  $L^s$ . Thus, taking the limit in measure of both sides of our formula, we obtain  $u_i = \sum_j R_i R_j u_j$ .

In other words, setting  $f = \sum_j R_j \frac{\partial u}{\partial x_j} \in L^p$ , we have  $u_i = R_i f$ .

Now for  $f \in L^p$ , we know that  $\frac{\partial}{\partial x_i} (I^1 f) = R_i f$  in the sense of tempered distributions. (It's trivial for  $f \in C_0^\infty$ ; in general we express  $f$  as the limit in  $L^p$  of smooth compactly supported  $f_N$  as  $N \rightarrow \infty$ .  $R_i f_N \rightarrow R_i f$  in  $L^p$  and hence weakly as distributions, and  $I^1 f_N \rightarrow I^1 f$  in  $L^s$  and hence weakly, so that  $\frac{\partial}{\partial x_i} (I^1 f_N)$  also converges weakly to  $\frac{\partial}{\partial x_i} (I^1 f)$ .)

On the other hand, the Riesz transforms or 1<sup>st</sup> fractional integral of a smooth function in  $L^p$  are smooth functions, so that  $\frac{\partial}{\partial x_i} (u - I^1 f) = R_i f - R_i f = 0$ , not just as distributions, but as pointwise derivatives of smooth functions. Thus  $u = \text{constant} + I^1 f$ . So to prove the claim, it will be enough to show that

$$\lim_{x_1 \rightarrow \infty} I^1 f(x_1, x') = 0$$

for almost all  $x'$ , whenever  $f$  is a smooth function in  $L^p$ .

Now

$$\begin{aligned} I^1 f(x_1, x') &= \int_{\mathbb{R}^n} \frac{f(x) dy}{|x - y|^{n-1}} = \int_{|x-y|<1} \frac{f(y) dy}{|x - y|^{n-1}} + \\ &\quad + \int_{|x-y|\geq 1} \frac{f(y) dy}{|x - y|^{n-1}} = I + II. \end{aligned}$$

Then II is easy ; it tends to zero as  $x \rightarrow \infty$ . (In fact, we just break up  $f$  into  $f = g + b$  where  $\|g\|_p \leq \|f\|_p$  and  $g$  lives on a bounded set, while  $\|b\|_p < \epsilon$ . The contribution of  $g$  to II clearly tends to zero, while  $b$  produces at most

$$\begin{aligned} \|b\|_p \cdot \left( \int_{|x-y|\geq 1} |x - y|^{-(n-1)q} dy \right) \left[ \frac{1}{p} + \frac{1}{q} = 1 \right] &\leq \\ &\leq C \|b\|_p \leq C \epsilon, \quad \text{since } p < n. \end{aligned}$$

So the main problem is term I.

The main step in dealing with term I is to prove the

**LEMMA.** — Let  $f \in L^p(Q)$  where  $Q$  is a cube of side 10, and  $1 < p < n$ . Say  $Q = [0, 10] \times Q' \subseteq \mathbb{R}^1 \times \mathbb{R}^{n-1}$ . Then

$$M(x') = \sup_{0 \leq x_1 \leq 10} \left| \int_Q \frac{f(y_1, y') dy_1 dy'}{((y_1 - x_1)^2 + (y' - x')^2)^{\frac{n-1}{2}}} \right|$$

belongs to  $L^p(Q')$ , and  $\|M\|_p \leq C \|f\|_p$ .

*Proof.* — Say  $f \geq 0$  and  $f = 0$  outside  $Q$ . Let  $f^\downarrow(\cdot, y')$  be the decreasing re-arrangement of  $f(\cdot, y')$  on  $[0, 10]$ , for each fixed  $y' \in Q'$ . Of course  $\|f\|_p = \|f^\downarrow\|_p$ . We claim that  $Mf(x') \leq 2Mf^\downarrow(x')$  pointwise. In fact, fix  $x', x_1, y'$ , and consider

$$\int_{y_1 > x_1} \frac{f(y_1, y') dy_1}{((y_1 - x_1)^2 + (y' - x')^2)^{\frac{n-1}{2}}}.$$

Since  $x_{\{y_1 | y_1 > x_1\}} (y_1) \cdot ((y_1 - x_1)^2 + (y' - x')^2)^{-\frac{(n-1)}{2}}$  is monotone decreasing in the interval  $(x_1, \infty)$  in which it is supported, we know that

$$\int_{y_1 > x_1} \frac{f(y_1, y') dy_1}{((y_1 - x_1)^2 + (y' - x')^2)^{\frac{n-1}{2}}} \leq \int_0^{10} \frac{f^\downarrow(y_1, y') dy_1}{((y_1)^2 + (y' - x')^2)^{\frac{n-1}{2}}}.$$

Integrating over  $y'$  yields

$$\int_{\substack{y_1 > x_1 \\ y' \in Q'}} \frac{f(y_1, y') dy_1}{((y_1 - x_1)^2 + (y' - x')^2)^{\frac{n-1}{2}}} \leq \int_Q \frac{f^\downarrow(y_1, y') dy_1}{(y_1^2 + (y' - x')^2)^{\frac{n-1}{2}}}.$$

Similarly,

$$\int_{y_1 < x_1} \frac{f(y_1, y') dy_1 dy'}{((y_1 - x_1)^2 + (y' - x')^2)^{\frac{n-1}{2}}} \leq \int_Q \frac{f^\downarrow(y_1, y') dy_1 dy'}{(y_1 + (y' - x')^2)^{\frac{n-1}{2}}}.$$

Adding and taking the sup over  $x_1$  gives us

$$Mf(x') \leq 2 \int_Q \frac{f^\downarrow(y_1, y') dy_1 dy'}{(y_1^2 + (y' - x')^2)^{\frac{n-1}{2}}}. \quad (*)$$

which is stronger than the claim.

We shall use what we proved in full strength. For a fixed  $y_1$ , the function  $M_{y_1}(x') = \int_{y' \in Q'} \frac{f^\downarrow(y_1, y') dy'}{(y_1^2 + (y' - x')^2)^{\frac{n-1}{2}}}$  is just the convolution of  $f^\downarrow(y_1, .)$  with  $(y_1^2 + (y' - x')^2)^{-\frac{n-1}{2}}$  on the cube  $Q' \subseteq \mathbb{R}^{n-1}$ . Thus

$$\begin{aligned} \|M_{y_1}(\cdot)\|_p &\leq \|\text{convolution kernel}_{y_1}\|_{L^1(Q')} \|f(y_1, .)\|_p \sim \\ &\sim C \left( \log \frac{1}{|y_1|} \right) \|f^\downarrow(y_1, .)\|. \end{aligned}$$

Therefore by estimate (\*),  $Mf(x') \leq \int_0^{10} M_{y_1}(x') dy'$ , and

$$\begin{aligned} \|Mf\|_p &\leq \int_0^{10} C \left( \log \frac{1}{|y_1|} \right) \|f^\downarrow(y_1, .)\|_p dy_1 \leq \\ &\leq \left( \int_0^{10} \left( C \log \frac{1}{|y_1|} \right)^q dy_1 \right)^{1/q} \left( \int_0^{10} \|f(y_1, .)\|_p dy_1 \right)^{1/p} \\ (\text{Note : } q < \infty \text{ since } p > 1) &\leq C \|f\|_p = C \|f\|_p. \end{aligned}$$

Q.E.D.

Now we return to term I above. Divide  $\mathbb{R}^{n-1}$  into a mesh of cubes of side 2,  $\mathbb{R}^{n-1} = \bigcup_j Q'_j$ , and write  $\mathbb{R}^n = \bigcup_{i,j} Q_{ij}$  where

$$Q_{ij} = \{(x_1, x') \in \mathbb{R}^n \mid x' \in Q_j, 2i \leq x_1 < 2(i+1)\} (-\infty < i < \infty).$$

Then let  $Q^*$  be the cube concentric with  $Q_{ij}$  but with side 10, and set  $f_{ij} = f \chi_{Q_{ij}}^*$ . For

$$g_{ij}(x') = \begin{cases} \sup_{2i < x_1 < 2(i+1)} \int_{|x-y|<1} \frac{|f(y_1, y')| dy_1 dy'}{|x-y|^{n-1}}, \\ 0 \text{ otherwise} \end{cases}$$

where  $x = (x_1, x')$  if  $x' \in Q'_j$  we have  $\|g_{ij}\|_{L^p(\mathbb{R}^{n-1})} \leq C \|f_{ij}\|_{L^p(\mathbb{R}^n)}$  by the lemma. For

$$\begin{aligned} x' \in Q_j, \sup_{x_1 \in \mathbb{R}^1} \int_{\substack{|x-y|<1 \\ \text{where } x=(x_1, x')}} \frac{|f(y)| dy}{|x-y|^{\frac{n-1}{2}}} &= \\ &= \sup_i g_{ij}(x') \leq \left( \sum_i g_{ij}^p(x') \right)^{1/p}. \end{aligned}$$

So certainly

$$\begin{aligned} \int_{\mathbb{R}^{n-1}} \left[ \sup_{x_1 \in \mathbb{R}^1} \int_{|(x_1, x') - y| < 1} \frac{|f(y)| dy}{|x-y|^{n-1}} \right]^p dx' &= \\ &= \sum_j \int_{x' \in Q'_j} [\sup (\text{etc. etc.})]^p dx' \leq \sum_j \int_{\mathbb{R}^{n-1}} \left( \sum_i g_{ij}^p(x') \right)^{p/p} dx' = \\ &= \sum_{i,j} \int_{\mathbb{R}^{n-1}} g_{ij}^p(x') dx' \leq C \sum_{i,j} \|f_{ij}\|_p^p \leq C \|f\|_{L^p(\mathbb{R}^n)}^p. \end{aligned}$$

In other words, if

$$N(x') = Nf(x') = \sup_{x_1 \in \mathbb{R}^1} \int_{|(x_1, x') - y| < 1} \frac{|f(y)| dy}{|x-y|^{n-1}},$$

then

$$\|Nf\|_{L^p(\mathbb{R}^{n-1})} \leq C \|f\|_{L^p(\mathbb{R}^n)}.$$

Now term I becomes easy. Again split up if  $f = g + b$  where  $g$  is supported in a bounded set and  $\|b\|_p < \epsilon$ .

In evaluating the claim

$$\lim_{x_1 \rightarrow \infty} \int_{|(x_1, x') - y| < 1} \frac{|f(y)| dy}{|x - y|^{n-1}} = 0, \quad ((x_1, x') = x)$$

we find that  $g$  makes no contribution at all to the last integral for  $x_1$  large enough. On the other hand, for  $b$  we know that

$$\left| \limsup_{x_1 \rightarrow \infty} \int_{|(x_1, x') - y| < 1} \frac{|f(y)| dy}{|(x_1, x') - y|^{n-1}} \right|_{L^p(dx')} < C \epsilon \quad \text{by } (**)$$

Since  $\epsilon > 0$  is arbitrary, the proof is complete.

Q.E.D.

## BIBLIOGRAPHIE

- [1] L.D. KUDRJAVCEV, Svočtba graničnyh značenij funkcij iz vesovyh prostranstv i ih priloženija k kraevym zadačam. Mehanika Splošnoj sredy i rodstvennye problemy analiza. Moskva 1972.
- [2] S.V. USPENSKIĬ, O teorema vloženija dlja vesovyh klassov, *Trudi Mat. Insta AN SSSR*, 60 (1961), 282-303.
- [3] V. PORTNOV, *Doklady AN SSSR*, to appear.

Manuscrit reçu le 9 juillet 1973  
accepté par J.P. Kahane.

Charles FEFFERMAN,  
Departement of Mathematics  
University of Chicago  
5734 University Avenue  
Chicago, Ill. 60637.