

SURJIT SINGH KHURANA

**Radon-Nikodym property for vector-valued
integrable functions**

Annales de l'institut Fourier, tome 28, n° 3 (1978), p. 203-208

http://www.numdam.org/item?id=AIF_1978__28_3_203_0

© Annales de l'institut Fourier, 1978, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

RADON-NIKODYM PROPERTY FOR VECTOR-VALUED INTEGRABLE FUNCTIONS

by Surjit Singh KHURANA

It is proved in ([6], Theorem 1) that if a Banach space E possesses Radon-Nikodym (R - N) property, then the Banach space $L_p(E, \lambda)$, $1 < p < \infty$, of Bochner p -integrable functions also possesses this property. In this paper we give a new proof of the corresponding result when E is a Frechet space (i.e. a complete metrizable locally convex space [5]).

Let (Y, \mathcal{B}, ν) be a positive measure which is non-trivial (i.e., there exists a $B \in \mathcal{B}$ such that $0 < \nu(B) < \infty$) with $\mathcal{R} = \{A \in \mathcal{B} : \nu(A) < \infty\}$, E a Frechet space with $\{P_i\}$ an increasing sequence of semi-norms on E generating the topology of E , and $L_p(E, \lambda)$ the equivalence classes of strongly p -power integrable functions $Y \rightarrow E$, $1 \leq p < \infty$. (A function $f: X \rightarrow E$ is called strongly p -power integrable if there exists a sequence $\{f_n\}$ of \mathcal{R} -simple E -valued functions of Y such that (i) $f_n \rightarrow f$ a.e. $[\nu]$, and (ii) $\int [P_i(f_n - f)]_i^p d\nu \rightarrow 0$, $\forall i$. The increasing sequence of semi-norms

$$N_{i,p}, N_{i,p}(f) = \left[\int [P_i(f)]^p d\nu \right]^{1/p}$$

makes $L_p(E, \lambda)$ a Frechet space. We use the definition of [4] for a Frechet space to have R - N property.

$E = K$, we denote $L_p(E, \nu)$ by $L_p(\nu)$ and the corresponding norm by $\|\cdot\|_p$.

THEOREM. — *Suppose E is a Frechet space with R - N property and (Y, \mathcal{B}, ν) a non-trivial positive measure space. Then $L_p(E, \nu)$ has R - N property for $1 < p < \infty$.*

Proof. – Using ([3], Theorem 5 (iv)) it is sufficient to prove the R – N property for every separable closed subspace; this means we can assume that E is separable ([3], Theorem 5). Let $(X, \mathcal{U}, \lambda)$ be a finite measure space, $\mu: \mathcal{U} \rightarrow L_p(E, \nu)$ a measure of finite variation (i.e., $\forall i$, the variation of μ relative to $N_{p,i}$ is finite, [4]), absolutely continuous with respect to λ . Assume first that $\nu(Y) < \infty$ and let $\lambda \times \nu$ be the product of λ and ν on the σ -algebra $\mathcal{U} \times \mathcal{B}$.

For an $A \in \mathcal{U}, B \in \mathcal{B}$, define $\omega(A \times B) = \int_B \mu(A) d\nu \in E$ (since $\nu(Y) < \infty, P_i(\mu(A)) \in L_p(\nu), \forall i$, implies $P_i(\mu(A)) \in L_1(\nu)$). Take $\{A_i \times B_i\}$ a disjoint sequence in $X \times Y$ ($A_i \in \mathcal{U}, B_i \in \mathcal{B}$) and let $\cup A_i \times B_i = A \times B$ ($A \in \mathcal{U}, B \in \mathcal{B}$). Fix an $f \in E'$. $f \circ \mu: \mathcal{U} \rightarrow L_p(\nu)$ is of bounded variation and absolutely continuous relative to λ . Since $L_p(\nu)$ has R – N property, there exists a function $\phi: X \times Y \rightarrow K$ such that

$$f \circ \mu(A) = \int_A \phi(x, y) d\lambda(x), \forall A \in \mathcal{U};$$

it is routine verification that $\phi(x, y) \in L_1(\lambda \times \nu)$. Thus

$$\int_{B_i} f \circ \mu(A_i) d\nu = \int_{A_i \times B_i} \phi(x, y) d(\lambda \times \nu)$$

(Fubini's theorem) and so $\sum \int_{B_i} f \circ \mu(A_i) d\nu = \int_B f \circ \mu(A) d\nu$ (un-

conditional convergence). Since $\langle f, \int_{B_i} \mu(A_i) d\nu \rangle = \int_{B_i} f \circ \mu(A_i) d\nu$, (simple verification), $\forall f \in E'$, by Pettis-Orlicz theorem,

$$\sum \int_{B_i} \mu(A_i) d\nu = \int_B \mu(A) d\nu.$$

Also for a finite disjoint collection $\{C_i \times D_i\}$ in $X \times Y$ ($C_i \in \mathcal{U}, D_i \in \mathcal{B}$),

$$\sum \int_{D_i} f \circ \mu(C_i) d\nu = \int_{\cup C_i \times D_i} \phi(x, y) d(\lambda \times \nu) \text{ (previous notation) and}$$

so $\left| f \circ \sum \int_{D_i} \mu(C_i) d\nu \right| \leq \int |\phi(x, y)| d(\lambda \times \nu)$. Combining these

results we see that ω can be uniquely extended to a finitely additive set function $\omega: \theta \rightarrow E$, θ being the algebra generated by $\{A \times B: A \in \mathcal{U}, B \in \mathcal{B}\}$. ω is countably additive, and $\omega(\theta)$ is bounded in E . Since E has R – N property it cannot contain a subspace isomorphic to c_0 ([1]; [3], Theorem 5). From this it easily follows that ω is exhaustive ([2], II; [7], Theorem 4). Thus ω can be uniquely extended to a countably additive measure on the σ -algebra $\mathcal{U} \times \mathcal{B}$ ([2], III). We claim that $\omega \ll \lambda \times \nu$. For an

$f \in E', A \in \mathcal{U}, B \in \mathcal{B}, f \circ \omega(A \times B) = \int_{A \times B} \phi(x, y) d(\lambda \times \nu)$ (pre-

vious notations) and so $f \circ \omega(H) = \int_H \phi d(\lambda \times \nu)$, $\forall H \in \mathcal{U} \times \mathcal{B}$.
 If $(\lambda \times \nu)(H) = 0$ we get $f \circ \omega(H) = 0$ and so $\omega(H) = 0$.

We now prove that ω is of finite variation. Fix $i \in \mathbb{N}$ and let $\lambda_0 =$ the finite variation of μ relative to the semi-norm $N_{i,p}$.

$$H = \{f \in E', |f(x)| \leq P_i(x), \forall x \in E\}$$

is a metrizable compact subset of $(E', \sigma(E', E))$ and so have a countable dense subset $\{f_j\}$. Let $\phi_j \in L_1(\lambda_0 \times \nu)$ such that

$$f_j \circ \mu(A) = \int_A \phi_j(x, y) d\lambda_0(x)$$

(same reasoning as before).

Let $\varphi_0 = \sup(|\varphi_1|, |\varphi_2|)$ and fix $x \in X$. Take

$$B_1 = \{y \in Y : |\varphi_1(x, y)| = \varphi_0(x, y)\} \text{ and } B_2 = Y \setminus B_1.$$

We claim the variation of $\xi = \chi_{B_1} f_1 \circ \mu + \chi_{B_2} f_2 \circ \mu$, in $L_p(\nu)$, does not exceed λ_0 . From

$$|\xi(A)|^p = |(\chi_{B_1} f_1 \circ \mu + \chi_{B_2} f_2 \circ \mu)(A)|^p \leq \chi_{B_1} (P_i(\mu(A)))^p + \chi_{B_2} (P_i(\mu(A)))^p = (P_i(\mu(A)))^p,$$

we get $\|\xi(A)\|_p \leq N_{i,p}(\mu(A)) \leq \lambda_0(A)$ and so the claim is established.

Now $\xi(A) = \int_A (\chi_{B_1} \varphi_1 + \chi_{B_2} \varphi_2) d\lambda_0$. If $|\xi|$ is the variation of ξ relative to $L_p(\nu)$, then $|\xi|(A) = \int_A \|\chi_{B_1} \varphi_1 + \chi_{B_2} \varphi_2\|_p d\lambda_0$.

If $\|\chi_{B_1} \varphi_1 + \chi_{B_2} \varphi_2\|_p \geq 1 + \eta$ for some $\eta > 0$ on $A \in \mathcal{U}$, then $\lambda_0(A) \geq |\xi|(A) \geq (1 + \eta)\lambda_0(A)$ which means $\lambda_0(A) = 0$ and so $\|\chi_{B_1} \varphi_1 + \chi_{B_2} \varphi_2\|_p \leq 1$ a.e. $[\lambda_0]$. Now

$$\|\chi_{B_1} \varphi_1 + \chi_{B_2} \varphi_2\|_1 \leq \|\chi_{B_1} \varphi_1 + \chi_{B_2} \varphi_2\|_p (\lambda_0(X))^{1/q} \leq (\lambda_0(X))^{1/q}$$

(Holder's inequality with $\frac{1}{p} + \frac{1}{q} = 1$) means

$$\int \varphi_0(x, y) d\nu(y) \leq (\lambda_0(X))^{1/q}, \text{ a.e. } [\lambda_0].$$

Since $x \in X$ was arbitrary we see $\varphi_0(x, y) \in L_1(X \times Y, \lambda_0 \times \nu)$.

If $\varphi = \sup(|\varphi_1|, |\varphi_2|, \dots)$, then proceeding in the same way we prove that $\int \varphi(x, y) d\nu(y) \leq (\lambda_0(X))^{1/q}$, a.e. $[\lambda_0]$ and so

$\varphi \in L_1(X \times Y, \lambda_0 \times \nu)$ (the set where φ takes values $+\infty$ has measure zero; we put $\varphi \equiv 0$ on that set). Fix $\epsilon > 0$ and let $\{H_j\}$ be a finite disjoint collection in $\mathcal{U} \times \mathcal{B}$.

$$\begin{aligned} \sum_j P_i(\omega(H_j)) - \epsilon &\leq \sum_j |f_{k(j)} \circ \omega(H_j)| \leq \sum_{H_j} |\varphi_{k(j)}| d(\lambda_0 \times \nu) \\ &\leq \sum_{H_j} \varphi d(\lambda_0 \times \nu) \leq \int \varphi d(\lambda_0 \times \nu) \end{aligned}$$

for some finite sequence $\{k(j)\} \subset \mathbb{N}$. This proves ω is of finite variation. Since E has $R - N$ property we get a $g \in L_1(E, \lambda \times \nu)$ such that $\int_B \mu(A) d\nu = \int_B \int_A g(x, y) d\lambda(x) d\nu(y)$.

Put $\psi = \mu(A) - \int_A g(x, y) d\lambda(x)$. We get $\int_B \psi d\nu = 0, \forall B \in \mathcal{B}$. Fix $i \in \mathbb{N}$ and let $\{f_j\}$ be a countable dense set in the compact metric space

$$H = \{f \in E' : |f(x)| \leq P_i(x), \forall x \in E\} \subset (E', \sigma(E', E)).$$

We get $\int_B f_j \circ \psi = 0, \forall B \in \mathcal{B}$ and so $P_i(\psi) = 0$ a.e. $[\nu]$. Thus $\psi = 0$ a.e. $[\nu]$. Thus $\mu(A) = \int_A g(x, y) d\lambda(x)$. It is easy to verify that $g(\cdot, x) \in L_1(L_p(E, \nu), \lambda)$.

Now we consider the case when $\nu(Y) = +\infty$. By ([3], Theorem 5) it is enough to prove the result for every closed separable subspace of $L_p(E, \nu)$. Let F be a closed separable subspace of $L_p(E, \nu)$. It is a simple verification that there exists a $B \in \mathcal{B}$ with σ -finite ν -measure such that $f = 0$ a.e. $[\nu]$ outside $B, \forall f \in F$. Thus ([3], Theorem 5) we can assume that ν is σ -finite. Let $\{K_n\}$ be a \mathcal{B} -measurable partition of Y , such that $0 < \nu(K_n) < \infty, \forall n$. Define $\nu_n = \chi_{K_n} \nu, \nu_n: \mathcal{B}_n \rightarrow [0, \infty), \mathcal{B}_n = \mathcal{B} \cap K_n$. Given μ as before, we get $\mu_n: \mathcal{U} \rightarrow L_p(E, \nu_n), \mu_n(A) = \chi_{K_n} \mu(A) \in L_p(E, \nu_n)$. It is easy to verify that μ_n is of finite variation relative to $L_p(E, \nu_n)$ and absolutely continuous relative to λ . Proceeding as before we get $g_n: X \times K_n \rightarrow E$ such that

$$\mu_n(A) = \int_A g_n(x, y) d\lambda(x) = \int_A \chi_{K_n} g_n(x, y) d\lambda(x).$$

Define $g(x, y) = g_n(x, y), y \in K_n$, we claim

$$\mu(A) = \int_A g(x, y) d\lambda(x).$$

If $\mu(A) = f \in L_p(E, \nu)$, then

$$(P_i(\mu(A) - \sum_{j=1}^k \mu_j(A)))^p \leq (P_i(f))^p$$

and so by dominated convergence theorem $\sum_{j=1}^k \mu_j(A)$ converges to $\mu(A)$ in $L_p(E, \nu)$. Let $|\mu|$ and $\left| \sum_{j=1}^k \mu_j \right|$ be the variations of μ and $\sum_{j=1}^k \mu_j$ relative to $N_{i,p}$. Then

$$|\mu|(A) \geq \left| \sum_{j=1}^{\infty} \mu_j \right| (A) = \int_A N_{i,p} \left(\sum \chi_{K_j} g_j \right) d\lambda.$$

By monotone convergence theorem $N_{i,p}(g) < \infty$, a.e. $[\lambda]$ and $\int N_{i,p}(g) d\lambda < \infty$. On the set where $N_{i,p}(g) = +\infty$ we change its value and the value of each of g_n to 0 and so $g(\cdot, x) \in L_p(E, \nu)$, $\forall x \in X$. Now it is easy to verify that $h_n = \sum_{j=1}^n \chi_{K_j} g_j$ converges to g in $L_p(E, \nu)$, a.e. $[\lambda]$ and $N_{i,p}(g - h_n)$, as a function of x , is decreasing as n increases. By monotone convergence theorem, $\int N_{i,p}(g - h_n) d\lambda \rightarrow 0$. Thus

$$N_{i,p} \left(\int_A g d\lambda - \int_A h_n d\lambda \right) \leq \int N_{i,p}(g - h_n) d\lambda \rightarrow 0$$

and so $\int_A h_n d\lambda \rightarrow \int_A g d\lambda$ in $L_p(E, \lambda)$. But

$$\int_A h_n d\lambda = \sum_{j=1}^n \mu_j(A) \rightarrow \mu(A)$$

and so $\mu(A) = \int_A g d\lambda$. The result now follows easily.

BIBLIOGRAPHY

- [1] J. DIESTEL, J.J. UHL, Jr., The Radon-Nikodym property for Banach space valued measures, *Rocky Mountain J. Math.*, 6 (1976), 1-46.
- [2] L. DREWNOWSKI, Topological rings of sets, continuous set functions, integration I, II, III, *Bull. Acad. Polon. Sci., Ser. Math. Astron. Phys.*, 20 (1972), 269-276, 277-286, 439-445.
- [3] E. SAAB, Dentabilité, points extrémaux et propriété de Radon-Nikodym, *Bull. Soc. Math.*, 99 (1975), 129-134.
- [4] E. SAAB, Dentabilité, points extrémaux et propriété de Radon-Nikodym, *C.R. Acad. Sci., Paris*, 280 (1975), 575-577.
- [5] H.H. SCHAEFER, Topological vector spaces, Macmillan, New York (1971).
- [6] K. SUNDARESAN, The Radon-Nikodym theorem for Lebesgue-Bochner function spaces, *J. Func. Anal.*, 24 (1977), 276-279.

- [7] Ju. B. TUMARKIN, On locally convex spaces with basis, *Doklady Acad. Sci. USSR*, 195 (1970), 1278-1281, English Translation: *Soviet Math.*, 11 (1970), 1672-1675.
- [8] P. TURPIN, Convexité dans les espaces vectoriels topologiques généraux, *Disser. Math.*, 131 (1976).

Manuscrit reçu le 21 septembre 1977

Proposé par G. Choquet.

Surjit Singh KHURANA,
Department of Mathematics
University of Iowa
Iowa City, Iowa 52242 (USA).