

ANNALES DE L'INSTITUT FOURIER

K. S. SARKARIA

Non-degenerescence of some spectral sequences

Annales de l'institut Fourier, tome 34, n° 1 (1984), p. 39-46

http://www.numdam.org/item?id=AIF_1984__34_1_39_0

© Annales de l'institut Fourier, 1984, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

NON-DEGENERESCENCE OF SOME SPECTRAL SEQUENCES

by K.S. SARKARIA

1. Introduction.

Given a Lie algebra \mathfrak{F} of vector fields (smooth sections of the complexified tangent bundle) of a smooth manifold M^m one has a decreasing filtered complex

$$A(M) = A_0(\mathfrak{F}) \supseteq A_1(\mathfrak{F}) \supseteq \dots \supseteq A_m(\mathfrak{F}) \supseteq A_{m+1}(\mathfrak{F}) = 0; \quad (1)$$

here $A(M)$ is the differential graded algebra of all smooth complex valued forms on M (the de Rham complex of M) and, for each $i > 0$, $A_i(\mathfrak{F})$ denotes the i th power of the ideal generated by 1-forms which vanish on \mathfrak{F} . Like any filtered complex, (1) has an associated spectral sequence which will be denoted by $E_k(\mathfrak{F})$. (We will follow Griffiths and Harris [2] regarding terminology for spectral sequences.) Two special cases of this construction will be important for us. (a) If M^{2n} is a complex manifold and \mathfrak{F} consists of vector fields which in local complex coordinates z_1, z_2, \dots, z_n are of the form $\sum_T \varphi_j \frac{\partial}{\partial \bar{z}_j}$, then $E_k(\mathfrak{F})$ is the well known *Fröhlicher spectral sequence* [1] of the complex manifold M . The usual notation for $E_1^{p,q}(\mathfrak{F})$ is $H^q(M, \Omega^p)$: the q th cohomology of M with coefficients in the sheaf Ω^p of germs of holomorphic p -forms on M . (b) If M carries a smooth foliation and \mathfrak{F} consists of all vector fields tangent to the leaves, then $E_k(\mathfrak{F})$ will be called the *spectral sequence of the foliated manifold* M . It has been considered e.g. in [3].

THEOREM. — (A) For all integers $m \geq 1$, $0 \leq c \leq m$, the m -torus T^m admits a codimension c real analytic foliation such that $E_\mu^{\mu,0}(\mathfrak{F})$ is infinite dimensional; here $\mu = \min(c, m - c + 1)$. (B) For all integers $n \geq 1$, $T^{2n-1} \times \mathbb{R}$ admits a complex structure such that $E_n^{n,0}(\mathfrak{F})$ is infinite dimensional. (C) For all integers $m \geq 1$, \mathbb{R}^m admits a compactly supported Lie algebra \mathfrak{F} of vector fields such that $E_m^{m,0}(\mathfrak{F})$ is infinite dimensional.

This theorem (which will be proved in § 2) is best possible in the sense that dimension considerations show that $E_k(\mathfrak{F})$ is isomorphic to the de Rham cohomology for $k \geq \mu + 1$ (resp. $k \geq n + 1$, resp. $k \geq m + 1$) and so is finite dimensional for these values of k . We note that the cohomology considered by Schwarz [4] is precisely the $E_2^{*,0}(\mathfrak{F})$ of a foliated manifold. In response to a question posed by Bott, he constructed smooth (non-analytic) foliations of compact manifolds for which $E_2^{*,0}(\mathfrak{F})$ is infinite dimensional. Thus (A) can be considered as an improvement on [4]; besides our construction is different and simpler and leads to foliations which are of a natural and non-pathological kind. Regarding (B) we remark that the degenerescence problem for compact complex manifolds is much more delicate; we hope to give some results about it in a subsequent paper.

2.

In each of the examples (A)–(C) we will check, for $k = \mu, n$ and m respectively, that $\text{Im} \{H^k(A_k(\mathfrak{F})) \rightarrow H^k(A_1(\mathfrak{F}))\}$ is infinite dimensional. The infinite dimensionality of $E_k^{k,0}(\mathfrak{F})$ (see [2], p. 441 for definition) would follow as an immediate consequence.

Proof of (A) (We ignore the trivial cases $c = 0, m$). — The torus T^m will be considered as the quotient $\mathbb{R}^m / (2\pi\mathbb{Z})^m$. Furthermore we put $\nu = m - (2\mu - 1)$ and identify \mathbb{R}^m with $\mathbb{R}^{\mu-1} \times \mathbb{R}^{\mu-1} \times \mathbb{R} \times \mathbb{R}^\nu$; thus a point of T^m will have coordinates $(\theta_{1,0}, \dots, \theta_{\mu-1,0}; \theta_{1,1}, \dots, \theta_{\mu-1,1}; r; t_1, \dots, t_\nu)$. Note that $\mu = \ell + 1$ (here $\ell = m - c$) or $\mu = c$; correspondingly $\mu - 1 = \ell$ or $\mu - 1 + \nu = \ell$. We define $\mu - 1$ real analytic vector fields on T^m by

$$X_i = \frac{\partial}{\partial \theta_{i,0}} + \sin r \cdot \frac{\partial}{\partial \theta_{i,1}}, \quad 1 \leq i \leq \mu - 1. \quad (2)$$

In case $\ell > \mu - 1$ we also take the vector fields $\frac{\partial}{\partial t_1}, \dots, \frac{\partial}{\partial t_\nu}$.

Taken together these ℓ vector fields span an Abelian ℓ dimensional Lie algebra \mathcal{A} of analytic vector fields. At each point of T^m these vector fields are linearly independent; so they determine a real analytic ℓ dimensional tangent plane field on T^m . By Frobenius theorem this involutive plane field is tangent to a real analytic ℓ dimensional foliation of T^m . We will prove that $E_\mu^{\mu,0}(\mathcal{F})$ is infinite dimensional for this foliation. (Note that \mathcal{A} and \mathcal{F} have the same filtration (1) and so have the same spectral sequence.)

For each smooth function $\varphi(r)$ of period 2π we define a closed form Ω_φ of degree μ on T^m by

$$\Omega_\varphi = \sum_{(\alpha_1, \dots, \alpha_{\mu-1})} \varphi(r) (-\sin r)^{\mu-1-\alpha} dr \wedge d\theta_{1,\alpha_1} \wedge \dots \wedge d\theta_{\mu-1,\alpha_{\mu-1}}; \quad (3)$$

here the summation is over all multi-indices $(\alpha_1, \dots, \alpha_{\mu-1})$ with entries 0 or 1, and $\alpha = \alpha_1 + \dots + \alpha_{\mu-1}$. Using (2) it follows

that the interior products $\iota_{X_i}(\Omega_\varphi)$ and $\iota_{\frac{\partial}{\partial t_k}}(\Omega_\varphi)$ vanish for $1 \leq i \leq \mu - 1$ and $1 \leq k \leq \nu$. Thus the forms Ω_φ constitute an infinite dimensional subspace of closed degree μ forms of $A_\mu(\mathcal{F})$. We assert that if $\Omega_\varphi = d\omega$ where $\omega \in A_1(\mathcal{F})$, then the form Ω_φ is the zero form (i.e. $\varphi \equiv 0$). This will suffice to prove the infinite dimensionality of $\text{Im} \{H^\mu(A_\mu(\mathcal{F})) \longrightarrow H^\mu(A_1(\mathcal{F}))\}$.

Let $T^{2\mu-2}$ denote the subtorus of T^m obtained by putting the last $\nu + 1$ coordinates equal to zero. For each $\theta \in T^{2\mu-2}$ we have the translation $L_\theta : T^m \longrightarrow T^m$ given by $L_\theta(u) = \theta + u$. From (3) we see that Ω_φ is preserved by this action of $T^{2\mu-2}$ on T^m . Since the infinitesimal generators of this action are $\frac{\partial}{\partial \theta_{i,\alpha_i}}$ and $\left[\frac{\partial}{\partial \theta_{i,\alpha_i}}, X_k \right] = 0 = \left[\frac{\partial}{\partial \theta_{i,\alpha_i}}, \frac{\partial}{\partial t_k} \right]$, we conclude also that our Lie algebra \mathcal{A} is preserved by this action. Hence if $\omega \in A_1(\mathcal{F})$ is such that $d\omega = \Omega_\varphi$ we can assume in addition that ω is also preserved by this action; for, otherwise, we can replace ω by the form $\int_{T^{2\mu-2}} L_\theta^*(\omega) d\theta$ obtained by averaging with respect to the normalized Haar measure $d\theta$ of $T^{2\mu-2}$. Let us now write

the expression of ω in the chosen coordinate system. We have

$$\omega = \sum_{(\alpha_1, \dots, \alpha_{\mu-1})} \omega_{\alpha_1 \dots \alpha_{\mu-1}} d\theta_{1, \alpha_1} \wedge \dots \wedge d\theta_{\mu-1, \alpha_{\mu-1}} + \bar{\omega} \quad (4)$$

where $\bar{\omega}$ denotes all terms other than those written out in the beginning. Since ω is preserved by the action of $T^{2\mu-2}$ every coefficient of ω in (4) is a function only of r, t_1, \dots, t_p . But $d\omega = \Omega_\varphi$ where Ω_φ is given by (3). This shows that $d\bar{\omega} = 0$ and that the coefficients $\omega_{\alpha_1 \dots \alpha_{\mu-1}}$ are functions only of r .

Since ω is a degree $\mu - 1$ form in $A_1(\mathcal{F})$ we should get zero when the operator $\sim_{X_{\mu-1}} \circ \dots \circ \sim_{X_1}$ is applied to (4). It is clear that on applying this to $\bar{\omega}$ we get zero. On applying it to the first part we get the condition

$$\sum_{(\alpha_1, \dots, \alpha_{\mu-1})} (\sin r)^\alpha \omega_{\alpha_1 \dots \alpha_{\mu-1}} = 0. \quad (5)$$

Next we use the conditions $\sim_{X_i}(d\omega) = 0, 1 \leq i \leq \mu - 1$ on (4). We see that for each multi-index $(\alpha_1, \dots, \alpha_{\mu-1})_k$ of length $\mu - 2$ obtained by omitting the k th entry we have

$$\omega'_{\alpha_1 \alpha_2 \dots 0 \dots \alpha_{\mu-1}} + \sin r \cdot \omega'_{\alpha_1 \alpha_2 \dots 1 \dots \alpha_{\mu-1}} = 0 \quad (6)$$

where primes denote differentiation with respect to r and 0 and 1 are placed at the k th place. This in turn implies the following identities

$$\sum_{\substack{(\alpha_1, \dots, \alpha_{\mu-1}) \\ s.t. \alpha > s}} \frac{\alpha!}{(\alpha - s)!} (\sin r)^{\alpha - s} \omega'_{\alpha_1 \alpha_2 \dots \alpha_{\mu-1}} = 0, \quad 0 \leq s \leq \mu - 2. \quad (7)_s$$

(To check $(7)_s$ use (6) and the binomial identities

$$\sum_{\alpha=s}^{\mu-1} \frac{\alpha!}{(\alpha - s)!} \binom{\mu - 1}{\alpha} (-1)^\alpha = 0.)$$

We now differentiate (5) with respect to r and use $(7)_0$ to get

$$\cos r \cdot \sum_{\substack{(\alpha_1, \dots, \alpha_{\mu-1}) \\ s.t. \alpha > 1}} \alpha (\sin r)^{\alpha - 1} \cdot \omega_{\alpha_1 \alpha_2 \dots \alpha_{\mu-1}} = 0$$

which implies —since $\cos r$ is non-zero on an open dense subset of T^m — that

$$\sum_{\substack{(\alpha_1, \dots, \alpha_{\mu-1}) \\ s.t. \alpha > 1}} \alpha (\sin r)^{\alpha - 1} \cdot \omega_{\alpha_1 \alpha_2 \dots \alpha_{\mu-1}} = 0.$$

Next we differentiate this and use $(7)_1$, etc., etc.; finally we get $\omega_{11\dots 1} = 0$. Now by using (6) it follows that $\omega'_{\alpha_1\alpha_2\dots\alpha_{\mu-1}} = 0$ for all multi-indices $(\alpha_1, \dots, \alpha_{\mu-1})$ which means that $d\omega = \Omega_\varphi = 0$.

Proof of (B). — We identify $T^{2n-1} \times \mathbf{R}$ with the quotient $(\mathbf{C}^{n-1} \times \mathbf{C})/\Gamma$; here Γ is the subgroup of $\mathbf{C}^{n-1} \times \mathbf{C}$ consisting of elements $(w_1, \dots, w_{n-1}; u)$ with u and each $\operatorname{Re} w_k, \operatorname{Im} w_k$ an integral multiple of 2π . Now let \mathcal{L} be the Abelian n dimensional Lie algebra of vector fields of $T^{2n-1} \times \mathbf{R}$ spanned by $\frac{\partial}{\partial \bar{u}}$ and

$$X_k = \frac{\partial}{\partial \bar{w}_k} + 2e^{iu} \frac{\partial}{\partial w_k}, \quad 1 \leq k \leq n-1. \quad (8)$$

We note that at each point x of $T^{2n-1} \times \mathbf{R}$ these n vector fields are linearly independent; so they span an n dimensional subspace $F(x)$ of the complexified tangent space $T(x)$. Furthermore $F(x) + \overline{F(x)} = T(x)$ (because $\frac{\partial}{\partial \bar{w}_k} + 2e^{iu} \frac{\partial}{\partial w_k}$ is always

linearly independent from $\frac{\partial}{\partial w_k} + 2e^{-i\bar{u}} \frac{\partial}{\partial \bar{w}_k}$). By the well known complex Frobenius theorem (of Newlander and Nirenberg) this involutive almost complex structure is integrable i.e. we can choose local complex coordinates z_1, z_2, \dots, z_n so that the smooth sections of F are precisely those vector fields which are of the type $\sum_j \varphi_j \frac{\partial}{\partial \bar{z}_j}$. We assert that $E_n^{n,0}(\mathcal{F})$ is infinite dimensional for this complex structure.

For each entire function $\varphi(u)$ of period 2π we define a closed form of degree n on $T^{2n-1} \times \mathbf{R}$ by

$$\Omega_\varphi = \sum_{(\alpha_1, \dots, \alpha_{n-1})} \varphi(u) (-2e^{iu})^{n-1-\alpha} du \wedge dw_{1,\alpha_1} \wedge \dots \wedge dw_{n-1,\alpha_{n-1}}. \quad (3)'$$

Here the summation is over all multi-indices $(\alpha_1, \dots, \alpha_{n-1})$ with entries 0, 1 and $\alpha = \alpha_1 + \dots + \alpha_{n-1}$; furthermore dw_{k,α_k} denotes $d\bar{w}_k$ if $\alpha_k = 0$ and dw_k if $\alpha_k = 1$. We can now prove, by a method entirely analogous to that in part (A), that these Ω_φ constitute an infinite dimensional subspace of closed forms in $A_n(\mathcal{F})$ and that no non-zero Ω_φ can be the boundary of a form $\omega \in A_1(\mathcal{F})$.

Proof of (C). – Case $m = 1$ is trivial (take $\mathfrak{F} = 0$); so we assume $m \geq 2$. Let us take the closed disk D in \mathbf{R}^2 with origin as centre and radius 1. In its interior we choose a countably infinite number of disjoint closed disks D_β . Let us choose polar coordinates (r_β, ϕ_β) for each D_β : so D_β is given by

$$0 \leq \phi_\beta < 2\pi, \quad 0 \leq r_\beta \leq \rho_\beta$$

where ρ_β is the radius of D_β . On the torus $T^{m-2} = \mathbf{R}^{m-2}/\mathbf{Z}^{m-2}$ we choose coordinates $\theta_1, \theta_2, \dots, \theta_{m-2}$. Now we define $m-1$ smooth vector fields on $T^{m-2} \times D^2$ by

$$\begin{aligned} X_i &= \sum_{\beta} c_{\beta} f_{\beta}(r_{\beta}) \frac{\partial}{\partial \theta_i} & \text{if } 1 \leq i < m-1 \\ &= \sum_{\beta} c_{\beta} f_{\beta}(r_{\beta}) \frac{\partial}{\partial \phi_{\beta}} & \text{if } i = m-1. \end{aligned} \quad (9)$$

Here $f_{\beta}(r_{\beta})$ is a smooth function on \mathbf{R}^+ , positive on $(\frac{\rho_{\beta}}{2}, \rho_{\beta})$ and zero outside this interval, while the positive constants c_{β} are so chosen that the sums in (9) converge in the C^{∞} topology. One notes that $[X_i, X_j] = 0 \forall i, j$ and that each vector field X_i vanishes on the boundary $T^{m-2} \times \partial D^2$ of our solid torus $T^{m-2} \times D^2$. The latter can be smoothly imbedded in \mathbf{R}^m and we can extend these vector fields to all of \mathbf{R}^m by defining them to be zero outside the solid torus. This gives us an Abelian $m-1$ dimensional Lie algebra \mathfrak{F} of compactly supported vector fields on \mathbf{R}^m ; we will prove that $E_m^{m,0}(\mathfrak{F})$ is infinite dimensional.

For each β , let us choose a smooth m -form Ω_{β} supported inside the region of $T^{m-2} \times D_{\beta}$ given by $r_{\beta} < \frac{\rho_{\beta}}{2}$ and such that

$$\int_{T^{m-2} \times D_{\beta}} \Omega_{\beta} \neq 0. \quad (10)$$

Note that each Ω_{β} lies in $A_m(\mathfrak{F})$. If possible let us suppose that Ω is a non-trivial finite linear combination of the Ω_{β} such that $\Omega = d\omega$ where ω is a smooth $m-1$ form lying in $A_{m-1}(\mathfrak{F})$. This last condition implies that $\omega(X_1, \dots, X_{m-1}) = 0$ and so in $T^{m-2} \times D_{\beta}$ we have

$$(c_{\beta} f_{\beta}(r_{\beta}))^{m-1} \omega \left(\frac{\partial}{\partial \theta_1}, \dots, \frac{\partial}{\partial \theta_{m-2}}, \frac{\partial}{\partial \phi_{\beta}} \right) = 0. \quad (11)$$

In the region $\frac{\rho_\beta}{2} < r_\beta < \rho_\beta$ the coefficient of (11) is non-zero ; so in this domain we must have $\omega\left(\frac{\partial}{\partial\theta_1}, \dots, \frac{\partial}{\partial\theta_{m-2}}, \frac{\partial}{\partial\phi_\beta}\right) = 0$ which remains valid on $r_\beta = \rho_\beta$ by continuity. Hence ω induces, on the boundary of each $T^{m-2} \times D_\beta$, the zero form. By Stoke's theorem this contradicts (10). Thus we have proved that

$$\text{Im} \{H^m(A_m(\mathcal{F})) \longrightarrow H^m(A_1(\mathcal{F}))\}$$

has infinite dimension.

3. Remarks.

The proof of (C) given above is inspired by the work of Schwarz [4]. We note that it is possible to use analogous ideas to construct some more smooth (non-analytic) codimension c foliations on T^m with $E_\mu^{\mu, \circ}(\mathcal{F})$ infinite dimensional. One can also ensure that the singular set of such foliations is nowhere dense. (The *singular set* is made up of those points where the infinitesimal transformations of the foliation fail to span the tangent space. In [3] it is proved that for a compact foliated manifold with empty singular set, $E_2(\mathcal{F})$ is finite dimensional.) In another direction we can prove that if a smooth manifold M^m admits one smooth codimension c foliation, then it admits another with $E_\mu^{\mu, \circ}(\mathcal{F})$ infinite dimensional.

Acknowledgement. – This work was done in 1979 while I was visiting France ; I would like to thank R. Barre and the University of Lille I for having made this pleasant trip possible.

BIBLIOGRAPHY

- [1] A. FROHLICHER, Relations between the cohomology groups of Dolbeault and topological invariants, *Proc. Nat. Acad. Sci. U.S.A.*, 41(1955), 641-644.
- [2] P. GRIFFITHS and J. HARRIS, *Principles of Algebraic Geometry*, John Wiley and Sons, New York, 1978.
- [3] K.S. SARKARIA, A finiteness theorem for foliated manifolds, *Jour. Math. Soc. Japan*, 30 (1978), 687-696.
- [4] G. SCHWARZ, On the de Rham cohomology of the leaf space of a foliation, *Topology*, 13 (1974), 185-187.

Manuscrit reçu le 4 octobre 1982.

K.S. SARKARIA,
213, 16A
Chandigarh 160016 (India).